Septate junction proteins are required for egg elongation and border cell

migration during oogenesis in Drosophila

Haifa Alhadyian[1], Dania Shoiab[1], and Robert E. Ward IV[1,2]

[1]Department of Molecular Biosciences, University of Kansas, Lawrence, Kansas 66045

[2]Current address and Corresponding Author information: Robert E. Ward IV, Department of

Biology, Case Western Reserve University, 2080 Adelbert Rd, Cleveland, Ohio 44106

E-mail: rew130@case.edu

Phone: (216) 368-6707

Running title: SJ proteins in oogenesis

Keywords: Septate junctions, oogenesis, Drosophila, tissue morphogenesis, egg elongation

Abstract

Protein components of the invertebrate occluding junction - known as the septate junction (SJ) - are required for morphogenetic developmental events during embryogenesis in *Drosophila melanogaster*. In order to determine whether SJ proteins are similarly required for morphogenesis during other developmental stages, we investigated the localization and requirement of four representative SJ proteins during oogenesis: Contactin, Macroglobulin complement-related, Neurexin IV, and Coracle. A number of morphogenetic processes occur during oogenesis, including egg elongation, formation of dorsal appendages, and border cell migration. We found that all four SJ proteins are expressed in the egg throughout oogenesis, with the highest and most sustained levels in the follicular epithelium (FE). In the FE, SJ proteins localize along the lateral membrane during early and mid-oogenesis, but become enriched in an apical-lateral domain (the presumptive SJ) by stage 10b. SJ protein relocalization requires the expression of other SJ proteins, as well as rab5 and rab11 in a manner similar to SJ biogenesis in the embryo. Knocking down the expression of these SJ proteins in follicle cells throughout oogenesis results in egg elongation defects and abnormal dorsal appendages. Similarly, reducing the expression of SJ genes in the border cell cluster results in border cell migration defects. Together, these results demonstrate an essential requirement for SJ genes in morphogenesis during oogenesis, and suggests that SJ proteins may have conserved functions in epithelial morphogenesis across developmental stages.

Article Summary

Septate junction (SJ) proteins are essential for forming an occluding junction in epithelial tissues

of *Drosophila melanogaster*. SJ proteins are also required for morphogenetic events during

embryogenesis prior to the formation of an occluding junction. To determine if SJ proteins

function in morphogenesis at other developmental stages, we examined their function during

oogenesis, and found that SJ proteins are expressed in the follicular epithelium of the egg

chamber and are required for egg elongation, dorsal appendages formation, and border cell

migration. Additionally, we found that the formation of SJs in oogenesis is similar to that in

embryonic epithelia.

Introduction

The septate junction (hereafter referred to as SJ) provides an essential paracellular barrier to

epithelial tissues in invertebrate animals (Noirot-timothée *et al.* 1978). As such, the SJ is

functionally equivalent to the tight junction in vertebrate tissues, although the molecular

components and ultrastructure of these junctions differ (reviewed in Izumi and Furuse 2014).

Studies in *Drosophila* have identified more than 20 proteins that are required for the organization

or maintenance of the SJ (Fehon *et al.* 1994; Baumgartner *et al.* 1996; Behr *et al.* 2003; Paul *et*

al. 2003; Genova and Fehon 2003; Faivre-Sarrailh *et al.* 2004; Wu *et al.* 2004; Wu *et al.* 2007;

Tiklová *et al.* 2010; Nelson *et al.* 2010; Ile *et al.* 2012; Bätz *et al.* 2014; Hall *et al.* 2014). Given

that some of these genes have clear developmental functions (e.g. *coracle*'s name derives from

its dorsal open embryonic phenotype; (Fehon *et al.* 1994), we previously undertook an

examination of the developmental requirements for a set of core SJ genes (Hall and Ward 2016).

We found that all of the genes we analyzed (9 in all) are required for morphogenetic

developmental events during embryogenesis including head involution, dorsal closure and

salivary gland organogenesis. Interestingly, these embryonic developmental events occur prior to

the formation of an intact SJ, suggesting that these proteins have a function independent of their

role in creating the occluding junction (Hall and Ward 2016). Since strong loss of function

mutations in every SJ gene are embryonic lethal (due to these morphogenetic defects and/or a

failure in establishing a blood-brain barrier in glial cells; Baumgartner *et al.* 1996), only a few

studies have examined the role of SJ proteins in morphogenesis at a later stages of development

(Lamb *et al.* 1998; Venema *et al.* 2004; Moyer and Jacobs 2008; Banerjee *et al.* 2008). These

studies have revealed roles for SJ proteins in planar polarization of the wing imaginal disc, for

epithelial rotations in the eye and genital imaginal discs, and ommatidia integrity.

To further explore the role of SJ proteins in morphogenesis beyond the embryonic stage, we

set out to examine the expression and function of a subset of SJ genes in the *Drosophila* egg

chamber during oogenesis. Each of the two *Drosophila* ovaries is comprised of approximately

16-20 ovarioles, which are organized into strings of progressively developing egg chambers

(Figure 1A). Each egg chamber forms in a structure called the germarium, where the germline

and somatic stem cells reside. Once the egg chamber is formed, it leaves the germarium as 16-

cell germline cyst consisting of 15 nurse cells and an oocyte surrounded by a layer of somatic

follicle cells (FCs) (Figure 1B). An egg chamber undergoes 14 developmental stages ending in a

mature egg that is ready for fertilization (reviewed in Horne-Badovinac and Bilder 2005).

Interfollicular cells called stalk cells connect egg chambers to each other. During oogenesis, the

follicular epithelium (FE) undergoes several morphogenetic events including border cell

migration, dorsal appendage formation and egg elongation (reviewed in Horne-Badovinac and

Bilder 2005; reviewed in Duhart *et al.* 2017).

Previous studies have revealed that a few core components of the SJ are expressed in the

ovary, including Macroglobulin complement-related (Mcr), Neurexin IV (Nrx-IV), Contactin

(Cont), Neuroglian (Nrg), and Coracle (Cora) (Wei *et al.* 2004; Schneider *et al.* 2006; Maimon *et*

al. 2014; Hall *et al.* 2014; Ben-Zvi and Volk 2019), although the developmental expression

pattern and subcellular localizations of these proteins have not been thoroughly investigated.

Furthermore, ultrastructural analysis has revealed the presence of mature SJs in the FE by stage

10/10b of oogenesis, while incipient SJ structures have been observed in egg chambers as early

as stage 6 (Mahowald 1972; Müller 2000). The biogenesis of SJs in embryonic epithelia is a

multistep process in which SJ proteins are initially localized along the lateral membrane, but

become restricted to an apical-lateral region (the SJ) in a process that required endocytosis and

recycling of SJ proteins (Tiklová *et al.* 2010). How SJ maturation occurs in the FE is unknown.

Here, we analyzed the expression and subcellular localization of the core SJ proteins Mcr,

Cont, Nrx-IV, and Cora throughout oogenesis. We find that all of these SJ proteins are expressed

in the FE throughout oogenesis. Interestingly, Mcr, Cont, Nrx-IV, and Cora become enriched at

the most apical-lateral region of the membrane in stage 10b/11 egg chambers, coincident with

the formation of the SJ revealed by electron microscopy (Mahowald 1972; Müller 2000) .

Similar to the biogenesis of SJs in the embryo, this enrichment of SJ proteins to the presumptive

SJ requires the function of other SJ genes, as well as *Rab5* and *Rab11*. Functional studies using

RNA interference (RNAi) of SJ genes in FCs results in defects in egg elongation, dorsal

appendage morphogenesis and border cell migration. Together, these results reveal a strong

similarity in the biogenesis of SJ between embryonic and follicular epithelia, and demonstrate

that at least some components of the SJs are required for morphogenesis in the ovary, and

suggest that these roles may be independent of their role in forming an occluding junction.

Material and methods

Fly stocks

All *Drosophila* stocks were maintained on media consisting of corn meal, sugar, yeast, and agar in incubators maintained at a constant temperature of 25°. GAL4 lines used in this study are as follows: *GR1-GAL4* (Bloomington Drosophila Stock Center (BDSC) #36287), *Slbo-GAL4, UAS-mCD8-GFP* (BDSC#76363), and *C306-GAL4; GAL80^{ts}/Cyo* (a gift from Jocelyn McDonald, Kansas State University, Manhattan, Kansas). RNAi stocks used for these studies are as follows: *UAS-Mcr-RNAi* (BDSC#65896 and Vienna Drosophila Resources Center (VDRC)#100197), *UAS-cora-RNAi* (BDSC#28933 and VDRC#9787), *UAS-Nrx-IV-RNAi* (BDSC#32424 and VDRC#9039), *UAS-Cont-RNAi* (BDSC#28923), *UAS-mCherry-RNAi* (BDSC#35787), *UAS-Lac-RNAi* (BDSC#28940), and *UAS-Sinu-RNAi* (VDRC#44929). *UAS-Rab5^{DN}* (BDSC#9771) was used to inhibit normal Rab5 function and *UAS-Rab11-RNAi* (BDSC#27730) was used to knock down Rab11 in the follicle cells. *UAS-GAL80^{ts}* (BDSC#7108) was used to deactivate GR1-GAL4 activity in Rab11-RNAi experiment. *UAS-GFP* (BDSC#1521) was crossed to *GR1-GAL4* as a control for the egg shape experiments, whereas *Slbo-GAL4, UAS-mCD8-GFP* was crossed to *UAS-mCherry-RNAi* as a control for border cell migration studies. Additionally, *C306-GAL4; GAL80^{ts}/Cyo* was crossed to *UAS-Dcr* (BDSC#24646) as a control for border cell migration studies. *w^{1118}* (BDSC# 5905) was used as the wild type stock for determining the expression of Mcr, Cont, Nrx-IV and Cora in the follicle cells. *Rab5-YFP* (BDSC#62543) and *Rab11YFP*

(BDSC#62549) were used to determine their co-localization with Mcr in the follicle cells at stage

10.

Fly staging

w^{1118} 1-2-day-old females and males were collected and reared at 25°C on fresh food sprinkled

with yeast for five to six days before the females were dissection for antibody staining. For egg

elongation analyses, crosses were maintained at 25°C, and 1-2-day-old females (control and

UAS-RNAi-expressing) were mated with sibling males and maintained at 29-30°C for 3 days

before dissection. For border cell migration analyses, *Slbo-GAL4* crosses were kept at 25°C,

whereas *C306-GAL4/UAS-Dcr; GAL80^{ts}/SJ-RNAi* crosses were kept at 18°C to prevent GAL4

activation. 1-2-day-old flies with the appropriate genotype (*Slbo-GAL4, UAS-mCD8-GFP/UAS-*

RNAi or *C306-GAL4 /UAS-Dcr;UAS-RNAi;GAL80^{ts}*) were shifted to 29-30°C for 48 hours

before dissection. It should be noted that by crossing *UAS-GFP* to *C306-GAL4*, we observed the

expression of GFP in polar cells in stage 10, but not stage 9 egg chambers (data not shown). For

Rab11-RNAi experiment, crosses were maintained at 18°C and 2-3-day-old males and females

with the appropriate genotype (*GR1-GAL4>UAS-mCherry-RNAi, UAS-GAL80ts or GR1-*

GAL4>UAS-Rab11-RNAi, UAS-GAL80ts) were collected and reared at 29°C-30°C overnight

before dissection. *Rab5-YFP* or *Rab11-YFP* 1-2-day old females and males were collected and

reared at 25°C on fresh food sprinkled with yeast for five days before dissection. For the *Rab5^{DN}*

experiment, crosses were maintained at 25°C, and 1-2-day-old females were mated to sibling

males and maintained at 29-30°C for 3 days before dissection.

Egg aspect ratio measurements

Stage 14 egg chambers were selected for analysis based on the overall morphology of the egg

and the absence of nurse cells nuclei by DAPI staining. Stage 14 egg chambers that have

irregular edges or touch other egg chambers were excluded from the analysis to prevent

inaccurate measurements. Egg length (anterior-posterior) and width (dorsal-ventral) were

measured using the ImageJ/Fiji (http://fiji.sc) (Schindelin *et al*. 2012) straight-line tool, and

aspect ratio was calculated as length divided by width using Excel Microsoft.

Border cell migration quantification

Stage 10 egg chambers were identified based on the morphology of the egg (oocyte occupies half

the egg chamber, whereas the other half is occupied by the nurse cells and centripetal cells). We

used the GFP signal in Slbo-GAL4 crosses and DAPI and/or Fas3 staining in c306-GAL4

crosses to identify the location of the cluster in stage 10 egg chambers. The location of the border

cell cluster was quantified and grouped into four categories - complete, incomplete, failed

migration - based on the location of the cluster relative to the oocyte in a stage 10 egg chamber

(Figure 6) and disassociated cluster. In some cases, border cell clusters display two phenotypes

such as complete and dissociated. In this case, we quantified both phenotypes in one egg

chamber.

Immunostaining and image acquisition

Ovaries were dissected in 1X Phosphate-buffered saline (PBS), fixed in 4% Paraformaldehyde

for 20 minutes, washed three times in 1X PBS, and then permeabilized in a block solution (1X

PBS + 0.1% Triton + 1% Normal Donkey Serum) for 30 minutes before incubation with primary

antibodies either overnight at 4°C or 2-4 hours at room temperature (~23-25°C). The following

antibodies were used at the given dilutions: guinea pig (gp) anti-Cont 1:2000 *(Faivre-Sarrailh et*

al. 2004) and rabbit (rab) anti-Nrx-IV 1:500 (Stefan Baumgartner *et al*. 1996) obtained from

Manzoor Bhat, University of Texas Health Science Center, San Antonio, TX, gp anti-Mcr

1:1000 (Hall *et al*. 2014), rat anti- *D*E-cad (DCAD2) obtained from the Developmental Studies

Hybridoma Bank (DSHB) at the University of Iowa, Iowa City, IA 1:27, mouse (m) anti-Cora

(C566.9 and C615.16 mixed 1:1, DSHB) 1:50, m anti-Fas3 (7G10, DSHB) 1:260, and m anti-Sxl

(M18, DSHB) 1:10. DAPI (1mg/ml) was used at a dilution of 1:1000. Rab anti-GFP (290-50,

abcam) was used at 1:1000. Secondary antibodies were obtained from Jackson ImmunoResearch

Laboratories (West Grove, Pennsylvania, USA) and were used at 1:500.

Images were acquired using an Olympus FV1000 confocal microscope equipped with

Fluoview software (version 4.0.3.4). Objectives used included an UPLSAPO 20X Oil (NA:0.85),

a PLAPON 60X Oil (NA: 1.42), and an UPLSAPO 100X Oil (NA:1.40). Stage 14 egg chambers

were imaged using on Nikon Eclipse 80*i* compound microscope. Raw images were rotated and

cropped in ImageJ/Fiji. Micrographs were adjusted for brightness using Adobe Photoshop 21.1.1

(San Jose, CA). Adobe Illustrator 24.1 was used to compile the figures.

Statistical Analysis

An unpaired *t*-test was used to calculate the P values in egg chamber aspect ratio between control

and SJ mutant stage 14 egg chambers using GraphPad Prism 8 (https://www.graphpad.com)

(version 8.4.2).

Data Availability

Fly stocks are available upon request. Supplemental files are available at FigShare. Figure

S1shows the expression of Contactin during border cell migration. Figure S2 shows the

expression of Nrx-IV during border cell migration. Figure S3 shows the expression of Coracle

during border cell migration. The authors affirm that all the data necessary for confirming the

conclusions of the article are present within the article and figures.

Results

Septate junction proteins are expressed in follicle cells throughout oogenesis

While a few SJ *proteins* have previously been reported to be expressed in the *Drosophila* ovary (Wei *et al.* 2004; Schneider *et al.* 2006; Hall *et al.* 2014; Maimon *et al.* 2014; Felix *et al.* 2015; Ben-Zvi and Volk 2019), a thorough analysis of their tissue distribution and subcellular localization throughout oogenesis is lacking. We therefore examined the spatial and temporal expression of four SJ proteins: Mcr, Cont, Nrx-IV and Cora (Fehon *et al.* 1994;Baumgartner *et al.* 1996; Faivre-Sarrailh *et al.* 2004; Bätz *et al.* 2014; Hall *et al.* 2014). These four proteins are core components of the junction for which well-characterized antibodies are available.

At early stages of oogenesis (stage 1-8), Mcr, Cont, and Nrx-IV all localize in puncta at the lateral membrane of follicle cells and nurse cells, and show a punctate distribution in these cells (Figure 1D-F). Mcr, Cont, and Nrx-IV are also more strongly expressed in polar cells (PCs) than the surrounding follicle cells (asterisks in Figure 1D-F). Cora is more uniformly localized along the lateral membrane of the FCs, including the PCs (Figure 1G and data not shown). We also find that all of these SJ proteins are expressed in stalk cells (arrowheads in Figure 1 D and E and data not shown). At stage 10b, Mcr, Nrx-IV, Cont and Cora become gradually enriched at the apical-lateral membrane of the FCs just basal to the AJ (arrows in Figure 2B, D, F, and H). The apical-lateral enrichment of Mcr, Cont, Nrx-IV and Cora coincides with the maturation of the SJ in the FCs based upon ultrastructural analysis (Mahowald 1972; Müller 2000), and so we will

refer to this region as the presumptive SJ. Finally, all of these SJ proteins continue to be

expressed in the FCs until stage 14 of oogenesis (Figure 2C, E, G and I).

Septate junction proteins have distinct expression patterns in the germarium

The most anterior region of each ovariole is the germarium. The germarium consists of three

regions (only region 1 and 2 are shown in Figure 3A). Region 1 is occupied by 2-3 germline

stem cells (GSCs) (Figure 3A). Each GSC is in close contact with two subpopulations of somatic

cells - cap cells and terminal filament cells. A GSC undergoes an asymmetric cell division giving

rise to a daughter cell that retains its stem cell characteristics and a cystoblast cell capable of

differentiation (Figure 3A). The cystoblast cell divides four times forming a 2-, 4-, 8-,and 16-cell

cysts while being encapsulated by escort cells (ESCs). As the 16-cell cyst progresses posteriorly

through region 2b of the germarium, it acquires pre-follicle cells derived from a pair of follicle

stem cells, which will generate the FE of the newly formed egg chamber (Figure 3A) (Spradling

et al. 1997; Kirilly and Xie 2007).

Consistent with our previous findings, Mcr is expressed at high levels in the anterior region of

the germarium (Hall *et al.* 2014). To determine which cell types Mcr is expressed in, we co-

stained the germarium with an antibody against Sex lethal (Sxl), which is an RNA-binding

protein expressed in the GSCs, cystoblasts, and 2-cell cysts (Bopp *et al.* 1991; Bopp *et al.*1993).

Co-localization reveals that Mcr is expressed in both GSCs (asterisks in Figure 3B-B'),

cystoblasts (arrow in Figure 3B-B'), and the developing cystocytes (arrowhead in Figure 3B-B').

Cont, in contrast, shows puncta staining throughout the germarium and is not specifically

enriched in any cell type (Figure 3C-C'). Interestingly, Nrx-IV outlines the outside membrane of

the terminal filaments (arrow in Figure 3D), as well as region 1 and 2a of the germarium

(arrowheads in Figure 3D-D') in 5 out of 7 germaria examined. Cora is found in the escort cells

(yellow arrows in Figure 3 E-E'), consistent with previous studies (Maimon *et al.* 2014; Ben-Zvi

and Volk 2019). Cora is also expressed in the terminal filaments (white arrow) and cap cells

(arrowhead) (Figure 3 E-E').

SJ proteins are required for egg elongation and dorsal appendage morphogenesis

Given our findings that these four SJ proteins are expressed in the FE throughout oogenesis,

and our previous studies indicating a role for SJ proteins in morphogenesis, we wondered

whether SJ proteins might be required for morphogenetic processes in the FE. The FE plays

critical roles in shaping the egg chamber and producing the dorsal appendage, while a subset of

FE cells participates in border cell migration to form the micropyle (Montell 2003; S. Horne-

Badovinac 2020). Because SJ mutant animals die during embryogenesis, we used the *GAL4*

UAS-RNAi system to knock-down the expression of SJ proteins in the FCs (Brand and Perrimon

1993). To knock down expression of SJ proteins throughout the majority of oogenesis, we used

GR1-GAL4, which is expressed in the FCs from stage 4 to 14 of oogenesis (Gupta and

Schüpbach 2003; Wittes and Schüpbach 2018). In all, we tested six different SJ genes (*Cont*,

cora, *Mcr*, *lac*, *Nrx-IV*, and *sinu*). To examine overall egg chamber shape, we dissected stage 14

egg chambers from females expressing *SJ-RNAi* under the control of *GR1-GAL4*, imaged them

on a compound microscope, and determined the aspect ratio of the egg chambers using

measurements of egg chamber length and width using ImageJ/Fiji. Control stage 14 egg

chambers (*GR1-GAL4>UAS-GFP*) had a mean aspect ratio of 2.3 (Figure 4A and J). In contrast,

the aspect ratio of stage 14 egg chambers from all *GR1-GAL4>SJ-RNAi* is statistically

significant than control egg chambers (aspect ratios from 1.7 to 2.1; Figure 4B-G and J). All *SJ-*

RNAi egg chambers are significantly shorter than control (Figure 4H), and all but *Mcr-RNAi* and

Cont-RNAi are also wider than control egg chamber (Figure 4I).

Further examination of stage 14 SJ mutant egg chambers revealed defects in dorsal appendage

morphogenesis. Dorsal appendages are tubular respiratory structures that form from two

populations of the dorsal FE known as floor and roof cells (Figure 1A) (Duhart *et al.* 2017). The

primary phenotypes we observed in the *SJ-RNAi*-expressing egg chambers were either missing

dorsal appendages, or appendages that appeared to be broken or excessively short (Figure 4L). In

quantifying these phenotypes, we found that 52% of *Mcr-RNAi-,* 15% of *Cora-RNAi-*, 19% of

Cont-RNAi-, and 13% of *Lac-RNAi*-expressing egg chambers have either missing or broken

dorsal appendages (Figure 4M). We did not observe these phenotypes in either *Nrx-IV-RNAi-* or

Sinu-RNAi-expressing egg chambers (Figure 4M).

SJ proteins are expressed in polar and border cells and are required for effective border cell

migration

The observation that Mcr, Cont, and Nrx-IV are strongly expressed in PCs and in all FCs

(Figure 1D-F), motivated us to investigate their expression during the process of border cell

migration. Border cell migration occurs during stages 9-10 of oogenesis (Figure 5A). During

stage 9, signaling from the anterior PCs recruits a group of 4-8 adjacent FCs to form a cluster

and delaminate apically into the egg chamber. The border cell cluster is organized with a pair of

PCs in the center surrounded by border cells (BCs). This cluster of cells migrates between the

nurse cells until they reach the anterior side of the oocyte (Figure 5A). This process takes

approximately 4-6 hours and is complete in wild type egg chambers by stage 10 of oogenesis

(Prasad and Montell 2007). The border cell cluster, along with the migratory centripetal cells,

collaborate to form the micropyle, a hole through which the sperm enters the egg (reviewed in

Montell 2003; Horne-Badovinac 2020). Previous studies show that Cora and Nrg are expressed

in the PCs during their migration (Wei *et al.* 2004; Felix *et al.* 2015). To determine if other SJ

proteins are also expressed during border cell migration, we stained stage 9-10 wild type egg

chambers with antibodies against Mcr, Cont, and Nrx-IV. We also examined the subcellular

localization of Cora in the border cell cluster. To mark the PCs, we co-stained the egg chambers

with Fasciclin 3 (Fas3; (Snow *et al.* 1989; Khammari *et al.* 2011). We find Mcr, Cont, Nrx-IV

and Cora are all expressed in border cell clusters throughout their migration (Figure 5B-D and

Supplemental Figures 1-3). Interestingly, the expression of these SJ proteins in PCs appears

highest at the interfaces between PCs-BCs (Figure 5B-D and Figures S1-3). SJ protein

expression is also asymmetric in the border cell cluster, with higher expression along border cells

closest to the oocyte, suggesting that they may mark the leading edge of the migrating cluster

(red arrows in Figure 5B"-D").

Given that Mcr, Cont, Nrx-IV, and Cora are expressed in border cell clusters throughout

border cell migration, we wondered if they are also required for some aspect of this process. To

address this issue, we used *Slbo-GAL4* (Ogienko *et al.* 2018) to express RNAi against individual

SJ genes specifically in border cells and analyzed the border cell clusters at stage 10 in these

ovaries. We noticed a range of defective migration phenotypes in these egg chambers and

classified them into three non-exclusive groups: failed, incomplete and dissociated clusters.

Complete migration (e.g. arrow in Figure 6C) is characterized by having the entire border cell

cluster near or physically touching the oocyte by the end of stage 10. A failed cluster (e.g. arrow

in Figure 6D) is characterized by the inability of the border cell cluster from delaminating from

the FE by stage 10. An incomplete migration phenotype (e.g. arrow in Figure 6E, two dashed

lines indicate the range of distance at which border cell cluster was categorized as incomplete) is

seen as an intact cluster that has not reached the oocyte by the end of stage 10. Finally, a

dissociated cluster phenotype (e.g. arrow in Figure 6F) is noted as a cluster that has broken into a

linear string of border cells or that has one or more border cells that remain between NCs that are

not connected to the larger border cell cluster (compare border cell cluster in Figure 6H to an

intact control cluster in Figure 6H).

In control (*Slbo-GAL4; UAS-mCD8-GFP/UAS-mCherry-RNAi*) stage 10 egg chambers, 86%

(n=81) of border cell clusters completed their migration, with the remainder showing incomplete

migration (Figure 6I). In contrast, stage 10 egg chambers expressing RNAi against *Cora*, *Nrx-IV,*

or *Cont* in the border cells resulted in 58% (n=67), 50% (n=74), and 40% (n=85) of border cell

clusters completing migration, respectively (Figure 6I). The remaining *Cora-RNAi-* and *Nrx-IV-*

RNAi-border cell clusters showed a combination of incomplete migration or failed to delaminate

(Figure 6I). *Cont-RNAi*-border cell clusters also showed 35% of incomplete border cell

migration, but additionally had 17% of the clusters disassociating during their migration (Figure

6I). *Mcr-RNAi*-border cell clusters were indistinguishable from controls with 86% (n=94)

completing their migration and the remainder showing only 13% incomplete migration (Figure

6I).

To extend these studies, we examined border cell migration in egg chambers expressing

RNAi against SJ genes using the *C306-GAL4* driver that is expressed in the border cells

throughout the process of border cell migration (Murphy and Montell 1996) and in PCs just at

stage 10 (H.A., unpublished observation). In control (*C306-GAL4/UAS-Dcr; GAL80^{ts}/+*) egg

chambers, 78% (n=91) of BC clusters completed their migration and 19% display incomplete

migration (Figure 6J). Stage 10 egg chambers expressing RNAi against *Mcr* in border and polar

cells resulted in 66% (n=98) complete border cell migration with 30% showing incomplete

migration, 5% dissociated and 3% failed border cell cluster (Figure 6J). Similarly, egg chambers

expressing *Nrx-RNAi* in border and polar cells show 66% (n=59) complete border cell migration

with 30% showing incomplete migration and 3% failing to delaminate (Figure 6J). Finally, 78%

(n=70) of stage 10 *Cora-RNAi*-expressing border and polar cells show complete border cell

migration, whereas 20% show incomplete migration and 3% fail to delaminate (Figure 6J).

SJ biogenesis in the follicular epithelium

The redistribution of SJ proteins in the FCs of later stage egg chambers resembles the

dynamic relocalization of SJ proteins during the biogenesis of the junction during embryogenesis

(Tiklová *et al*. 2010). In embryonic epithelia, SJ protein enrichment at the apical-lateral domain

(presumptive SJ) requires endocytosis and recycling of SJ proteins to the membrane, and is

interdependent on the presence of all core SJ proteins (Ward *et al.* 1998; Hall *et al.* 2014).

Coincident with the strong localization of SJ proteins to the presumptive SJ at stage 16 of

embryogenesis, ladder-like electron-dense intermembrane septae are visible by electron

microscopy that become progressively organized by stage 17 (Schulte *et al.* 2003; Hildebrandt *et*

al. 2015). We therefore sought to determine if similar processes occurred in the FE.

To test for the interdependence of SJ proteins for localization, we examined Cora and Mcr

expression in wild type, *Mcr-RNAi*, and *Nrx-IV-RNAi* stage 12 FCs. Whereas Cora is strongly

co-localized with Mcr at the apical-lateral domain of the FCs of wild type stage 12 egg chambers

(n=95 cells, 31 egg chambers) (Figure 7A), Cora and Mcr are mislocalized along the lateral

domain in *Nrx-IV-RNAi* FCs (Figure 7B), much like they do in earlier stage FCs (Figure 1D).

Specifically, we observed 6 out of 20 cells examined in 19 egg chambers that showed complete

mislocalization and 13 out of these 20 cells showing lateral localization with some degree of

apical enrichment. Similarly, in stage 12 *Mcr-RNAi* FCs, Cora is mislocalized along the lateral

membrane in 39 out of 47 cells examined from 19 egg chambers, with the remaining 8 cells

showing some enrichment of Cora at the apical lateral domain (Figure 7C). Notably, cells that

showed apical enrichment of Cora also retained small foci of Mcr expression that overlaps with

the enriched Cora (Figure 7D), suggesting the perdurance of Mcr in these cells may have allowed

for normal SJ organization. Together, these experiments indicate that SJ biogenesis in the

follicular epithelium of late stage egg chambers requires the expression of at least some core SJ

proteins.

We next wanted to investigate whether the relocalization of SJ proteins to the presumptive SJ

required endocytosis and recycling. In the embryonic hindgut, Cora, Gliotactin (Glio), Sinu, and

Melanotransferrin (Mtf) localize with the early endosomal marker, Rab5, and partially localize

with the recycling marker, Rab11 during SJ biogenesis (Tiklová *et al.* 2010). Moreover, blocking

Rab5 or Rab11 function prevents Cora, Glio, Sinu, and Mtf apical-lateral localization (Tiklová *et*

al. 2010), and thus SJ formation. To determine if the redistribution of SJ proteins in the FCs is

mediated by Rab5 and Rab11, we first asked whether Mcr co-localized with Rab5 and

Rab11compartments in stage 10 FCs. We chose to analyze stage 10 egg chambers because this is

the stage that immediately precedes Mcr's re-localization to the presumptive SJ (similar to stage

13/14 of embryogenesis). During stage 10 of oogenesis, we find that Mcr still localizes

uniformly along the lateral (and to the apical membrane), but is also found in punctate

intracellular compartments (Figure 8A-A''). Notably, some of these Mcr positive puncta co-

localize with Rab5-YFP puncta, suggesting that these Mcr molecules are in early endosomes. We

noted less colocalization of Mcr puncta with Rab11-YFP-positive puncta, although we did find

some examples of co-localization (Figure 8B-B''). To determine if the co-localization of Mcr

and Rab5 have functional significance, we expressed a dominant negative Rab5 (*UAS-Rab5^{DN}*)

in the FCs using *GR1-GAL4* and examined the expression of Mcr and Cora in stage 11 FCs. In

wild-type FCs at that stage, Mcr and Cora are enriched at the apical-lateral membrane (n=91

cells, 15 egg chambers) (arrows in Figure 8 C), whereas both Mcr and Cora remains localized

along the lateral membrane in the Rab5^{DN}-expressing FCs (n=97 cells, 15 egg chambers) (Figure

8D). Similarly, Cora and Mcr co-localize at the apical-lateral membrane of the FCs of stage 12

egg chambers (n=64 cells, 15 egg chambers) (arrows in Figure 8E), whereas knocking down the

expression of Rab11 in stage 12 FCs results in the lateral localization of Mcr and/or Cora (n=23

cells out of 44, 16 egg chambers) (arrow in Figure 8F). Interestingly, we observed *Rab11-RNAi*

expressing cells in which Cora is mislocalized laterally and Mcr is completely missing from the

plasma membrane (n=21 cells out of 44, 16 egg chambers) (arrowhead in Figure 8F-F''),

suggesting that Mcr may have been endocytosed and shunted for degradation since it could not

be recycled. Taken together, these results suggest that similar to embryonic epithelia, the

maturation of SJs in the follicular epithelium requires Rab5-mediated endocytosis and Rab11-

mediated recycling.

Discussion

In this study, we have demonstrated that a subset of SJ proteins are expressed in egg chambers

throughout oogenesis and are required for critical morphogenetic processes that shape the egg,

including egg chamber elongation, dorsal appendage formation and border cell migration

(required to form the micropyle). Interestingly, the subcellular localization of these SJ proteins in

the ovarian FCs changes coincident with the establishment of the occluding junction in much the

same way that they do during embryogenesis in ectodermal epithelial cells (Tiklová *et al.* 2010),

suggesting that a similar maturation process for the SJ occurs in this tissue.

Biogenesis of the SJ in the FE

The mechanisms of SJ biogenesis in embryonic epithelia has been well-studied and involves

several steps in which transmembrane SJ proteins are first localized all along lateral plasma

membranes (by stage 12 of embryogenesis), but then must be endocytosed and recycled back to

the plasma membrane prior to aggregating in the region of the presumptive SJ (between stages

13 and 16; (Tiklová *et al.* 2010)). Prior to this relocalization step, SJ proteins show high mobility

in the plane of the membrane by Fluorescence Recovery After Photobleaching (FRAP) analysis,

but become strongly immobile as the relocalization is occurring (Oshima and Fehon 2011). As

these steps are occurring (e.g. stage 14 of embryogenesis), electron-dense material begins to

accumulate between adjacent cells in the presumptive SJ that takes on the appearance of ladder-

like septae by stage 17 of embryogenesis (Tepass and Hartenstein 1994). Functional studies

indicate that the occluding function of the junction is established late in embryogenesis, near the

end of stage 15 (Paul *et al.* 2003). Importantly, the process of SJ biogenesis is interdependent on

the function of all core components of the junction and several accessory proteins including Rab

5 and Rab 11 (Ward *et al.* 1998; Genova and Fehon 2003; Tiklová *et al.* 2010).

Here, we observe that many of these same steps occur in the maturation of SJs in the FE. We

first show that SJ proteins are expressed in the FE beginning early in ovarian development where

they localize all along the lateral membrane (Figure 1). These proteins appear to be endocytosed

later in egg chamber development (Figure 8) prior to becoming enriched at the presumptive SJ

by stage 10b/11 (Figure 2). The relocalization of SJ proteins to the SJ requires core SJ proteins

including Mcr and Nrx-IV, and accessory proteins including Rab 5 and Rab 11 (Figures 7 and 8).

Prior studies indicate the presence of electron dense material between FE cells as early as stage 6

(Müller 2000), with a ladder-like appearance of extracellular septae by stage 10B (Mahowald

1972). A recent study of the occluding function in the FE show a similar pattern of protein

localization for endogenously tagged Neuroglian-YFP and Lachesin-GFP, and demonstrates that

an intact occluding junction is formed during stage 11 of oogenesis (Isasti-Sanchez *et al.* 2020).

It is interesting to note the FE is a secondary epithelium initiated by a mesenchymal to epithelial

transition (Tepass *et al.* 2001), and yet SJ biogenesis appears to be very similar to that observed

in the primary epithelia found in the embryo.

SJ proteins are required for morphogenetic events during oogenesis

The similarities in the dynamic expression of SJ proteins in the FE and embryonic epithelia,

coupled with the observation that SJ proteins are required for numerous embryonic

developmental events ((Hall *et al.* 2014) and references therein) motivated us to explore whether

SJ proteins have similar roles in morphogenetic processes that shape the egg. Using a targeted

RNAi approach, we show that reducing the expression of Mcr, Nrx-IV, Cont, Cora, Sinu, or Lac

throughout oogenesis in the FCs results in significantly rounder stage 14 egg chambers, with

many showing additional defects in dorsal appendages (Figure 4). The initiation and

maintenance of egg elongation is achieved at various stages throughout oogenesis (Gates 2012;

Bilder and Haigo 2012; Cetera and Horne-Badovinac 2015). From stage 3-8, a gradient of JAK-

STAT signaling is required at each pole of the egg chamber to promote Myosin II-based apical

cell contractions (Alégot *et al.* 2018). From stage 6-8, the formation of a robust planar polarized

molecular corset - consisting of basal actin cytoskeleton and basement membrane protein fibrils -

is required for egg elongation, and requires collective FC migration over a static basement

membrane (Gutzeit *et al.* 1991; Frydman and Spradling 2001; Bateman *et al.* 2001; Viktorinová

et al. 2009; Haigo and Bilder 2011; Horne-Badovinac *et al.* 2012; Cetera *et al.* 2014; Isabella

and Horne-Badovinac 2016; Campos *et al.* 2020). During the middle stages of oogenesis (stages

9 and 10), basal actin stress fibers undergo actomyosin contractions, which contribute to egg

elongation (He *et al.* 2010; Qin *et al.* 2017). Finally, later in oogenesis, the maintenance of a

planar polarized molecular corset is required to retain an elongated egg chamber shape (Haigo

and Bilder 2011; Cha *et al.* 2017; Campos *et al.* 2020). Future studies are needed to determine if

SJ proteins are required for the establishment and/or maintenance of egg shape. Since many of

the events involved in egg elongation occur prior to the formation of a functional (and

ultrastructurally intact) occluding junction, it raises the possibility that SJ proteins have a

function in morphogenesis that is independent of their role in forming a tight occluding junction,

much as they do in the embryo.

Our observation that specifically knocking down the expression of several SJ proteins in

border cells (including polar cells using *C306-GAL4*) results in defective border cell migration

(Figure 6) supports a role for SJ proteins in morphogenesis, independent of their role in forming

an occluding junction. The phenotypes we observed include failing to complete migration by the

end of stage 10 and partial disassociation of the complex, although it is possible that these

phenotypes underestimate the full requirement of SJ proteins in border cell migration since this

process takes a relatively short time (4-6 hours) (Prasad and Montell 2007), while SJ proteins are

thought to be very stable in the membrane (Oshima and Fehon 2011). Nevertheless, these

phenotypes may indicate a role for SJ proteins in cell adhesion or cell polarity during migration.

Specifically, we note that Mcr appears to be enriched in border cells along membranes that are

closest to the oocyte (Figure 5). Perhaps the incomplete migration defect results from a

meandering migration through the nurse cells. A potential role for SJ proteins in cell adhesion in

the ovary has been reported previously. Reducing the level of Nrg in FCs results in the failure of

newly divided FCs to reintegrate into the FE, indicating a role for *Nrg* in lateral cell adhesion

(Bergstralh *et al.* 2015). In addition, expressing a null allele of *Nrg* in FCs enhances the invasive

tumor phenotype of a *Discs Large* (*Dlg*) mutation (Wei *et al.* 2004).

Unique roles for SJ proteins in the germarium

Unlike the polarized FE, the germarium consists of numerous unique cell types with distinct

functions. Cap cells, terminal filament cells and escort cells act as a niche for germ stem cell

renewal and differentiation (Eliazer and Buszczak 2011; Xie and Spradling 2000). Follicle stem

cells also reside in the germarium to produce the FE needed to encapsulate newly formed germ

cell cysts. It is interesting that Mcr, Cont, Nrx-IV, and Cora all show distinct expression pattern

within the germarium (Figure 3). For example, Cora is enriched in cap cells, terminal filament

cells and escort cells, whereas Nrx-IV outlines the outside membrane of region 1 and 2a of the

germarium. Mcr is highly expressed in the germ line stem cells, the cystoblasts and cystocytes

(Figure 3). This location-specific expression of SJ proteins provides a useful tool to identify

different cell types using antibodies against SJ proteins as markers.

In the *Drosophila* testes, Nrx-IV and Cora are expressed and are required for germ stem cell

differentiation by forming a permeability barrier at a late stage of spermatogenesis (Fairchild *et*

al. 2015). Additionally, Duby *et al.* showed that a total of five SJ proteins, including Nrx-IV and

Cora, co-localize with Dlg at the germline-somatic cell interfaces during the early and late

spermatogenesis (Dubey *et al*. 2019). It is unclear whether a similar permeability barrier is

established in the germarium, but the non-overlapping expression of SJ proteins in the

germarium argues against a SJ-like occluding function and instead suggests unique roles for

these proteins in aspects of early egg chamber development. For example, Nrx-IV and Cora are

expressed in cells/regions that are known to provide a niche for GSCs, suggesting a role for these

proteins in germline maintenance and differentiation, whereas Mcr may play a role in

maintaining cell-cell adhesion to regulate germline cell division and differentiation. Thus, future

studies could be aimed at identifying these unique functions as well as uncovering common

mechanisms by which SJ proteins facilitate morphogenetic processes during development.

Acknowledgements

We thank Jocelyn McDonald, the Bloomington *Drosophila* Stock Center, and the Vienna

Drosophila RNAi Center for fly stocks. We thank Manzoor Bhat and the Developmental Studies

Hybridoma Bank (created by the NICHD of the NIH and maintained at The University of Iowa,

Department of Biology, Iowa City, IA 52242) for antibodies used in this study. We thank Brian

Ackley for the use of his Olympus FV1000 confocal microscope. We thank Lindsay Ussher for

preliminary studies on border cell migration and thoughtful discussions on the project. We also

thank Sally Horne-Badovinac, Jocelyn McDonald, Yujun Chen, and members of the Ward lab

for helpful discussion about the project and manuscript.

References:

Alégot, H., P. Pouchin, O. Bardot, and V. Mirouse. 2018. Jak-Stat pathway induces Drosophila

follicle elongation by a gradient of apical contractility. *ELife* 7: 1–21.

https://doi.org/10.7554/elife.32943.

Banerjee, S, R. J. Bainton, N. Mayer, R Beckstead, and M. A. Bhat. 2008. Septate junctions are

required for ommatidial integrity and blood-eye barrier function in Drosophila. *Nature* 23:

1–7. https://doi.org/10.1038/jid.2014.371.

Bateman, Jack, R. Srekantha Reddy, Haruo Saito, and David Van Vactor. 2001. The receptor

tyrosine phosphatase dlar and integrins organize actin filaments in the Drosophila follicular

epithelium. *Curr Biol* 11: 1317–27. https://doi.org/10.1016/S0960-9822(01)00420-1.

Bätz, T., D. Förster, and S. Luschnig. 2014. The transmembrane protein macroglobulin

complement-related is essential for septate junction formation and epithelial barrier function

in Drosophila. *Development* 141: 899–908. https://doi.org/10.1242/dev.102160.

Baumgartner, S., J. T. Littleton, K. Broadie, M. A. Bhat, R. Harbecke *et al.*, 1996. A Drosophila

neurexin is required for septate junction and blood-nerve barrier formation and function.

Cell 87: 1059–68. https://doi.org/10.1016/S0092-8674(00)81800-0.

Behr, M., D. Riedel, and R. Schuh. 2003. The claudin-like megatrachea is essential in septate

junctions for the epithelial barrier function in Drosophila. *Dev Cell* 5: 611–620.

https://doi.org/10.1016/S1534-5807(03)00275-2.

Ben-Zvi, D. S., and T. Volk. 2019. Escort cell encapsulation of Drosophila germline cells Is

maintained by irre cell recognition module proteins. *Biol Open* 8.

https://doi.org/10.1242/bio.039842.

Bilder, D., and S. L. Haigo. 2012. Expanding the morphogenetic repertoire: rerspectives from the

Drosophila egg. *Dev Cell* 22: 12–23. https://doi.org/10.1016/j.devcel.2011.12.003.

Bopp, D., L. R. Bell, T. W. Cline, and P. Schedl. 1991. Developmental distribution of female-

specific sex-lethal proteins in Drosophila melanogaster. *Genes Dev* 5: 403–15.

https://doi.org/10.1101/gad.5.3.403.

Bopp, D., J. I. Horabin, R. A. Lersch, T. W. Cline, and P. Schedl. 1993. Expression of the sex-

lethal gene Is controlled at multiple levels during Drosophila oogenesis. *Development* 118:

797–812.

Brand, A. H., and N. Perrimon. 1993. Targeted gene expression as a means of altering cell fates

and generating dominant phenotypes. *Development* 118: 401–15.

Campos, F. C., C. Dennis, H. Alégot, C. Fritsch, A. Isabella *et al*., 2020. Oriented basement

membrane fibrils provide a memory for f-actin planar polarization via the Dystrophin-

Dystroglycan complex during tissue elongation. *Development* 147 (7).

https://doi.org/10.1242/dev.186957.

Cetera, M., and S. Horne-Badovinac. 2015. Round and round gets you somewhere: collective

cell migration and planar polarity in elongating Drosophila egg chambers. *Curr Opin Genet*

Dev 32: 10–15. https://doi.org/10.1016/j.gde.2015.01.003.

Cetera, M., G. R. Ramirez-San Juan, P. W. Oakes, L. Lewellyn, M. J. Fairchild *et al*., 2014.

Epithelial rotation promotes the global alignment of contractile actin bundles during

Drosophila egg chamber elongation. *Nat Commun* 5: 5511.

https://doi.org/10.1038/ncomms6511.

Cha, I. J., J. H. Lee, K. S. Cho, and S. B. Lee. 2017. Drosophila tensin plays an essential role in

cell migration and planar polarity formation during oogenesis by mediating integrin-

dependent extracellular signals to actin organization. *Biochem and Biophys Res Commun*

484: 702–9. https://doi.org/10.1016/j.bbrc.2017.01.183.

Dubey, P., T. Kapoor, S. Gupta, S. Shirolikar, and K. Ray. 2019. Atypical septate junctions

maintain the somatic enclosure around maturing spermatids and prevent premature sperm

release in Drosophila testis. *Biol Open* 8: bio036939. https://doi.org/10.1242/bio.036939.

Duhart, J. C., T. T. Parsons, and L. A. Raftery. 2017. The repertoire of epithelial morphogenesis

on display: progressive elaboration of Drosophila egg structure. *Mech Dev*. 148: 13-39.

https://doi.org/10.1016/j.mod.2017.04.002.

Eliazer, S., and M. Buszczak. 2011. Finding a niche: studies from the Drosophila ovary. *Stem*

Cell Res Ther. 2:45. https://doi.org/10.1186/scrt86.

Fairchild, M. J., C. M. Smendziuk, and G. Tanentzapf. 2015. A somatic permeability barrier

around the germline is essential for Drosophila spermatogenesis. *Development* 142: 268–81.

https://doi.org/10.1242/dev.114967.

Faivre-Sarrailh, C., S. Banerjee, J. Li, M. Hortsch, L. Monique *et al*., 2004. Drosophila contactin,

a homolog of vertebrate contactin, is required for septate junction organization and

paracellular barrier function. *Development* 131: 4931–42.

https://doi.org/10.1242/dev.01372.

Fehon, R. G., I. A. Dawson, and S. Artavanis-Tsakonas. 1994. A *Drosophila* homologue of

membrane-skeleton protein 4.1 is associated with septate junctions and is encoded by the

coracle gene. *Development* 120: 545–57. https://doi.org/10.1083/jcb.120.1.129.

Felix, M., M. Chayengia, R. Ghosh, A. Sharma, and M. Prasad. 2015. Pak3 regulates apical-

basal polarity in migrating border cells during Drosophila oogenesis. *Development* 142:

3692–3703. https://doi.org/10.1242/dev.125682.

Frydman, H. M., and A. C. Spradling. 2001. The receptor-like tyrosine phosphatase lar is

required for epithelial planar polarity and for axis determination within Drosophila ovarian

follicles. *Development* 128: 3209–20.

Gates, J. 2012. Drosophila egg chamber elongation: insights into how tissues and organs are

shaped." *Fly* 6: 213–27. https://doi.org/10.4161/fly.21969.

Genova, J. L., and R. G. Fehon. 2003. Neuroglian, gliotactin, and the Na+/K+ ATPase are

essential for septate junction function in Drosophila. *J of Cell Biol* 161: 979–89.

https://doi.org/10.1083/jcb.200212054.

Gupta, T., and T. Schüpbach. 2003. Cct1, a phosphatidylcholine biosynthesis enzyme, is required

for Drosophila oogenesis and ovarian morphogenesis." *Development* 130: 6075–87.

https://doi.org/10.1242/dev.00817.

Gutzeit, H. O, W. Eberhardt, and E. Gratwohl. 1991. Laminin and basement membrane-

associated microfilaments in wild-type and mutant Drosophila ovarian follicles. *J Cell Sci*

100: 781–88.

Haigo, S. L., and D. Bilder. 2011. Global tissue revolutions in a morphogenetic movement

controlling elongation. *Science* 331: 1071–74. https://doi.org/10.1126/science.1199092.

Hall, S., C. Bone, K. Oshima, L. Zhang, M. McGraw *et al.*, 2014. Macroglobulin complement-

related encodes a protein required for septate junction organization and paracellular barrier

function in Drosophila. *Development* 141: 889–98. https://doi.org/10.1242/dev.102152.

Hall, S., and R. E. Ward. 2016. Septate junction proteins play essential roles in morphogenesis

throughout embryonic development in Drosophila." *G3* 6: 2375–84.

https://doi.org/10.1534/g3.116.031427.

He, L., X. Wang, H. L. Tang, and D. J. Montell. 2010. Tissue elongation requires oscillating

contractions of a basal actomyosin network. *Nat Cell Biol* 12:1133-1142.

https://doi.org/10.1038/ncb2124.

Hildebrandt, A., R. Pflanz, M. Behr, T. Tarp, D. Riedel, and R. Schuh. 2015. Bark beetle

controls epithelial morphogenesis by septate junction maturation in Drosophila. *Dev Biol*

400: 237–47. https://doi.org/10.1016/j.ydbio.2015.02.008.

Horne-Badovinac, S. 2020. The Drosophila micropyle as a system to study how epithelia build

complex extracellular structures. *Phil Trans R Soc* 375: 20190561.

https://doi.org/10.1098/rstb.2019.0561.

Horne-Badovinac, S., and D. Bilder. 2005. Mass transit: epithelial morphogenesis in the

Drosophila egg chamber. *Dev Dyn* 232: 559–74. https://doi.org/10.1002/dvdy.20286.

Horne-Badovinac, S., J. Hill, G. Gerlach, W. Menegas *et al*., 2012. A Screen for round egg

mutants in Drosophila identifies tricornered, furry, and misshapen as regulators of egg

chamber elongation. *G3* 2: 371–78. https://doi.org/10.1534/g3.111.001677.

Ile, K. E., R. Tripathy, V. Goldfinger, and A. D. Renault. 2012. Wunen, a Drosophila lipid

phosphate phosphatase, is required for septate junction-mediated barrier function.

Development 139: 2535–46. https://doi.org/10.1242/dev.077289.

Isabella, A. J., and S. Horne-Badovinac. 2016. Rab10-mediated secretion synergizes with tissue

movement to build a polarized basement membrane architecture for organ morphogenesis.

Dev Cell 38: 47–60. https://doi.org/10.1016/j.devcel.2016.06.009.

Isasti-Sanchez, J., F. Munz-Zeise, and S. Luschnig. 2020. Transient opening of tricellular

vertices controls paracellular transport through the follicle epithelium during Drosophila

oogenesis. *BioRxiv*, no. CiM: 2020.02.29.971168.

https://doi.org/10.1101/2020.02.29.971168.

Ward, R. E., R. S. Lamb, and R. G. Fehon. 1998. A conserved functional domain of Drosophila

coracle is required for localization at the septate junction and has membrane-organizing

activity. *Cell* 140: 1463–73.

Izumi, Y., and M. Furuse. 2014. Molecular organization and function of invertebrate occluding

junctions. *Semin Cell Dev Bio* 36: 186–93. https://doi.org/10.1016/j.semcdb.2014.09.009.

Khammari, A., F. Agnès, P. Gandille, and A. M. Pret. 2011. Physiological apoptosis of polar

cells during Drosophila oogenesis is mediated by hid-dependent regulation of diap1. *Cell*

Death and Differ. 18: 793–805. https://doi.org/10.1038/cdd.2010.141.

Kirilly, D., and T. Xie. 2007. The Drosophila ovary: an active stem cell community. *Cell Res* 17:

15–25. https://doi.org/10.1038/sj.cr.7310123.

Lamb, R. S., R. E. Ward, L. Schweizer, and R. G. Fehon. 1998. Drosophila coracle, a member of

the protein 4.1 superfamily, has essential structural functions in the septate junctions and

developmental functions in embryonic and adult epithelial cells. *Mol Bio Cell* 9: 3505–19.

https://doi.org/10.1091/MBC.9.12.3505.

Mahowald, A. P. 1972. Ultrastructural observations on oogenesis in Drosophila. *J Morph* 137:

29–48. https://doi.org/10.1002/jmor.1051370103.

Maimon, I., M. Popliker, and L. Gilboa. 2014. Without children is required for stat-mediated

zfh1 transcription and for germline stem cell differentiation. *Development* 141: 2602–10.

https://doi.org/10.1242/dev.109611.

Montell, D. J. 2003. Border-Cell Migration: The Race Is On. *Nat Rev Mol Cell Bio* 4: 13–24.

https://doi.org/10.1038/nrm1006.

Moyer, K. E., and J. R. Jacobs. 2008. Varicose: a MAGUK required for the maturation and

function of Drosophila septate junctions. *BMC Dev. Bio*. 8;99. https://doi.org/10.1186/1471-

213X-8-99.

Müller, H. J. 2000. Genetic control of epithelial cell polarity: lessons from Drosophila. *Dev Dyn*

218: 52–67. https://doi.org/10.1002/(sici)1097-0177(200005)218:1<52::aid-

dvdy5>3.3.co;2-c.

Murphy, A. M., and D. J. Montell. 1996. Cell type-specific roles for cdc42, rac, and rho1 in

Drosophila oogenesis. *J Cell Bio* 133: 617–30. https://doi.org/10.1083/jcb.133.3.617.

Nelson, K. S., M. Furuse, and G.J. Beitel. 2010. The Drosophila claudin kune-kune is required

for septate junction organization and tracheal tube size control. *Genetics* 185: 831–39.

https://doi.org/10.1534/genetics.110.114959.

Noirot-timothée, C., D. S. Smith, M. L. Cayer, and C. Noirot. 1978. Septate junctions in insects:

comparison between intercellular and intramembranous structures." *Tissue Cell* 10: 125–36.

https://doi.org/10.1016/0040-8166(78)90011-3.

Ogienko, A. A., L. A. Yarinich, E. V. Fedorova, M. O. Lebedev *et al.*, 2018. New slbo-gal4

driver lines for the analysis of border cell migration during Drosophila oogenesis.

Chromosoma 127: 475–87. https://doi.org/10.1007/s00412-018-0676-7.

Oshima, K., and R. G. Fehon. 2011. Analysis of protein dynamics within the septate junction

reveals a highly stable core protein complex that does not include the basolateral polarity

protein discs large." *J Cell Sc* 124: 2861–71. https://doi.org/10.1242/jcs.087700.

Paul, S. M., M. Ternrt, P. Salvaterra, and G. Beitel. 2003. The Na+/K+ ATPase is required for

septate junction function and epithelial tube-size control in the Drosophila tracheal system.

Development 130: 4963–74. https://doi.org/10.1242/dev.00691.

Prasad, M., and D. J. Montell. 2007. Cellular and molecular mechanisms of border cell migration

analyzed using time-lapse live-cell imaging. *Dev Cell* 12: 997–1005.

https://doi.org/10.1016/j.devcel.2007.03.021.

Qin, X., B. O. Park, J. Liu, B. Chen, V. Choesmel-Cadamuro *et al*., 2017. Cell-matrix adhesion

and cell-cell adhesion differentially control basal myosin oscillation and Drosophila egg

chamber elongation. *Nat Commun*. 8: 14708. https://doi.org/10.1038/ncomms14708.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al*., 2012. Fiji: an open-

source platform for biological-image analysis. *Nat Methods* 9: 676–82.

https://doi.org/10.1038/nmeth.2019.

Schneider, M., A.K. Khalil, J. Poulton, C. Castillejo-Lopez, D. Egger-Adam *et al*., 2006.

Perlecan and dystroglycan act at the basal side of the Drosophila follicular epithelium to

maintain epithelial organization. *Development* 133: 3805–15.

https://doi.org/10.1242/dev.02549.

Schulte, J., U. Tepass, and V. J. Auld. 2003. Gliotactin, a Novel Marker of Tricellular Junctions,

Is Necessary for Septate Junction Development in Drosophila. *Journal Cell Bio* 161 (5):

991–1000. https://doi.org/10.1083/jcb.200303192.

Snow, Peter M., Allan J. Bieber, and Corey S. Goodman. 1989. Fasciclin III: a novel homophilic

adhesion molecule in Drosophila. *Cell* 59: 313–23. https://doi.org/10.1016/0092-

8674(89)90293-6.

Spradling, A. C., M. De Cuevas, D. Drummond-Barbosa, L. Keyes, M. Lilly *et al*., 1997. The

Drosophila germarium: stem cells, germ line cysts, and oocytes. *Cold Spring Harb Symp*

Quant Biol 62: 25–34.

Tepass, U., G. Tanentzapf, R. Ward, and R. Fehon. 2001. Epithelial cell polarity and cell

junctions in Drosophila. *Annu Rev Gene*. 35: 747–84.

https://doi.org/10.1146/annurev.genet.35.102401.091415.

Tepass, Ulrich, and Volker Hartenstein. 1994. The development of cellular junctions in the

Drosophila embryo. *Dev Biol* 161:563-96. https://doi.org/10.1006/dbio.1994.1054.

Tiklová, K., K. A. Senti, S. Wang, A. GräCurrency Signslund, and C. Samakovlis. 2010.

Epithelial septate junction assembly relies on melanotransferrin iron binding and

endocytosis in Drosophila. *Nature Cell Biol*. 12: 1071–77. https://doi.org/10.1038/ncb2111.

Venema, D. R., T. Zeev-Ben-Mordehai, and V. J. Auld. 2004. Transient apical polarization of

gliotactin and coracle is required for parallel alignment of wing hairs in Drosophila. *Dev*

Biol 275: 301–14. https://doi.org/10.1016/j.ydbio.2004.07.040.

Viktorinová, I., T. König, K. Schlichting, and C. Dahmann. 2009. The cadherin fat2 is required

for planar cell polarity in the Drosophila ovary. *Development* 136: 4123–32.

https://doi.org/10.1242/dev.039099.

Ward, R. E., R. S. Lamb, and R. G. Fehon. 1998. A conserved functional domain of Drosophila

coracle is required for localization at the septate junction and has membrane-organizing

activity. *Cell* 140: 1463–73.

Wei, J., M. Hortsch, and S. Goode. 2004. Neuroglian Stabilizes Epithelial Structure during

Drosophila Oogenesis. *Dev*. *Dyn*. 230: 800–808. https://doi.org/10.1002/dvdy.20108.

Wittes, J., and T. Schüpbach. 2018. A gene expression screen in Drosophila melanogaster

identifies novel JAK/STAT and EGFR targets during oogenesis. *G3* 9: 47–60.

https://doi.org/10.1534/g3.118.200786.

Wu, V. M., J. Schulte, A. Hirschi, U. Tepass, and G.J. Beitel. 2004. Sinuous is a Drosophila

claudin required for septate junction organization and epithelial tube size control. *J Cell*

Biol. 164: 313–23. https://doi.org/10.1083/jcb.200309134.

Wu, V. M., M. H. Yu, R. Paik, S. Banerjee, Z. Liang *et al*., 2007. Drosophila varicose, a member

of a new subgroup of basolateral MAGUKs, is required for septate junctions and tracheal

morphogenesis. *Development* 134: 999–1009. https://doi.org/10.1242/dev.02785.

Xie, T., and A. C. Spradling. 2000. A niche maintaining germ line stem cells in the Drosophila

ovary. *Science* 290: 328–30. https://doi.org/10.1126/science.290.5490.328.

Figure Legends

Figure 1. Mcr, Cont, Nrx-IV, and Cora expression during early stages of oogenesis. (A)

Diagram of a female *Drosophila* ovariole. Egg chambers are formed in the most anterior region

of the ovariole called the germarium (Germ). Each egg chamber undergoes 14 developmental

stages while connecting to each other through stalk cells (SCs) to form a mature stage 14 egg.

(B) Diagram of a stage 8 egg chamber. An egg chamber consists of 15 nurse cells (NCs) and one

oocyte (Oo.), which are surrounded by a monolayer of follicle cells (FCs). At the anterior and

posterior ends of an egg chamber resides a pair of differentiated FCs called polar cells (PCs). (C)

Diagram of the follicular epithelium in a stage 10b egg chamber. FCs have defined apical-basal

polarity with the apical surface facing the germline. By stage 10b the septate junction (SJ) has

formed along the lateral membrane. At that stage, the junctional complex consists of a marginal

zone (MZ), an adherens junction (AJ), and a SJ. (D-G) Confocal optical sections of wild type

early stages egg chambers stained with antibodies against Mcr (D and single channel in D'),

Cont (E and E'), Nrx-IV (F and F'), and Cora (G and G'), and co-stained with antibodies against

Ecad (green) and labeled with DAPI. All four SJ proteins are expressed throughout the egg

chamber along FC membranes and in the NCs. Mcr, Cont, and Nrx-IV (D-F) are found along the

membrane and in puncta, whereas Cora is found predominantly at the membrane (G). In

addition, Mcr, Cont, and Nrx-IV are highly expressed in the PCs (asterisks in D-F), whereas

Cora is expressed in these cells with same level of expression relative to the follicle cells (data

not shown). Anterior is to the left in each ovariole. Scale bar= 25µm.

Figure 2. Mcr, Cont, Nrx-IV, and Cora localization at later stages of oogenesis. (A) Diagram

of a stage 12 egg chamber, showing the area where confocal images were taken in panels B, D, F

and H (red box) which is the boundary between the oocyte (Oo.) and nurse cells (NCs). Confocal

optical sections of wild type stage 11 – 12 egg chambers (B, D, F, and H) or stage 14 (C, E, G,

and I) stained with antibodies against Mcr (B and C), Cont (D and E), Nrx-IV (F and G), or Cora

(H and I) and co-stained with antibodies against Ecad (green) and labeled with DAPI (blue).

Individual channels for SJ protein staining are shown in (B'-H'). Note that Mcr, Cont, Nrx-IV,

and Cora become enriched at the apical-lateral region of FCs membrane (arrow in B, D, F, and

H). The expression of all of these proteins persists in stage 14 egg chambers (C, E, G, and I).

Anterior is to the left. Scale bar = 25µm in B, D, F, and H and 100µm in C, E, G, and I.

Figure 3. Mcr, Cont, Nrx-IV, and Cora expression in the germarium. (A) Diagram of a

germarium showing regions 1 and 2 (the full germarium has 3 regions). Terminal filament (TF)

and cap cells (CPCs) reside at the most anterior tip of the germarium. In region 1, germline stem

cells (GSCs) divide to give rise to a cystoblast (CB), which undergoes four rounds of cell

division forming a 16-cell cyst (CCs). In region 2a of the germarium, CCs move posteriorly as

they are encapsulated by the prefollicle cells in region 2b. (B-E) Confocal optical sections of

wild type germaria stained with antibodies against Mcr (B and single channel in B'), Cont (C and

C'), Nrx-IV (D and D'), and Cora (E and E') and co-stained with antibodies against Sxl (B) or

Ecad (green) and labeled with DAPI (blue) (C-E). (B) Mcr is highly expressed in GSCs

(asterisks), CBs (arrow), and developing cysts (arrowhead), whereas Cont is found in puncta

throughout the germarium region (C). Nrx-IV is found in puncta throughout the germarium and

outlines the TFs (arrow in D) and the outside membrane of region 1 and region 2a (arrowheads

in D). However, the localization of Nrx-IV at the outside membrane in the germarium varies as it

was observed in 5 out of 7 germaria examined. (E) Cora is expressed throughout the germarium

with high level in the TF (arrow), CPCs (arrowhead) and ESCs (yellow arrows). Anterior is to

the left. Scale bar= 25μm.

Figure 4. *SJ* genes are required for egg elongation. (A-G) Brightfield photomicrographs of

stage 14 egg chambers. (A) *GR1-GAL4>UAS-GFP*, (B) *GR1-GAL4>UAS-Cont-RNAi*, (C) *GR1-*

GAL4>UAS-Nrx-IV-RNAi, (D) *GR1-GAL4>UAS-Mcr-RNAi*, (E) *GR1-GAL4>UAS-Cora-RNAi*,

(F) *GR1-GAL4>UAS-Sinu-RNAi*, and (G) *GR1-GAL4>UAS-Lac-RNAi*. Images in this figure

represent average phenotypes observed in each genotype. (H and I) Quantification of length and

width of stage 14 egg chambers from control and *SJ-RNAi* egg chambers. (J) Quantification of

the aspect ratio of length (orange line in A) to width (yellow line in A) from control and *SJ-RNAi*

stage 14 egg chambers. Note that the aspect ratio of all *SJ-RNAi* expressing egg chambers are

statistically significant from the control egg chambers (unpaired T-test; $P<0.0001$). (K) Zoom in

images of *GR1-GAL4>UAS-GFP* and *GR1-GAL4>UAS-Lac-RNAi* stage 14 egg chambers

showing the dorsal appendages (L). In the control, dorsal appendages are formed, whereas the

dorsal appendages in eggs expressing *Lac-RNAi* are either deformed or absent (arrows). (M)

Quantification of dorsal appendage defects from control and *SJ-RNAi* stage 14 egg chambers. *n*,

total number of egg chambers measured per genotype. Data represents individual values with

mean ± s.d. *P* value <0.0001. Anterior is to the left. Scale bar = 100μm.

Figure 5. Mcr expression during border cell migration. (A) Diagram of border cell migration.

At early stage 9 of oogenesis, a group of 4-6 follicle cells are specified by the polar cells (PCs)

(yellow) to become border cells (BCs) (red). The BC/PC cluster delaminates apically and

migrates between the nurse cells (NCs) until it reaches the oocyte (Oo.) by stage 10 of oogenesis.

(B-D) Confocal optical images of wild type stage 9-10 egg chambers (insets show border cell

clusters) stained with antibodies against Mcr (red and in " channel), Fas3 (green and in "'

channel), and labeled with DAPI (blue and in '"" channel). Mcr is expressed in the PCs and BCs

with the highest expression at the interface between the PCs and BCs. Mcr is also enriched at the

leading edge of migration (red arrows in B"-D"). Anterior is to the left. Scale bar = 25μm in (B-

D) and 10μm in (B'-B'''' through D'-D'''').

Figure 6. Mcr, Cont, Nrx-IV, and Cora are required for effective border cell migration. (A-

C) Border cell migration in control egg chambers. Confocal optical section of *Slbo>mCD8-GFP*

egg chambers immunostained with anti-Fas3 (red) to mark the polar cells, GFP (green) to mark

border cells and labeled with DAPI (blue). Arrow indicates a border cell cluster that has

completed migration to the oocyte by stage 10. (D-F) Stage 10 egg chambers expressing *Cont-*

RNAi in border cells immunostained with anti-Cont (red) and anti-Fas3 (blue) as examples of BC

cluster migration defects in *SJ-RNAi* egg chambers. Failed (arrow in D), incomplete (arrow in E;

note two dashed lines indicate the range of distance at which border cell cluster was categorized

as incomplete), or disassociated border cell migration (arrow in F). Higher magnification of wild

type (G) and Cont-RNAi-expressing (H) border cell cluster showing the dissociation of border

cell cluster observed in *SJ-RNAi* clusters. (I and J) Quantification of border cell cluster

phenotypes at stage 10 egg chambers in control and *SJ-RNAi* driven by either Slbo-GAL4 (I) or

C306-GAL4 (J) line. Anterior is to the left. Scale bar = 25μm.

Figure 7. SJ localization of Cora and Mcr depends on the expression of other SJ proteins.

(A-D) Confocal optical sections of stage 12 follicle cells (FCs) stained with Mcr (red and '

channel) and Cora (green and " channel) and labeled with DAPI (blue). In wild type FCs, Mcr

and Cora co-localize in the region of the SJ (A), whereas Mcr and Cora are mislocalized along

the lateral membrane of *NrxIV-RNAi* expressing FCs (B). In *Mcr-RNAi* expressing FCs, Cora is

mislocalized laterally (C), however, when Cora appears to maintain its enrichment at the SJ, we

often see residual Mcr protein (arrow, D). Scale bar = 10µm (A-D) and 0.5µm (E and F).

Figure 8. SJ localization of Cora and Mcr requires Rab5 and Rab11. (A and B) Confocal

optical sections of stage 10 follicle cells (FCs) expressing Rab5-YFP (A) or Rab11-YFP (B)

stained with antibody against GFP (green and ' channel) and Mcr (red and " channel). Mcr

cytoplasmic puncta partially co-localize with Rab5 (white circles) (A-A'') and Rab11 (B-B'').

(C and D) Confocal optical sections of stage 11 control (C) and Rab5^{DN} expressing (D) FCs

expressing Mcr (red and in ' channel) and Cora (green and in " channel) and labeled with DAPI

(blue). Note that Mcr and Cora co-localize at the region of the SJ in control FCs (arrows), but are

mislocalized along the lateral membrane in Rab5^{DN} FCs. (E and F) Confocal optical sections of

stage 12 control (E) and Rab11-RNAi expressing (F) FCs expressing Mcr (red and in ' channel)

and Cora (green and in " channel) and labeled with DAPI (blue). Mcr and Cora are mislocalized

in Rab11-RNAi expressing FCs (compare arrows in E and F). In some instances, it appears that

Mcr is lost from the membrane in Rab11-RNAi expressing FCs (arrowhead in F). Anterior is to

the left. Scale bars = 10μm.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure S1. Cont expression during border cell migration. (A-C) Confocal optical images of wild type stage 9-10 egg chambers (insets show border cell clusters) stained with antibodies against Cont (red and in " channel), Fas3 (green and in "' channel), and labeled with DAPI (blue and in '"' channel). Cont is expressed in the border cell cluster throughout oogenesis, with high level at the polar cells-border cells interfaces (asterisks in A', B' and C'). Anterior is to the left. Scale bar = 25μm.

Figure S2. Nrx-IV expression during border cell migration. (A-C) Confocal optical images of wild type stage 9-10 egg chambers (insets show border cell clusters) stained with antibodies against Nrx-IV (red and in " channel), Fas3 (green and in "' channel), and labeled with DAPI (blue and in "" channel). Nrx-IV is expressed in both polar cells (PCs) and border cells (BCs) throughout border cell migration, with high level at the PCs-BCs interfaces (asterisks in A', B' and C'). Anterior is to the left. Scale bar = 25μm.

Figure S3. Cora expression during border cell migration. (A-C) Confocal optical images of wild type stage 9-10 egg chambers (insets show border cell clusters) stained with antibodies against Cora (red and in ” channel), Fas3 (green and in ”’ channel), and labeled with DAPI (blue and in ”” channel). (A) At early stage 9, Cora is expressed in both polar cells (PCs) and border cells (BCs). (B) During middle stage 9 of oogenesis, Cora continues to be expressed in the cluster until stage 10 with the highest level in the PCs relative to the BCs (C) (asterisks in A', B' and C'). Anterior is to the left. Scale bar in A-C = 25μm and 10μm in A'-A'''', B'-B'''', and C'-C''''.