The *Thermosynechococcus* genus: wide environmental distribution, but a highly

conserved genomic core

A. Paulina Prondzinsky[1,2]*, Sarah J. Berkemer[3,4], Lewis M. Ward[2,5], Shawn E. McGlynn[2]*

[1]Department of Chemical Science and Engineering, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8550 Japan

[2]Earth-Life Science Institute, Ota-ku, Tokyo 145-0061, Japan

[3]Bioinformatics Group, Department of Computer Science, University Leipzig, Leipzig, Germany

[4]Competence Center for Scalable Data Services and Solutions, Dresden/Leipzig, Germany

[5]Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA

*prondzinsky@elsi.jp / mcglynn@elsi.jp

Abstract

Cyanobacteria thrive in very diverse environments. In Earth history however, delayed oxygenation has

raised questions of growth limitation in ancient environmental conditions. As a single genus, the

Thermosynechococcus are known to be cosmopolitan and live in chemically diverse habitats. To

understand the genetic basis for this, we compared the protein coding component of

Thermosynechococcus genomes. Supplementing the known genetic diversity of *Thermosynechococcus*,

we report draft metagenome-assembled genomes of two *Thermosynechococcus* recovered from ferrous

carbonate hot springs in Japan. We find that as a genus, *Thermosynechococcus* is genomically

conserved, having a small pan-genome with few accessory genes per individual strain and only 18

protein clusters appearing in all *Thermosynechococcus* but not in any other cyanobacteria in our

analysis. Furthermore, by comparing orthologous protein groups, including an analysis of genes

encoding proteins with an iron related function (uptake, storage or utilization), no clear differences in

genetic content, or adaptive mechanisms could be detected between genus members, despite the range

of environments they inhabit. Overall, our results highlight a seemingly innate ability for

Thermosynechococcus to inhabit diverse habitats without having undergone substantial genomic

adaptation to accommodate this. The finding of *Thermosynechococcus* in both hot and high iron

environments without adaptation recognizable from the perspective of protein coding genes has

implications for understanding the basis of thermophily within this clade, and also suggests that ferrous

iron in ancient oceans may not have inhibited the proliferation of Cyanobacteria on Earth. The conserved

core genome may be indicative of an allopatric lifestyle – or reduced genetic complexity of hot spring

habitats relative to other environments.

Keywords: cyanobacteria, great oxygenation event, hot springs, comparative genomics, pan-genome,

Thermosynechococcus

Introduction

Water oxidizing cyanobacteria fundamentally altered the distribution of carbon and electrons on Earth (Canfield, 1998; Raven, 2009; Ward and Shih, 2019). A marker of this reorganization is in the Great Oxygenation Event (GOE), which marks a major transition in the evolution of life on Earth ~2.3 billion years ago (Lyons *et al.*, 2014; Fischer *et al.*, 2016). While the GOE is widely accepted to have been driven by the production of O_2 by oxygenic photosynthesis performed by members of the Cyanobacteria, the timing and proximal trigger for the GOE is debated. At least six hypotheses for the timing of the GOE can be considered i) the evolution of oxygenic photosynthesis by Cyanobacteria just before the GOE (Fischer *et al.*, 2016; Shih *et al.*, 2017), ii) an earlier evolution of Cyanobacteria with O_2 accumulation delayed due to the transition of cyanobacteria from small-scale freshwater to large-scale marine environments (Sánchez-Baracaldo, 2015), iii) the transition from unicellular to multicellular organisms for increased evolutionary success (Schirrmeister *et al.*, 2015), iv) the inhibition of early cyanobacteria due to high iron concentrations (Swanner, Mloszewska, *et al.*, 2015; Swanner, Wu, *et al.*, 2015), v) a possible nitrogen throttle on cyanobacterial growth (Fennel *et al.*, 2005; Shi and Falkowski, 2008; Allen *et al.*, 2019), vi) or depressed Archaean productivity due to phosphate availability (Hao *et al.* 2020). Although cyanobacterial taxonomy continues to be refined (Knoll 2006; Tomitani *et al.*, 2006; Shih *et al.*, 2013; Soo *et al.*, 2019; Parks *et al.*, 2020), understanding the ecological distribution of cyanobacteria may help us generate, or reject hypotheses about their evolutionary trajectories, and the factors which led to the observed timing of the GOE.

Name	Strain name (Thermosynechococcus)	Max. Temperature [°C]	pH	Sulfate [mM]	Iron [mM]	References
Kangding Geothermal Field (Sichuan, China)	*T. elongatus* PKUAC-SCTE54	94	6.35 – 8.84	1.2	0.0104	Mei-hua *et al.*, 2015; Guo *et al.*, 2017; Tang *et al.*, 2018; Wang *et al.*, 2019
Okuoku-hachikurou Hot Spring (Akita, Japan)	*T. nakabusensis* OHK	44	6.8	6.5	0.114	Ward *et al.*, 2017
Yunomine Hot Spring (Wakayama, Japan)	*T. vulcanus* NIES2134	91	8	0.06 – 0.229	< 0.0018	Ichikuni and Kikuchi, 1972 Onsen information sheet*
Nakabusa Hot Spring (Nagano, Japan)	*T. nakabusensis* NK55a	76	8.5 – 9	0.218 – 0.246	0.0004	Nakagawa and Fukui, 2002; Nakamura *et al.*, 2002; Stolyar *et al.*, 2014 Onsen information sheet*
Jinata Hot Spring (Shikinejima,	*T. nakabusensis* Jinata	63	5.4	17.4	0.261	Ward *et al.*, 2019

Tokyo, Japan)						
Chung-Lun Hot Spring (Taiwan)	*T.* CL1	62	9.3	1.35 – 1.39	0.0006	Yeh *et al.*, 2005; Hsueh *et al.*, 2007; Maity *et al.*, 2011; Cheng *et al.*, 2020
Beppu Hot Spring Kamegawa Shinoyu (Oita, Japan)	*T. elongatus* BP1	78	6.8	1.09	0.0036	Onsen information sheet*

Table 1: Geochemical parameters for each hot spring. References are related to the first publication of the strains

or the geochemistry of the hot spring. *information available online through local governments, last accessed in

11/2019.

Cyanobacteria are found in a wide range of environments – the knowledge of which continues to expand

(Whitton, 1992; Puente-Sánchez *et al.*, 2018; Callieri *et al.*, 2019). As a single phylogenetic group,

members of the *Thermosynechococcus* genus have been documented to inhabit a range of chemical

environments, some of which might be similar to places and times on the early Earth. Those

Thermosynechococcus with genomes available are from hot springs which vary in temperature from 44

– 94 °C, pH ranging from 5.4 – 9.3, sulfate concentrations between 0.06 mM and 17.4 mM and iron

concentrations between 0.0004 mM and 0.261 mM (Table 1) at their source. It is noteworthy that for two

of the genomes - those from Okuoku-hachikurou and Jinata hot springs - the ferrous iron concentrations

at the source by far exceed those at the other springs with 114 µM at OHK and 261 µM at Jinata (Ward

et al., 2019; Cheng *et al.*, 2020).

Seminal work by Papke et al. on phylotype:geographical relationships of *Thermosynechococcus* posed

questions as to how these organisms could be so widely distributed: in their analysis, the distribution of

Thermosynechococcus could not be explained by measured geochemical parameters.

Thermosynechococcus thus appear to be cosmopolitan, but the basis for this remains unresolved.

Motivated by the finding of *Thermosynechococcus* members in ferrous iron carbonate hot springs that

we have been studying (Ward *et al.*, 2017, 2019), and in an attempt to find a genetic basis for the

geochemically wide distribution of the organisms, we took a comparative genomics approach to search

for resolvable traits which may underlie environmental adaptations of *Thermosynechococcus* members.

Figure 1: *Thermosynechococcaceae* phylogeny built with concatenated ribosomal proteins. Branch supports are derived from bootstrapping with BOOSTER and the tree scale bar indicates substitutions per nucleotide site.

Methods

Genome recovery

The OHK43 genome was recovered from genome-resolved metagenomic sequencing of samples from Okuoku-hachikurou Onsen (OHK) in Akita Prefecture, Japan, following methods described previously (Ward *et al.*, 2019, 2020) and described briefly here. Samples of thin biofilms in the outflow of the hot spring were sampled for metagenomic sequencing in September 2016. DNA was preserved in the field with a Zymo Terralyzer BashingBead Matrix and Xpedition Lysis Buffer (Zymo Research, Irvine, CA) after disruption of cells in polyethylene sample tubes via attachment to and operation of a cordless reciprocating saw (Makita JR101DZ). Microbial DNA was extracted and purified after return to the lab with a Zymo Soil/Fecal DNA extraction kit (Zymo Research, Irvine, CA Quantification of DNA was performed with a Qubit 3.0 fluorimeter (Life Technologies, Carlsbad, CA). DNA was submitted to SeqMatic LLC (Fremont, CA) for library preparation using an Illumina Nextera XT DNA Library Preparation Kit prior to 2x100bp paired-end sequencing via Illumina HiSeq 4000 technology. Raw sequence reads were quality controlled with BBTools (Bushnell 2014) and assembled with MegaHit v. 1.02 (Li *et al.*, 2016). The OHK43 genome bin was recovered via differential coverage binning with MetaBAT (Kang *et al.*, 2015). Completeness and contamination/redundancy were determined with CheckM v1.1.2 (Parks *et al.*, 2015). The genome was uploaded to RAST v2.0 for annotation and characterization (Aziz *et al.*, 2008). Presence or absence of metabolic pathways of interest was

predicted using MetaPOAP v1.0 (Ward *et al.*, 2018). Taxonomic assignment was determined with

GTDB-Tk v1.2 (Parks *et al.*, 2018, 2020; Chaumeil *et al.*, 2019). Genome assembly and quality statistics

for OHK43 and other *Thermosynechoccus* genomes are reported in Supplemental Table 4.

Organismal Phylogenies

Concatenated ribosomal phylogenies were constructed following methods from Hug *et al.* (2016).

Members of the *Thermosynechococcaceae* and outgroups were identified using GTDB (Chaumeil *et al.*,

2019) and their genomes downloaded from the NCBI WGS and Genbank databases. Ribosomal protein

sequences were extracted using the tblastn function of BLAST+ (Camacho *et al.*, 2009) and aligned

with MUSCLE (Edgar, 2004). Trees were built with RAxML v.8.2.12 (Stamatakis, 2014) on the Cipres

science gateway (Miller *et al.*, 2010). Transfer bootstrap support values were determined with

BOOSTER (Lemoine *et al.*, 2018). Visualization of trees was performed with the Interactive Tree of Life

Viewer (Letunic and Bork, 2016).

Genome comparison

We compared the core- and pangenomes of *Thermosynechococcus* at the genus level, family level and

with a sub-sample of Cyanobacteria across the GTDB defined class Cyanobacteriia. ProteinOrtho

(Lechner *et al.*, 2011) was used for the identification of Conserved Likely Orthologous Groups (CLOG)

and analysis.

At the genus level 7 *Thermosynechococcus* strains from varying hot spring environments (Table 1 and

2) were compared. The genomic data from five available sequences of *Thermosynechococcus* strains

T. sp. CL1/1-2178 (CL1), *T. elongatus* BP1/1-2178 (*elongatus* BP1)*, T. vulcanus* NIES2134/1-2178

(*vulcanus*)*, T.* sp. NK55a/1-2022 (NK55a), *T. elongatus* PKUAC-SCTE542 (*elongatus* PKUAC) and two

unnamed strains from Jinata Onsen (Jinata) and Oku-Okuhachikurou Onsen (OHK) in Japan were

compared using ProteinOrtho (Lechner *et al.*, 2011), BLAST (Altschul *et al.*, 1990), and FeGenie (Garber

et al., 2019). Phylogenetic relationships between these strains were established using concatenated

ribosomal protein phylogenies following Hug *et al.* (2016), taxonomic classifications with GTDB-tk

(Chaumeil *et al.*, 2019) and average nucleotide identities (Rodriguez-R and Konstantinidis, 2017).

For the analysis at the family level, we included 6 more species that appear as representative strains at

the *Thermosynochoccaceae* family level in the Genome Taxonomy Data Base (GTDB). The

representative GTDB strains include *Acaryochloris marina* MBIC11017, *Cyanothece* sp. PCC 7425,

Acaryochloris sp. CCMEE 5410, *Synechococcus* sp. PCC 6312, *Synechococcus lividus* PCC 6715 and

Acaryochloris sp. RCC1774. This resulted in a total of 13 strains for family level analysis.

At the class level we used the 7 genus level strains and additionally 16 species, which do not cover all

of the family level species, 15 after (Beck *et al.*, 2012) and one more *Gloeobacter* species (*G. kilaueensis*

JS-1). Here the goal was to test the coherence of our analysis parameters when compared to previous

studies of cyanobacterial core- and pangenomes (Beck *et al.*, 2012, 2018). We acknowledge that the

distribution of species within the families of the class is not even, however this comparison was used to

verify our methods and the stability of the class level core. This resulted in a total of 23 genomes for the

class level analysis.

It is important to note that although the spring source water properties differ, the source water is not

necessarily where the DNA originated. Thus there is some uncertainty in the precise

geochemistry:genome relationships discussed here. We also acknowledge that not all genomes in this

study reach 100 % completeness. We might therefore be missing some gene clusters and some

genomes might cluster differently if they were complete. We compared genomic data as of August 4[th],

2020.

To compare the appearance of genes related to iron uptake and regulation we used FeGenie (Garber

et al., 2019) with standard parameters. We used results from ProteinOrtho (Lechner *et al.*, 2011) for

further analysis of the core, shared, unique, TS core and TS shared clusters. Proteinortho was applied

such that the output included also singleton clusters (only containing a single protein) and with an

algebraic connectivity of 0 as a measure for the structure of the orthologous clusters. We did not obtain

many differences when running ProteinOrtho on our data using a value of 0 or 0.1 (default) for the

algebraic connectivity, however, a value of 0 resulted in slightly larger clusters for the core and a few

less singletons. A comparison of results from FeGenie and ProteinOrtho resulted in similar gene clusters

for iron related genes such that our results here are based on both program outputs.

159 Results and Discussion

Genome Size	T. nakabusensis Jinata	T. nakabusensis OHK	T. elongatus BP-1	T. elongatus PKUAC-SCTE542	T. sp CL1	T. nakabusensis NK55a	T. vulcanus NIES2134
T. nakabusensis Jinata	2.31 MB	99.70	92.65	86.82	87.53	99.68	92.59
T. nakabusensis OHK		2.25 MB	92.58	86.85	87.52	99.84	92.55
T. elongatus BP-1			2.59 MB	86.65	87.34	92.53	99.14
T. elongatus PKUAC-SCTE542				2.64 MB	89.95	86.81	86.637
T. sp CL1					2.64 MB	87.50	87.32
T. nakabusensis NK55a						2.51 MB	92.45
T. vulcanus NIES2134							2.57 MB

Genome Sizes for GTDB representative family level species:
Acaryochloris marina MBIC11017	8.36 MB
Acaryochloris sp. CCMEE 5410	7.87 MB
Cyanothece sp. PCC 7425	5.78 MB
Synechococcus lividus PCC 6715	2.65 MB
Synechococcus sp. PCC 6312	3.72 MB
Acarychloris RCC 1774	5.93 MB

160
Table 2: Genus level average nucleotide identities (ANI) and genome sizes (diagonal). Additionally, genomes sizes for family level species are added in the lower left.

Phylogeny of the Thermosynechococcus and the species proposal T. nakabusensis

The _Thermosynechococcus_ genus is phylogenetically coherent within the Cyanobacteria (Figure 1) and

the genome sizes of genus members are similar to one another (Table 2). Based on similarity observed

with ANI and GTDB-tk (Table 2, supplementary Table 4), 4 species are present within the

Thermosynechococcus genus: _T. elongatus_ BP1 and _T. vulcanus_ belonging to one species, _T._ NK55a,

the Jinata and OHK genomes belonging to a second species, and _T._ CL1 and _T. elongatus_ PKUAC-

SCTE542 as one species each. For the species including the _T._ NK55a, Jinata, and OHK genomes we

propose the name _Thermosynechococcus nakabusensis_ as the first and so far only isolated organism

which originates from Nakabusa hot spring in Nagano Prefecture, Japan.

Genus and family level comparison of _Thermosynechococcus_ and

Thermosynochococcaceae

Comparing the conserved likely orthologous groups (CLOGS), we analyzed i) the core-genomes: those

CLOGS shared by all genomes in an analysis, ii) the shared CLOGs: those shared by at least 2 but not

all of the genomes in the analysis, and iii) unique CLOGs: those CLOGs that are unique to a single

genome (Table 3). The *Thermosynechococcus* genus specific core (**core TS)** comprises CLOGs shared by all 7 genus level genomes that are not present in any other species, and the *Thermosynechococcus* genus-specific shared CLOGs (**shared TS)** corresponds to CLOGs that are shared by at least 2 and at most 6 genus level genomes.

	Genus (7 genomes)	Family (13 genomes)	Class (23 genomes)
Core of all genomes analyzed	1878	1283	742
Core of 5 Thermosynechococcus (not including Jinata/OHK)	27	3	3
Jinata and OHK only	28	25	24
Protein Clusters Unique to Thermosynechococcus genus	-	18	79*

Table 3 - Numbers of CLOGs per grouping and phylogenetic level. *note that the number of genus level core CLOGs increases due to the exclusion of some family level genomes at the class level.

Comparing the genomes of the seven genus members, the protein core is made up of 1878 CLOGs and contains 80 to 89 % of the putative protein coding genes in a genome. This is higher than in other comparisons of organisms, for example in Reno *et al.* (2009) 69 – 79 % of genes from *S. islandicus* strains made up the core. Wu *et al.* (2018) found that the core genes of species belonging to the genus *Comamonas* account for 18 – 33 % of all genes, and Barajas *et al.* (2019) investigated the core genome within the genus *Streptococcus* which ranges in size from 9.6 % to 24 %. The high proportion of CLOGs which make up the core leaves few unique CLOGs for each genome and between genus members, and only slight variations in genome content between genome are observed: for example 27 CLOGs are specific to the 5 genomes that do not include Jinata and OHK and 28 CLOGs are specific to Jinata and OHK (Figure 2, supplementary Figure 1).

Golicz *et al.* (2020) suggested that the size of a pangenome is related to organismal lifestyles, with sympatric organisms having open pangenomes with many accessory genes and allopatric (isolated from other organisms) organisms having more closed and conserved pangenomes. The *Thermosynechococcus* genus could thus be considered as allopatric as they have a comparatively large core and few shared and unique genes. Chen *et al.* (2020) also noted that differences in horizontal gene transfer (HGT) are related to genome size with smaller genomes showing less HGT and bigger genomes having a larger probability that HGT occurred. They also found that hot spring cyanobacteria specifically have smaller genome sizes and less HGT into the genome. Excluding gene loss, we suggest that the conserved core might be indicative of a more ancient gene repertoire in hot spring cyanobacteria, with

other cyanobacteria gaining more functionality through HGT over evolutionary timescales. Furthermore

thermotolerance is phylogenetically scattered across the cyanobacterial tree of life, mostly in organisms

comprising smaller genomes. Since all *Thermosynechococcus* in this study are found in hot springs –

which typically have reduced microbial diversity in comparison to other environments such as soils

(Ward *et al.,* 1998) – this finding provides support for the still tentative hypothesis that hot spring

environments may provide more limited opportunity for lateral gene transfer, which in turn could lead to

less opportunity for lateral gene transfer and smaller genomes.

In contrast to the genus level where genome size varies from 2.25 MB to 2.64 MB, at the family level it

varies to 8.36 MB (*Acaryocholoris marina* MBIC11017, Table 2). Running proteinortho analyses with the

seven genus level *Thermosynechococcus* and the six family level sequences, the overall core is reduced

Figure 2: (A) Thermosynechococcaceae family level comparison of core- and pangenomes, (B) number of CLOGs observed at the family level in relationship to the number of genomes in the analysis. coreTS indicates CLOGs found in all genus level genomes, sharedTS are CLOGs found in at least 2 but not all genus level genomes and no other genomes. Full species names as mentioned above.

by just over 30%, from 1878 to 1283 CLOGs. This number of ~1900 core CLOGs within the genus is

different from a recent analysis (Cheng *et al.*, 2020), with the higher number observed here being due

to the inclusion of the newly available and revised *T. elongatus* PKUAC-SCTE542 genome. Shared

genes, not core genes, comprise a larger percentage of the genomes for smaller genomes, while unique

genes are abundant in larger genomes (Figure 2a).

Including the added diversity at the family level, the number of CLOGs shared between the 5

Thermosynechococcus genus members that do not include Jinata and OHK is reduced from 27 to 3,

showing that these CLOGs are found in other closely related members at the family level. However, the

number of CLOGs found only in Jinata and OHK changes from 28 to 25 CLOGs, highlighting that these

are unique. Also observed when comparing the *Thermosynechococcus* genus across the

Thermosynochococcaceae family, the specific CLOGs that appear in only, and all of the 7 genus level

genomes is made up of 18 CLOGs, showing a low number of conserved *Thermosynechococcus* genus

only CLOGs. From the CLOG perspective, the seven members share much in common with their closest

relatives, but they appear to lack major distinguishing qualities.

Genomic differentiation of the *Thermosynechococcus* genus from other cyanobacteria

At the class level the number of *Thermosynechococcus* genus level core CLOGs increases when compared

to the family level due to the exclusion of some family level species at this level (see Methods and Table 3).

The overall core decreases by almost 60% when class level representatives are added (1878 core

CLOGs at the genus level and 742 core CLOGs at the class level). This is expected as Beck *et al.* (2018)

suggested that the clustering of CLOGs depends on the variability in genome size as well as

phylogenetic distance between the analyzed genomes as well as the total amount of genomes analyzed .

Moreover this is also in line with the larger analyses of microbial genomes which have shown that the

continued addition of taxonomic diversity in an analysis in an analysis leads to increasingly smaller cores

(Charlebois and Doolittle, 2004; Lapierre and Gogarten, 2009).

Overall these results conform with Beck *et al.* (2012), with variations attributed to the different methods

and parameters used for each analysis. Additionally, we confirm that the class level core is stable when

adding a higher diversity of organisms, similar to Beck *et al.* (2018), as the number of core family CLOGs

only slightly decreases when adding more species at the class level.

At the class level, we looked at specific cases among the shared, but not unique CLOGs, to investigate

gene content which may differentiate the *Thermosynechococcus* genus from other cyanobacteria. In the

case of CLOGs which appear in all *Thermosynechococcus* but not in any other cyanobacteria in our

analysis, 79 CLOGs were found, 47 of which are hypothetical proteins of unknown functions and could

be of interest to further biochemical studies to understand the adaptations of the genus. CLOGs unique

to the genus appear to be of known functionalities account for a variety of processes with genes involved

in transcriptional regulation, transporters and membrane proteins (e.g. acetyltransferases, glycosidases

and ATPases; more information in supplementary Table 2). Since all the *Thermosynechococcus*

analyzed are from hot springs, these genes potentially provide some basis for that lifestyle.

Adaptation of Thermosynechococcus to their respective environments

We were especially interested in those CLOGs that are shared between the strains from environments

with elevated iron concentrations (Jinata and OHK) but which are not present in any other cyanobacteria.

Previous studies have shown that some cyanobacteria express higher levels of genes involved in iron

ion homeostasis which are expressed in iron limiting conditions (Cheng and He, 2014), and here we

looked at the presence or absence of iron related gene products in *Thermosynechococcus* compared

to other cyanobacteria using FeGenie and BLAST comparisons. 24 CLOGs are specific to Jinata and

OHK at the class level, some of which show high partial identity but low coverage matches with genes

from other *Thermosynechococcus* (supplementary Table 4). It is notable that these 24 CLOGs do not

appear in the third strain of the same species (NK55a). With our current understandings after

considering results from BLAST and FeGenie, none of the 24 CLOGs comprises genes that could

explain the organisms adaptation to elevated iron concentrations. 3 unique CLOGs are found in the 5

Thermosynechococcus that do not include Jinata and OHK, and we confirmed that none of those

CLOGs are related to iron regulation as best as can be assessed from the sequence, similar to the

Jinata and OHK CLOGs (supplementary Table 4). There is no sequence resolvable genomic signature

specific to Jinata and OHK related to iron tolerance.

These genomes lack genes coding ferrous iron transport and uptake proteins EfeB, EfeO and EfeU, the

metal transport gene ZupT, cellular iron storage proteins Bfr and the iron regulator under iron limiting

conditions PfsR (Table 4). Genes encoding these proteins are found in other species of cyanobacteria,

but not in the *Thermosynechococcus* genus members with currently available genomes. Within the

genus there are no differences with regard to genes encoding proteins involved in iron regulation (PfsR),

and in all cases the same genes encoding proteins related to ferrous iron uptake (FeoA, FeoB, YfeA

and YfeB), ferric iron uptake or transport (ExbD, FutA, FutB and FutC), siderophore iron acquisition

(FpvD), metal ion binding (Ho1 and Ho2) and iron starvation acclimation (IsiA) are present (Table 4).

Protein	Function	BP-1	PKUAC	CL1	NK55a	vulcanus	Jinata	OHK	Present in Themosyn	Reference
FeoA	ferrous iron uptake	X	X	X	X	X	X	X	All	Lau *et al.*, 2016
FeoB	ferrous iron uptake	X	X	X	X	X	X	X	All	Lau *et al.*, 2016
YfeA	ferrous iron uptake	X	X	X	X	X	X	X	All	Toulza *et al.*, 2012
YfeB	ferrous iron uptake	X	X	X	X	X	X	X	All	Toulza *et al.*, 2012
FpvD	siderophore iron acquisition	X	X	X	X	X	X	X	All	Brillet *et al.*, 2012
ExbD	ferric iron uptake	X	X	X	X	X	X	X	All	Jiang *et al.*, 2015
PfsR	iron regulator under iron limiting conditions	Not found when Blasting Synechocystis sequence							Synechocystis and others	Cheng and He, 2014
FutA	ABC-type ferric iron transport	X	X	X	X	X	X	X	All	Katoh *et al.*, 2001
FutB	ABC-type ferric iron transport	X	X	X	X	X	X	X	All	Katoh et al., 2001
FutC	ABC-type ferric iron transport	X	X	X	X	X	X	X	All	Katoh *et al.*, 2001
EfeB	ferrous iron transport	Not found when Blasting Synechococcus sequence							Synechococcus and others	Lau *et al.*, 2016
EfeO	ferrous iron transport	Not found when Blasting Synechococcus sequence							Synechococcus and others	Lau *et al.*, 2016
EfeU	ferrous iron uptake	Not found when Blasting Microcystis sequence							Microcystis aeruginosa	Lau *et al.*, 2016
ZupT	metal transport (including ferrous iron)	Not found when Blasting with Synechococcus sequence as query							Synechococcus, J055 and J083 and others	Lau *et al.*, 2016
Bfr	cellular iron storage	Not found when Blasting with Synechococcus sequence as query							Synechococcus and others	Keren *et al.*, 2004; Cheng and He, 2014
IsiA	iron starvation acclimation	X	X	X	X	X	X	X	Some	Cheng and He, 2014
Ho1	metal ion binding	X	X	X	X	X	X	X	Some	Cheng and He, 2014
Ho2	metal ion binding	X	X	X	X	X	X	X	Some	Cheng and He, 2014

275 Table 4: Genes known to be involved in iron regulation within the class Cyanobacteriia and their presence in the

276 *Thermosynechococcus* genus. Presence or absence of genes was confirmed with BLAST searches.

The conserved genomic core of *Thermosynechococcus* in relationship to environmental distribution is unique

In addition to resolving a genetic basis for widespread environmental distribution, our work is also relevant to historical proliferation of cyanobacteria, since some modern-day hot springs and their biogeochemistries can be used as historical process analogues (Brown *et al.*, 2005, 2007; Ward *et al.*, 2019). Considering contemporary environments, the analysis of *Thermosynechococcus* also provide insight into island biogeography of microbes. Ionescu *et al.* (2010) observed that the speciation patterns of microorganisms are shaped by local community structures and environmental influences, and Bahl *et al.* (2011) additionally suggest a positive correlation between geographic and genetic distance. Papke *et al.* (2003) found that isolated environments such as geothermal springs may lead to evolutionary divergence of closely related *Thermosynechococcus* strains due to an island effect. Our analysis suggests that geographically widespread organisms belonging to the genus inhabit hot springs with varying geochemistries without genomically recognizable adaptations specific to their site of origin. Instead, the finding of highly conserved genomes within the genus, and furthermore, that the genetic content of the genus is not markedly different from other cyanobacteria, implies that the genus is inherently flexible and viable in the geochemical regimes studied. The large portion of shared genes within the genus provides a genetic basis for the lack of correlation between geographic and genetic distances within the genus found by Papke *et al.* (2003).

This genomic coherence at the genus level is in contrast to other studies, such as Reno *et al.* (2009; Wu *et al.* (2018) or Barajas *et al.* (2019)as mentioned above. Furthermore the organisms tested by Reno *et al.* (2009) are obligate thermoacidophilic archaea at the species level which are environmentally restricted similar to organisms here. They found that the core and pan genomes of these organisms are shaped by their geographical distribution and relatedness within and across different environments. Apparently, *Thermosynechococcus* is environmentally promiscuous, and have fewer restrictive requirements concerning their distribution.

Outlook

One aim of this study was to identify those proteins that make hot-spring inhabiting *Thermosynechococcus* genus members unique. Although the genus *Thermosynechococcus* is rather genomically conserved, 47 out of the 79 CLOGs unique to these genomes are identified as hypothetical proteins of unknown function and thus biochemical studies on these proteins is warranted.

Based on the genome comparisons presented here, a viability test of isolates in environments other

than those of their origin is suggested as future work. For example, since *Thermosynechococcus* is

lacking genes that are involved in iron regulation under limiting conditions, genus level iron tolerance

experiments are proposed to test if strains from low-iron environments can also withstand elevated iron

without any adaptations in their genome. In a similar way thermotolerance of these organisms could

also be investigated. This could help us understand if *Thermosynechococcus* is indeed less restrictive

or to identify mechanisms unresolvable by the CLOG approach that account for the geographical

distribution.

Traits which are unresolvable with an orthology comparison approach like that employed here include

amino acid substitutions or sequence changes that lead to variations in enzyme activity, structure, and

regulation. Furthermore our analysis is not sensitive to differences in gene regulation or gene copy

number. Access to solvents, solubility, active side peptides and differential folding can influence a

protein's functionality and therefor may have implications for adaptability or infer evolutionary history

(Lesk and Chothia, 1986; Pandey and Braun, 2020). If the orthology approach employed here is correct,

the geochemical parameters that are often thought to be important in microbial selection might simply

be not as important as we assume them to be in the environment. Here, orthology between genes was

assessed based on sequence comparison (proteinortho, BLAST) and sequence and structure

comparison (FeGenie, HMMer). However, there exists a diverse set of methods and approaches to

determine orthology between genes (Lechner *et al.*, 2011; Forslund *et al.*, 2018), and as orthology

definition techniques change, our results could be interpreted.

From an Earth history perspective, the presence of cyanobacteria in high-iron environments today could

indicate that ancient cyanobacteria may not have been limited by elevated iron concentrations on early

Earth, whereas previous studies have reported on the toxicity of high iron concentrations to

cyanobacteria (Swanner, Mloszewska, *et al.*, 2015; Swanner, Wu, *et al.*, 2015). We could not find

striking adaptive mechanisms on a genome level that can explain the tolerance of strains found in

environments with elevated iron concentrations, however the presence and perseverance of

cyanobacteria in these environments have important implications when considering the earliest

organisms capable of oxygenic photosynthesis on early Earth. Ionescu et al. proposed that simply by

increasing photosynthetic rate and oxygen production, cyanobacteria might protect themselves from

ferrous iron by promoting its precipitation at some distance from the cell (Ionescu *et al.*, 2014). In line

with this observation, is worthwhile to note that the biomass accumulation in high iron environments like

Jinata hot spring is appreciable, with co-occurring visible biomass and super saturated dissolved oxygen

concentrations from cyanobacterial activity (Ward *et al.*, 2019).

Data availability for newly described strains

The Whole Genome Shotgun project for OHK43 has been deposited at DDBJ/ENA/GenBank under the

accession JACOMP000000000. The version described in this paper is version JACOMP010000000.

Supplemental Material

SI Figure 1 – class and genus level comparison of core- and pan-genomes; distribution of number of

CLOGs with number of genomes at class and genus level. Core TS corresponds to the CLOGs found

in all *Thermosynechococcus* genus members, and shared TS corresponds to CLOGs that are shared

by at least 2 and at most 6 genus level genomes.

SI Table 1 – *Thermosynechococcus* core CLOGs at class level and their hits in the BLAST protein

database.

SI Table 2 – Jinata and OHK specific CLOGs at class level with coverage and identity scores, 5 TS

specific CLOGs at class level

SI Table 3 – genome information for all family level genomes.

Acknowledgements

LMW was supported by a Simons Foundation Postdoctoral Fellowship in Marine Microbial Ecology.

Metagenomic sequencing of OHK was supported by NSF grant #OISE 1639454. SEM acknowledges

support from the Astrobiology Center Program of the National Institutes of Natural Sciences (grant no.

AB311013). We would also like to thank the group around M. Daroch for providing us with an updated

version for the *T. elongatus* PKUAC-SCTE542 genome which increased the strain's coherence

compared to previous versions of the genome.

References

Allen, J.F., Thake, B., and Martin, W.F. (2019) Nitrogenase Inhibition Limited Oxygenation of Earth's Proterozoic Atmosphere. *Trends in Plant Science* **24**: 1022–1031.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403–410.

Aziz, R.K., Bartels, D., Best, A., DeJongh, M., Disz, T., Edwards, R.A., et al. (2008) The RAST Server: Rapid annotations using subsystems technology. *BMC Genomics* **9**: 75.

Bahl, J., Lau, M.C.Y., Smith, G.J.D., Vijaykrishna, D., Cary, S.C., Lacap, D.C., et al. (2011) Ancient origins determine global biogeography of hot and cold desert cyanobacteria. *Nature Communications* **2**: 1–6.

Barajas, H.R., Romero, M.F., Martínez-Sánchez, S., and Alcaraz, L.D. (2019) Global genomic similarity and core genome sequence diversity of the Streptococcus genus as a toolkit to identify closely related bacterial species in complex environments. *PeerJ* **2019**: e6233.

Beck, C., Knoop, H., Axmann, I.M., and Steuer, R. (2012) The diversity of cyanobacterial metabolism: genome analysis of multiple phototrophic microorganisms. *BMC genomics* **13**: 56.

Beck, C., Knoop, H., and Steuer, R. (2018) Modules of co-occurrence in the cyanobacterial pan-genome reveal functional associations between groups of ortholog genes. *PLOS Genetics* **14**: e1007239.

Brillet, K., Ruffenach, F., Adams, H., Journet, L., Gasser, V., Hoegy, F., et al. (2012) An ABC Transporter with Two Periplasmic Binding Proteins Involved in Iron Acquisition in Pseudomonas aeruginosa. *ACS Chemical Biology* **7**: 2036–2045.

Callieri, C., Slabakova, V., Dzhembekova, N., Slabakova, N., Peneva, E., Cabello-Yeves, P.J., et al. (2019) The mesopelagic anoxic Black Sea as an unexpected habitat for Synechococcus challenges our understanding of global "deep red fluorescence." *ISME Journal* **13**: 1676–1687.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L. (2009) BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 1–9.

Canfield, D.E. (1998) A new model for Proterozoic ocean chemistry. *Nature* **396**: 450.

Charlebois, R.L. and Doolittle, W.F. (2004) Computing prokaryotic gene ubiquity: Rescuing the core from extinction. *Genome Research* **14**: 2469–2477.

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2019) GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**: 1925–1927.

Chen, M.Y., Teng, W.K., Zhao, L., Hu, C.X., Zhou, Y.K., Han, B.P., et al. (2020) Comparative genomics reveals insights into cyanobacterial evolution and habitat adaptation. *ISME J* 1–17.

Cheng, D. and He, Q. (2014) PfsR Is a Key Regulator of Iron Homeostasis in Synechocystis PCC 6803. *PLoS ONE* **9**: e101743.

Cheng, Y.-I., Chou, L., Chiu, Y.-F., Hsueh, H.-T., Kuo, C.-H., and Chu, H.-A. (2020) Comparative Genomic Analysis of a Novel Strain of Taiwan Hot-Spring Cyanobacterium Thermosynechococcus sp. CL-1. *Frontiers in Microbiology* **11**: 82.

Edgar, R.C. (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**:113

Fennel, K., Follows, M., and Falkowski, P.G. (2005) The co-evolution of the nitrogen, carbon and oxygen cycles in the proterozoic ocean. *American Journal of Science* **305**: 526–545.

Fischer, W.W., Hemp, J., and Valentine, J.S. (2016) How did life survive Earth's great oxygenation? *Current Opinion in Chemical Biology* **31**: 166–178.

Forslund, K., Pereira, C., Capella-Gutierrez, S., da Silva, A.S., Altenhoff, A., Huerta-Cepas, J., et al. (2018) Gearing up to handle the mosaic nature of life in the quest for orthologs. *Bioinformatics* **34**: 323–329.

Garber, A.I., Nealson, K.H., Okamoto, A., McAllister, S.M., Chan, C.S., Barco, R.A., and Merino, N. (2019) FeGenie: a comprehensive tool for the identification of iron genes and iron gene neighborhoods in genomes and metagenome assemblies. *bioRxiv* 777656.

Golicz, A.A., Bayer, P.E., Bhalla, P.L., Batley, J., and Edwards, D. (2020) Pangenomics Comes of Age: From Bacteria to Plant and Animal Applications. *Trends in Genetics* **36**: 132–145.

Guo, Q., Pang, Z., Wang, Y., and Tian, J. (2017) Fluid geochemistry and geothermometry applications of the Kangding high-temperature geothermal system in eastern Himalayas. *Applied Geochemistry* **81**: 63–75.

Hao, J., Knoll, A.H., Hazen, R.M., Daniel, I., Huang, F., and Schieber, J. (2020) Cycling Phosphorus on the Archean Earth: Part II. Phosphorus Limitation on Primary Production in Archean Ecosystems Pattern-forming abiotic reactions during organic diagenesis View project Deep Time Data Infrastructure View project Cycling phosphorus on the Archean Earth: Part II. Phosphorus limitation on primary production in Archean ecosystems ScienceDirect.

Hsueh, H.T., Chu, H., and Chang, C.C. (2007) Identification and Characteristics of a Cyanobacterium Isolated from a Hot Spring with Dissolved Inorganic Carbon. *Environmental Science & Technology* **41**: 1909–1914.

Hug, L.A., Baker, B.J., Anantharaman, K., Brown, C.T., Probst, A.J., Castelle, C.J., et al. (2016) A new view of the tree of life. *Nature Microbiology* **1**: 1–6.

Ichikuni, M. and Kikuchi, K. (1972) Retention of boron by travertines. *Chemical Geology* **9**: 13–21.

Ionescu, D., Buchmann, B., Heim, C., Häusler, S., de Beer, D., and Polerecky, L. (2014) Oxygenic photosynthesis as a protection mechanism for cyanobacteria against iron-encrustation in environments with high Fe2+ concentrations. *Frontiers in Microbiology* **5**: 459.

Jiang, H.-B., Lou, W.-J., Ke, W.-T., Song, W.-Y., Price, N.M., and Qiu, B.-S. (2015) New insights into iron acquisition by cyanobacteria: an essential role for ExbB-ExbD complex in inorganic iron uptake. *The ISME journal* **9**: 297.

Kang, D.D., Froula, J., Egan, R., and Wang, Z. (2015) MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* **2015**: e1165.

Katoh, H., Hagino, N., and Ogawa, T. (2001) Iron-Binding Activity of FutA1 Subunit of an ABC-type Iron Transporter in the Cyanobacterium Synechocystis sp. Strain PCC 6803, Oxford Academic.

Keren, N., Aurora, R., and Pakrasi, H.B. (2004) Critical roles of bacterioferritins in iron storage and proliferation of cyanobacteria. *Plant Physiology* **135**: 1666–1673.

Knoll, A.H. (2006) Cyanobacteria and Earth History Diverse aspects of testate amoebae evolution View project The Co-Evolution of the Geo-and Biospheres: An Integrated Program for Data-Driven Abductive Discovery in Earth Sciences View project.

Lapierre, P. and Gogarten, J.P. (2009) Estimating the size of the bacterial pan-genome. *Trends in Genetics* **25**: 107–110.

Lau, C.K.Y., Krewulak, K.D., and Vogel, H.J. (2016) Bacterial ferrous iron transport: the Feo system. *FEMS microbiology reviews* **40**: 273–298.

Lechner, M., Findeiß, S., Steiner, L., Marz, M., Stadler, P.F., and Prohaska, S.J. (2011) Proteinortho: detection of (co-) orthologs in large-scale analysis. *BMC bioinformatics* **12**: 124.

Lemoine, F., Entfellner, J.-B.D., Wilkinson, E., Correia, D., Felipe, M.D., De Oliveira, T., and Gascuel, O. (2018) Renewing Felsenstein's phylogenetic bootstrap in the era of big data. *Nature* **556**: 452–456.

Lesk, A.M. and Chothia, C.H. (1986) The response of protein structures to amino-acid sequence changes. *Philosophical Transactions of the Royal Society of London Series A, Mathematical and Physical Sciences* **317**: 345–356.

Letunic, I. and Bork, P. (2016) Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**: W242–W245.

Li, D., Luo, R., Liu, C.M., Leung, C.M., Ting, H.F., Sadakane, K., et al. (2016) MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies and community practices. *Methods* **102**: 3–11.

Lyons, T.W., Reinhard, C.T., and Planavsky, N.J. (2014) The rise of oxygen in Earth's early ocean and atmosphere. *Nature* **506**: 307.

Maity, J.P., Liu, C.-C., Nath, B., Bundschuh, J., Kar, S., Jean, J.-S., et al. (2011) Biogeochemical characteristics of Kuan-Tzu-Ling, Chung-Lun and Bao-Lai hot springs in southern Taiwan. *Journal of Environmental Science and Health, Part A* **46**: 1207–1217.

Mei-hua, W., Junhao, W., and Tingshan, T. (2015) Study of the Scaling Trend of Thermal Groundwater in Kangding County of Sichuan Province. *World Geothermal Congress 2015* 11.

Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *2010 gateway computing environments workshop (GCE)*. Ieee, pp. 1–8.

Nakagawa, T. and Fukui, M. (2002) Phylogenetic characterization of microbial mats and streamers from a Japanese alkaline hot spring with a thermal gradient. *The Journal of General and Applied Microbiology* **48**: 211–222.

Nakamura, Y., Kaneko, T., Sato, S., Ikeuchi, M., Katoh, H., Sasamoto, S., et al. (2002) Complete genome structure of the thermophilic cyanobacterium Thermosynechococcus elongatus BP-1. *DNA research* **9**: 123–130.

Pandey, A. and Braun, E.L. (2020) Phylogenetic Analyses of Sites in Different Protein Structural Environments Result in Distinct Placements of the Metazoan Root. *Biology* **9**: 64.

Papke, R.T., Ramsing, N.B., Bateson, M.M., and Ward, D.M. (2003) Geographical isolation in hot spring cyanobacteria. *Environmental Microbiology* **5**: 650–659.

Parks, D.H., Chuvochina, M., Chaumeil, P.A., Rinke, C., Mussig, A.J., and Hugenholtz, P. (2020) A complete domain-to-species taxonomy for Bacteria and Archaea. *Nature Biotechnology* 1–8.

Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarshewski, A., Chaumeil, P.A., and Hugenholtz, P. (2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nature Biotechnology* **36**: 996.

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., and Tyson, G.W. (2015) CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research* **25**: 1043–1055.

Puente-Sánchez, F., Arce-Rodríguez, A., Oggerin, M., García-Villadangos, M., Moreno-Paz, M., Blanco, Y., et al. (2018) Viable cyanobacteria in the deep continental subsurface. *Proceedings of the National Academy of Sciences of the United States of America* **115**: 10702–10707.

Raven, J. (2009) Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquatic Microbial Ecology* **56**: 177–192.

Reno, M.L., Held, N.L., Fields, C.J., Burke, P. v., and Whitaker, R.J. (2009) Biogeography of the Sulfolobus islandicus pan-genome. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 8605–8610.

Rodriguez-R, L.M. and Konstantinidis, K.T. (2017) Bypassing Cultivation To Identify Bacterial Species Culture-independent genomic approaches identify credibly distinct clusters, avoid cultivation bias, and provide true insights into microbial species.

Sánchez-Baracaldo, P. (2015) Origin of marine planktonic cyanobacteria. *Scientific Reports* **5**:.

Schirrmeister, B.E., Gugger, M., and Donoghue, P.C.J. (2015) Cyanobacteria and the Great Oxidation Event: evidence from genes and fossils. *Palaeontology* **58**: 769–785.

Shi, T. and Falkowski, P.G. (2008) Genome evolution in cyanobacteria: The stable core and the variable shell. *Proceedings of the National Academy of Sciences of the United States of America* **105**: 2510–2515.

Shih, P.M., Hemp, J., Ward, L.M., Matzke, N.J., and Fischer, W.W. (2017) Crown group Oxyphotobacteria postdate the rise of oxygen. *Geobiology* **15**: 19–29.

Shih, P.M., Wu, D., Latifi, A., Axen, S.D., Fewer, D.P., Talla, E., et al. (2013) Improving the coverage of the cyanobacterial phylum using diversity-driven genome sequencing. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 1053–1058.

Soo, R.M., Hemp, J., and Hugenholtz, P. (2019) Evolution of photosynthesis and aerobic respiration in the cyanobacteria. *Free Radical Biology and Medicine* **140**: 200–205.

Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**: 1312–1313.

Stolyar, S., Liu, Z., Thiel, V., Tomsho, L.P., Pinel, N., Nelson, W.C., et al. (2014) Genome sequence of the thermophilic cyanobacterium Thermosynechococcus sp. strain NK55a. *Genome Announc* **2**: e01060-13.

Swanner, E.D., Mloszewska, A.M., Cirpka, O.A., Schoenberg, R., Konhauser, K.O., and Kappler, A. (2015) Modulation of oxygen production in Archaean oceans by episodes of Fe(II) toxicity. *Nature Geoscience* **8**: 126–130.

Swanner, E.D., Wu, W., Hao, L., Wüstner, M.L., Obst, M., Moran, D.M., et al. (2015) Physiology, Fe(II) oxidation, and Fe mineral formation by a marine planktonic cyanobacterium grown under ferruginous conditions. *Frontiers in Earth Science* **3**: 60.

Tang, J., Jiang, D., Luo, Y., Liang, Y., Li, L., Shah, Md.M.R., and Daroch, M. (2018) Potential new genera of cyanobacterial strains isolated from thermal springs of western Sichuan, China. *Algal Research* **31**: 14–20.

Tomitani, A., Knoll, A.H., Cavanaugh, C.M., and Ohno, T. (2006) The evolutionary diversification of cyanobacteria: Molecular-phylogenetic and paleontological perspectives. *Proceedings of the National Academy of Sciences of the United States of America* **103**: 5442–5447.

Toulza, E., Tagliabue, A., Blain, S., and Piganeau, G. (2012) Analysis of the Global Ocean Sampling (GOS) Project for Trends in Iron Uptake by Surface Ocean Microbes. *PLoS ONE* **7**: e30931.

Wang, L., Liu, Q., Wu, X., Huang, Y., Wise, M.J., Liu, Z., et al. (2019) Bioinformatics Analysis of Metabolism Pathways of Archaeal Energy Reserves. *Scientific Reports* 1–12.

Ward, D.M., Ferris, M.J., Nold, S.C., and Bateson, M.M. (1998) A Natural View of Microbial Biodiversity within Hot Spring Cyanobacterial Mat Communities. *Microbiol Mol Biol Rev* **62**: 1353–1370.

Ward, L.M., Fischer, W.W., and Mcglynn, S.E. Candidatus Anthektikosiphon siderophilum OHK22, a New Member of the Chloroflexi Family Herpetosiphonaceae from Oku-okuhachikurou Onsen. *Microbes Environ* **35**: 2020.

Ward, L.M., Idei, A., Nakagawa, M., Ueno, Y., Fischer, W.W., and McGlynn, S.E. (2019) Geochemical and Metagenomic Characterization of Jinata Onsen, a Proterozoic-Analog Hot Spring, Reveals Novel Microbial

Diversity including Iron-Tolerant Phototrophs and Thermophilic Lithotrophs. *Microbes and environments*
34: 278–292.
Ward, L.M., Idei, A., Terajima, S., Kakegawa, T., Fischer, W.W., and McGlynn, S.E. (2017) Microbial diversity and
iron oxidation at Okuoku-hachikurou Onsen, a Japanese hot spring analog of Precambrian iron
formations. *Geobiology* **15**: 817–835.
Ward, L.M. and Shih, P.M. (2019) The evolution and productivity of carbon fixation pathways in response to
changes in oxygen concentration over geological time. *Free Radical Biology and Medicine* **140**: 188–199.
Ward, L.M., Shih, P.M., and Fischer, W.W. (2018) MetaPOAP: presence or absence of metabolic pathways in
metagenome-assembled genomes. *Bioinformatics (Oxford, England)* **34**: 4284–4286.
Whitton, B.A. (1992) Diversity, ecology, and taxonomy of the cyanobacteria. In *Photosynthetic prokaryotes*.
Springer, pp. 1–51.
Wu, Y., Zaiden, N., and Cao, B. (2018) The Core- and Pan-Genomic Analyses of the Genus Comamonas: From
Environmental Adaptation to Potential Virulence. *Frontiers in Microbiology* **9**: 3096.
Yeh, G.-H., Yang, T.F., Chen, J.-C., Chen, Y.-G., and Song, S.-R. (2005) Fluid geochemistry of mud volcanoes in
Taiwan. *Mud Volcanoes, Geodynamics and Seismicity Springer, Netherlands* 227–237.

553