Single-cell RNA-sequencing reveals dysregulation of molecular programs associated with SARS-CoV-2 severity and outcomes in patients with chronic lung disease

Linh T. Bui[1]*, Nichelle I. Winters[2]*, Mei-I Chung[1], Chitra Joseph[3], Austin J. Gutierrez[1], Arun C. Habermann[2], Taylor S. Adams[4], Jonas C. Schupp[4], Sergio Poli[5], Lance M. Peter[1], Chase J. Taylor[2], Jessica B. Blackburn[2], Bradley W. Richmond[2,6], Andrew G. Nicholson[7,8], Doris Rassl[9], William A. Wallace[10,11], Ivan O. Rosas[12], R. Gisli Jenkins[3], Naftali Kaminski[4], Jonathan A. Kropski[2,6,13]**, Nicholas E. Banovich[1]** and the Human Cell Atlas Lung Biological Network

* denotes equal contribution
** denotes equal contribution

Affiliations
[1] Translational Genomics Research Institute, Phoenix, AZ, USA
[2] Division of Allergy, Pulmonary and Critical Care Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA
[3] Respiratory Medicine NIHR Biomedical Research Centre, University of Nottingham, Nottingham, UK
[4] Section of Pulmonary, Critical Care and Sleep Medicine, Yale School of Medicine, New Haven, CT, USA
[5] Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA
[6] Department of Veterans Affairs Medical Center, Nashville, TN, USA
[7] National Heart and Lung Institute, Imperial College, London, SW3 6LY, UK
[8] Royal Brompton and Harefield NHS Foundation Trust, Department of Histopathology, Sydney Street, London, SW3 6NP, UK
[9] Pathology Research, Royal Papworth Hospital NHS Foundation Trust, Cambridge, UK
[10] Department of Pathology, Royal Infirmary of Edinburgh, Edinburgh, UK
[11] Division of Pathology, Edinburgh University Medical School, Edinburgh, UK
[12] Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, Baylor College of Medicine, Houston, TX, USA
[13] Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN, USA

Note: A full list of authors and affiliations of HCA Lung Biological Network members is included at the end of the manuscript.

Address correspondence to:
Linh T. Bui (lbui@tgen.org) or Nicholas E. Banovich (nbanovich@tgen.org)

ABSTRACT

Rationale: Patients with chronic lung disease have an increased risk for severe coronavirus disease-19 (COVID-19) and poor outcomes.

Objectives: To identify molecular characteristics of diseased lung epithelial and immune cells that may account for worse COVID-19 outcomes in patients with chronic lung diseases.

Methods: We analyzed the transcriptomes of 605,904 single cells isolated from healthy (79 samples) and diseased human lungs (31 chronic obstructive pulmonary disease (COPD), 82 idiopathic pulmonary fibrosis (IPF) and 18 non-IPF interstitial lung disease samples).

Measurements and Main Results: Cellular distribution and relative expression of SARS-CoV-2 entry factors (*ACE2*, *TMPRSS2*) was similar in disease and control lungs. Epithelial cells isolated from diseased lungs expressed higher levels of genes linked directly to efficiency of viral replication and the innate immune response. Unique *ACE2*-correlated gene sets were identified for each diagnosis group in the type II alveolar cells. Diseased lungs have a significant increase in the proportion of CD4, CD8 and NK cells compared to control lungs. Components of the interferon pathway, the IL6 cytokine pathway and the major histocompatibility complex (MHC) class II genes are upregulated in several diseased immune cell types. These differences in inflammatory gene expression programs highlight how chronic lung disease alters the inflammatory microenvironment encountered upon viral exposure to the peripheral lung.

Conclusions: Chronic lung disease is accompanied by changes in cell-type-specific gene expression programs that prime the lung epithelium for and influence innate and adaptive immune responses to SARS-CoV-2 infection.

INTRODUCTION

In December 2019, a respiratory disease associated with a novel coronavirus emerged in Wuhan, China (1–3). The syndrome, now called COVID-19, was caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and has since rapidly spread worldwide (4). As of October 19, 2020, a total of over 40.3 million confirmed COVID-19 cases and more than one million deaths have been reported around the globe (https://coronavirus.jhu.edu/).

The clinical manifestations of SARS-CoV-2 infection range from asymptomatic to fulminant cases of acute respiratory distress syndrome (ARDS) and life-threatening multi-system organ failure. Development of ARDS in patients with SARS-CoV-2 dramatically increases the risk of ICU admission and death (5–11). Risk factors for severe SARS-CoV-2 include age, smoking status, ethnicity and male sex (12–14). Baseline comorbidities including hypertension, diabetes and obesity, increase SARS-CoV-2 susceptibility and severity (9, 15–19). In addition, chronic lung disease has been identified as a risk factor for hospitalization and mortality in patients with COVID-19 (20–27). Patients with chronic obstructive pulmonary disease (COPD) and interstitial lung disease (ILDs), especially Idiopathic Pulmonary Fibrosis (IPF), have a significantly higher COVID-19 mortality rate compared to patients without chronic lung disease (28). However, the molecular mechanisms underlying the increased risk of SARS-CoV-2 severity and mortality in patients with pre-existing lung diseases are not well understood.

To investigate the molecular basis of SARS-CoV-2 severity and mortality risk in chronic lung disease patients, we performed an integrated analysis of four lung single cell RNA-sequencing (scRNA-seq) datasets (29–32) in addition to unpublished data, together including 79 control and 131 chronic lung disease samples (n=31 COPD, 82 IPF and 18 other interstitial lung diseases). We found that chronic lung disease is associated with baseline changes in cell-type specific expression of genes related to viral replication and the immune response, as well as evidence of immune exhaustion and altered inflammatory gene expression. Together, these data provide a molecular framework underlying the increased risk of SARS-CoV-2 severity and poor outcomes in patients with certain pre-existing lung diseases.

METHODS

The code for genomic analyses in this paper is available at https://github.com/tgen/banovichlab/

scRNA-seq samples: scRNA-seq data were obtained from published data with samples in the "VUMC/TGen" dataset from Habermann *et. al.* (2020) (GEO accession GSE135893), samples in the "Yale/BWH" dataset came from Adams *et. al.* (2020) (GEO accession number GSE136831), samples in the "Pittsburg" dataset from Morse *et. al.* (2019) (GEO accession GSE128033) and samples in the "Northwestern" dataset from Reyfman *et al.* (2019) (GEO accession GSE122960) (Table S1). Additionally, there are 39 unpublished scRNA-seq samples in the "VUMC/TGen" dataset that were collected under Vanderbilt IRB #'s 060165, 171657 and Western IRB # 20181836.

scRNA-seq data processing and analysis: Please see online data supplement.

Immunohistochemistry of ACE2 and anti-αvβ6 integrin: Please see online data supplement.

Data availability: the majority of the data used in this manuscript is publicly available from published paper: GEO accession GSE135893 (32), GEO accession GSE136831(31), GEO accession GSE128033 (30) and GEO accession GSE122960 (29). The unpublished data from VUMC/TGen (39 samples) are included in the supplementary data (Table S3) as a count matrix format containing all the genes being used in the manuscript.

RESULTS

Expression profile of SARS-CoV-2 associated receptors and factors in the diseased lung

To determine why COVID-19 patients with chronic lung disease have a higher risk of infection severity and poorer outcomes, we performed an integrated analysis of the transcriptomes from 605,904 single cells derived from healthy donors (79 samples), COPD (31 samples), IPF (82 samples) and Non-IPF ILD (Other ILD, 18 samples) (Table S1). Using published cell type specific markers (31, 32), we identified 38 distinct cell types in the dataset (Figure S1, Table S2).

SARS-CoV-2 utilizes the host *ACE2,* and other putative factors such as *BSG* and *HSPA5*, as entry factors and *TMPRSS2*, *CTSL* or *FURIN* as priming proteases for cellular entry. Consistent with prior reports analyzing normal lung tissue (33–35), *ACE2* and *TMPRSS2* are expressed predominantly in epithelial cell types (Figure 1A), while other putative SARS-CoV-2 entry receptors (*BSG*, *HSPA5*) and priming proteases (*CTSL*, *FURIN*) have substantially more widespread expression in nearly all cell types (Figure S2). The number of *ACE2+* cells is highest in type 2 alveolar cells (AT2) and secretory cells, while *TMPRSS2* is widely expressed in all epithelial cell types. There were no significant differences in the proportion of *ACE2+* cells in any cell-type in diseased versus control groups (Figure 1B). The proportion of *TMPRSS2+* AT2 cells are decreased in IPF lungs while *TMPRSS2+* *SCGB3A2*+/*SCGB1A1+* club cells are in significantly higher numbers in COPD and IPF patients compared to controls (Figure 1C). The putative priming protease *CTSL* is expressed in more ciliated cells, proliferating macrophages, fibroblasts and pericytes isolated from Other-ILD samples (Figure S2).

Next, we compared the number of double positive cells, (cells co-expressing a receptor and priming protease), in control and disease samples. Interestingly, a notable fraction of cells co-express all established and putative entry receptors (*ACE2*, *BSG*, *HSPA5*) and proteases (*TMPRSS2*, *CTSL*, *FURIN*) (Figure 1D, Figure S3). While the percentage of cells co-expressing *ACE2* and priming proteases was similar across disease subtypes (Figure 1E, Figure S3), there were significant differences in the number of cells co-expressing *BSG*, *HSPA5* with a priming protease in disease samples in multiple cell types (Figure 1F, Figure S3).

To examine whether chronic lung disease patients express higher levels of SARS-CoV-2 receptors and priming proteases, we performed differential expression analysis of those genes in the disease versus control samples. The two major SARS-CoV-2 cellular entry factors, *ACE2* and *TMPRSS2*, have similar expression profiles in the disease and control samples. *ACE2* expression is highest in AT2 and Ciliated Cells, but there were no significant differences in *ACE2* expression in disease groups compared to control (Figure S4A). *TMPRSS2* expression is also highest in AT2 cells, and is lower in the COPD samples (Figure S4B), similar to a recent publication demonstrating decreased *TMPRSS2* expression in severe COPD (36). The putative alternative receptor *HSPA5* is down-regulated in AT2, Ciliated Cells and Transitional AT2 cells in disease groups (Figure S4B, D). Two alternative priming proteases (*CTSL* and *FURIN*) showed no significant differences in expression between control and disease samples (Figure S4E, F). However, the SARS-CoV-2 entry gene score (calculated on the average expression levels of all SARS-CoV-2 entry factors over a random control gene set) is significantly increased in the disease samples in many epithelial cell types, including AT1, AT2, Basal and Club cells, but not in *KRT5-/KRT17+* cells, an ECM-producing epithelial cell type enriched in the fibrotic lung (31, 32), (Figure 2B, Figure S5A-B).

Figure 1. Percentage of cells expressing SARS-CoV-2 receptor genes in lung cell types in different diagnosis subgroups. (*A*) Percentage of cells expressing *ACE2* and *TMPRSS2* in all cell types. Total number of *ACE2+* or *TMPRSS2+* cells in the dataset (Percentage of *ACE2+* or *TMPRSS2+* cells). (*B-C*) Percentage of cells expressing *ACE2* and *TMPRSS2* in each diagnosis group, only cell types with at least 0.5% of cells expressing *ACE2* (*B*) and at least 10% of cells expressing *TMPRSS2* (*C*) are represented. (*D*) Venn diagram shows overlapping of cells co-expressing the proposed receptors (*ACE2*, *BSG* and *HSPA5*) and the protease *TMPRSS2*. (*E-F*) Percentage of cells co-expressing receptors and *TMPRSS2* split by cell type and diagnosis group. Significant differences between diagnosis groups were calculated using Tukey HSD test, p-value < 0.05: *, p-value < 0.01: **, p-value < 0.001: ***, p-value < 0.0001: ****.

Dysregulation of viral infection and innate immune response genes in disease epithelial cells

Given the relatively modest differences in SARS-CoV-2 entry factors in epithelial cells between diseased and control lungs (Figure S4), we hypothesized that rather than greatly increased cellular susceptibility to SARS-CoV-2 infection, patients with chronic lung are predisposed to severe lung injury due to underlying differences in epithelial gene expression in key pathways mediating the antiviral response. Epithelial cells isolated from diseased lungs expressed higher levels of genes thought to directly impact viral replication (*PIKFYVE*, *TPCN2*) and the immune response to viral infection (Figure 2A). The complement pathway genes *C3* and *C4B*, important components of the innate immune

response and previously found to be elevated in SARS patients (37), are upregulated in diseased samples. Interestingly, key regulators of the host viral response including cytokine and inflammatory response genes (IFN type I and type II receptors, *SOCS1/2, CCL2, CSF3, TRIM28, EIF2AK2*) are also elevated in diseased epithelial cells. Additionally, autophagy *(ATG5, ATG7, BECN1)* and ER stress response *(EIF2AK3, ATF6, ERN1)* genes which are well known to be aberrantly regulated in IPF lungs were also upregulated in the diseased epithelial cells; these pathways are important for propagating viral infection and the host response (38–40).

In the distal lung, AT2 cells have been proposed to be the primary targets of SARS-CoV-2 (33, 35, 41). Thus, we examined the gene expression profile of diseased AT2 cells in more detail. As described above, AT2 cells in all disease subgroups have significantly higher SARS-CoV-2 entry gene scores than control cells (Figure 2B). Additionally, genes involved in the complement pathway, immune response and autophagy pathway are upregulated in diseased AT2 cells (Figure 2A). COPD and Other-ILD, but not IPF, AT2 cells express higher levels of *CSF3*, an important cytokine in the regulation of granulocytes (Figure 2D). The epithelial integrin *ITGB6,* involved in wound healing and pathogenic fibrosis (42), and the suppressors of cytokine signaling-1 and -2 (*SOCS1, SOCS2*) are upregulated in all disease AT2 cells (Figure 2D, Figure S5). Gene correlation analysis showed strong positive correlation between *ADAM17*, the main enzyme responsible for *ACE2* shedding, and *ACE2* but negatively correlated with *BSG* and *HSPA5* in the COPD and IPF samples (Figure 2C). These data suggest that basal differences in the expression profiles of genes regulating viral infection and immune response in diseased epithelial cells, specifically AT2 cells, contribute to COVID-19 severity and poor outcomes.

Disease specific *ACE2* correlated gene profiles in AT2 cells

Since *ACE2* is the best-established SARS-CoV-2 entry factor, we sought to identify *ACE2*-correlated genes in the *ACE2+* AT2 cells in different disease groups. We performed Spearman correlation analysis with Benjamini-Hochberg adjusted p-values and identified distinct gene profiles significantly correlated with *ACE2* for each disease group (Figure 2E). There were only three *ACE2* correlated genes in the Control samples with a cut-off of 99th percentile Spearman rho values and q-value less than 0.03, none of those genes are associated with the immune response. In the disease samples, we identified 671 genes (COPD: 429 genes, IPF: 140 genes and Other-ILD: 102 genes) with significant correlation to *ACE2* (99th percentile rho values, q-value less than 0.03) (Table S6). Interestingly, many *ACE2*-correlated genes in the disease samples are associated with the innate and antiviral immune response. In the COPD samples, genes with strong correlation coefficients with *ACE2* include several interferon-induced genes (*IFI6, IFI27, IFIT1, IFIT2, IFIT3*), integrin beta 7 (*ITGB7*), a modulator of innate immune function (*OAS1*), the chemokine receptor *ACKR4* and a gene associated with West Nile viral infection (*OASL*). In the IPF samples, *ACE2* expression is strongly correlated with the interferon-induced gene *GBP6* (Guanylate Binding Protein Family Member 6), the interleukin receptor *IL31RA*, and *DDX17,* a gene downstream of the IFN pathway. n other ILD diseases (non-IPF related), the cytokine *IL10*, a member of the TGF-β superfamily (*NRTN*), and an innate immune pathway component (*ITLN1*) are among the genes with high correlation with *ACE2*. Interestingly, *IFIT3* and *GBP4* (a member in the same gene family of *GBP6*) were shown to correlate with *ACE2* in asthma patients (43), and *OAS1* was among the top 50 genes with a significant correlation coefficient with *ACE2* in a previous study (34). The presence of immune-associated genes in these gene correlation profiles suggests that in patients with chronic lung diseases, *ACE2+* AT2 cells are conditioned and primed to express these genes to cope with viral infection.

Figure 2. Expression profile of SARS-CoV-2 mediators and response genes in the epithelial cell population. (*A*) Binary heatmap with log₂FC converted into a 1 (genes upregulated in the Disease samples) and 0 (genes downregulated in the Disease samples) shows the majority of the SARS-CoV-2 response genes are upregulated in the Disease samples compared to Control, the upregulation pattern in the disease samples were tested with an Exact binomial test, p-value = 9.909e-07. Up-regulation: genes being up-regulated in Disease samples, Down-regulation: genes being down-regulated in Disease samples, Not detected: gene expression was not detected in either of the two tested populations (Disease vs. Control). (*B*) SARS-CoV-2 entry module score in different cell types, SARS-CoV-2 mediators included *ACE2, BSG (CD147), HSPA5 (GRP78), TMPRSS2, CTSL, ADAM17, FURIN*. The outliers were removed in this plot, please see Figure S5A with outliers included. *:

p-value < 0.05, **: p-value < 0.01, ***: p-value < 0.001, ****: p-value < 0.0001, Tukey_HSD post-hoc test. (*C*) Significant gene expression correlation in AT2 cells between *ADAM17* and *ACE2*, *BSG* (*CD147*) and *HSPA5* (*GRP78*) in COPD and IPF samples. (*D*) Boxplot shows differences in gene expression of selected Covid-19 response genes in the AT2 cell types, **: p_value_adj < 0.05 (negative binomial test, corrected for Age, Ethnicity, Smoking_status and Dataset). (*E*) Spearman gene correlation analysis identified genes correlated with *ACE2* expression in AT2 *ACE2+* cells in different diagnosis groups, p_value was adjusted using Benjamini-Hochberg corrections and dashed lines indicate the 99th percentile of Spearman rho values.

Elevated ACE2 protein expression level in the small airway in IPF lungs

To further study the expression of the major SARS-CoV-2 entry factor *ACE2* in the fibrotic lungs, we examined the protein level of ACE2 in different lung regions. In agreement with the transcript quantification above and previous immunohistology analysis (44), we detected overall low expression

Figure 3. ACE2 and ITGB6 protein expression in IPF lung sections. (*A-C*) IPF lung sections stained for ACE2**:** (*A*) small airway, (*B*) large airway and (*C*) lung parenchyma. (*D-F*): IPF lung sections stained for αvβ6: (*D*) small airway, (*E*) large airway and (*F*) lung parenchyma. (*G*) semi–quantitative evaluation of ACE2 scoring among control and IPF sections (both the percentage of staining and staining intensity of ACE2 expression;0–Negative; 1-0–⩽10%; 2-11–⩽25%; 3-⩽26%). Significant differences between IPF and control were calculated using Tukey HSD test, p-value < 0.05 *. Scale bar = 100μm.

level of ACE2 across all tissue types in both IPF (Figure 3A-C) and control lung sections (Figure S8A). Notably, semi-quantitative evaluation of ACE2 expression scoring showed elevated ACE2 expression in all IPF sections compared to control, reaching statistical significance in the IPF small airway sections (Figure 3A, G), suggesting patients with IPF may be more susceptible to SARS-CoV-2 infection. Upregulation of the epithelial integrin alpha-V beta-6 (αvβ6) plays an important role in enhanced fibrosis in response to lung injury (45), and enhances TGFβ activation which can suppress type I interferon responses from alveolar macrophages increasing susceptibility to viral infection (46). We detected a significant increase of αvβ6 integrin expression in all lung sections isolated from IPF patients (Figure 3D-F, H). While there was additional positive staining in the peripheral lung, αvβ6 expression is highest in the AT2 epithelial cells in the IPF samples compared to overall low expression level in the normal lung sections (Figure S8B), mirroring the expression data of *ITGB6* described above (Figure 2D).

Baseline differences in inflammatory response programs in chronic lung disease

Recent publications have suggested that immune dysregulation, including sustained cytokine production and hyper-inflammation, is associated with SARS-CoV-2 severity (47–50). We performed in-depth examination of the immune cell population to determine whether pre-existing immune dysregulation in chronic lung disease patients could contribute to SARS-CoV-2 severity and mortality. We analyzed a total of 418,891 cells from 13 immune cell types (Table S2, Figure S1) and found significant increases in the proportion of CD4, CD8, cDCs and NK cells in the disease groups, most notably in COPD samples (Figure 4A). Moreover, many immune cells isolated from disease samples expressed higher levels of genes in the interferon pathway, IL6-related cytokine pathway (*IL6ST, TGFB1, AREG*) and especially major histocompatibility complex (MHC) class II genes (HLA type II genes) (Figure 4B). Expression of HLA type II genes increased across all disease groups but especially in the Other-ILD samples, compared to controls (Figure 4D). Type I IFN response (IFNa score) is slightly elevated in the diseased macrophages and pDCs (Figure S9). IL6-associated tocilizumab responsive genes (IL6 score) are expressed at a higher level in the disease groups IPF and Other-ILD, but lower in the COPD samples (Figure S9). Previous studies demonstrated elevated exhaustion levels in CD8 T cells in severely affected COVID-19 patients (51, 52). Notably, diseased immune cells, especially CD8 T cells and NK cells, have higher expression levels of cytotoxicity and exhaustion genes compared to controls (Figure 4E, F). These perturbations in the T Cell population of diseased lungs may diminish the host immunity response to viral infection, leading to a higher risk of severe disease and poor outcomes in response to SARS-CoV-2 infection.

To further investigate differences in immune cell type-specific gene expression profiles, we examined expression levels of genes associated with viral infection in disease versus control samples. Amphiregulin (AREG), a ligand for epidermal growth factor receptor (EGFR), is known to have essential roles in wound repair and inflammation resolution; furthermore, upregulation of *AREG* is associated with viral infections of the lung (53). In COVID-19 patients, *AREG* is significantly upregulated in PBMCs (54), monocytes, CD4 T Cells, NK cells, neutrophils and DCs (50), suggesting that upregulation of *AREG* may be an attempt to ameliorate the severe injury induced by SARS-CoV-2 infection. We observed reduced expression of *AREG* in the cDCs and macrophages, but not in the monocytes, in the disease samples (Figure 4C, Table S5). *SOCS1*, a suppressor of cytokine signaling, was shown to reduce the type I IFN antiviral response in bronchial epithelial cells after influenza infection (55, 56). Expression of the chemokine *CXCL8* and the IL6 co-receptor *IL6ST* was elevated in COVID-19 patients (57, 58). In our study, *CXCL8* expression is lower in the disease samples in cDCs, macrophages and monocytes while *SOCS1* expression is elevated in disease samples in these cell types

(Figure 4C, Table S5). Interestingly, *IL6ST* expression level is elevated significantly in COPD but reduced dramatically in Other-ILD samples (average two fold down-regulation compared to Control samples in all three cell types) (Figure 4C, Table S5). These basal differences in inflammatory gene expression programs highlight how chronic lung disease alters the inflammatory microenvironment encountered upon viral exposure to the peripheral lung.

Figure 4. Analysis of SARS-CoV-2 candidate immune response genes in immune cells. (*A*) quantification of cell types as a percent of all immune cells in control versus diseased lungs. (*B*) Binary heatmap shows the majority of the SARS-CoV-2 response genes are upregulated in the Disease samples compared to Control; p-value = 8.298e-08 (Exact binomial test); Up-regulation: genes being up-regulated in Disease samples, Down-regulation: genes being down-regulated in Disease samples, Not detected: gene expression was not detected in either of the two tested populations (Disease vs. Control). (*C*) Differential expression analysis for SARS-CoV-2 immune candidate genes in cDCs, Macrophages and Monocytes. *: p_adjusted_value < 0.1, **: p_adjusted_value < 0.05, p_adjusted_value were Bonferroni adjusted from Seurat FindMarkers differential expression analysis using a negative binomial test and corrected for Age, Ethnicity, Smoking_status and Dataset. (*D*) Compared to the healthy control samples, HLA type II score is higher in all disease groups (especially Other-ILD). (*E-F*) In the T cell population, cytotoxicity scores (*E*) and exhaustion scores (*F*) are higher in the disease samples than in control samples. (*A*), (*D*), (*E*) and (*F*): Boxes: interquartile range, *: p-value < 0.05, **: p-value < 0.01, ***: p-value < 0.001, ****: p-value < 0.0001, Tukey_HSD post-hoc test. See Figure S8 for plots with outliers included for (*D*), (*E*), (*F*).

DISCUSSION

The COVID-19 pandemic, caused by the SARS-CoV-2 virus, has affected tens of millions of individuals around the globe in just the first nine months of 2020. Patients with chronic lung disease have an increased risk for severe SARS-CoV-2 infection: COPD patients have a five-fold increased risk of severe COVID-19 (23, 24, 59–61) and ILD patients have up to a four-fold increased odds of death from COVID-19 (28, 62). Here, we performed an integrated transcriptomic analysis of scRNA-seq data from healthy and chronic lung disease patients to identify potential molecular causative factors determining SARS-CoV-2 severity. To summarize the results: (1) *ACE2* and *TMPRSS2* are expressed predominantly in epithelial cells and there are no significant differences in the number of *ACE2+* cells in all cell types in disease compared to control samples; (2) a viral entry score including multiple entry factors is increased in cells isolated from diseased lungs; (3) diseased epithelial cells exhibit pre-existing dysregulation of genes involved in viral infection and the immune response; (4) a unique *ACE2* correlated gene profiles for each diagnosis group included anti-viral and immune regulatory genes; (5) ACE2 protein levels are elevated in the IPF airway sections; (6) there are baseline differences in the cellular immune population in disease compared to control samples.

Similar to other coronaviruses, SARS-CoV-2 utilizes cellular receptors (*ACE2* and putatively, *BSG* and *HSPA5* gene products) and priming proteases (*TMPRSS2, CTSL, FURIN*), for viral entry. These factors are expressed predominantly in the upper and lower airways, with *ACE2* being expressed highly in nasal goblet and ciliated cells, a subset of AT2 cells and the absorptive enterocytes in the gut (33–35, 41, 44). We observed a similar expression pattern of *ACE2* in our dataset, with AT2 cells having the highest number *ACE2+* cells. To our knowledge, publications investigating baseline expression of these SARS-CoV-2 entry factors in lung disease have been limited to asthma and COPD with variable results (43, 63)(36). Despite the diversity of disease processes in our study, we observed surprisingly little variation in individual viral entry genes (receptors or proteases) between diseased and control samples, although a composite score of SARS-CoV-2 cell entry genes was increased in several epithelial cell types including AT2 cells in diseased versus control lungs. These data suggest that patients with fibrotic lung disease and COPD may have increased susceptibility to SARS-CoV-2 infection on a cellular level; this observation alone is likely insufficient to explain the increased morbidity and mortality due to COVID-19 among these patient populations.

IPF lungs exhibit abnormal expansion of epithelial cell programs, specifically the presence of more proximal specific cell types in the distal lungs (31, 32). ACE2 protein expression level is elevated in all IPF lung sections, with the highest being in the airway cells, compared to healthy lungs. While the overall frequency of *ACE2+* cells and ACE2 expression level may be low, changes in the proportional cellular makeup of the disease lung epithelium may lead to a proportionate increase *ACE2+* "infectable" cells into the distal lung. Further, increased epithelial αvβ6 integrin expression was observed in IPF patients and has been associated with a poor prognosis (64). αvβ6 integrin is central to transforming-growth factor beta activation and could contribute to more severe/fibrotic SARS-CoV-2 pathology.

A balanced immune response is crucial to viral clearance and avoidance of excessive injury to the host, as evidenced by poor outcomes related both to immunosuppression as well as hyperinflammation in COVID-19 patients(65). COPD patients with severe COVID-19 had elevated serum levels of various inflammatory cytokines including IL-2R, IL-6, IL-8, IL-10 and TNF-α suggesting there may be global alterations in the immune response (27).We observed that COPD AT2 cells expressed elevated levels of immune-response related genes (*CSF3, ITGB6, SOCS1, SOCS2*). G-CSF (encoded by *CSF3)* is found at high levels in patients with severe COVID-19 and thought to play a role in the hyperinflammatory syndrome while SOCS1 and SOCS2 are part of a negative feedback

system that regulates the response to cytokines (66, 67). *ACE2* correlated genes in this cell population were enriched for regulators of the immune response (Figure 2E). In addition to alterations in the cytokine microenvironment, changes in cellular immune populations were also identified in the COPD samples, dysregulation of several genes in inflammatory pathways (*AREG, CCL3, CXCL8, SOCS1*), and high levels of cytotoxic and exhaustion-related genes in CD4 and CD8 T Cells from COPD lungs. Expression of cytotoxic and exhaustion genes was increased compared to controls but similar in IPF and Other-ILD immune cell types. Together, our data suggest that the immune microenvironment at both the molecular and cellular level in the fibrotic and COPD lung is dysregulated and may promote severe infection and poor outcomes in COVID-19.

One limitation of our study is that we focus mainly on the peripheral regions of the lungs, and did not analyze cells in the upper airways or trachea. It is possible that there are significant differences in SARS-CoV-2 entry gene expression between disease and control samples in the more proximal regions of the lungs. Our study is also limited to the expression profiles of SARS-CoV-2 infection in baseline conditions, without viral infection.

CONCLUSION

Analysis of scRNA-sequencing data from >210 unique lungs indicates that dysregulation of genes related to viral replication and innate immune response in the epithelial cells, and baseline differences in inflammatory gene expression programs are key factors leading to an increased risk of COVID-19 severity and poorer outcomes in chronic lung disease patients.

FUNDING SOURCES
This study was supported by the NIH/NHLBI R01HL145372 (NEB/JAK), the Department of Defense W81XWH1910415 (NEB/JAK), Doris Duke Charitable Foundation (JAK), T32HL094296 (NIW, JBB), the Department of Veterans Affairs IK2BX003841 (BWR), DoD W81XWH-19-1-0131 (JCS), R01HL127349 (NK), R01HL141852 (NK), U01HL145567 (NK), UH2 HL123886 (NK), and a generous gift from Three Lakes Partners to NK and IOR. The integrated data sets were funded by various sponsors as indicated in the original publications. RGJ is funded by and NIHR Research Professorship (RP-2017-08-ST2-014).

AUTHOR CONTRIBUTIONS
L.T.B, N.I.W., M.-I.C, N.E.B and J.A.K conceived and designed the analysis. A.J.G., L.T.B., N.E.B., and J.A.K. performed quality checks, data integration, and computational analyses. L.T.B., N.E.B., and J.A.K. analyzed and interpreted scRNA-seq data. C.J. performed the immunohistology and semi-quantification analysis. L.T.B., N.I.W., M.-I.C, N.E.B., and J.A.K. wrote and revised the manuscript, with significant input from I.O.R., G.J., N.K. and the HCA Lung Biological Network. All authors read and approved the manuscript before submission.

CONFLICT OF INTERESTS
JAK has received advisory board fees from Boehringer Ingelheim, Inc, Janssen Pharmaceuticals, is on the scientific advisory board of APIE Therapeutics, and has research contracts with Genentech. In the last 36 months, NK reported personal fees from Biogen Idec, Boehringer Ingelheim, Third Rock, Samumed, Numedii, AstraZeneca, Life Max, Teravance, RohBar, and Pliant and Equity in Pliant; collaboration with MiRagen, AstraZeneca; Grant from Veracyte, all outside the submitted work. In addition, NK has a patent for New Therapies in Pulmonary Fibrosis, and Peripheral Blood Gene Expression licensed to Biotech. AGN has received advisory board fees from Boehringer Ingelheim, Galapagos, Medical Quantitative Image Analysis and personal fees for educational material from Up to Date and Boehringer Ingelheim. RGJ reports grants from Astra Zeneca, grants from Biogen, personal fees from Boehringer Ingelheim, personal fees from Chiesi, personal fees Daewoong, personal fees from Galapagos, grants from Galecto, grants from GlaxoSmithKline, personal fees from Heptares, non-financial support from NuMedii, grants and personal fees from Pliant, personal fees from Promedior, non-financial support from Redx, personal fees from Roche, other from Action for Pulmonary Fibrosis, outside the submitted work.

CONSORTIUM MEMBERS

HCA Lung Biological Network Members

Alexander V. Misharin
Division of Pulmonary and Critical Care Medicine, Northwestern University, Chicago, Illinois
a-misharin@northwestern.edu

Alexander M. Tsankov
Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY USA
alexander.tsankov@mssm.edu

Avrum Spira
Department of Medicine, Boston University School of Medicine, Boston, MA, USA; Johnson & Johnson Innovation, Cambridge, MA, USA.
aspira@bu.edu

Pascal Barbry
Université Côte d'Azur, CNRS, IPMC, Sophia-Antipolis, 06560, France
barbry@ipmc.cnrs.fr

Alvis Brazma
European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome Trust Genome Campus, Hinxton, Cambridge, CB10 1SD, UK
brazma@ebi.ac.uk

Christos Samakovlis
SciLifeLab, Department of Molecular Biosciences, Stockholm University, Stockholm Sweden and Cardiopulmonary Institute, Justus Liebig University; Giessen Germany
christos.samakovlis@scilifelab.se

Douglas P Shepherd
Center for Biological Physics and Department of Physics, Arizona State University, Tempe, AZ USA
douglas.shepherd@asu.edu

Emma L Rawlins
Wellcome Trust/ CRUK Gurdon Institute and Department Physiology, Development and Neuroscience, University of Cambridge
e.rawlins@gurdon.cam.ac.uk

Fabian J Theis
Institute of Computational Biology, Helmholtz Zentrum München and Departments of Mathematics and Life Sciences, Technical University Munich, Germany
fabian.theis@helmholtz-muenchen.de

Jennifer Griffonnet

Nice University-Affiliated Hospital : Pneumology department, Nice, 06002 France
griffonnet.j@chu-nice.fr

Haeock Lee
Department of Biomedicine and Health Sciences, The Catholic University of Korea, Seoul, Korea
haeocklee@gmail.com

Herbert B Schiller
Comprehensive Pneumology Center (CPC) / Institute of Lung Biology and Disease (ILBD), Helmholtz Zentrum München, Member of the German Center for Lung Research (DZL), Munich, Germany
herbert.schiller@helmholtz-muenchen.de

Paul Hofman
Laboratory of Clinical and Experimental Pathology, Pasteur Hospital, University Côte d'Azur, Nice, France / Hospital-Related Biobank, Pasteur Hospital, University Côte d'Azur, Nice, France / FHU OncoAge, Pasteur Hospital, BP69, 06001 Nice cedex 01, France
hofman.p@chu-nice.fr

Joseph E Powell
Garvan-Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, Sydney, NSW, Australia; UNSW Cellular Genomics Futures Institute, University of New South Wales, Sydney, NSW, Australia
j.powell@garvan.org.au

Joachim L Schultze
Department for Genomics & Immunoregulation, LIMES-Institute, University of Bonn, 53115 Bonn, Germany
PRECISE Platform for Single Cell Genomics & Epigenomics, Germany Center for Neurodegenerative Diseases and University of Bonn, Bonn, Germany
j.schultze@uni-bonn.de

Jeffrey Whitsett
Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA
jeffrey.whitsett@cchmc.org

Jiyeon Choi
Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, 20892, USA.
jiyeon.choi2@nih.gov

Joakim Lundeberg
SciLifeLab, Department of Gene Technology, KTH Royal Institute of Technology
joakim.lundeberg@scilifelab.se

Jonathan A Kropski

Division of Allergy, Pulmonary and Critical Care Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN; Department of Veterans Affairs Medical Center, Nashville, TN; Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN
jon.kropski@vumc.org

Jose Ordovas-Montanes
Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA; Program in Immunology, Harvard Medical School, Boston, MA 02115, USA; Division of Gastroenterology, Hepatology, and Nutrition, Boston Children's Hospital, Boston, MA 02115, USA; Harvard Stem Cell Institute, Cambridge, MA 02138, USA.
jordovas@mit.edu

Jayaraj Rajagopal
Harvard Stem Cell Institute, Cambridge, Massachusetts; Center for Regenerative Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA.
jrajagopal@mgh.harvard.edu

Kerstin B Meyer
Cellular Genetics Programme, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, United Kingdom.
km16@sanger.ac.uk

Mark A Krasnow
Department of Biochemistry and Wall Center for Pulmonary Vascular Disease, Stanford University, Stanford, CA 94305, USA.
krasnow@stanford.edu

Kourosh Saeb-Parsy
Department of Surgery, University of Cambridge and NIHR Cambridge Biomedical Research Centre, UK
ks10014@cam.ac.uk

Kun Zhang
UCSD Department of Bioengineering, 9500 Gilman Drive, MC0412, PFBH402, La Jolla, CA 92093, USA.
kzhang@bioeng.ucsd.edu

Robert Lafyatis
Division of Rheumatology, Department of Medicine, University of Pittsburgh Medical Center, Pittsburgh, PA, USA.
lafyatis@pitt.edu

Sylvie Leroy
Université Côte d'Azur, CHU de Nice, FHU OncoAge, Department of Pulmonary Medicine and Allergology, Nice, France ; CNRS UMR 7275 - Institut de Pharmacologie Moléculaire et Cellulaire, Sophia Antipolis, France

leroy.s2@chu-nice.fr

Muzlifah Haniffa
Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, UK; Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne NE2 4HH, UK; Department of Dermatology and NIHR Newcastle Biomedical Research Centre, Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne NE2 4LP, UK.
m.a.haniffa@ncl.ac.uk

Muzlifah Haniffa
Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, U.K.
Department of Dermatology and Newcastle NIHR Biomedical Research Centre, Royal Victoria Infirmary, The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, U.K.
m.a.haniffa@newcastle.ac.uk

Martijn C Nawijn
Department of Pathology and Medical Biology, University of Groningen, GRIAC Research Institute, University Medical Center Groningen, the Netherlands
m.c.nawijn@umcg.nl

Marko Z Nikolić
UCL Respiratory, Division of Medicine, University College London, London, UK
m.nikolic@ucl.ac.uk

Maarten van den Berge
Department of Pulmonary Diseases, University Medical Center Groningen, and Groningen Research Institute for Asthma and COPD, University of Groningen, NL-9700-RB Groningen, Netherlands
m.van.den.berge@umcg.nl

Malte Kuhnemund
Cartana AB, Nobels vag 16, 17165 Stockholm, Sweden
malte@cartana.se

Charles-Hugo Marquette
Team 4, IRCAN, FHU OncoAge, University Côte d'Azur, CNRS, INSERM, 06107 Nice CEDEX 02, France. marquette.c@chu-nice.fr.
Department of Pneumology and Oncology, CHU Nice, FHU OncoAge, University Côte d'Azur, 06100 Nice, France. marquette.c@chu-nice.fr.
Marquette.c@chu-nice.fr

Michael Von Papen
Comma Soft AG, Bonn, Germany
michael.von.paper@comma-soft.com

Naftali Kaminski

Pulmonary, Critical Care and Sleep Medicine, Yale University School of Medicine
naftali.kaminski@yale.edu

Nicholas Banovich
Translational Genomics Research Institute
nbanovich@tgen.org

Oliver Eickelberg
oliver.eickelberg@ucdenver.edu

Orit Rosenblatt-Rosen
Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
orit@broadinstitute.org

Paul A Reyfman
Northwestern University Feinberg School of Medicine, Division of Pulmonary and Critical Care Medicine
paul.reyfman@northwestern.edu

Dana Pe'er
Computational and Systems Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, New York, USA
peerster@gmail.com

Peter Horvath
Biological Research Centre of the Hungarian Academy of Sciences, Temesvári krt. 62., 6726 Szeged Hungary.
Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Tukholmankatu.
peter.horvath@fimm.fi

Purushothama Rao Tata
Department of Cell Biology, Regeneration Next Initiative, Duke University School of Medicine, Durham, NC, USA, 27710
purushothamarao.tata@duke.edu

Aviv Regev
Broad Institute of MIT and Harvard, Cambridge, MA 02142
Howard Hughes Medical Institute
Department of Biology, MIT, Cambridge, MA 02140
Genentech, 1 DNA Way, South San Francisco, CA 94080
regev.aviv.5@gmail.com (aregev@broadinstitute.org)

Mauricio Rojas
Division of Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh
rojasm@upmc.edu

Max A Seibold
Department of Pediatrics; Center for Genes, Environment, and Health; National Jewish Health; Denver, CO 80206
seiboldm@njhealth.org

Alex K Shalek
Ragon Institute of MGH, MIT, and Harvard, Cambridge, MA, USA; Institute for Medical Engineering and Science (IMES), Koch Institute for Integrative Cancer Research, and Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, USA; Broad Institute of MIT and Harvard, Cambridge, MA, USA
shalek@mit.edu

Jason R Spence
Department of Internal Medicine, Gastroenterology, University of Michigan Medical School, Ann Arbor, MI 48109, USA; Department of Cell and Developmental Biology, University of Michigan Medical School, Ann Arbor, MI 48109, USA; Department of Biomedical Engineering, University of Michigan College of Engineering, Ann Arbor, MI 48109, USA.
spencejr@umich.edu

Sarah A Teichmann
Cellular Genetics Programme, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, United Kingdom.
Dept Physics/Cavendish Laboratory, University of Cambridge, JJ Thompson Ave, Cambridge CB3 0EH, United Kingdom.
st9@sanger.ac.uk

Stephen Quake
Chan Zuckerberg Biohub
steve@czibiohub.org

Thu Elizabeth Duong
Department of Pediatrics, Respiratory Medicine, University of California, San Diego, USA
taduong@ucsd.edu

Tommaso Biancalani
Broad Institute of MIT and Harvard, Cambridge, MA 02142
tbiancal@broadinstitute.org

Tushar Desai
Department of Medicine and Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA 94116
tdesai@stanford.edu

Xin Sun
Department of Pediatrics, University of California, San Diego, San Diego, CA 92093, USA.

Laboratory of Genetics, University of Wisconsin–Madison, Madison, WI 53706, USA.
xinsun@ucsd.edu

Laure Emmanuelle Zaragosi
Université Côte d'Azur, CNRS, IPMC, Sophia-Antipolis, 06560, France
zaragosi@ipmc.cnrs.fr

REFERENCES

1. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.

2. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, China Novel Coronavirus Investigating and Research Team. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020;382:727–733.

3. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet* 2020;

4. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, Si H-R, Zhu Y, Li B, Huang C-L, Chen H-D, Chen J, Luo Y, Guo H, Jiang R-D, Liu M-Q, Chen Y, Shen X-R, Wang X, Zheng X-S, Zhao K, Chen Q-J, Deng F, Liu L-L, Yan B, Zhan F-X, Wang Y-Y, Xiao G-F, Shi Z-L. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–273.

5. Marini JJ, Gattinoni L. Management of COVID-19 Respiratory Distress. *JAMA* 2020;doi:10.1001/jama.2020.6825.

6. Murthy S, Gomersall CD, Fowler RA. Care for Critically Ill Patients With COVID-19. *JAMA* 2020;doi:10.1001/jama.2020.3633.

7. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, Greninger AL, Pipavath S, Wurfel MM, Evans L, Kritek PA, West TE, Luks A, Gerbino A, Dale CR, Goldman JD, O'Mahony S, Mikacenic C. Covid-19 in Critically Ill Patients in the Seattle Region - Case Series. *N Engl J Med* 2020;382:2012–2022.

8. Wu C, Chen X, Cai Y, Xia J 'an, Zhou X, Xu S, Huang H, Zhang L, Zhou X, Du C, Zhang Y, Song J, Wang S, Chao Y, Yang Z, Xu J, Zhou X, Chen D, Xiong W, Xu L, Zhou F, Jiang J, Bai C, Zheng J, Song Y. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med* 2020;doi:10.1001/jamainternmed.2020.0994.

9. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, Guan L, Wei Y, Li H, Wu X, Xu J, Tu S, Zhang Y, Chen H, Cao B. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–1062.

10. Fan E, Beitler JR, Brochard L, Calfee CS, Ferguson ND, Slutsky AS, Brodie D. COVID-19-associated acute respiratory distress syndrome: is a different approach to management warranted? *The Lancet Respiratory Medicine* 2020;

11. Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S, Chiumello D. COVID-19 Does Not Lead to a "Typical" Acute Respiratory Distress Syndrome. *American Journal of Respiratory and Critical Care Medicine* 2020;

12. Zheng Z, Peng F, Xu B, Zhao J, Liu H, Peng J, Li Q, Jiang C, Zhou Y, Liu S, Ye C, Zhang P, Xing Y, Guo H, Tang W. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *J Infect* 2020;81:e16–e25.

13. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, Curtis HJ, Mehrkar A, Evans D, Inglesby P, Cockburn J, McDonald HI, MacKenna B, Tomlinson L, Douglas IJ, Rentsch CT, Mathur R, Wong AYS, Grieve R, Harrison D, Forbes H, Schultze A, Croker R, Parry J, Hester F, Harper S, Perera R, Evans SJW, Smeeth L, *et al.* Factors associated with COVID-19-related

death using OpenSAFELY. *Nature* 2020;584:430–436.

14. Chen L, Yu J, He W, Chen L, Yuan G, Dong F, Chen W, Cao Y, Yang J, Cai L, Wu D, Ran Q, Li L, Liu Q, Ren W, Gao F, Wang H, Chen Z, Gale RP, Li Q, Hu Y. Risk factors for death in 1859 subjects with COVID-19. *Leukemia* 2020;34:2173–2183.

15. Wortham JM. Characteristics of persons who died with COVID-19—United States, February 12--May 18, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69.:

16. Dowd JB, Andriano L, Brazel DM, Rotondi V, Block P, Ding X, Liu Y, Mills MC. Demographic science aids in understanding the spread and fatality rates of COVID-19. *Proc Natl Acad Sci U S A* 2020;117:9696–9698.

17. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.

18. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, and the Northwell COVID-19 Research Consortium, Barnaby DP, Becker LB, Chelico JD, Cohen SL, Cookingham J, Coppa K, Diefenbach MA, Dominello AJ, Duer-Hefele J, Falzon L, Gitlin J, Hajizadeh N, Harvin TG, Hirschwerk DA, Kim EJ, Kozel ZM, Marrast LM, Mogavero JN, Osorio GA, Qiu M, Zanos TP. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 2020;doi:10.1001/jama.2020.6775.

19. Feng Y, Ling Y, Bai T, Xie Y, Huang J, Li J, Xiong W, Yang D, Chen R, Lu F, Lu Y, Liu X, Chen Y, Li X, Li Y, Summah HD, Lin H, Yan J, Zhou M, Lu H, Qu J. COVID-19 with Different Severities: A Multicenter Study of Clinical Features. *Am J Respir Crit Care Med* 2020;201:1380–1388.

20. CDC. Coronavirus Disease 2019 (COVID-19). 2020;at <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>.

21. Southern BD. Patients with interstitial lung disease and pulmonary sarcoidosis are at high risk for severe illness related to COVID-19. *Cleve Clin J Med* 2020;doi:10.3949/ccjm.87a.ccc026.

22. George PM, Wells AU, Jenkins RG. Pulmonary fibrosis and COVID-19: the potential role for antifibrotic therapy. *Lancet Respir Med* 2020;8:807–815.

23. Lippi G, Henry BM. Chronic obstructive pulmonary disease is associated with severe coronavirus disease 2019 (COVID-19). *Respir Med* 2020;167:105941.

24. Alqahtani JS, Oyelade T, Aldhahir AM, Alghamdi SM, Almehmadi M, Alqahtani AS, Quaderi S, Mandal S, Hurst JR. Prevalence, Severity and Mortality associated with COPD and Smoking in patients with COVID-19: A Rapid Systematic Review and Meta-Analysis. *PLoS One* 2020;15:e0233147.

25. Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *The Lancet Respiratory Medicine* 2020;

26. Leung JM, Niikura M, Yang CWT, Sin DD. COVID-19 and COPD. *Eur Respir J* 2020;56.:

27. Song J, Zeng M, Wang H, Qin C, Hou H, Sun Z, Xu S, Wang G, Guo C, Deng Y, Wang Z, Ma J, Pan L, Liao B, Du Z, Feng Q, Liu Y, Xie J, Liu Z. Distinct effects of asthma and COPD comorbidity on disease expression and outcome in patients with COVID-19. *Allergy* 2020;doi:10.1111/all.14517.

28. Jenkins G, Drake T, Docherty AB, Harrison E, Quint J, Adamali H, Agnew S, Babu S, Barber C, Barratt S, Bendstrup E, Bianchi S, Castillo D, Chaudhuri N, Chua F, Coker R, Chang W, Cranshaw A, Crowley L, Dosanjh D, Fiddler C, Forrest IA, George P, Gibbons M, Groom K, Haney S, Hart

S, Heiden E, Henry M, *et al.* Outcome of hospitalisation for COVID-19 in patients with Interstitial Lung Disease: An international multicentre study. *Respiratory Medicine* 2020;doi:10.1101/2020.07.15.20152967.

29. Reyfman PA, Walter JM, Joshi N, Anekalla KR, McQuattie-Pimentel AC, Chiu S, Fernandez R, Akbarpour M, Chen C-I, Ren Z, Verma R, Abdala-Valencia H, Nam K, Chi M, Han S, Gonzalez-Gonzalez FJ, Soberanes S, Watanabe S, Williams KJN, Flozak AS, Nicholson TT, Morgan VK, Winter DR, Hinchcliff M, Hrusch CL, Guzy RD, Bonham CA, Sperling AI, Bag R, *et al.* Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. *Am J Respir Crit Care Med* 2019;199:1517–1536.

30. Morse C, Tabib T, Sembrat J, Buschur KL, Bittar HT, Valenzi E, Jiang Y, Kass DJ, Gibson K, Chen W, Mora A, Benos PV, Rojas M, Lafyatis R. Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. *Eur Respir J* 2019;54.:

31. Adams TS, Schupp JC, Poli S, Ayaub EA, Neumark N, Ahangari F, Chu SG, Raby BA, DeIuliis G, Januszyk M, Duan Q, Arnett HA, Siddiqui A, Washko GR, Homer R, Yan X, Rosas IO, Kaminski N. Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. *Sci Adv* 2020;6:eaba1983.

32. Habermann AC, Gutierrez AJ, Bui LT, Yahn SL, Winters NI, Calvi CL, Peter L, Chung M-I, Taylor CJ, Jetter C, Raju L, Roberson J, Ding G, Wood L, Sucre JMS, Richmond BW, Serezani AP, McDonnell WJ, Mallal SB, Bacchetta MJ, Loyd JE, Shaver CM, Ware LB, Bremner R, Walia R, Blackwell TS, Banovich NE, Kropski JA. Single-cell RNA sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Science Advances* 2020;

33. Ziegler CGK, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, Cao Y, Yousif AS, Bals J, Hauser BM, Feldman J, Muus C, Wadsworth MH 2nd, Kazer SW, Hughes TK, Doran B, Gatter GJ, Vukovic M, Taliaferro F, Mead BE, Guo Z, Wang JP, Gras D, Plaisant M, Ansari M, Angelidis I, Adler H, Sucre JMS, Taylor CJ, *et al.* SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* 2020;181:1016–1035.e19.

34. Sungnak W, Huang N, Bécavin C, Berg M, Queen R, Litvinukova M, Talavera-López C, Maatz H, Reichart D, Sampaziotis F, Worlock KB, Yoshida M, Barnes JL, HCA Lung Biological Network. SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med* 2020;26:681–687.

35. Muus C, Luecken MD, Eraslan G, Waghray A, Heimberg G, Sikkema L, Kobayashi Y, Vaishnav ED, Subramanian A, Smillie C, Jagadeesh K, Duong ET, Fiskin E, Triglia ET, Ansari M, Cai P, Lin B, Buchanan J, Chen S, Shu J, Haber AL, Chung H, Montoro DT, Adams T, Aliee H, Allon SJ, Andrusivova Z, Angelidis I, Ashenberg O, *et al.* Integrated analyses of single-cell atlases reveal age, gender, and smoking status associations with cell type-specific expression of mediators of SARS-CoV-2 viral entry and highlights inflammatory programs in putative target cells. *bioRxiv* 2020;doi:10.1101/2020.04.19.049254.

36. Aliee H, Massip F, Qi C, de Biase MS, van Nijnatten JL, Kersten ETG, Kermani NZ, Khuder B, Vonk JM, Vermeulen RCH, Others. Determinants of SARS-CoV-2 receptor gene expression in upper and lower airways. *medRxiv* 2020;at <https://www.medrxiv.org/content/10.1101/2020.08.31.20169946v1.abstract>.

37. Chen J-H, Chang Y-W, Yao C-W, Chiueh T-S, Huang S-C, Chien K-Y, Chen A, Chang F-Y, Wong C-H, Chen Y-J. Plasma proteome of severe acute respiratory syndrome analyzed by two-dimensional gel electrophoresis and mass spectrometry. *Proc Natl Acad Sci U S A* 2004;101:17039–17044.

38. Yang N, Shen H-M. Targeting the Endocytic Pathway and Autophagy Process as a Novel Therapeutic Strategy in COVID-19. *Int J Biol Sci* 2020;16:1724–1731.

39. So J-S. Roles of endoplasmic reticulum stress in immune responses. *Mol Cells* 2018;41:705.

40. Chan C-P, Siu K-L, Chin K-T, Yuen K-Y, Zheng B, Jin D-Y. Modulation of the unfolded protein response by the severe acute respiratory syndrome coronavirus spike protein. *J Virol* 2006;80:9279–9287.

41. Singh M, Bansal V, Feschotte C. A single-cell RNA expression map of human coronavirus entry factors. *bioRxiv* 2020;doi:10.1101/2020.05.08.084806.

42. Meecham A, Marshall JF. The ITGB6 gene: its role in experimental and clinical biology. *Gene X* 2020;5:100023.

43. Camiolo M, Gauthier M, Kaminski N, Ray A, Wenzel SE. Expression of SARS-CoV-2 receptor ACE2 and coincident host response signature varies by asthma inflammatory phenotype. *J Allergy Clin Immunol* 2020;146:315–324.e7.

44. Hikmet F, Méar L, Edvinsson Å, Micke P, Uhlén M, Lindskog C. The protein expression profile of ACE2 in human tissues. doi:10.1101/2020.03.31.016048.

45. Jenkins G. Demystifying pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2020;

46. Meliopoulos VA, Van de Velde L-A, Van de Velde NC, Karlsson EA, Neale G, Vogel P, Guy C, Sharma S, Duan S, Surman SL, Jones BG, Johnson MDL, Bosio C, Jolly L, Jenkins RG, Hurwitz JL, Rosch JW, Sheppard D, Thomas PG, Murray PJ, Schultz-Cherry S. An Epithelial Integrin Regulates the Amplitude of Protective Lung Interferon Responses against Multiple Respiratory Pathogens. *PLoS Pathog* 2016;12:e1005804.

47. Lucas C, Wong P, Klein J, Castro TBR, Silva J, Sundaram M, Ellingson MK, Mao T, Oh JE, Israelow B, Takahashi T, Tokuyama M, Lu P, Venkataraman A, Park A, Mohanty S, Wang H, Wyllie AL, Vogels CBF, Earnest R, Lapidus S, Ott IM, Moore AJ, Muenker MC, Fournier JB, Campbell M, Odio CD, Casanovas-Massana A, Yale IMPACT Team, *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* 2020;584:463–469.

48. Valle DMD, Del Valle DM, Kim-Schulze S, Huang H-H, Beckmann ND, Nirenberg S, Wang B, Lavin Y, Swartz TH, Madduri D, Stock A, Marron TU, Xie H, Patel M, Tuballes K, Van Oekelen O, Rahman A, Kovatch P, Aberg JA, Schadt E, Jagannath S, Mazumdar M, Charney AW, Firpo-Betancourt A, Mendu DR, Jhang J, Reich D, Sigel K, Cordon-Cardo C, *et al.* An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nature Medicine* 2020;doi:10.1038/s41591-020-1051-9.

49. Giamarellos-Bourboulis EJ, Netea MG, Rovina N, Akinosoglou K, Antoniadou A, Antonakos N, Damoraki G, Gkavogianni T, Adami M-E, Katsaounou P, Ntaganou M, Kyriakopoulou M, Dimopoulos G, Koutsodimitropoulos I, Velissaris D, Koufargyris P, Karageorgos A, Katrini K, Lekakis V, Lupse M, Kotsaki A, Renieris G, Theodoulou D, Panou V, Koukaki E, Koulouris N, Gogos C, Koutsoukou A. Complex Immune Dysregulation in COVID-19 Patients with Severe Respiratory Failure. *Cell Host Microbe* 2020;27:992–1000.e3.

50. Unterman A, Sumida TS, Nouri N, Yan X, Zhao AY, Gasque V, Schupp JC, Asashima H, Liu Y, Cosme C, Others. Single-Cell Omics Reveals Dyssynchrony of the Innate and Adaptive Immune System in Progressive COVID-19. *medRxiv* 2020;at <https://www.medrxiv.org/content/10.1101/2020.07.16.20153437v1>.

51. Zheng H-Y, Zhang M, Yang C-X, Zhang N, Wang X-C, Yang X-P, Dong X-Q, Zheng Y-T. Elevated exhaustion levels and reduced functional diversity of T cells in peripheral blood may predict severe progression in COVID-19 patients. *Cell Mol Immunol* 2020;

52. Zhang J-Y, Wang X-M, Xing X, Xu Z, Zhang C, Song J-W, Fan X, Xia P, Fu J-L, Wang S-Y, Xu

R-N, Dai X-P, Shi L, Huang L, Jiang T-J, Shi M, Zhang Y, Zumla A, Maeurer M, Bai F, Wang F-S. Single-cell landscape of immunological responses in COVID-19 patients. *bioRxiv* 2020;doi:10.1101/2020.07.23.217703.

53. Zaiss DMW, Gause WC, Osborne LC, Artis D. Emerging functions of amphiregulin in orchestrating immunity, inflammation, and tissue repair. *Immunity* 2015;42:216–226.

54. Xiong Y, Liu Y, Cao L, Wang D, Guo M, Jiang A, Guo D, Hu W, Yang J, Tang Z, Wu H, Lin Y, Zhang M, Zhang Q, Shi M, Liu Y, Zhou Y, Lan K, Chen Y. Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect* 2020;9:761–770.

55. Pothlichet J, Chignard M, Si-Tahar M. Cutting edge: innate immune response triggered by influenza A virus is negatively regulated by SOCS1 and SOCS3 through a RIG-I/IFNAR1-dependent pathway. *J Immunol* 2008;180:2034–2038.

56. Sallenave J-M, Guillot L. Innate Immune Signaling and Proteolytic Pathways in the Resolution or Exacerbation of SARS-CoV-2 in Covid-19: Key Therapeutic Targets? *Front Immunol* 2020;11:1229.

57. Coperchini F, Chiovato L, Croce L, Magri F, Rotondi M. The cytokine storm in COVID-19: An overview of the involvement of the chemokine/chemokine-receptor system. *Cytokine Growth Factor Rev* 2020;53:25–32.

58. Zhu L, Yang P, Zhao Y, Zhuang Z, Wang Z, Song R, Zhang J, Liu C, Gao Q, Xu Q, Wei X, Sun H-X, Ye B, Wu Y, Zhang N, Lei G, Yu L, Yan J, Diao G, Meng F, Bai C, Mao P, Yu Y, Wang M, Yuan Y, Deng Q, Li Z, Huang Y, Hu G, *et al.* Single-Cell Sequencing of Peripheral Mononuclear Cells Reveals Distinct Immune Response Landscapes of COVID-19 and Influenza Patients. *Immunity* 2020;53:685–696.e3.

59. Banerjee A, Pasea L, Harris S, Gonzalez-Izquierdo A, Torralbo A, Shallcross L, Noursadeghi M, Pillay D, Sebire N, Holmes C, Pagel C, Wong WK, Langenberg C, Williams B, Denaxas S, Hemingway H. Estimating excess 1-year mortality associated with the COVID-19 pandemic according to underlying conditions and age: a population-based cohort study. *Lancet* 2020;395:1715–1725.

60. Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, Holden KA, Read JM, Dondelinger F, Carson G, Merson L, Lee J, Plotkin D, Sigfrid L, Halpin S, Jackson C, Gamble C, Horby PW, Nguyen-Van-Tam JS, Ho A, Russell CD, Dunning J, Openshaw PJ, Baillie JK, Semple MG, ISARIC4C investigators. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;369:m1985.

61. Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, Liu L, Shan H, Lei C-L, Hui DSC, Du B, Li L-J, Zeng G, Yuen K-Y, Chen R-C, Tang C-L, Wang T, Chen P-Y, Xiang J, Li S-Y, Wang J-L, Liang Z-J, Peng Y-X, Wei L, Liu Y, Hu Y-H, Peng P, Wang J-M, Liu J-Y, *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382:1708–1720.

62. Esposito AJ, Menon AA, Ghosh AJ, Putman RK, Fredenburgh LE, El-Chemaly SY, Goldberg HJ, Baron RM, Hunninghake GM, Doyle TJ. Increased Odds of Death for Patients with Interstitial Lung Disease and COVID-19: A Case-Control Study. *Am J Respir Crit Care Med* 2020;doi:10.1164/rccm.202006-2441LE.

63. Kermani N, Song W-J, Lunt A, Badi Y, Versi A, Yike GUO, Sun K, Bhavsar P, Howarth P, Dahlen S-E, Others. Airway expression of SARS-CoV-2 receptor, ACE2, and proteases, TMPRSS2 and furin, in severe asthma. *medRxiv* 2020;at <https://www.medrxiv.org/content/10.1101/2020.06.29.20142091v1.abstract>.

64. Saini G, Porte J, Weinreb PH, Violette SM, Wallace WA, McKeever TM, Jenkins G. αvβ6 integrin may be a potential prognostic biomarker in interstitial lung disease. *Eur Respir J* 2015;46:486–494.

65. Fung M, Babik JM. COVID-19 in Immunocompromised Hosts: What We Know So Far. *Clin Infect Dis* 2020;doi:10.1093/cid/ciaa863.

66. Lu C, Li S, Liu Y. Role of immunosuppressive therapy in rheumatic diseases concurrent with COVID-19. *Ann Rheum Dis* 2020;79:737–739.

67. Yoshimura A, Naka T, Kubo M. SOCS proteins, cytokine signalling and immune regulation. *Nat Rev Immunol* 2007;7:454–465.