Restructuring of genomic provinces of surface ocean plankton under climate change

Paul Frémont[1,2*], Marion Gehlen[3*], Mathieu Vrac[3], Jade Leconte[1,2], Patrick Wincker[1,2], Daniele Iudicone[4], Olivier Jaillon[1,2*]

[1]Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Université d'Evry, Université Paris-Saclay, 91057 Evry, France. [2]Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara Oceans, Paris, France. [3]LSCE-IPSL, CEA/CNRS/Université Paris-Saclay, Gif-sur-Yvette, France. [4]Stazione Zoologica Anton Dhorn. Villa Comunale, 80121, Naples, Italy.

*Corresponding authors: pfremont@genoscope.cns.fr; marion.gehlen@lsce.ipsl.fr; ojaillon@genoscope.cns.fr

Abstract

The impact of climate change on diversity, functioning and biogeography of marine plankton is a major unresolved scientific issue. Here, niche theory is applied on plankton metagenomes sampled during the *Tara* Oceans expedition to derive pan-ocean geographical structuring in *climato-genomic* provinces characterized by signature genomes for 6 size fractions, from viruses to meso-zooplankton. Assuming a high warming scenario (RCP8.5), the identified tropical provinces would expand and temperate provinces would shrink. Poleward shifts are projected for 96% of provinces in five major basins leading to their reorganization over ~50% of the surface ocean south of 60°N, of which 3% correspond to novel assemblages of provinces. Sea surface temperature is identified as the main driver and accounts only for ~51 % of the changes followed by phosphate (11%) and salinity (10.3%). These results demonstrate the potential of integration of genomics with physico-chemical data for higher scale modeling and understanding of ocean ecosystems.

Planktonic communities are composed of complex and heterogeneous assemblages of small animals, small single-celled eukaryotes (protists), prokaryotes and viruses - that drift with currents. They contribute to the regulation of the Earth system notably through primary production via photosynthesis[1], carbon export to the deep oceans[2,3] and form the base of the food webs that sustain the whole trophic chain in the oceans and beyond[4].

The composition of communities is known to vary over time at a given site with daily[5] to seasonal fluctuations[6] following environmental variability[7,8]. Overlying these relatively

short scale spatio-temporal variations, a more macroscale partitioning of the ocean was

evidenced by different combinations of biological and physico-chemical data[9-11], and

recently at the resolution of community genomics[12]. The basin scale biogeographical

structure was proposed to result from a combination of multiple bio-physico-chemical

processes named the seascape[7,8], including both abiotic and biotic interactions[13], neutral

genetic drift[14], natural selection[15-17], temperature variations, nutrient supply but also

advection and mixing along currents[12,14].

Today, knowledge of global scale plankton biogeography at the DNA level is in its infancy.

We lack understanding and theoretical explanations for the emergence and maintenance of

biogeographical patterns at genomic resolution. Omics data (*i.e.* the DNA/RNA sequences

representative of the variety of coding and non-coding sequences of organisms) provide

the appropriate resolution to track and record global biogeographical features[12],

modulation of the repertoire of expressed genes in a community in response to

environmental conditions[2,18,19] as well as eco-evolutionary processes[14,16,17]. Importantly,

metagenomic sequencing can be consistently analyzed across plankton organisms as

recently demonstrated by global expeditions[20-23]. Furthermore, the strong links between

plankton and environmental conditions suggest potentially major consequences of climate

change on community composition and biogeography[24,25]. Time series observations have

highlighted recent changes in the planktonic ecosystem attributed to this anthropogenic

pressure, such as changes in community composition[26-28] or poleward shifts of some

species[29,30]. These changes are expected to intensify with ongoing climate warming and

could lead to major reorganizations in plankton community composition[24,25], with a

potential decline in diversity[31-33]. Another major consequence of a global reorganization of

the seascape on biological systems (e.g. growth, grazing) would be a decrease of primary

production at mid-latitudes and an increase at higher latitudes[34].

Here we report a global structure of plankton biogeography based on metagenomic data

using niche models and its putative modifications under climate change. First, we show

that environmental niches[35], *i.e.* the envelope of environmental parameters suitable for an

organism or a population, can be defined at the scale of genomic provinces across 6

organism size fractions representing major plankton groups from nano- (viruses) to meso-

zooplankton (small metazoans). Then, we spatially extrapolate their niches into *climato-*

genomic provinces to depict the structure of plankton biogeography of all but arctic regions

for each size fraction and for all combined. Then, considering the same niches, we assess

putative spatial reorganization of the same provinces and their associated environmental

drivers under climate change at the end of the century.

Niche models and signature genomes from genomic provinces

We use 38 previously defined genomic provinces[12] containing at least 4 sampling sites;

they correspond to 595 metagenomes for 6 size fractions (ranging from 0 to 2000 μm) and

sampled in 95 sites from all oceans except the Arctic (Supplementary Figs. 1-2).

To compute and test the validity of realized environmental niches, we train four machine

learning techniques to probabilistically associate genomic provinces with environmental

data: sea surface temperature, salinity, three macronutrients (dissolved silica, nitrate and

phosphate), one micronutrient (dissolved iron) plus a seasonality index of nitrate. A valid

environmental niche is obtained for 27 out of 38 initial provinces (71%) comforting their

definition and covering 529 samples out of 595 (89%, Supplementary Fig. 2). Rejected

provinces contain relatively few stations (mean of 6 ± 2.6 versus 19 ± 15.3 for valid

provinces, p-value<10^{-3} Wilcoxon test). For spatial and temporal extrapolations of the

provinces presented below, we use the ensemble model approach[36] that considers mean

predictions of machine learning techniques.

The signal of ocean partitioning is likely due to abundant and compact genomes whose

geographical distributions closely match provinces. Within a collection of 523 prokaryotic

and 713 eukaryotic genomes[37,38] from *Tara* Oceans samples, we find signature genomes for

all but 4 provinces. In total, they correspond to respectively 96 and 52 of the genomes and

their taxonomies are coherent with the size fractions (Fig. 1 for eukaryotes and

Supplementary Fig. 3 for prokaryotes). Some of them correspond to unexplored lineages

highlighting the gap of knowledge for organisms that structure plankton biogeography and

the strength of a rationale devoid of any *a priori* on reference genomes or species.

Structure of present day biogeography of plankton

To extrapolate to a global ocean biogeography for each size fraction, we define the most

probable provinces, named hereafter as *dominant* and assigned to a climatic annotation

(Supplementary Table 1), on each 1°x1° resolution grid point using 2006-13 WOA13

climatology[39] (Supplementary Fig. 4 and Fig. 3).

Overall, in agreement with previous observations[12], provinces of large size fractions (>20

µm) are wider and partially decoupled from those of smaller size fractions, probably due to

differential responses to oceanic circulation and environmental variations, different life

cycle constraints, lifestyles[7,8,12] and trophic networks positions[40]. Biogeographies of small

metazoans that enrich the largest size fractions (180-2000 µm and 20-180 µm) are broadly

aligned with latitudinal bands (tropico-equatorial, temperate and (sub)-polar) dominated

by a single province (Fig. 2a,b). A more complex oceanic structuring emerges for the

smaller size fractions (<20 µm) (Fig. 2c-f) with several provinces per large geographical

region. Taking size fraction 0.8-5 µm enriched in small protists (Fig. 2d) as an example,

distinct provinces are identified for oligotrophic gyres in the Atlantic and Pacific Oceans,

and one for the nutrient-rich equatorial upwelling region. A complex pattern of provinces,

mostly latitudinal, is also found for the bacteria enriched size class (Fig. 2e, 0.22-3 µm) and

the virus enriched size class (Fig. 2f, 0-0.2 µm) though less clearly linked to large-scale

oceanographic regions. A single province extending from temperate to polar regions

emerges from the size fraction 5-20 µm enriched in protists (Fig. 2c), for which a smaller

number of samples is available (Supplementary Fig. 2b-c), which probably biases this

result. Finally, we use PHATE[41] dimension reduction algorithm to combine all provinces for

all size classes into a single consensus biogeography revealing 4 or 7 robust clusters (Fig.

2g,h). The 4 cluster consensus biogeography is mainly latitudinally organized

distinguishing polar, subpolar, temperate and tropico-equatorial regions. The 7 cluster

consensus biogeography distinguishes the equatorial pacific upwelling biome and three

subpolar biomes that most likely reflect the chemico-physical structuring of the Southern

Ocean and known polar fronts (red lines Fig. 2h). However, learning data are scarcer south

of 60°S so these extrapolations need to be taken with caution.

Previous ocean partitioning either in biomes[9–11] or biogeochemical provinces (BGCPs)[9,11]

are based on physico-biogeochemical characteristics including SST[9–11], chlorophyll a[9–11],

salinity[9–11], bathymetry[9,11], mixed layer depth[10] or ice fraction[10]. Considering three of these

partitions as examples we notice differences with our partitions (Supplementary Fig. 7-8)

for example in terms of number of regions in the considered oceans (56 for 2013

Reygondeau et al. BGCPs[11], 17 for Fay and McKingley[10]) and structure (the coastal biome

for 2013 Reygondeau et al. biomes[11]). Numerical comparison of our partitions with others

(*Methods*), reveals low similarity between them, the highest being with Reygondeau biomes

(Supplementary Figs. 7-9). However, biomes and BGCP frontiers closely match our

province frontiers in many cases. Near the frontiers, *dominant* provinces have smaller

probabilities in agreement with smooth transitions instead of sharp boundaries as already

proposed[9] and with a seasonal variability of the frontiers[11] (Supplementary Fig 7). Some of

these transitions are very large and match entire BGCPs, for example in subtropical North

Atlantic and subpolar areas where high annual variations are well known[11].

Future changes in plankton biogeography structure

We assess the impacts of climate change on plankton biogeography at the end of the 21st

century following the Representative Concentration Pathway 8.5 (RCP8.5)[42] greenhouse

gas emission scenario. To consistently compare projections of present and future

biogeographies with coherent spatial structures, we use a bias-adjusted mean of 6 Earth

System Model (ESM) climatologies (Supplementary Table 2, Supplementary Fig. 10,

Materials and Methods). The highest warming (7.2°C) is located off the east coast of Canada

in the North Atlantic while complex patterns of salinity and nutrient variations are

projected in all oceans (Supplementary Figure 11). According to this scenario, future

temperature at most sampling sites will be higher than the mean and maximum

contemporary temperature within their current province (Supplementary Fig. 12).

Our projections indicate multiple large-scale changes in biogeographical structure

including plankton organism size-dependent province expansions or shrinkages and shifts

(Supplementary Fig. 13-15). A change in the *dominant* province in at least one size fraction

would occur over 60.1 % of the ocean surface, ranging from 12 % (20-180 μm) to 31%

(0.8-5 μm) (Table 1).

Centroids of provinces with *dominance* areas larger than 10^6 km^2 within a basin would be

moved at least 200 km away for 77% of them, 96 % of which move poleward

(Supplementary Figs. 14 and 14). Most longitudinal shift distances are smaller (50% <190

km) but a few are larger than 1000 km while the distribution of latitudinal shifts is more

concentrated around the mean (290 km) with no shifts superior to 1000 km

(Supplementary Fig. 15b). These important longitudinal shifts corroborate existing

projections[24,43,44] and differ from trivial poleward shifts due to temperature increase,

reflecting more complex spatial rearrangements of the other environmental drivers

(Supplementary Fig. 11). We find an average displacement speed of the provinces'
centroids of 76 ± 79 km.dec^{-1} (latitudinally mean of 34 ± 82 km.dec^{-1} and 59 ± 82 km.dec^{-1}
longitudinally; median of 47 km.dec^{-1}, 23 km.dec^{-1} latitudinally and 27 km.dec^{-1}
longitudinally). We project phytoplankton provinces displacements similar to previously
published shifts for the North Pacific (poleward shift of 118 km.dec^{-1} for province C4
versus 100 km.dec^{-1} for the subtropical biome of Polovina et al.[44] and eastward shift of 195
km.dec^{-1} for province C9 versus 200 km.dec^{-1} for the equatorial biome of Polovina et al.[44]).
For all size fractions, climate change would lead to a poleward expansion of tropical and
equatorial provinces at the expense of temperate provinces (Table 1 and Supplementary
Fig. 13). This is illustrated for the size fraction 180-2000 by the example of the temperate
province F5 (Supplementary Fig. 16) for which significant shrinkage is projected in the five
major ocean basins. In the North Atlantic, its centroid would move approximately 800 km
to the northeast (Supplementary Fig. 16c).
To simplify comparisons between future and present biogeographies, we combine all
projections into two comparable consensus maps (Fig. 2e,f). Some particularly visible
patterns of geographical reorganization are common to several or even all size fractions
and visible on the consensus maps (Fig. 2e,f compared to Fig. 2a-d and Supplementary Fig.
13). For example, the tropico-equatorial and tropical provinces expand in all size fractions
and the provinces including the pacific equatorial upwelling shrink for size fractions
smaller than 20 µm.
To further quantify patterns of expansion or shrinkage, we calculate the surface covered by
the *dominant* provinces weighted by probabilities of presence (Supplementary Fig. 17,
Supplementary Table 1). In this way, *dominant* provinces are defined on 100% of the
surface ocean (327 millions of km^2) but their presence probabilities correspond to the
equivalent of 45 to 74% (due to sampling variability and niche overlaps) of the surface
ocean depending on plankton size fraction (Table 1). Overall, our results indicate
expansions of the surface of tropical and tropico-equatorial provinces but in very different
ways depending on the size fractions of organisms. The surface area of temperate
provinces is ~22 million km^2 on average (from 10 Mkm^2 for 0-0.2 µm to 49 Mkm^2 for 20-
180 µm) and should decrease by 36% on average (from -20 % for 5-20 µm up to -54% for
0.8-5 µm, -12 million km^2 on average, +6 % for 0.22-3 µm). Tropical provinces cover ~118

million km² on average (from 86 Mkm² for 0.8-5 µm up to 169 Mkm² for 180-2000 µm) and

their coverage should increase by 32% on average (from +13% for 0-0.2 µm up to +75%

for 0.8-5 µm, +25 million km² on average) (Supplementary Fig. 17 and Supplementary

Table 1).

We calculate at each grid point a single dissimilarity index (*Materials and Methods*)

between probabilities of future and present *dominant* provinces for all size fractions

combined (Fig. 4a). Areas located between future and present borders of expanding and

shrinking provinces would be the most subject to replacements by other contemporary

provinces, as exemplified by the poleward retraction of the southern/northern edges of the

temperate provinces (red arrows on Fig. 4a, Table 1). High dissimilarities are obtained

over northern (25° to 60°) and symmetrically southern (-25 to -60°) temperate regions

(mean of 0.29 and 0.24 respectively). Despite important environmental changes in austral

and equatorial regions (Supplementary Fig. 11) and projected change in diversity[31–33] and

biomass[45], the contemporary provinces remain the most probable at the end of the century

using our statistical models (mean dissimilarities of 0.18 and 0.02 respectively) as no

known contemporary provinces could replace them.

To further study the decoupling between provinces of different size fractions in the future

we considered the assemblages of *dominant* provinces of each size fraction. By using two

differently stringent criteria, from 45.3 to 57.1% of ocean surface, mainly located in

temperate regions, would be inhabited in 2090-99 by assemblages that exist elsewhere in

2006-15 (Fig. 4b versus Fig. 4c). Contemporary assemblages would disappear on 3.5 to

3.8% of the surface, and, conversely, novel assemblages, not encountered today, would

cover 2.9 to 3.0% of the surface. These areas appear relatively small but they include some

important economic zones (Fig 4b, Supplementary Fig. 18). On 41.8% to 51.8% of the

surface of the main fisheries and 41.2% to 54.2% of Exclusive Economic Zones (*Materials*

and Methods), future assemblages would differ from those present today (Supplementary

Fig. 18).

Drivers of plankton biogeography reorganization

We quantify the relative importance of considered environmental parameters

(temperature, salinity, dissolved silica, macronutrients and seasonality of nitrate) into

niche definition and in driving future changes of the structure of plankton biogeography.

Among environmental properties that define the niches, temperature is the first influential parameter (for 19 niches out of 27) but only at 22.6% on average. Particularly, the distribution of relative influences of temperature is spread over a much wider range than that of other parameters (Supplementary Fig. 18a).

The relative impact of each environmental parameter is calculated[24] for each site presenting a significant dissimilarity between 2006-15 and 2090-99 (Fig 5a). Overall, SST would be responsible for reorganization of half of the provinces followed by Phosphate (11 %) and Salinity (10.3%) (Supplementary Fig. 20). SST is the primary driver over the majority of the ocean (Fig. 5a). In some regions, salinity (e.g. eastern North Atlantic) and Phosphate (e.g. equatorial region) dominate (Fig. 5a) and excluding the effect of SST, they are the primary drivers of global reorganization of the provinces (Fig. 5 b). The impact of SST varies across size classes with a significantly higher contribution in large size classes (>20 μm) compared to the small ones (mean of ~73% versus ~49%; Fig. 5c). Even though the contribution of combined nutrients to niche definition is similar for small and large size classes under present day conditions (mean of ~56% versus ~61%, Supplementary Fig. 19, Supplementary Table 3), their future projected variations have a higher relative impact on the reorganization of small organisms' biogeographies (mean of ~39% versus ~20 %, t-test p-value < 0.05, Supplementary Fig. 19, Supplementary Table 3). For instance, in the tropical zone, the shrinkage of the equatorial province (province C9, size fraction 0.8 – 5 μm, Fig. 2b,d, Supplementary Fig. 16e) is driven at 24 % by reduction of dissolved phosphate concentrations and at 25% by SST increase. In contrast, SST and Salinity would drive respectively at 56% and 16% the shrinkage of the temperate province F5 of size fraction 180-2000 μm (Supplementary Fig. 16d versus e). Non-poleward shifts are found only within small size fractions (<20 μm) (Supplementary Fig. 14, 15) highlighting differential responses to nutrients and SST changes between large and small size classes, the latter being enriched in phytoplankton that directly rely on nutrient supplies.

Discussion

We propose a novel partitioning of the ocean in plankton size dependent *climato-genomic* provinces, complementing pioneer and recent efforts based on other bio-physico-chemical data[9–11]. Though they are initially built at genomic scale, our biogeographies paradoxically reveal basin scale provinces that are larger than BGCPs[11] and biomes[10], and probably

relatively stable across seasons evoking limited effects of seasonality on frontiers positions
of BGCPs provinces[11]. We propose that this apparent paradox emerges from the
combination of the scale, nature and resolution of sampling. First, two proximal samples
from the *Tara* Oceans expedition are separated by ~300 km on average sampled over three
years; this relatively large spatio-temporal scale overlies shorter scale compositional
variations previously observed[5,6]. Second, our estimates of plankton community
dissimilarities are highly resolutive; they are computed at genomic scale with billions of
small DNA fragments[12,20] thus smoothing the more discrete species level signal. Together,
from these combinations of processes and patterns occurring at multiple scales emerge
basin scale provinces associated with coherent environmental niches and signature
genomes.

These *climato-genomic* provinces are structured in broad latitudinal bands with smaller
organisms (<20 μm) displaying more complex patterns and partially decoupled from larger
organisms. This decoupling is the result of distinct statistical links between provinces
based on organism size fractions and environmental parameters and could reflect their
respective trophic modes[40].

Complex changes of the parameters defining the niches are projected under climate change
leading to size-dependent modifications of biogeographical patterns, as for example
smaller organisms being more sensitive to nutrient changes. Assuming the maintenance of
environmental characteristics that define the *climato-genomic* provinces, climate change is
projected to restructure them over approximately 50% of surface oceans south of 60°N by
the end of the century (Fig. 4). The largest reorganization is detected in subtropical and
temperate regions in agreement with other studies[32,44] and is accompanied by appearance
and disappearance of size-fractionated provinces' assemblages. For tropico-equatorial and
austral regions, out of contemporary range and novel environmental conditions are
projected. While some studies extrapolate important diversity and biomass changes in
these zones[31–33,45], here we project shifts of their boundaries and maintain their climatic
label. However, the present approach does not account for putative changes in community
composition nor the emergence of novel niches over these regions for which novel
environmental selection pressure is expected.

Overall, our projections for the end of the century do not take into account possible future

changes of major bio-physico-chemical factors such as dynamics of community mixing,

trophic interactions through transport[46], the potential dynamics of the genomes[14,16,17]

(adaptation or acclimation) and biomass variations[45]. New sampling in current and future

expeditions[47], as well as ongoing technological improvements in bio-physico-chemical

characterization of seawater samples[38,47,48] will soon refine functional[18,49], environmental

(micronutrients[50]) and phylogenetic[16,17] characterization of plankton ecosystems for

various biological entities (genotypes, species or communities) and spatio-temporal

scales[47]. Ultimately, crossing this varied information will allow a better understanding of

the conditions of emergence of ecological niches in the seascape and their response to a

changing ocean.

References

1. Field, C. B., Behrenfeld, M. J., Randerson, J. T. & Falkowski, P. Primary production of the
biosphere: Integrating terrestrial and oceanic components. *Science (80-.).* (1998).
doi:10.1126/science.281.5374.237

2. Guidi, L. *et al.* Plankton networks driving carbon export in the oligotrophic ocean. *Nature*
(2016). doi:10.1038/nature16942

3. Henson, S. A., Sanders, R. & Madsen, E. Global patterns in efficiency of particulate organic
carbon export and transfer to the deep ocean. *Global Biogeochem. Cycles* (2012).
doi:10.1029/2011GB004099

4. Azam, F. *et al.* The Ecological Role of Water-Column Microbes in the Sea. *Mar. Ecol. Prog. Ser.*
(1983). doi:10.3354/meps010257

5. Saab, M. A. abi. Day-to-day variation in phytoplankton assemblages during spring blooming
in a fixed station along the Lebanese coastline. *J. Plankton Res.* (1992).
doi:10.1093/plankt/14.8.1099

6. Djurhuus, A. *et al.* Environmental DNA reveals seasonal shifts and potential interactions in a
marine community. *Nat. Commun.* (2020). doi:10.1038/s41467-019-14105-1

7. Kavanaugh, M. T. *et al.* Seascapes as a new vernacular for pelagic ocean monitoring,
management and conservation. *ICES J. Mar. Sci.* (2016). doi:10.1093/icesjms/fsw086

8. Steele, J. H. The ocean 'landscape'. *Landsc. Ecol.* (1989). doi:10.1007/BF00131537

9. Longhurst, A. R. *Ecological Geography of the Sea. Ecological Geography of the Sea* (2007).
doi:10.1016/B978-0-12-455521-1.X5000-1

10.   Fay, A. R. & McKinley, G. A. Global open-ocean biomes: Mean and temporal variability. *Earth*
312   *Syst. Sci. Data* (2014). doi:10.5194/essd-6-273-2014

313   11.   Reygondeau, G. *et al.* Dynamic biogeochemical provinces in the global ocean. *Global*
314   *Biogeochem. Cycles* (2013). doi:10.1002/gbc.20089

315   12.   Richter, D. J. *et al.* Genomic evidence for global ocean plankton biogeography shaped by
316   large-scale current systems. *bioRxiv* 867739 (2019). doi:10.1101/867739

317   13.   Dutkiewicz, S. *et al.* Dimensions of marine phytoplankton diversity. *Biogeosciences* (2020).
318   doi:10.5194/bg-17-609-2020

319   14.   Hellweger, F. L., Van Sebille, E. & Fredrick, N. D. Biogeographic patterns in ocean microbes
320   emerge in a neutral agent-based model. *Science (80-.).* (2014). doi:10.1126/science.1254421

321   15.   Sauterey, B., Ward, B., Rault, J., Bowler, C. & Claessen, D. The Implications of Eco-Evolutionary
322   Processes for the Emergence of Marine Plankton Community Biogeography. *Am. Nat.* (2017).
323   doi:10.1086/692067

324   16.   Laso-Jadart, R. *et al.* Investigating population-scale allelic differential expression in wild
325   populations of Oithona similis (Cyclopoida, Claus, 1866). *Ecol. Evol.* (2020).
326   doi:10.1002/ece3.6588

327   17.   Delmont, T. O. *et al.* Single-amino acid variants reveal evolutionary processes that shape the
328   biogeography of a global SAR11 subclade. *Elife* (2019). doi:10.7554/eLife.46497

329   18.   Carradec, Q. *et al.* A global ocean atlas of eukaryotic genes. *Nat. Commun.* (2018).
330   doi:10.1038/s41467-017-02342-1

331   19.   Salazar, G. *et al.* Gene Expression Changes and Community Turnover Differentially Shape the
332   Global Ocean Metatranscriptome. *Cell* (2019). doi:10.1016/j.cell.2019.10.014

333   20.   Alberti, A. *et al.* Viral to metazoan marine plankton nucleotide sequences from the Tara
334   Oceans expedition. *Sci. Data* (2017). doi:10.1038/sdata.2017.93

335   21.   Pesant, S. *et al.* Open science resources for the discovery and analysis of Tara Oceans data.
336   *Sci. Data* (2015). doi:10.1038/sdata.2015.23

337   22.   Karsenti, E. *et al.* A holistic approach to marine Eco-systems biology. *PLoS Biol.* (2011).
338   doi:10.1371/journal.pbio.1001177

339   23.   Duarte, C. M. Seafaring in the 21st century: The Malaspina 2010 circumnavigation
340   expedition. *Limnology and Oceanography Bulletin* (2015). doi:10.1002/lob.10008

341   24.   Barton, A. D., Irwin, A. J., Finkel, Z. V. & Stock, C. A. Anthropogenic climate change drives shift
342   and shuffle in North Atlantic phytoplankton communities. *Proc. Natl. Acad. Sci.* (2016).
343   doi:10.1073/pnas.1519080113

25. Benedetti, F., Guilhaumon, F., Adloff, F. & Ayata, S. D. Investigating uncertainties in
zooplankton composition shifts under climate change scenarios in the Mediterranean Sea.
Ecography (Cop.). (2018). doi:10.1111/ecog.02434

26. Beaugrand, G. *et al.* Prediction of unprecedented biological shifts in the global ocean. *Nature*
Climate Change (2019). doi:10.1038/s41558-019-0420-1

27. McMahon, K. W., McCarthy, M. D., Sherwood, O. A., Larsen, T. & Guilderson, T. P. Millennial-
scale plankton regime shifts in the subtropical North Pacific Ocean. *Science (80-.).* (2015).
doi:10.1126/science.aaa9942

28. Rivero-Calle, S., Gnanadesikan, A., Del Castillo, C. E., Balch, W. M. & Guikema, S. D.
Multidecadal increase in North Atlantic coccolithophores and the potential role of rising CO_2.
Science (80-.). (2015). doi:10.1126/science.aaa8026

29. Beaugrand, G. Decadal changes in climate and ecosystems in the North Atlantic Ocean and
adjacent seas. *Deep. Res. Part II Top. Stud. Oceanogr.* (2009). doi:10.1016/j.dsr2.2008.12.022

30. Pinsky, M. L., Worm, B., Fogarty, M. J., Sarmiento, J. L. & Levin, S. A. Marine taxa track local
climate velocities. *Science (80-.).* (2013). doi:10.1126/science.1239352

31. Thomas, M. K., Kremer, C. T., Klausmeier, C. A. & Litchman, E. A global pattern of thermal
adaptation in marine phytoplankton. *Science (80-.).* (2012). doi:10.1126/science.1224836

32. Busseni, G. *et al.* Large scale patterns of marine diatom richness: Drivers and trends in a
changing ocean. *Glob. Ecol. Biogeogr.* (2020). doi:10.1111/geb.13161

33. Ibarbalz, F. M. *et al.* Global Trends in Marine Plankton Diversity across Kingdoms of Life. *Cell*
(2019). doi:10.1016/j.cell.2019.10.008

34. Bopp, L. *et al.* Multiple stressors of ocean ecosystems in the 21st century: Projections with
CMIP5 models. *Biogeosciences* (2013). doi:10.5194/bg-10-6225-2013

35. Hutchinson, G. E. Concludig remarks. *Cold Spring Harb. Symp. Quant. Biol.* (1957).

36. Jones, M. C. & Cheung, W. W. L. Multi-model ensemble projections of climate change effects
on global marine biodiversity. *ICES J. Mar. Sci.* (2015). doi:10.1093/icesjms/fsu172

37. Delmont, T. O. *et al.* Nitrogen-fixing populations of Planctomycetes and Proteobacteria are
abundant in surface ocean metagenomes. *Nat. Microbiol.* (2018). doi:10.1038/s41564-018-
0176-9

38. Delmont, T. O. *et al.* Functional repertoire convergence of distantly related eukaryotic
plankton lineages revealed by genome-resolved metagenomics. *bioRxiv* 2020.10.15.341214
(2020). doi:10.1101/2020.10.15.341214

39. Boyer, T. P. *et al.* WORLD OCEAN DATABASE 2013, NOAA Atlas NESDIS 72. *Sydney Levitus,*

Ed.; Alexey Mishonoc, Tech. Ed. (2013). doi:10.7289/V5NZ85MT

40.   Sunagawa, S. *et al.* Tara Oceans: towards global ocean ecosystems biology. *Nature Reviews*
Microbiology (2020). doi:10.1038/s41579-020-0364-5

41.   Moon, K. R. *et al.* Visualizing structure and transitions in high-dimensional biological data.
Nat. Biotechnol. (2019). doi:10.1038/s41587-019-0336-3

42.   van Vuuren, D. P. *et al.* The representative concentration pathways: An overview. *Clim.*
Change (2011). doi:10.1007/s10584-011-0148-z

43.   Marinov, I. *et al.* North-South asymmetry in the modeled phytoplankton community
response to climate change over the 21st century. *Global Biogeochem. Cycles* (2013).
doi:10.1002/2013GB004599

44.   Polovina, J. J., Dunne, J. P., Woodworth, P. A. & Howell, E. A. Projected expansion of the
subtropical biome and contraction of the temperate and equatorial upwelling biomes in the
North Pacific under global warming. *ICES J. Mar. Sci.* (2011). doi:10.1093/icesjms/fsq198

45.   Flombaum, P., Wang, W. L., Primeau, F. W. & Martiny, A. C. Global picophytoplankton niche
partitioning predicts overall positive response to ocean warming. *Nat. Geosci.* (2020).
doi:10.1038/s41561-019-0524-2

46.   Iudicone, D. Some may like it hot. *Nature Geoscience* (2020). doi:10.1038/s41561-020-0535-
z

47.   Gorsky, G. *et al.* Expanding Tara Oceans Protocols for Underway, Ecosystemic Sampling of
the Ocean-Atmosphere Interface During Tara Pacific Expedition (2016–2018). *Front. Mar.*
Sci. (2019). doi:10.3389/fmars.2019.00750

48.   Istace, B. *et al.* de novo assembly and population genomic survey of natural yeast isolates
with the Oxford Nanopore MinION sequencer. *Gigascience* (2017).
doi:10.1093/gigascience/giw018

49.   Busseni, G. *et al.* Meta-Omics Reveals Genetic Flexibility of Diatom Nitrogen Transporters in
Response to Environmental Changes. *Mol. Biol. Evol.* (2019). doi:10.1093/molbev/msz157

50.   Grand, M. M. *et al.* Developing autonomous observing systems for micronutrient trace
metals. *Frontiers in Marine Science* (2019). doi:10.3389/fmars.2019.00035

Acknowledgments

406 PF was supported by a CFR doctoral fellowship and the *NEOGEN impulsion* grant from the
407 Direction de la recherche fondamentale (DRF) of the CEA. We thank the LSCE (Laboratoire
408 des sciences du climat et de l'environnement, CEA) for providing Earth System Models
409 outputs, Tilla Roy for preparation of the data, LAGE (Laboratoire d'Analyses Génomiques

des Eucaryotes, CEA) members for stimulating discussions on this project, Mahendra Mariadassou, Sakina Dorothée Ayata and Bruno Hay Mele for discussions on statistics and climate envelop models, Laurent Bopp for initial discussions on this project and on climate models, Tom Delmont for providing Metagenome-Assembled Genomes data and Noan Le Bescot (TernogDesign) for artwork on Figures. We thank all members of the *Tara* Oceans consortium for maintaining a creative environment and for their constructive criticism. *Tara* Oceans would not exist without the *Tara* Ocean Foundation and the continuous support of 23 institutes (https://oceans.taraexpeditions.org/).

This article is contribution number XX of *Tara* Oceans.

Competing interests

The authors declare no competing interests.

Author contributions

PF, OJ and MG conceived the study. MV wrote the bias correction algorithm. PF computed the results, compiled and analyzed the data. PF wrote the initial draft of the paper. JL, OJ and MG conducted a preliminary study. PF, OJ, MG, MV, DI, and PW discussed the results and contributed to write the paper.

Online content

Supplementary information, additional references, source data and codes are available at www.doi.xx.com/

Materials and methods

Genomic provinces of plankton

Environmental niches are computed for trans-kingdom plankton genomic provinces from Richter et al.[12]. They consist of the clustering of metagenomic dissimilarity matrices from 6 available size fractions with sufficient metagenomic data from the *Tara* Oceans dataset. The six size fractions (0-0.2, 0.22-3, 0.8-5, 5-20, 20-180 and 180-2000 μm) represent major plankton groups. Two large size classes (180-2000 μm and 20-180 μm) are enriched in zooplankton dominated by arthropods (mainly copepods) and cnidarians. They are expected to directly depend on smaller eukaryotes as they feed on them. Size classes 5-20 μm and 0.8-5 μm are enriched in smaller eukaryotic algae, such as dynophytes (5-20 μm),

pelagophytes and haptophytes (0.8-5 μm). The distribution of these photoautotrophs presumably depends on nutrient availability. Finally, size classes 0.22-3 μm and 0-0.2 μm are respectively enriched in bacteria and viruses. Bacteria are characterized by a wide range of trophisms including autotrophy (cyanobacteria), mixotrophy and heterotrophy, while viruses are mainly parasites. Within each size fraction (from large to small), there are respectively 8, 8, 11, 6, 6 and 8 (48 in total) provinces defined in Richter et al.[12] formed by *Tara* Oceans metagenomes (644 metagenomes sampled either at the surface (SUR) or at the Deep Chlorophyll Maximum (DCM) across 102 sites). The clustering of individual size fractions is independent.

Genome signature of the provinces

We analyzed the distribution of 713 eukaryotic and 523 prokaryotic genomes[37,38] within the genomic provinces. These genomes are Metagenome-Assembled Genomes (MAGs) obtained from *Tara* Oceans metagenomes. For each size class, we select MAGs that are present (according to a criteria defined in Delmont et al.[38]) in at least 5 samples. We computed an index of presence enrichment of MAGs within provinces as the Jaccard index[51], defined as follows:

$$J = \frac{M_{11}}{M_{11} + M_{01} + M_{10}}$$

M_{11} is the number of samples where the MAG is present and match a sample of the province. M_{01} and M_{10} are respectively the number of samples where the MAG is not present in a sample of the province and inversely. A MAG is considered to be signature of a province if the Jaccard index is superior to 0.5 with this province and inferior to 0.1 for all other provinces of the given size class (Fig. 1 and Supplementary Fig. 3).

World Ocean Atlas data

Physicochemical parameters proposed to have an impact on plankton genomic provinces[12] are used to define environmental niches: sea surface temperature (SST), salinity (Sal); dissolved silica (Si), nitrate (NO_3), phosphate (PO_4), iron (Fe), and a seasonality index of nitrate (SI NO_3). With the exception of Fe and SI NO_3, these parameters are extracted from the gridded World Ocean Atlas 2013 (WOA13)[39]. Climatological Fe fields are provided by the biogeochemical model PISCES-v2[52]. The seasonality index of nitrate is defined as the

range of nitrate concentration in one grid cell divided by the maximum range encountered

in WOA13 at the *Tara* Oceans sampling stations. All parameters are co-located with the

corresponding stations and extracted at the month corresponding to the *Tara* Oceans

sampling. To compensate for missing physicochemical samples in the *Tara* Oceans *in situ*

data set, climatological data (WOA) are preferred. The correlation between in situ samples

and corresponding values extracted from WOA are high (r^2: SST: 0.96, Sal: 0.83, Si: 0.97,

NO_3: 0.83, PO_4: 0.89). In the absence of corresponding WOA data, a search is done within 2°

around the sampling location and values found within this square are averaged.

Nutrients, such as NO_3 and PO_4, display a strong collinearity when averaged over the global

ocean (correlation of 0.95 in WOA13) which could complicate disentangling their

respective contributions to niche definition. However, observations and experimental data

allow identification of limiting nutrients at regional scale characterized by specific plankton

communities[53]. The projection of niches into future climate would yield spurious results

when the present-day collinearity is not maintained[54] but there is up to now no evidence

for large scale changes in global nutrient stoichiometry[55].

Earth System Models and bias correction

Outputs from 6 Earth System Models (ESM) (Supplementary Table 2) are used to project

environmental niches under greenhouse gas emission scenario RCP8.5[42]. Environmental

drivers are extracted for present day (2006-2015) and end of century (2090-2099)

conditions for each model and the multi-model mean is computed. A bias correction

method, the Cumulative Distribution Function transform, CDFt[56], is applied to adjust the

distributions of SST, Sal, Si, NO_3 and PO_4 of the multi-model mean to the WOA database.

CDFt is based on a quantile mapping (QM) approach to reduce the bias between modeled

and observed data, while accounting for climate change. Therefore, CDFt does not rely on

the stationarity hypothesis and present and future distributions can be different. CDFt is

applied on the global fields of the mean model simulations. By construction, CDFt preserves

the ranks of the simulations to be corrected. Thus, the spatial structures of the model fields

are preserved.

Environmental niche models: training, validation and projections

Provinces with similar metagenomic content are retrieved from Richter et al.[12]. From a total of 48 initial provinces, 10 provinces are removed either because they are represented by too few samples (7 out of 10) or they are found in environments not resolved by ESMs (e.g. lagoons of Pacific Ocean islands, 3 out of 10). Four machine learning methods are applied to compute environmental niches for each of the 38 provinces: Gradient Boosting Machine (gbm)[57], Random Forest (rf)[58], fully connected Neural Networks (nn)[59] and Generalized Additive Models (gam)[60]. Hyper parameters of each technique (except gam) are optimized. These are (1) for gbm, the interaction depth (1, 3 and 5), learning rate (0.01, 0.001) and the minimum number of observations in a tree node (1 to 10); (2) for rf, the number of trees (100 to 900 with step 200 and 1000 to 9000 with step 2000) and the number of parameters used for each tree (1 to 7); (3) for nn, the number of layers of the network (1 to 10) and the decay (1.10^{-4} to 9.10^{-4} and 1.10^{-5} to 9.10^{-5}). For gam the number of splines is set to 3, respectively 2 only when not enough points are available (for fraction 0-0.2, 65 points). R packages gbm (2.1.3), randomForest (4.6.14), mgcv (1.8.16) and nnet (7.3.12) are used for gbm, rf, nn and gam models.

To define the best combination of hyper parameters for each model, we perform 30 random cross-validations by training the model on 75% (85 % for gbm and gam) of the dataset randomly sampled and by calculating the Area Under the Curve[61] (AUC) on the 25% (15 % for gbm and gam) remaining points of the dataset. The best combination of hyper parameters is the one for which the mean AUC over the 30 cross-validation is the highest. A model is considered valid if at least 3 out of the 4 techniques have a mean AUC superior to 0.65, which is the case for 27 out of the 38 provinces (Supplementary Fig. 2a). A climatic annotation is given to the 27 validated niches (Supplementary Table 1). Final models are trained on the full dataset and only the techniques that have a mean AUC higher than 0.65 are considered to make the projections. The majority (23) of the 27 validated niches is validated by all four models and 4 by only 3 models. Relative influences of each parameter in defining environmental niches are calculated using the feature_importance function from the DALEX R package[62] for all four statistical methods (Supplementary Fig. 19a). To evaluate the consistency and coherence of environmental niche models, we first make global projections on the 2006-13 WOA2013 climatology. Projections are consistent with sampling regions for provinces encompassing vast oceanic areas. For example, the

genomic province sampled in temperate Atlantic regions of size fraction 180-2000 μm is

projected to be present in the north and south temperate Atlantic but also other temperate

regions (Supplementary Fig. 4). For model training and projections, physicochemical

variables are scaled to have a mean of 0 and a variance of 1. For this scaling, the mean and

standard deviation of each WOA13 variable (+ PISCES-v2 Fe) co-localized with *Tara* Oceans

stations with a value available are used. This standardization procedure allows for better

performance of nn models. Finally, as statistical models often disagree on projection sets

whereas they give similar predictions on the training set (Supplementary Fig. 5, 6), we use

the ensemble model approach for global-scale projections of provinces[36] *i.e.* the mean

projections of the validated machine learning techniques.

Combined size classes provinces and ocean partitioning comparisons

To combine all size classes' provinces, we use the PHATE algorithm[41,63] from R package

phateR. This algorithm allows visualization of high dimensional data in the requested

number of dimensions while best preserving the global data structure[63]. We choose to train

PHATE separately on WOA13 projections and present day and end of century projections

including presence probabilities of *non dominant* provinces. We use 3 dimensions and set

hyper parameter k-nearest neighbors (knn) and decay respectively to 1000 for WOA13 and

2000 for model data as in this case there are twice as many points. The hyper parameter

knn reflects the degree to which the mapping of PHATE from high to low dimensionality

should respect the global features of the data. We argue that 1000 and 2000 are good

choices as it will be sufficient to have a highly connected graph, conserve global structure,

allow visualization of structures of the size of the provinces (mean number of points in a

province: 4867) and have a reasonable computational time. Decay is set to 20 in both cases.

Then we cluster the resulting distance matrix using the k-medoïds algorithm[64] and the

silhouette average width criteria[65] is used as an indicator of good fit. The silhouette

criterion is maximal for 2, 3 and 4 clusters and 2 peaks are found at 7 and 14 clusters (the

peak at 7 is slightly less high than the one at 14, data not shown). We choose to present the

4 and 7 clusters geographical patterns as they seem more relevant with respect to the

resolutions of our environmental datasets (WOA13 and climate models). We compare the

three polar clusters of the 7 cluster geographical patterns with Antarctic Circumpolar

Currents fronts[66] by overlying them on the map (red lines Fig. 2h).

To visualize the global biogeography structure, the resulting 3 vectors of PHATE are plotted using an RGB color code. Each coordinate of each vector is respectively assigned to a given degree of color component between 0 and 255 (8 bits red, green or blue) using the following formula (Fig. 1 g,h; Fig 2 e,f; Supplementary Fig. 7):

$$C\_col(i) = \frac{C(i) - \min\{C(1), C(2), C(3)\}}{\max(C(1), C(2), C(3)) - \min\{C(1), C(2), C(3)\}} * 255$$

$C(i)$ is the ith component of the PHATE axes. Respectively, components 1, 2 and 3 are assigned to red, green and blue.

To compare the six size fraction provinces, the combined size class with existing biogeochemical partitions of the oceans[10,11] and with each other, we use the adjusted rand index[67] (Supplementary Fig. 7-9) and overlay their masks above our partitions. In this case, presence probabilities of *dominant* provinces are not used anymore. Instead, each ocean grid point is assigned to the *dominant* provinces or the phate clusters.

Bray-Curtis dissimilarity index

Climate change impact on global projections is calculated at each grid point as the Bray-Curtis dissimilarity index[68,69] defined as follows:

$$BC = \frac{\sum_n |P_n^{future} - P_n^{present}|}{\sum_n |P_n^{future} + P_n^{present}|}$$

Where ($P_n^{present}$ and P_n^{future}) are respectively the probability of presence of the province n in present day and at the end of the century. Only the probabilities of *dominant* provinces are non-null and all others are set to zero. The mask of main fisheries[70] (chosen as the first 4 deciles) and Exclusive Economical Zones[71] is overlaid on the Bray-Curtis map.

Change in province assemblages

A province assemblage is defined as the assemblage of *dominant* provinces of each size fraction at a given grid point of the considered ocean. We consider two criteria of change in province assemblage between present day and end of the century conditions. The first one, more straightforward and less stringent, considers that a province assemblage occurs when a change of *dominant* province is found in at least one size fraction. In a more stringent way, a change of assemblage is considered significant for $BC > \frac{1}{6}$ (previous section). This threshold corresponds to an idealized case where each *dominant* province

has a probability of one and a change of *dominant* province is found in only one size

fraction. For example, the *dominant* province assemblage goes from vector

(F5,E6,D3,C8,B7,A7) (with the size fractions in decreasing order) corresponding to all

temperate provinces to vector (F8,E6,D3,C8,B7,A7). This example corresponds to the

replacement of the temperate province of size fraction 180-2000 μm (F5) by the tropico-

equatorial province (F8). This criterion allows us to discard assemblage changes for which

the changes in probability of presence of *dominant* provinces are very low. With this

criterion, only on a small oceanic area is found to have no changes of assemblage (Fig. 4c

light blue zones).

Centroids and migration shifts

The centroid of each province is defined as the average latitude and longitude for which the

probability of presence is superior to 0.5 and weighted by both the probability of presence

at each grid point and the grid cell area. It is calculated for both present day conditions and

end of the century conditions. The migration shift is calculated as the distance between the

present day and the end of the century centroids considering the earth as a perfect sphere

of radius 6371 km. For consistency (*i.e.* avoid long distance aberrant shifts), it is only

calculated for provinces with an area of dominance larger than 10^6 km^2 in the given basin.

Driver analysis

To assess the relative importance of each driver in province changes, the methodology

from Barton et al.[24] is adopted. For a set N of n provinces (individual provinces or all

provinces together), the probability of presence of each province is recalculated for present

day conditions except for driver d (from the set of drivers D) for which end of the century

condition is used ($P_n^{future\ for\ dth\ driver\ only}$). The set of driver D can be either all drivers

(Fig. 5a,c) or all drivers except SST (Fig. 5b). The relative importance of driver d at a given

grid point for the set N of provinces is computed as follow:

612
$$RI(d) = \frac{\sum_{n \in N} \left| P_n^{future\ for\ dth\ driver\ only} - P_n^{present} \right|}{\sum_{d \in D} \sum_{n \in N} \left| P_n^{future\ for\ dth\ driver\ only} - P_n^{present} \right|}$$

613 $RI(d)$ is computed at grid cells where $BC > \frac{1}{6}$, BC being calculated with either the set of all

614 drivers (Fig. 5a,c) or all drivers except SST (Fig. 5b). When RI(d) is calculated for individual

615 provinces (Fig. 5c and Supplementary Fig. 16d,e), it is computed only at grid cells where

616    $BC > \frac{1}{6}$ and the concerned province is either *dominant* in present day and/or end of

617   century conditions.

References

619   51.   Jaccard, P. Distribution comparée de la flore alpine dans quelques régions des Alpes
620   occidentales et orientales. *Bull. la Murithienne* (1902).

621   52.   Aumont, O., Ethé, C., Tagliabue, A., Bopp, L. & Gehlen, M. PISCES-v2: An ocean biogeochemical
622   model for carbon and ecosystem studies. *Geosci. Model Dev.* (2015). doi:10.5194/gmd-8-
623   2465-2015

624   53.   Moore, C. M. *et al.* Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*
625   (2013). doi:10.1038/ngeo1765

626   54.   Brun, P., Kiørboe, T., Licandro, P. & Payne, M. R. The predictive skill of species distribution
627   models for plankton in a changing climate. *Glob. Chang. Biol.* (2016). doi:10.1111/gcb.13274

628   55.   Redfield, A. C. On the Proportions of Organic Derivatives in Sea Water and Their Relation to
629   the Composition of Plankton. in *James Johnstone Memorial Volume* 1767–192 (Liverpool
630   Univ. Press, Liverpool, U.K., 1934).

631   56.   Michelangeli, P. A., Vrac, M. & Loukos, H. Probabilistic downscaling approaches: Application
632   to   wind   cumulative   distribution   functions.   *Geophys.   Res.   Lett.*   (2009).
633   doi:10.1029/2009GL038401

634   57.   Ridgeway, G. gbm: Generalized Boosted Regression Models. *R Packag. version 1.6-3.1* (2010).

635   58.   Breiman, L. & Cutler, A. Breiman and Cutler's random forests for classification and
636   regression. *Packag. 'randomForest'* (2012). doi:10.5244/C.22.54

637   59.   Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S Fourth edition by. World*
638   (2002). doi:10.2307/2685660

639   60.   Wood, S. N. Stable and efficient multiple smoothing parameter estimation for generalized
640   additive models. *J. Am. Stat. Assoc.* (2004). doi:10.1198/016214504000000980

641   61.   Fawcett,   T.   An   introduction   to   ROC   analysis.   *Pattern   Recognit.   Lett.*   (2006).
642   doi:10.1016/j.patrec.2005.10.010

643   62.   Biecek, P. DALEX: explainers for complex predictive models. *J. Mach. Learn. Res.* **19**, 1–5
644   (2018).

645   63.   Vallejos, C. A. Exploring a world of a thousand dimensions. *Nature Biotechnology* (2019).
646   doi:10.1038/s41587-019-0330-9

647   64.   L., K. & P., R. Clustering by means of Medoids. in *Statistical Data Analysis Based on the L1*
648   *Norm and Related Methods* (1987).

65. Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and validation of cluster
analysis. *J. Comput. Appl. Math.* (1987). doi:10.1016/0377-0427(87)90125-7

66. Orsi, A. H., Whitworth, T. & Nowlin, W. D. On the meridional extent and fronts of the Antarctic
Circumpolar Current. *Deep. Res. Part I* (1995). doi:10.1016/0967-0637(95)00021-W

67. Hubert, L. & Arabie, P. Comparing partitions. *J. Classif.* (1985). doi:10.1007/BF01908075

68. Somerfield, P. J. Identification of the Bray-Curtis similarity index: Comment on Yoshioka
(2008). *Marine Ecology Progress Series* (2008). doi:10.3354/meps07841

69. Bloom, S. Similarity Indices in Community Studies: Potential Pitfalls. *Mar. Ecol. Prog. Ser.*
(1981). doi:10.3354/meps005125

70. Watson, R. A. A database of global marine commercial, small-scale, illegal and unreported
fisheries catch 1950-2014. *Sci. Data* (2017). doi:10.1038/sdata.2017.39

71. Flanders Marine Institute (2018). Maritime Boundaries Geodatabase: Maritime Boundaries
and Exclusive Economic Zones (200NM), version 10. (2018).
doi:https://doi.org/10.14284/313.

Fig. 1 | Signature genomes of provinces of eukaryotes enriched size classes. For each plankton size class, indexes of presence enrichment for 713 genomes of eukaryotic plankton[38] in corresponding provinces are clustered and represented in a color scale. Signature genomes (see *Methods*) are found for almost all provinces, their number and taxonomies are summarized (detailed list in Supplementary Table 6).

Fig. 2 | Global biogeographies of size-structured plankton provinces projected on WOA2013 dataset. (**a**) Metazoans enriched (180-2000 μm) (**b**) Small metazoans enriched (20-180 μm) (**c**) Small eukaryotes enriched (5-20 μm) (**d**) Small eukaryotes enriched (0.8-5 μm) (**e**) Bacteria enriched (0.22-3 μm) (**f**) Viruses enriched (0-0.2 μm). (**a-f**) Dotted areas represent uncertainty areas where the delta of presence probability of the dominant province and an other (from the same size fraction) is inferior to 0.5. (**g**) Combined size class biogeography using PHATE algorithm

partitioned in 4 clusters. Areas of uncertainty are highlighted with dotted lines. (**h**) Combined size

class biogeography using PHATE algorithm partitioned in 7 clusters overlaid with Antarctic

Circumpolar Current boundaries (red). Simple biogeographies are observed in large size fractions

(>20 µm) with a partitioning in three major oceanic areas: tropico-equatorial, temperate and polar.

More complex geographic patterns and patchiness are observed in smaller size fractions with the

distinction of pacific equatorial provinces and provinces associated to oligotrophic tropical gyres.

Fig. 3 | **Global biogeography of small metazoans enriched size fraction (180-2000 μm), protists enriched (0.8-5 μm) and combined size classes in modeled present day (a, c, e) and end of the century (b, d, f).** (a-d) The dominant niche *i.e.* the one predicted to have the highest probability of presence is represented at each grid point of the map. The color transparency is the probability of presence of the dominant niche. A simple biogeography is observed for size fraction 180-2000 μm (**a, b**) with a polar niche, a temperate and a tropico-equatorial niche. Biogeography of size fraction 0.8-5 μm size fraction is more complex and patchy with several temperate and tropical niches (**c, d**). Biogeography of large size plankton and small size plankton are therefore decoupled. Climate change impacts tropical niches which are expanding towards the poles in both size fractions with a more complex behavior in size fraction 0.8-5 μm (**b** and **d**). (**e, f**) Present and end of century combined size classes using PHATE algorithm.

Size fraction (μm)	Present day covered area (%)	End of century covered area (%)	% area with a change of dominant province	Most frequent transition	%	2[nd] most frequent transition	%
180-2000	74	74	13	temperate->tropico-equatorial (5->8)	67	polar->temperate (1->5)	14
20-180	78	77	12	temperate->tropico-equatorial (6->5)	67	polar->temperate (1->6)	29
5-20	45	49	22	temperate -> equatorial (3->4)	47	equatorial -> tropical (4->6)	25
0.8-5	56	59	31	equatorial -> tropical (9->11)	22	temperate -> equatorial (8->9)	15
0.22-3	60	61	15	temperate-> tropical (7->5)	22	polar -> temperate (8->6)	16
0-0.2	64	66	16	equatorial -> tropical (6->3)	32	temperate -> equatorial (8->6)	31
Total			60				

697 **Table 1 | Global statistics of covered areas and provinces' changes and transitions.** From 12 %

698 to 31% of the total covered area is estimated to be replaced by a different province at the end of the

699 century compared to present day depending on the size fraction. In total, considering all size

700 fractions this represents 60 % of the total covered area with at least one predicted change of

701 dominant province across the six size fractions.

Fig. 4 | (a) Bray-Curtis dissimilarity index map comparing present day with end of the century projections of dominant provinces. Maps of trans-kingdom assemblages reorganization of dominant provinces (b) and with a criterion of significance (c). (a) Bray-Curtis dissimilarity index is calculated by integrating all the dominant provinces presence probabilities over the six size fraction. Most important changes appear in subtropical, temperate and subpolar regions. These changes are due to the displacement of tropical and temperate provinces towards the pole but also the geographical decoupling between large and small size plankton. The mean change in niche dissimilarity index is 0.25. (b) An assemblage is the combined projected presence of the dominant province of each size class. Assemblage reorganization (present day versus end of the century) is either mapped on all considered oceans or with a criterion on the Bray-Curtis dissimilarity index (BC>1/6, see *Materials and Methods*) (c). Depending on the criterion from 60.1 % (b, dark blue) to 48.7 % (c, dark blue) of the oceanic area is projected to change of assemblage. New assemblage are expected to appear in 2090 (purple+blue) whereas some 2006 specific assemblages are projected to disappear (red+blue). New assemblages as well as lost assemblages are mostly found in temperate, subtropical and tropical regions where most of the rearrangements are projected.

Fig. 5 | Map of most impacting drivers on dominant province changes (a), most impacting driver without considering temperature change (b) and relative importance of the drivers in the different size fractions (c). (**a**) Temperature appears as the top impacting driver on the majority of the projected ocean with a significant change of province (Fig. 4). (**b**) Salinity and dissolved phosphate are found to be the second and third driver of province reorganization notably at tropical and subpolar latitudes. Note the importance of nitrate at temperate southern latitudes. (**c**) Temperature is found to be the most important driver for all size classes but has a more important impact in large size classes (>20 μm). Nutrients have on average a relative more important impact in small size classes in driving provinces reorganization.

729
Supplementary Fig. 1 | Study pipeline. Metagenomic data from the 2009-2013 *Tara Oceans*
expedition and *in situ* measurements of physicochemical variables (*World Ocean Atlas 2013*,
WOA13)[1] are combined to define environmental niches at the plankton community level across 6
size fractions of the plankton realm. Bias corrected outputs from a mean model of 6 Earth System
Models[2-7] and WOA13 data are then used to project global plankton provinces for present day
conditions and end of the century conditions under a high emission climate change scenario
(RCP8.5)[8]. Physico-chemical variables are Sea Surface Temperature (SST), Salinity (Sal), Dissolved
silica (Si), Nitrate (NO_3), Phosphate (PO4), Iron (Fe) and a seasonality index of nitrate (SI NO_3).

Supplementary Fig. 2 | (a) Barplot of mean AUC over 30-fold cross validation process of the 38 initial metagenomic provinces. (b,c) Map of validated and non validated stations across the six size fractions in surface samples (b) and DCM samples (c). (a) Mean AUC (Area Under the receiver operating Characteristic)[9] is plotted for the best hyperparameter combination of the 4 machine learning techniques used for each of the 38 metagenomic provinces. General Additive Models[10] (gam) are shown in red (no optimization), Gradient Boosting Machines[11] (gbm) in green, fully connected Neural Networks[12] (nn) in blue and Random Forest[13] models in purple (rf). A star for valid models is drawn at the top of each considered valid model. A model is validated when at least 3 out of the four models have a mean AUC superior to 0.65. A valid model is found for 27 out of 38 initial provinces. Out of the 27 validated models 4 are not valid for the gbm method. (b-c) For each Tara sample present in the dataset at surface (b) or Deep Chlorophyll Maximum (DCM) (c) and for each size fraction, the filter (one size fraction at one location and one depth) belongs either to a validated niche (green), a non validated niche (red), an excluded filter (sea Materials & Methods) (black) or no data is available (white). Non validated niches represent only 11 % of filters present in the dataset (66 out of 595).

Supplementary Figure 3 | Signature genomes of provinces of the prokaryotes enriched size.
Indexes of presence enrichment[14] for 523 genomes of prokaryotic plankton[15] in corresponding
provinces are clustered and represented in a color scale. Signature genomes (see *Methods*) are
found for almost all provinces, their number and taxonomies are summarized (detailed list in
Supplementary Table 6).

Supplementary Fig. 4 | Sampling and projection maps on WOA13 climatological data of two
example provinces from size fraction 180-2000 μm. (**a**) Sampling map of province F5. (**b**)
Province F8. (**c**) Projection map of province F5 on WOA13[1]. (**d**) Province F8. Qualitatively,
projection maps are coherent with sampling maps of the two provinces with the highest probability
of presence projected in the sampling regions. Other presence zones are also predicted by the
projection. Sampling of these zones might be interesting to confirm our approach and projections
such as South of Australia where a high probability of presence for province F5 is predicted.

Supplementary Fig. 5 | Pairwise Pearson correlation coefficient heat maps between outputs of the 4 machine learning models. rf: Random Forest, gbm: gradient boosting machines, gam: general additive models, nn: neural networks. In rows are the provinces of the different size fractions. In columns are the pairwise comparisons of each machine learning technique. (a) Training set outputs. (b) WOA13 average data (except for Iron, PISCES-v2[16] is used) projection outputs. (c) Present day data projection outputs (bias corrected mean model of 6 Earth System Models). (d) End of the century projection outputs. On the training set, models are in agreement with most of the correlation coefficients superior to 0.9. A drop of correlation is observed for modeled data (c, d) especially in small size fractions. This shows modeled data are more distant from the training set than WOA data. Random Forest and Gradient Boosting Machine are in very good agreement (first columns) which could be expected as they are both based on multiple decision trees.

Supplementary Fig. 6 | Probability range map (WOA data) of province F5 of size fraction 180-
2000 μm. At each point of the grid, the maximum delta probability of presence between the 4
machine learning projections is calculated. In black are the zones where two models completely
disagree: one model predicts presence with certainty whereas the other predicts absence with
certainty. Disagreement appears mostly in uncertain presence areas (Supplementary Fig. 3c)
whereas models are generally in good agreement in absence areas (blue zones).

 Supplementary Fig. 7 | Three existing oceanic partitions overlaid on top of plankton

provinces for the six fractions and combined size fractions. The three oceanic partitions are

Reygondeau et al. Biogeochemical Provinces (BGCP)[17]; Reygondeau et al. Biomes[17]; Fay and

McKingley Biomes[18]. Each partitioning mask is overlaid in the above order on top of plankton

provinces for the six size fractions. (**a-c**) 180-2000 µm (**d-f**) 20-180 µm (**g-i**) 5-20 µm (**j-l**) 0.8-5

µm(**m-o**) 0.22-3 µm and (**p-r**) 0-0.2 µm. Above all maps, the adjusted rand index (ARI), an index

used to compare partitions, comparing mapped biogeographies with the black lines mask is shown.

Supplementary Fig. 8 | Three existing oceanic partitions overlaid on top of plankton provinces for the combined size fractions. The three oceanic partitions are Reygondeau et al.[17] Biogeochemical Provinces (BGCP); Reygondeau et al. Biomes[17]; Fay and McKingley Biomes[18]. Each partitioning mask is overlaid in the above order on top of size fraction plankton provinces combined by the PHATE algorithm in (**a-c**) 4 clusters and (**d-f**) 7 clusters.

802   **Supplementary Fig. 9 | Pairwise comparisons of ocean partitions based on plankton**
803   **provinces and existing biogeochemical partitions.** The three oceanic partitions are Reygondeau
804   et al.[17] Biogeochemical Provinces (BGCP); Reygondeau et al. Biomes[17]; Fay and McKingley Biomes[18].

Supplementary Fig. 10 | Taylor diagram exhibiting statistical comparison of WOA 2013 observations and present day ESM model mean of the different drivers. Each variable is centered and scaled according to the mean and standard deviation of the observed variable (black circle point at standard deviation 1 on the x-axis). Dark color points are the ESM model mean without Cumulative Distribution Function transform (CDFt[19]) bias correction. They are to be compared with light color points for which the bias correction is performed. Overall, good spatial correlations are found between the model mean and the observations. CDFt bias correction performs well by bringing standard deviations of the model to the observed standard deviations without decreasing correlations.

Supplementary Fig. 11 | Differences in drivers' intensity (2090/99-2006/15) in the bias corrected ESM model-mean under RCP8.5. (**a**) Sea Surface Temperature (SST). (**b**) Dissolved silica. (**c**) Nitrate. (**d**) Iron. (**e**) Phosphate. (**f**) Salinity (Sal). (**g**) Seasonality Index of Nitrate (SI NO_3). Note that the scale for dissolved silica variations is restricted to visualize small variations. Regions for which out of range values (*i.e.* inferior to the minimum or superior to the maximum found in

821   2006-15) are reached at the end of the century are highlighted with small stars reflecting high

822   uncertainty zones for machine learning approaches.

Supplementary Fig. 12 | Distribution of deltas between future temperature at each sampling site minus either the mean or maximum temperature within their contemporary genomic province. For most of the sites and across size fractions the future temperature projected by the bias adjusted ESM ensemble model is higher than both the maximum and mean contemporary temperature of their genomic province.

Supplementary Fig. 13 | Global geographical patterns for 20-180, 5-20, 0.22-3, 0-0.2 μm

plankton size fractions in present day (a, c, e , g) and at the end of the century (b, d, f, h). The

dominant province *i.e.* the one predicted to have the highest probability of presence is represented

at each grid point of the map. The color transparency is the probability of presence of the dominant

province. Expansion of tropical provinces and shrinkage of temperate communities is consistently

projected in all size fractions.

Supplementary Fig. 14 | Projected migration shifts of the 27 provinces between present day and end of the century. Predicted migration shifts are presented in 5 major ocean basins: (a) North Atlantic (b) North Pacific (c) South Atlantic (d) South Pacific and (e) Indian Ocean. 96 % of migration shifts (larger than 200 km) are oriented towards the pole. Mean shift is 641 km (76 ± 79 km.dec[-1]) and median shift is 394 km (47 km.dec[-1]). Few provinces are projected to shift more than thousands of kilometers toward suitable environmental conditions with a maximum shift of 4325 km.

Supplementary Fig. 15 | (a) Projected migration shifts of provinces on the world map. (b)
Latitudinal shift distribution (red bars) and longitudinal shift distribution (blue bars). (a)
Migration shifts are represented as arrows pointing at the end of century centroid. Arrows are
colored according to the size fraction. Some shifts seem to correlate with each other (exemplified
with circled arrows). For instance, parallel shifts are projected in the southern pacific equatorial
communities of size fractions 0.8-5 μm and 5-20 μm (blue and green circled arrows). All non-
poleward arrows belong to small size classes (<20μm) showing differential responses to climate
change depending on the size class. (**b**) Some longitudinal shifts are more important than
latitudinal shifts with 14 longitudinal shifts superior to 1000 kms. Mean longitudinal shift (around
500 kms) is significantly higher (Student t-test p-value<0.01) than mean latitudinal shift (around
290 kms) while medians (longitudinal 190 kms vs latitudinal 230 kms) are not significantly
different (Wilcoxon test).

Supplementary Fig. 16 | Projection maps of province F5 of size fraction 180-2000 μm in present day (a) and at the end of the century (b). At each grid point, the probability of presence of the province is computed as the average of the predicted probability of each of the four machine learning techniques (gbm, nn, rf and gam). Red color indicates a high probability of presence. (**a**)The projected province corresponds to the sampled province (North Atlantic and South Atlantic) but several other places have high probabilities of presence such as South Australia where no sampling is available. (**b**) At the end of the century, the province is projected to reduce significantly in size. (**c**) Delta probability of presence map (2006/15 – 2090/99) and core range shift in the 5 major oceanic basins of province F5 of size fraction 180-2000 μm. In all the basins, the centroid of the province is projected to migrate poleward. (**d**) Main drivers associated with the projected changes. Changes are mainly driven by sea surface temperature (55%) followed by salinity (14%). (**e**) Main drivers associated with the shrinkage of the equatorial cluster C9 of size fraction 0.8-5.

Considering only latitudes between the two tropics, changes are mainly driven by decreases in PO_4

(24 %) in addition to SST (27 %) (overall 34 % STT and 20 % PO_4).

Supplementary Fig. 17 | Probabilistic covered areas of the provinces projected in present
day and at the end of the century. (**a**) 180-2000 μm (**b**) 20-180 μm (**c**) 5-20 μm (**d**) 5-20 μm (**e**)
0.8-5 μm (**f**) 0-0.2 μm. The covered area by a province is defined as the area in which this province
is dominant and weighted by its probability of presence at each point and grid cell area. Areas not
covered by the provinces are represented in white.
877

878 **Supplementary Fig. 18 | Bray-Curtis dissimilarity index and assemblage changes maps**
879 **comparing present day with end of the century projections of *dominant* provinces in** (**a**)
880 principal fisheries (4 last deciles[20]) and (**b**) Exclusive Economic Zones[21]. Assemblage changes in (**c**)
881 Principal fisheries (**d**) Exclusive Economic Zones. Assemblage changes in (**e**) Principal fisheries (**f**)
882 Exclusive Economic Zones with a Bray-Curtis dissimilarity index superior to 1/6.
883

Supplementary Fig. 20 | Maps of environmental drivers' relative influences in driving province reorganization. Each environmental parameter relative influence is quantified by considering only the variation of each parameter individually between present day and end of the century as defined in Barton et al.[23] (*Materials and Methods*) and where a significant change is projected. Importantly the mean impact of SST (51.5 %) is largely the highest. PO_4 is the second most impacting drivers (11.0 %). Contrary to supplementary Fig. 16, relative influence is calculated here by combining all provinces together. Therefore, mean relative influences slightly differ especially for dissolved silica (Si).

Model	Reference
CESM1-BGC	Gent et al., 2011
GFDL-ESM2G	Dunne et al., 2013
GFDL-ESM2M	Dunne et al., 2013
HadGEM2-ES	Collins et al., 2011
IPSL-CM5A-LR	Dufresne et al., 2013
IPSL-CM5A-MR	Dufresne et al., 2013
MPI-ESM-LR	Giorgetta et al., 2013
MPI-ESM-MR	Giorgetta et al., 2013
NorESM1-ME	Bentsen et al., 2013

905 **Supplementary Table 1 | Earth System models used to compute the mean model.**
906

Fraction	Province	Climatic annotation	Area 2006-15 (MKm^2)	Area 2090-99 (MKm^2)	Delta area (MKm^2 (%))
180-2000	F1	polar	82	73	-9 (-11%)
180-2000	F5	temperate	44	26	-18 (-41%)
180-2000	F8	tropico-equatorial	140	169	29 (+21%)
20-180	E1	polar	97	84	-13 (-13%)
20-180	E5	tropico-equatorial	119	145	+26 (+22%)
20-180	E6	temperate	65	49	15 (-23%)
5-20	D3	temperate	51	41	-10 (-20%)
5-20	D4	equatorial	46	37	-8 (-18%)
5-20	D6	tropical	65	97	32 (+48%)
0.8-5	C1	polar	71	64	-6 (-8%)
0.8-5	C3	subtropical	3,6	1,4	-2,2 (-61%)
0.8-5	C4	subtropical	6	19	+13 (+217%)
0.8-5	C8	temperate	33	15	-18 (-54%)
0.8-5	C9	equatorial	35	19	-16 (-45%)
0.8-5	C10	subtropical	3.8	7.4	+3,6 (+95%)
0.8-5	C11	tropical	49	86	37 (+75%)
0.22-3	B2	subtropical	5,1	6,7	+1,6 (+31 %)
0.22-3	B3	equatorial	33	32	-1 (-0.3%)
0.22-3	B5	tropical	88	102	+13 (+15%)
0.22-3	B6	temperate	30	35	+5 (+17%)
0.22-3	B7	temperate	8	2	-6 (-75%)
0.22-3	B8	polar	51	44	-7 (-14%)
0-0.2	A3	tropical	101	114	+13 (+13 %)
0-0.2	A4	subtropical	6	19	+13 (+216 %)
0-0.2	A6	equatorial	40	36	-4 (-10%)
0-0.2	A7	temperate	1,7	1	-0,7 (-41%)
0-0.2	A8	polar	81	67	-14 (-17%)

908 **Supplementary Table 2 | Genomic provinces climatic annotations and oceanic surfaces they**
909 **cover in present day (2006-15) and at the end of the century (2090-990).**

	Fraction (μm)	SST	Sal	Si	NO3	PO4	Fe	SI NO3
Niche definition	180-2000	27,7	16,8	10,1	10,3	11,4	14	9,7
	20-180	29,7	13,3	8,9	10,7	16,3	13,2	7,9
	5-20	21,5	14,2	8,4	12,7	13,7	12,6	17
	0,8-5	22,6	17,3	9,2	12,3	13,4	14,5	10,8
	0,22-3	23,9	13,7	10,5	13	12,5	14,2	12,2
	0-0,2	27,1	16,4	9,3	9,5	17,9	12,1	7,6
	all	25	15,5	9,5	11,6	14,2	13,6	10,8
Climate change	180-2000	75,2	8,4	2,5	2,6	4,1	5,8	1,4
	20-180	71,7	4,2	7,3	3,9	8,7	3,3	1,0
	5-20	57,9	6,1	4,4	6,6	9,0	6,5	1,0
	0,8-5	54,9	13,4	7,2	5,5	9,7	5,9	3,6
	0,22-3	42,6	9,9	13,5	6,4	9,1	15,2	3,3
	0-0,2	42,3	18,9	6,6	6,4	12,1	10,8	2,9
	all	51,5	10,3	8,7	6	11	8,7	3,8

Supplementary Table 3 | Summary table of relative importance of each environmental
driver in niche definition and in driving geographical reorganization in response to climate
change. Note that in both cases (niche definition and climate change), the row 'all' is not the mean
over the size fractions. In the case of niche definition, this is due to a different number of niches in
each size class. In the case of climate change, relative influence is either calculated for single
provinces at a given grid point then recalculated for individual size class or calculated for all
provinces together (row 'all').

References

1. Boyer, T. P. et al. WORLD OCEAN DATABASE 2013, NOAA Atlas NESDIS 72. Sydney Levitus, Ed.; Alexey Mishonoc, Tech. Ed. (2013). doi:10.7289/V5NZ85MT

2. Gent, P. R. et al. The community climate system model version 4. J. Clim. (2011). doi:10.1175/2011JCLI4083.1

3. Dunne, J. P. et al. GFDL's ESM2 global coupled climate-carbon earth system models. Part II: Carbon system formulation and baseline simulation characteristics. J. Clim. (2013). doi:10.1175/JCLI-D-12-00150.1

4. Collins, W. J. et al. Development and evaluation of an Earth-System model - HadGEM2. Geosci. Model Dev. (2011). doi:10.5194/gmd-4-1051-2011

5. Dufresne, J. L. et al. Climate change projections using the IPSL-CM5 Earth System Model: From CMIP3 to CMIP5. Clim. Dyn. (2013). doi:10.1007/s00382-012-1636-1

6. Giorgetta, M. A. et al. Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5. J. Adv. Model. Earth Syst. (2013). doi:10.1002/jame.20038

7. Bentsen, M. et al. The Norwegian Earth System Model, NorESM1-M – Part 1: Description and basic evaluation of the physical climate. Geosci. Model Dev. (2013). doi:10.5194/gmd-6-687-2013

8. van Vuuren, D. P. et al. The representative concentration pathways: An overview. Clim. Change (2011). doi:10.1007/s10584-011-0148-z

9. Fawcett, T. An introduction to ROC analysis. Pattern Recognit. Lett. (2006). doi:10.1016/j.patrec.2005.10.010

10. Wood, S. N. Stable and efficient multiple smoothing parameter estimation for generalized additive models. J. Am. Stat. Assoc. (2004). doi:10.1198/016214504000000980

11. Ridgeway, G. gbm: Generalized Boosted Regression Models. R Packag. version 1.6-3.1 (2010).

12. Venables, W. N. & Ripley, B. D. Modern Applied Statistics with S Fourth edition by. World (2002). doi:10.2307/2685660

13. Breiman, L. & Cutler, A. Breiman and Cutler's random forests for classification and regression. Packag. 'randomForest' (2012). doi:10.5244/C.22.54

14.   Jaccard, P. Distribution comparée de la flore alpine dans quelques régions des Alpes
occidentales et orientales. Bull. la Murithienne (1902).

15.   Delmont, T. O. et al. Nitrogen-fixing populations of Planctomycetes and Proteobacteria are
abundant in surface ocean metagenomes. Nat. Microbiol. (2018). doi:10.1038/s41564-018-0176-9

16.   Aumont, O., Ethé, C., Tagliabue, A., Bopp, L. & Gehlen, M. PISCES-v2: An ocean
biogeochemical model for carbon and ecosystem studies. Geosci. Model Dev. (2015).
doi:10.5194/gmd-8-2465-2015

17.   Reygondeau, G. et al. Dynamic biogeochemical provinces in the global ocean. Global
Biogeochem. Cycles (2013). doi:10.1002/gbc.20089

18.   Fay, A. R. & McKinley, G. A. Global open-ocean biomes: Mean and temporal variability. Earth
Syst. Sci. Data (2014). doi:10.5194/essd-6-273-2014

19.   Michelangeli, P. A., Vrac, M. & Loukos, H. Probabilistic downscaling approaches: Application
to wind cumulative distribution functions. Geophys. Res. Lett. (2009). doi:10.1029/2009GL038401

20.   Watson, R. A. A database of global marine commercial, small-scale, illegal and unreported
fisheries catch 1950-2014. Sci. Data (2017). doi:10.1038/sdata.2017.39

21.   Flanders Marine Institute (2018). Maritime Boundaries Geodatabase: Maritime Boundaries
and Exclusive Economic Zones (200NM), version 10. (2018). doi:https://doi.org/10.14284/313.

22.   Biecek, P. DALEX: explainers for complex predictive models. J. Mach. Learn. Res. 19, 1–5
(2018).

23.   Barton, A. D., Irwin, A. J., Finkel, Z. V. & Stock, C. A. Anthropogenic climate change drives shift
and shuffle in North Atlantic phytoplankton communities. Proc. Natl. Acad. Sci. (2016).
doi:10.1073/pnas.1519080113