Asymmetries around the visual field:
From retina to cortex to behavior

Authors

Eline R. Kupers[1,2,3]*, Noah C. Benson[1,2,4], Marisa Carrasco[1,2], Jonathan Winawer[1,2]

Affiliations

[1] Dept. of Psychology, New York University, New York, NY 10003

[2] Center for Neural Sciences, New York University, New York, NY 10003

[3] Current affiliation: Department of Psychology, Stanford University, Stanford, CA, 94305

[4] Current affiliation: eScience Institute, University of Washington, Seattle, WA, 98195

* Corresponding author. Email to: eline.kupers@nyu.edu

Funding

This work was supported by NIH R01-EY027401 to MC and JW.

Acknowledgements

We thank Michael Landy and Brian Wandell for their useful comments.

1 Abstract

Visual performance varies around the visual field. It is best near the fovea compared to the

periphery, and at iso-eccentric locations it is best on the horizontal, intermediate on the lower,

and poorest on the upper meridian. The fovea-to-periphery performance decline is linked to the

decreases in cone density, retinal ganglion cell (RGC) density, and V1 cortical magnification

factor (CMF) as eccentricity increases. The origins of polar angle asymmetries are not well

understood. Optical quality and cone density vary across the retina, but recent computational

modeling has shown that these factors can only account for a small percentage of behavior.

Here, we investigate how visual processing beyond the cone photon absorptions contributes to

polar angle asymmetries in performance. First, we quantify the extent of asymmetries in cone

density, midget RGC density, and V1 CMF. We find that both polar angle asymmetries and

eccentricity gradients increase from cones to mRGCs, and from mRGCs to cortex. Second, we

extend our previously published computational observer model to quantify the contribution of

phototransduction by the cones and spatial filtering by mRGCs to behavioral asymmetries.

Starting with photons emitted by a visual display, the model simulates the effect of human

optics, cone isomerizations, phototransduction, and mRGC spatial filtering. The model performs

a forced choice orientation discrimination task on mRGC responses using a linear support

vector machine classifier. The model shows that asymmetries in a decision-maker's

performance across polar angle are greater when assessing the photocurrents than when

assessing isomerizations and are greater still when assessing mRGC signals. Nonetheless, the

polar angle asymmetries of the mRGC outputs are still considerably smaller than those

observed from human performance. We conclude that cone isomerizations, phototransduction

and the spatial filtering properties of mRGCs contribute to polar angle performance differences,

but that a full account of these differences will entail additional contribution from cortical

representations.

Introduction

Visual performance is not uniform across the visual field. The most well-known effect is a decrease in visual acuity as a function of eccentricity: we see more poorly in the periphery compared to the center of gaze [1-4]. This observed difference in visual performance has been attributed to several physiological factors, starting as early as the distribution of photoreceptors [5, 6]. In the human fovea, the cones are tightly packed such that visual input is encoded at high spatial resolution. In peripheral retinal locations, cones are larger and interspersed among rods, resulting in a drastically lower density [7-10]; hence a decrease in spatial resolution.

Visual performance also differs as a function of polar angle. At matched eccentricity, performance is better along the horizontal than vertical visual meridian (horizontal-vertical anisotropy or "HVA", *e.g.*, [11-16]) and better along the lower than upper vertical visual meridian (vertical-meridian asymmetry or "VMA", *e.g.*, [12-18]). These polar angle asymmetries are observed in many different visual tasks, such as those mediated by contrast sensitivity [12-15, 19-31] and spatial resolution [11, 16, 17, 19, 20, 32-34], contrast appearance [35], visual search [36-44], crowding [44-47], and tasks that are thought to recruit higher visual areas such as visual working memory [34]. Covert spatial attention improves performance similarly at all iso-eccentric stimulus locations, thus it does not eliminate the polar angle asymmetries [12, 13, 48, 49].

These polar angle effects can be large. For instance, for a Gabor patch at 4.5° eccentricity with a spatial frequency of 4 cycles per degree, contrast thresholds are close to double for the upper vertical meridian compared to the horizontal meridian [12, 13, 15]. This is an effect size similar to doubling stimulus' eccentricity from 4.5° to 9° on the horizontal axis [15, 20]. Additionally, these performance differences are retinotopic, shifting in line with the retinal location of the stimulus rather than its location in space [14].

The visual system has polar angle asymmetries from its earliest stages, including in the optics and cone density. In a computational observer model that tracked information from the photons in the scene through the optics and cone isomerizations, variations in optical quality and cone density accounted for less than 10% of the observed polar angle asymmetries in a contrast threshold task [50]. This leads to the question, what additional factors later in the visual processing stream give rise to visual performance differences with polar angle?

One possibility is that even without additional asymmetries in cell density, later processing could exacerbate the earlier asymmetries. For example, the larger cone apertures

observed at lower cone densities result in greater downregulation of the cone photocurrent [51],

hence this decrease in signal-to-noise ratio might exacerbate polar angle asymmetries.

A second –not mutually exclusive– possibility is that there are additional polar angle

asymmetries in the distribution of other downstream cell types. In the human retina, the best

described retinal ganglion cells (RGCs) are the midget and parasol cells. Both of these cell

types show a decrease in density as a function of eccentricity and vary in density as a function

polar angle in humans [52-58] and monkeys [59-62]. Because midget RGCs are the most

numerous ganglion cells in primates (*i.e.*, 80% of ~1 million RGCs compared to 10% parasols

and 10% other types) and have small cell bodies and small dendritic field trees that increase

with eccentricity [60, 61, 63], they are often hypothesized to set an anatomical limit on high

resolution spatial vision such as acuity and contrast sensitivity at mid to high spatial frequencies

[55, 61].

Interestingly, in the range of eccentricities used for many psychophysical tasks (0–10°),

cone density shows an HVA (greater density on the horizontal than vertical meridian), but not a

VMA, inconsistent with behavior (there is a slightly greater density on the upper than lower

vertical visual meridian, opposite what one would predict to explain behavior) [8-10]. Midget

RGC density, in contrast, shows both an HVA and a VMA, making their distribution patterns

more similar to behavioral patterns [52-54, 57, 64].

Here, we investigate how asymmetries in the visual system vary across processing

stages. First, we quantify asymmetries in spatial sampling around the visual field in three early

visual processing stages: cones, mRGCs, and V1 cortex. We do so because it is important to

first identify *if* there are any differences in spatial encoding across these processing stages, and

if so, *how* these differences relate to differences in behavior. Then we extend our previously

published computational observer model, which included optics and cone sampling, by adding a

model of conversion from photon absorptions to photocurrent, and then mRGC-like spatial

filtering. We compare this observer model to our previous model (no RGC layer) and to human

performance on a 2-AFC orientation discrimination task. By comparing the predicted

performance to human observers, we can quantify the contribution of mRGCs to visual

performance differences around the visual field.

Results

We quantify the asymmetries in cone density, midget retinal ganglion cells (mRGCs) density
and V1 cortical magnification factor (CMF)—both as a function of eccentricity and for the four
cardinal meridians. In the next two sections, we first show that both eccentricity gradients and
polar angle asymmetries are amplified from cones to mRGCs and from mRGCs to early visual
cortex. Then we implement the observed variations in mRGC density in a computational
observer model to test whether biologically plausible differences in mRGC sampling across the
cardinal meridians can quantitatively explain psychophysical performance differences as a
function of polar angle.

Fovea-to-periphery gradient is amplified from retina to mRGCs to early visual cortex

A hallmark of the human retina is the sharp drop in cone density from fovea to periphery [8-10].
Within the central one degree, cone density decreases dramatically (on average by 3.5-fold).
Beyond the fovea, cone density continues to decrease by 10-fold between 1° and 20°
eccentricity (**Fig 1A**, left panel). This decrease in cone density is due to an increase in cone
spacing caused by the presence of rods and by the increase in cone diameter [9].

The second processing stage we focus on are the midget RGCs. The mRGC cell bodies
are laterally displaced from their receptive fields by the foveal cones. Therefore, we use a
computational model by Watson [64] that combines cone density [9], mRGC density [53] and
displacement [57] to infer the mRGC density referred to the visual field, rather than the cell body
positions. Throughout, we refer to mRGC density with respect to receptive fields. Like the
cones, midget RGCs sample the visual field differentially as a function of eccentricity. At the
central one degree, mRGC density is greater than cone density. The fovea-to-periphery gradient
is steeper for mRGCs than for cones (**Fig 1A,** middle panel compared to left panel). This
divergence results in a cone:mRGC ratio of 0.5 (**Fig 1B**, left panel), indicating a 'direct line'
between a single cone and a pair of on- and off-center mRGCs. In the periphery, mRGC density
falls off at a faster rate than cones. For example, cone density decreases by 10-fold between 1°
and 20° eccentricity, whereas mRGC density decreases by 80-fold. This *convergence* can also
be expressed in the cone:mRGC ratio, which increases as a function of eccentricity (**Fig 1B**, left
panel).

Fig 1. Foveal over-representation is amplified from cones to mRGCs to cortex. (A) Cone density, mRGC
receptive field density and V1 cortical magnification factor as a function of eccentricity. Left panel: Cone data from
Curcio *et al*. [9]. Middle panel: midget RGC RF density data from Watson [64]. Both cone and mRGC data are the
average across cardinal retinal meridians of the left eye using the publicly available toolbox ISETBIO [65-67]. Right
panel: V1 CMF is predicted by the areal equation published in Horton and Hoyt [68]. **(B) Transformation ratios from
cones to mRGCs and mRGCs to V1.** The cone:mRGC ratio is unitless, as both cone density and mRGC density are
quantified in cells/deg^2. The increasing ratio indicates higher convergence of cone signals by the mRGCs. For
mRGC:V1 CMF ratio units are defined in cells/mm^2. The ratio increase in the first 20 degrees indicates an
amplification of the foveal over-representation in V1 compared to mRGCs.

Third, we quantify the amount of V1 surface area devoted to a portion of the visual field,

also known as the cortical magnification factor (**Fig 1A**, right panel). There have been claims

that V1 CMF is proportional to retinal ganglion cell density [69-72] and see Discussion).

However, when comparing human mRGCs density [64] to V1 CMF [68], we find that the ratio is

not constant: The foveal magnification is even more accentuated in V1 up to 20° eccentricity

(**Fig 1B**, right panel). These results are consistent with the findings in squirrel monkey [73]; owl

monkey [74], and macaque [75], all of which show that the cortical magnification function falls

off with eccentricity more steeply in V1 than would be predicted by mRGC density alone.

Beyond 20° eccentricity, the mRGC to V1 CMF ratio declines slowly. This effect is driven by V1

CMF falling off slightly more steeply than mRGC density. The relative compression of V1 CMF

138 vs mRGC density in the far periphery has been reported in owl monkey [74]. However, given

139 that this result has not been confirmed in human cortex, we cannot exclude the possibilities that

140 in the far periphery Watson's formula [64] overpredicts mRGC density, Horton and Hoyt's

141 formula [68] underpredicts V1 CMF, or a combination of both.

Polar angle asymmetries are amplified from cones to mRGCs

Cone density differs as a function of polar angle. It is higher along the horizontal visual field

meridian (average of nasal and temporal retina meridians) than the upper and lower vertical

visual field meridians (representing the inferior and superior retinal meridians) (**Fig 2A**, left

panel). This horizontal-vertical asymmetry is around 20% and relatively constant with

eccentricity. There is no systematic difference between the cone density in the upper and lower

visual field meridians. If anything, there is a slight 'inverted' vertical-meridian asymmetry in the

central five degrees: cones are more densely packed along the upper vertical visual meridian.

Assuming greater density leads to better performance, this would predict better performance on

the upper vertical meridian in the central three degrees, opposite of the typical asymmetry

reported in behavior, which has been found up to 1.5° eccentricity in a study on contrast

sensitivity [30]. All of these patterns of cone density asymmetries are found using two different

datasets with different methods: a post-mortem retinal dataset [9] and an *in vivo* dataset [10],

indicating reproducibility of the biological finding. All of the patterns are also consistent when

computed using two different analysis toolboxes (ISETBIO [65-67] and rgcDisplacementMap

[76], **Supplemental Fig 1**, top row), indicating computational reproducibility.

The polar angle asymmetries in density are larger in the mRGC distribution. The

horizontal visual field meridian (average of nasal and temporal retina) contains higher cell

densities (after correction for cell body displacement) than the upper and lower visual field

meridians (**Fig 2A**, middle panel). This horizontal-vertical asymmetry increases with eccentricity.

For example, at 3.5° eccentricity, the average horizontal visual field density is ~20% higher than

the average of upper and lower visual field meridians. By 40° eccentricity, this density difference

increases to ~60%. Beyond 10° eccentricity, this horizontal-vertical asymmetry is mostly driven

by the nasal retina, as it contains higher mRGC density than the temporal retina. This finding is

in line with earlier histology reports in macaque [62] and positively correlated with spatial

resolution tasks (*e.g.,* [77]). This nasal-temporal asymmetry, although interesting, is beyond the

focus of this paper, as the asymmetries in performance we observe are found in both binocular

and monocular experiments [12, 16]. Overall, the emphasis on the horizontal is substantially

greater in the mRGCs than the cones.

Unlike the cones, mRGC receptive fields show a consistent asymmetry along the vertical

meridian: The lower visual meridian (superior retinal meridian) contains a higher mRGC density

than the upper visual meridian (inferior retinal meridian). This is consistent with the

psychophysical VMA, showing better performance on the lower vertical meridian [12-15, 19-31].

This asymmetry increases with eccentricity. For example, the lower vertical meridian (superior

retina) has ~15% higher density compared to upper vertical (inferior) at 3.5°, and ~50% higher

density at 40° eccentricity. This interaction between retinal meridian and eccentricity is

summarized in the cone-to-mRGC transformation plot (**Fig 2B**, left panel), where the

convergence ratio from cones to mRGCs increases more rapidly along the upper than the lower

vertical and the horizontal visual meridians (see also **Supplemental Fig 2**).

Polar angle asymmetries are amplified from mRGCs to early visual cortex

Because the areal V1 CMF calculation by Horton and Hoyt [68] does not make separate

predictions for the cardinal meridians, we used the publicly available retinotopy dataset from the

Human Connectome Project (HCP) analyzed by Benson *et al.* [79] to calculate the CMF along

the meridians (see also [78]). As a first check on agreement between the two datasets, we

found that the V1 CMF data measured in 163 subjects with functional MRI [78], pooled across

all polar angles, was a close match to Horton and Hoyt's [68] prediction based on lesion case

studies from three decades ago. We then used the HCP dataset to compute CMF along the

separate meridians.

We find that polar angle asymmetries in cortical magnification factors are yet larger than

those found in mRGC density (**Fig 2A**, right panel), where V1 CMF is higher on the horizontal

than vertical meridian, and the V1 CMF is higher for the lower than the upper vertical meridian.

For example, at 3.5° eccentricity CMF is ~52% higher on the horizontal than vertical meridians

and ~41% higher for the lower than upper vertical meridian. These polar angle asymmetries

show a 2x increase within the first three degrees of eccentricity before flattening (**Fig 2B**, right

panel) and are mostly driven by the upper vertical meridian (**Supplemental Fig 2**). This

indicates that the mapping of the visual field in early visual cortex is not simply predicted from

the distribution of midget retinal ganglion cells, but rather the cortex increases the retinal polar

angle asymmetries.

A computational observer model from stimulus to mRGCs to behavior

To understand how polar angle asymmetries in visual field representations might affect visual
performance, we added a photocurrent transduction and retinal ganglion cell layer to our
computational observer model [50]. In this observer model, we used the publicly available
ISETBIO toolbox [65-67] to simulate the first stages of visual pathway including the stimulus
scene, fixational eye movements, chromatic and achromatic optical aberrations, and
isomerization by the cone array. Combining model output with a linear support vector machine
classifier allowed us to simulate performance on a 2-AFC orientation discrimination task given
information available in the cones. When matching stimulus parameters in the model to a
previously published psychophysical experiment [13], we showed that biologically plausibly
variations in optical quality and cone density together would contributed no more than ~10% to
the observed polar angle asymmetries in contrast sensitivity.

Given the inability of cone density to quantitatively explain differences in visual
performance, we extended our model further into the retina to include temporal and spatial
filtering, and noise at two later processing stages. First, we added temporal filtering and noise in
the conversion of cone isomerizations to photocurrent in the cone outer segments. Second, we
added spatial filtering and noise in a model of midget RGCs. The mRGCs are especially
interesting because they show a systematic asymmetry between the upper and lower visual
field (where the cones did not), and an amplification of the horizontal-vertical asymmetry. The
mRGC computational stage is implemented after cone isomerizations and photocurrent and
before the model performs the discrimination task. We provide a short overview of the modeled
stages that precede the mRGC layer, as details of these stages can be found in our previous
paper [50], followed by a discussion of the implementation details of the photocurrent
transduction and mRGC layer.

Scene radiance

The first stage of the model comprises the photons emitted by a visual display. This results in a
time-varying scene defined by the spectral radiance of an achromatic low contrast Gabor
stimulus (**Fig 3**, panel 1). The Gabor was oriented 15° clockwise or counter-clockwise from
vertical with a spatial frequency of 4 cycles per degree. These stimulus parameters were
chosen to match a recent psychophysical experiment [15] to later compare model and human
performance.

Retinal irradiance

The second stage simulates the effect of emitted photons passing through the human cornea, pupil, and lens. This computational step results in time-varying retinal irradiance (**Fig 3**, panel 2). Optics are modeled as a typical human wavefront with a 3-mm diameter pupil without defocus, and contain a spectral filter that reduces the fraction of short wavelengths (due to selective absorption by the lens). We do not vary the optics across the different simulations.

Cone absorptions

The third stage implements a rectangular cone mosaic with L-cones only (2x2° field-of-view). For each cone, we compute the number of photons absorbed in each 2-ms bin, resulting in a 2D time-varying cone absorption image (**Fig 3**, panel 3). The number of absorptions depends on the photoreceptor efficiency and on the wavelengths of light and on Poisson sampling due to the quantal nature of light. This stage differs in two ways from our previous model. First, we use an L-cone-only retina, and second, we exclude fixational eye movements. We make these two simplifications to keep the model tractable and the calculations to reasonable size. As we describe in the Methods, the number of trials is much larger than in our previous work (to ensure that the classifier has sufficient information to learn the best classification), the number of conditions simulated is much larger (because we vary both cone density and mRGC:cone ratios), and the noise level is substantially higher (because we add noise at phototransduction and mRGC stages). The lack of eye movements enables us to average time points across trials, greatly speeding up processing, as well as simplifying the interpretation of how the new stages contributed to performance.

Cone photocurrent

The fourth stage converts photon absorptions to photocurrent, incorporating the recently added phototransduction functionality in ISETBIO by Cottaris *et al*. [51], Here, phototransduction is implemented as a temporal filter followed by gain control and additive noise (**Fig 3**, panel 4). The result is a continuous time-varying signal in units of current (picoamps). While we use the same photocurrent model for all cones irrespective of size or location, the effect of the photocurrent depends on properties of the cones, due to the additive noise. Specifically, the signal-to-noise decreases more for larger cones than smaller cones, because large cones capture more photons and are subject to more downregulation before the additive noise.

Fig 3. Overview of computational observer model with additional mRGC layer. A 1-ms frame of a 100% contrast Gabor stimulus is used at each computational step for illustration purposes. **(1) Scene radiance.** Photons emitted by the visual display, resulting in a time-varying scene spectral radiance. Gabor stimulus shows radiance summed across 400-700 nm wavelengths. **(2) Retinal irradiance.** Emitted photons pass through simulated human cornea, pupil, and optics, indicated by the schematic point spread function (PSF) in the top right-side box, resulting in time-varying retinal irradiance. Gabor stimulus shows irradiance with wavelengths converted to RGB values for illustration purposes. **(3) Cone absorptions.** Retinal irradiance is isomerized by a rectangular cone mosaic, resulting in time-varying photon absorption rates for each L-cone with Poisson noise. **(4) Cone photocurrent.** Absorptions are converted to photocurrent via temporal integration, gain control, followed by adding Gaussian white noise. This results in time-varying photocurrent for each cone. **(5) Midget RGC responses.** Time-varying cone photocurrents are convolved with a 2D Difference of Gaussians spatial filter (DoG), followed by additive Gaussian white noise and subsampling (see also Fig 4). **(6) Behavioral inference.** A linear support vector machine (SVM) classifier is trained on the RGC outputs to classify stimulus orientation per contrast level. With 10-fold cross-validation, left-out data are tested, and accuracy is fitted with a Weibull function to extract the contrast threshold at ~80%.

Midget RGC responses

The fifth stage is spatial filtering by the mRGCs. We model the mRGCs in a rectangular array
with each mRGC receptive field centered on a cone. We do not add further temporal filtering
beyond that inherited from the photocurrent stage. We do not explicitly model spiking and its
associated noise, but instead add independent Gaussian white noise to each RGC output at
each time point. Unlike the photocurrent, where the noise is implemented in ISETBIO according
to a physiologically informed model [80], the noise added in the mRGC layer is not constrained
by a physiological model because the noise added by mRGCs (after accounting for noise
inherited from prior stages) is less well known. For this reason, in additional simulations, we
explore the effect of noise level in RGCs, and find that while the mean performance declines
with increasing noise (as expected), the differences between conditions are largely unaffected
by noise level (**Supplemental Fig 4**). In the Discussion, we elaborate on the possible
contribution of other aspects of retinal processing to polar angle asymmetries such as spatial
subunits and spiking.

The mRGC layer has the same field-of-view as the cone array. Because we do not
model rectification or spiking non-linearities, we do not separately model on- and off-cells. Our
mRGC receptive fields are 2D difference of Gaussian (DoG) models, approximating the shape
of receptive fields measured with electrophysiology [81, 82] (**Fig 3**, panel 5), based on
parameters from macaque [83]. The width of the center Gaussian (σ_c, 1 sd) is ⅓ of the spacing
between neighboring cones, and the surround Gaussian (σ_s) is 6x the width of the center. This
creates an mRGC array with one mRGC per cone and where mRGC RFs overlap at 1.3
standard deviations from their centers, which matches the overlap of dendritic fields reported in
human retina [55]. We compute the mRGC responses by convolving the cone absorptions with
this mRGC DoG receptive field. Because the ratio of mRGCs to cones varies across the retina,
we simulate differences in this ratio by subsampling the mRGC array (**Fig 4**). Thus, the mRGC
density (cells/deg^2) is determined by both the cone array density and the cone-to-mRGC ratio,
creating a 2D space of simulations.

Fig 4. Difference of Gaussians filters used to model mRGC layer. Two mRGCs are illustrated for a 2x2° field-of-view mRGC array centered at 4.5° and 40° eccentricity. **(A) 1D representation of two example mRGC layers in visual space.** The mRGC responses are computed by convolving the cone image with the mRGC DoG RF, followed by adding noise (not shown), and subsampling the cone array to the corresponding mRGC density. Width for Gaussian center (σ_c) and surround (σ_s) are converted to units of degree. As the mRGC filters in our model are not rectified, they respond to both increments and decrements. Physiologically, this would require two cells (an ON and OFF cell), so we count each modeled mRGC location as two cells. Both panels show a mRGC:cone ratio of 2:1. **(B) 1D representation of Difference of Gaussians in Fourier space.** The Fourier representation illustrates the band-pass and unbalanced nature of the DoGs (*i.e.*, non-zero amplitude at DC). Depending on the width/subsample rate, DoGs attenuate different spatial frequencies. However, at our peak stimulus frequency (4 cycles per degree, indicated with red dashed line) the two DoGs filters vary a relatively small amount, preserving most stimulus information. Fourier amplitudes are normalized. Note that y-axis is truncated for illustration purposes. **(C) 2D representation of two example mRGC layers shown in panel (A).** Midget RGC DoG filters are zoomed into a 1x1° field-of-view cone array (black raster) centered at 4.5° (red center with purple surround) and 40° eccentricity (red center with yellow surround), corresponding to the 1D examples in panel A. Centers and surrounds are plotted at 2 standard deviations. For illustration purposes, only one mRGC is shown; the mRGC array in our computational observer model tiles the entire cone array.

Behavioral inference

The final stage of the computational observer model is the decision maker. For the main analysis, we use a linear support vector machine (SVM) classifier to discriminate stimulus orientation (clockwise or counter-clockwise from vertical) given the cone absorptions, cone photocurrent, or mRGC responses. We compute a weighted average across time for the output of each cell before running the classifier. This greatly reduces the dimensionality of the classifier input, and therefore speeds up computation time and reduces the number of trials needed for

the classifier to learn optimal classification boundary. The weighting is proportional to the

temporal filter in the photocurrent simulation, such that the time points with the highest weight in

the filter has the largest contribution to the weighted average. Because we do not simulate eye

movements or vary the phase of the stimulus, the only changes over time arise from the noise

and temporal filtering by the photocurrent, and hence there is little to no loss of signal from

averaging. The classifier trains and tests on the averaged responses for each stimulus contrast

separately, where each contrast level results in a percent correct identified stimulus. The

accuracy results are then fitted with a Weibull function to extract the contrast threshold at ~80%.

The cone photocurrent and mRGCs have a large effect on orientation discrimination

We find large effects on performance of the computational observer when adding the cone

photocurrent and the mRGC layers. For comparison, we ran the SVM decision maker either on

the cone absorptions, the cone photocurrent, or the mRGC outputs while varying the cone

density and the stimulus contrast. Consistent with our prior model [50], thresholds are low (~0.1-

0.2%) when analyzed on the cone absorptions, and show only a small effect of cone density

(**Fig 5A**). Thresholds increase sharply, about 5-10x, after the absorptions are converted to

photocurrent (**Fig 5B**). This increase is due to noise in the photocurrent, consistent with prior

results [51]. Surprisingly, the effect of cone density is also substantially increased, as seen in

the greater spread of the psychometric functions. This is because the cones in the lower density

retinal patches have larger apertures, resulting in greater photon capture, and hence more

downregulation when converted to photocurrent. Over the 10-fold range of retinal densities,

threshold vary by only about 1.4:1 for the absorptions, much less in contrast to about 5:1 for the

photocurrent. The spatial filtering and late noise from the mRGCs further elevate thresholds, but

at a fixed mRGC:cone ratio there is little change in the effect of cone density: the threshold vs

density plot shows a vertical shift compared to the cone photocurrent, with about the same

slope (**Fig 5c**).

Fig 5. Model performance for different computational stages. Left column shows classifier accuracy as function of stimulus contrast. Data are from simulated experiments with 1,000 trials per stimulus class, using a model with a L-cone only mosaic varying in cone density. Data are fitted with a Weibull function. Contrast thresholds are plotted separately as a function of cone density in the right column. **(A) Cone absorptions.** Applying a linear SVM classifier to cone absorptions averaged across stimulus time points. **(B) Cone photocurrent.** Applying a linear SVM classifier to cone outer segment photocurrent responses, averaged across time weighted by a temporally delayed stimulus time course. This transformation of cone absorptions into photocurrent causes a ~10x increase in contrast thresholds, interacting with cone density (*i.e.*, Weibull functions are spaced out compared to cone absorptions). **(C) RGC responses.** Applying a linear SVM classifier to spatially filtered photocurrent with added white noise. This transformation causes an additional increase in contrast thresholds for all cone densities. Data show results for a fixed subsampling ratio of 2 mRGCs per cone.

We next quantified the effect of the mRGC:cone ratio on computational observer performance. We find that as the ratio increases, contrast thresholds decline (**Fig 6A**). The effect of the mRGC:cone ratio is largely independent of the cone density. For example, at any cone density, downsampling the mRGC density by 4x elevates thresholds by about 70% to 80%. The better model performance with more mRGCs comes from higher SNR, which arises because the signal is correlated across mRGCs (due to spatial pooling), whereas the noise added in the mRGC layer is independent. To visualize the space of predicted contrast thresholds as a function of cone density and mRGC:cone ratio, we plot model thresholds as a

function of both independent variables (**Fig 6B**). This surface plot confirms the observation from

the line plots (**Fig 6A**) that the effects of these two retinal factors—cone density and

mRGC:cone ratio—have approximately independent, additive effects on model contrast

threshold.

Fig 6. The effect of spatial filtering properties by mRGCs on full model performance. (A) Contrast thresholds
as a function of cone density and mRGC:cone ratio. Data points are contrast thresholds for cone absorptions,
cone photocurrent, and each mRGC:cone ratio separately (for psychometric functions see **Supplemental Fig 3**).
Individual mRGC fits are slices of the 3D mesh fit shown in panel B. **(B) Mirrored views of combined effect of cone**
density and mRGC:cone ratio on contrast sensitivity. The mesh is fitted with a locally weighted regression to 3D
data: log cone density (x-axis) by log mRGC:cone ratio (y-axis) by log contrast thresholds (z-axis). Individual dots
represent the predicted model performance for nasal retina or horizontal visual (red star), superior retina or lower

visual (blue star), temporal retina or horizontal visual (green star) and inferior or upper visual (black star) meridian
locations at 4.5° eccentricity (matched to stimulus eccentricity in [15]). Contour lines show possible cone densities
and mRGC:cone ratios that would predict the same horizontal-vertical and upper/lower vertical-meridian asymmetry
as observed in psychophysical data at 4.5° eccentricity. To do so, we scaled the difference in contrast threshold
between the lower (blue) and upper (black) vertical visual meridian relative to the horizontal meridian to match the
difference in behavior. Goodness of fit of 3D mesh fit is $R^2 = 0.96$.

Comparison between model and human contrast sensitivity

To compare model performance to human observers, we evaluate the model outputs for cone

densities and mRGC:cone ratios that match the values on the different meridians according to

the literature. We then compare these predicted thresholds to those obtained in a recent

psychophysical experiment [15]. We also compare both the human data and the mRGC model

data to two simplified models, one which omits the mRGCs and one which omits mRGCs and

the conversion from isomerizations to photocurrent.

According to Curcio *et al.* [9], cone density at 4.5° eccentricity is ~1,575 cones/deg^2 on

the horizontal retinal meridian (nasal: 1590 cones/deg^2, temporal: 1560 cones/deg^2), 1300

cones/deg^2 on the superior retinal meridian, and 1382 cones/deg^2 on the inferior retinal

meridian. We combine these cone density values with the mRGC:cone ratios from the

computational model by Watson [64], which ranges between 0.84 mRGCs per cone on the

horizontal meridian (nasal: 0.87, temporal: 0.82), to 0.81 on the superior retina and 0.68 on the

inferior retina.

Consistent with our previous report [50], we find that a model in which the pattern of

photon absorptions is fed into the linear SVM classifier shows only a small effect of cone density

(**Fig 7A**, left). Given the expected cone densities at the different polar angles at 4.5°

eccentricity, the model predicts only about 5% higher sensitivity for the horizontal than vertical

visual meridians, much less than the 40% difference found in behavioral experiments [15] (**Fig

7B**). The model also predicts almost no difference between upper and lower vertical visual

meridian, whereas human sensitivity was found to be about 20% higher on the lower than upper

vertical visual meridian. The overall sensitivity of the model observer (800-900) is considerably

higher than human sensitivity (~30-50).

The conversion from cone absorptions to cone photocurrent reduces the sensitivity by

about 4- to 5-fold, and increases the asymmetries. The linear SVM classifier performance based

on the cone photocurrent shows about 15% higher sensitivity for horizontal than vertical visual

meridian, an asymmetry that is 3 times larger than that found in a model up to cone

isomerizations. It also predicts about 9% higher sensitivity for upper vertical than lower vertical

visual meridian (opposite to the pattern in human data). This is because the cone density is

slightly higher for the upper than lower vertical visual meridian at this eccentricity (4.5 degrees).

Finally, the mRGC model brings overall performance closer to behavior, with sensitivity

of about 70-90, and ~18% higher sensitivity for the horizontal than vertical visual meridian,

predicting almost half the asymmetry found in behavior (~40%). The mRGC model also

eliminates the advantage for upper over lower vertical visual meridian (now predicting slightly

higher performance for the lower vs upper vertical), which is the same direction as the pattern

observed in the human data.

Fig 7. Comparison of model performance to human performance. (A) Contrast sensitivity predicted by
computational observer model up to isomerizations in cones (blue), up to cone outer segment
phototransduction (turquoise), up to spatial filtering and subsampling in mRGCs (red), and behavior
observed (purple) by Himmelberg *et al*. (2020) using matching stimulus parameters. HM: horizontal meridian,
UVM: upper visual meridian, LVM: lower visual meridian. Model prediction shows contrast sensitivity (reciprocal of
contrast threshold) for stimuli at 4.5° eccentricity, with a spatial frequency of 4 cycles per degree. HM is the average
of nasal and temporal meridians. Model error bars indicate simulation results allowing for uncertainty in the cone or
mRGC density along each meridian (see Methods for details). Behavioral plots show group average results (*n*=9)
from Himmelberg *et al.* [15], and error bars represent standard error of the mean across observers. **(B) Polar angle**
asymmetries for cone absorptions, photocurrent, mRGCs, and behavior. HVA: horizontal-vertical asymmetry.
VMA: vertical-meridian asymmetry. Blue, turquoise, red, and purple bars match panel (A) and correspond to model
prediction up to cone absorptions, cone photocurrent, mRGCs, human behavior. Error bars represent the HVA and
VMA when using the upper/lower bound of predicted model error from panel A.

Overall, our models show that although including an mRGC layer predicts polar angle

asymmetries closer to behavior than a model up to cone absorptions or up to photocurrent, the

biological variations in the spatial properties of mRGCs are not sufficient to fully explain

differences in behavior. For example, the measured cone densities for the upper and lower

vertical visual meridians are about 12% and 19% lower than for the horizontal. To predict the

horizontal-vertical and vertical-meridian asymmetries as observed in human performance, and

without further changing the mRGC:cone ratios, the cell densities would instead have to be

~37% and 30% lower than the horizontal. Alternatively, one could keep the cone densities fixed

at the levels estimated by Curcio *et al*. [9], and instead vary the mRGC:cone ratio as observed

by Watson [64]. In this case, the ratios would have to decrease from 0.81 to 0.52 for the lower

vertical and 0.68 to 0.32 for the upper vertical visual meridian. If one decreased both the cone

densities and the mRGC:cone ratios by tracing out the values along the nasal retinal meridian,

one would need to increase eccentricity of a stimulus from 4.5° to 7.3° (upper vertical) or 6.3°

(lower vertical) to match the behavioral asymmetries.

497 Discussion

The visual system, from retina to subcortex to cortex, is organized in orderly maps of the visual

field. But within each particular processing stage, the retinotopic map is distorted. Here we

investigated the polar angle asymmetries in these spatial representations across three stages of

the early visual pathway: cones, mRGCs and V1 cortex. Our study revealed that both the

eccentricity gradient (foveal bias) and polar angle asymmetries (HVA and VMA) in spatial

representations are amplified from cones to mRGCs, and further amplified from mRGCs to early

visual cortex. Additionally, we showed that although mRGC density has considerably polar

angle asymmetries in the directions predicted by psychophysical studies, they are insufficient to

explain observed differences in human's contrast sensitivity around the visual field.

507 Linking behavior to eccentricity and polar angle asymmetries in visual field
508 representations

For over a century, limits in retinal sampling were hypothesized to cause the fovea-to-periphery

gradient in human visual performance [1, 5, 6]. Initial tests of this idea showed that the fall-off in

cone density could explain some, but not all of the observed decrease in visual acuity [2, 3, 84-

87]. Later, more detailed computational models, reported that mRGCs come closer in predicting

the eccentricity-dependent decrease in achromatic contrast sensitivity and resolution, and

conclude that mRGCs are sufficient to explain some aspects of behavior, such as spatial

resolution and contrast sensitivity [88-94]. Similar to the retina, the cortical magnification factor

in V1 has been linked to visual performance as a function of eccentricity, for example,

explaining differences in acuity [92, 95, 96], contrast sensitivity and resolution [20], visual search

[97, 98], and the strength of some visual illusions [99].

Conversely, polar angle asymmetries have rarely been considered. For instance, all

above-mentioned studies either ignored the stimulus polar angle for analysis or limited

measurements to a single meridian, usually the horizontal. Despite the fact that the existence of

polar angle asymmetries in human early visual cortex was predicted based on behavior in the
late 70's [19, 20], further reports on polar angle differences have been scarce. One fMRI study
reported a higher V1 BOLD amplitude for stimuli on the lower than the upper visual meridian
[100] and two studies found more cortical surface area devoted to the horizontal than the
vertical meridian [101, 102]. Our recent studies suggest that V1 surface area is highly correlated
to spatial frequency thresholds [78] and contrast sensitivity [103]. Yet several studies have
assumed little to no polar angle differences in macaque V1 CMF [104, 105] or did not account
for polar angle differences in human V1 CMF [46, 96] to explain differences in behavior.
Computational models that include retinal and/or V1 sampling across visual space generally
exclude polar angle asymmetries (*e.g.,* [106, 107]). A few cases do incorporate polar angle
asymmetries in the retinal ganglion cell distribution, but they assume that these asymmetries
are not amplified in cortex [108-110].

Early visual cortex does not sample the retina uniformly

It is well documented that the convergence of cones to retinal ganglion cells varies with
eccentricity (*e.g.,* see [91]). In the fovea of both primates and humans, there is one cone per
pair of bipolar cells and pair of midget RGCs, with pairs comprised of an "on" and an "off" cell. In
contrast, in the periphery, there are many cones per pair of bipolar cells and midget RGCs, with
the ratio depending on the eccentricity. In the far periphery, there can be dozens of cones per
ganglion cell [9].

It has been long debated whether V1 further distorts the visual field representation, or if
V1 samples uniformly from RGCs, as reviewed previously [71, 72]. Our analysis showed more
cortical surface area devoted to the fovea than the parafovea and to the horizontal than vertical
meridian, supporting previous findings using retinotopy informed by anatomy [101] and
functional MRI [78, 102, 103, 111]. Importantly, these eccentricity and polar angle non-
uniformities are larger in V1 than they are in mRGC density, in agreement with findings from
monkey [61, 73-75, 112, 113]. Whether these non-uniformities arise in cortex, or depend on the
mapping from retina to LGN, LGN to V1, or both, is a question of interest in both human [114,
115] and monkey [116-120], but beyond the scope of this paper. The implication of the
increased spatial non-uniformities in the cortical representation is that cortex cannot be
understood as a canonical wiring circuit from the retina repeated across locations.

Because visual field distortions are larger as a function of eccentricity than polar angle,
one might surmise that polar angle asymmetries contribute little to visual performance. Even

though the polar angle asymmetries are smaller than the eccentricity effects, they can in fact be

large. For example, within the central eight degrees, the surface area in V1 is about 60% larger

for the horizontal meridian than the vertical meridian [78]. Given that virtually all visual tasks

must pass through V1 neurons, these cortical asymmetries are likely to have a large effect on

perception. The number of cortical cells could be important for extracting information quickly

[121], for increasing the signal-to-noise ratio, and for tiling visual space and visual features (*e.g.*,

orientation, spatial frequency) more finely [122]. To know how the number of V1 neurons affect

performance, there is a need for a computational model that explicitly links cortical resources to

performance around the visual field.

563 Temporal summation in cone photocurrent accentuates polar angle
564 asymmetries

We found one physiological factor in the retina—gain control in the cone photocurrent—that

appears to accentuate the polar angle asymmetries. This is because at matched eccentricities,

cone density varies with polar angle (*i.e.*, cone density is higher on the horizontal meridian), and

cone aperture size varies inversely with density. Specifically, at lower densities, the apertures

are larger, capturing more photons per receptor. As a result of the higher absorption rates, there

is greater downregulation of the photocurrent gain. Cottaris *et al*. [51] observed in their modeling

work that the lower gain in the photocurrent for larger cones caused a reduction in the signal-to-

noise ratio. In their simulations, this resulted in sensitivity loss for stimuli that extended further

into the periphery. In our simulations, lower density results in lower sensitivity, therefore

contributing to the difference in performance as a function of polar angle.

Overall, while adding a photocurrent stage decreases overall thresholds, bringing them

closer to human performance (especially for simulations with low cone density mosaics), it still

leaves a large gap between the predicted and observed psychophysical asymmetries as a

function of polar angle. Moreover, the photocurrent model does not explain any of the vertical

meridian asymmetry, as cone density, and presumably aperture size, do not differ between

lower and upper vertical meridian in a way that matches behavior.

581 Model limitations

Despite implementing known facts about the eye, our model, like any model, is a simplification.

The lack of comprehensiveness trades off with interpretability. For this model, we make the

trade-off between complexity and understanding by treating a local patch of mRGCs as a linear,

shift-invariant system (*i.e.*, a spatial filter). As several components of the model here are

identical to our previous model, we will focus on the limitations of those components that are

different (addition of cone photocurrent and mRGC layer, and exclusion of eye movements),

and refer to Kupers, Carrasco and Winawer [50] for model limitations related to the pathways

from scene to cone absorptions and the inference engine.

Spatial properties: Uniform sampling within a patch and subunits

Hexagonal cone arrays that include within-patch density gradients have been implemented in

ISETBIO by Cottaris *et al.* (e.g., [51, 67]). Nonetheless, our mRGC layer is implemented as a

rectangular patch of retina, initially with the same size as the cone mosaic. This allows for

filtering by convolution and then linear subsampling to account for mRGC density, making the

model computationally efficient. We do not incorporate several known complexities of RGC

sampling in the retina: (i) density gradients within a patch, (ii) irregular sampling, and (iii) spatial

RGC subunits. (i) Given our relatively small patch size (2x2° field-of-view) in the parafovea

(centered at 4.5°), the change in density across the patch would be small (~10%). We found

that a much larger change in mRGC density (spanning a 5-fold range) had only a modest effect

on performance of our observer model, so it is unlikely that accounting for a small gradient

within a patch would have significantly influenced our results. (ii) Given the relatively low spatial

frequency content of our test stimulus (4 cycles per degree), it is unlikely that irregular sampling

would have resulted in a substantial difference from the regular sampling we implemented. (iii)

Our low spatial frequency test stimuli also reduce concerns of omitting spatial subunits [123-

126], as these non-linearities are most likely to be important for stimuli at high spatial

frequencies (reviewed by [127]). Moreover, we showed for our linear RGC filters that sensitivity

differences are only large at high spatial frequencies (around 8 cycles per degree and higher);

even when receptive field sizes differ by a factor of 3 (as shown in **Fig 4B**). Hence for the

relatively low spatial frequency stimuli modeled here, the detailed spatial properties that we

excluded would likely not have large enough effects to make up the difference between the

predicted model performance and human behavior.

Temporal properties and eye movements

In contrast to our previous work [50], our current model includes temporal integration but omits

fixational eye movements and multiple cone types. The omission of eye movements made the

model more tractable and the computations more efficient. We think this omission is unlikely to

have a large effect on our results. In recent related work, it was shown that fixational eye

movements combined with temporal integration resulted in spatial blur and degraded

performance, causing a loss in contrast sensitivity up to a factor of 2.5 [51]. However, the

largest losses were for stimulus spatial frequencies over 8 cycles per degree, with little loss from

eye movements for stimuli with lower peak spatial frequency (2-4 cycles per degree). Given that

the spatial frequency of our test stimulus falls within this range, the influence of fixational eye

movements on the computational observer performance would have been modest.

Noise implementation

Our expectation was that the largest effect of mRGCs on performance as a function of polar

angle would arise from variation in cell density: where mRGC density is higher, SNR will be

higher, thus performance will be better. This effect of density on performance emerged in our

simulations from the noise added after spatial filtering, before subsampling: without this

additional noise component, the spatial filtering of the mRGC would just be a linear transform of

the cone outputs, which would have little or no effect on performance of a linear classifier. We

simulated this late noise as additive Gaussian noise rather than the stochastic nature of spiking,

as we were not trying to fit spiking data but rather predict behavior. While we also did not build

in correlated noise between RGCs (*e.g.*, [128]), there is nonetheless some shared noise in our

mRGCs due to common inputs from cones, which is the major source of noise correlations in

RGCs [129]. Moreover, we found that the general pattern of model performance was unchanged

over a large range of noise levels (up to an overall scale factor in performance), suggesting that

the effect of density is likely to hold in many noise regimes.

Other retinal cell types

Midgets are not the only retinal ganglion cells that process the visual field. Parasol (pRGCs) and

bistratified retinal ganglion cells are less numerous but also cover the entire retina. pRGCs are

the next most common retinal ganglion cells, and have generally larger cell bodies and dendritic

field sizes than mRGCs, both increasing with eccentricity [54]. These differences cause

parasols to be more sensitive to relative contrast changes and have higher temporal resolution,

with the consequence of losing spatial resolution [130]. For this reason, the small mRGCs are

much more likely to put a limit on spatial vision, and thus our model does not include pRGCs.

The discussion above raises the question, had we incorporated more known features of

the retina in our model, would the model make predictions more closely matched to human

performance? We think it is unlikely that doing so would fully explain the observed asymmetries

in behavior, because we measured substantially larger asymmetries in cortex than in retina. If
the retinal simulations entirely accounted for behavior, this would leave no room for the
additional cortical asymmetries on behavior.

A case for cortical contributions to visual performance asymmetries

Recent retinal modeling of contrast sensitivity in the fovea showed that very little information
used for behavior seems to be lost from the retinal output [51]. This may not be the case for the
parafovea and periphery. Incorporating temporal properties of phototransduction and spatial
properties of mRGC followed by additive noise could explain about half the differences in
behavior of HVA and ~1/6 of VMA. These differences indicate a contribution from downstream
processing, such as early visual cortex. V1 cortex has several characteristics that suggest a
tight link between cortical topography and polar angle visual performance asymmetries. Hence
a model that incorporates properties of early visual cortex is likely to provide a substantially
better account of polar angle asymmetries in behavior than one that only incorporates properties
of the eye. We have not developed such a model but outline some of the reasons that cortex-
specific properties are important for explaining polar angle asymmetries.

First, the representation of the visual field is split across hemispheres in visual cortex
along the vertical, but not horizontal meridian. This split may require longer temporal integration
windows for visual input that spans the vertical meridian, as information needs to travel between
hemispheres. For example, the response in the left visual word form area is delayed by ~100
ms compared to the right visual word form area when presenting a stimulus in the left visual
field [131]. Longer integration windows may in turn impair performance on some tasks, as eye
movements during integration will blur the representation. Longer integration time of visual
information spanning the vertical meridian is consistent with behavior, as accrual time is slower
when stimuli are presented at the vertical than the horizontal meridian [38]. Interestingly, the
hemispheric split is not precise: there is some ipsilateral representation of the visual field along
the vertical meridian in early visual cortex. The amount of ipsilateral coverage is larger along the
lower than upper vertical meridian and increases from 1-6° eccentricity [132]. It is possible that
the split representation affects performance for stimuli on the vertical meridian (contributing to
the HVA), and that the asymmetry in ipsilateral coverage between the lower and upper vertical
meridian contributes to the VMA.

Second, there is good correspondence between the angular patterns of asymmetries in
V1 cortex and behavior. Polar angle asymmetries in the CMF of early visual cortex are largest

along the cardinal meridians (*i.e.,* horizontal vs vertical and upper vertical vs lower vertical). The

asymmetries gradually fall-off with angular distance from the meridians [78]. This gradual

decrease in polar angle asymmetry in cortex parallels the gradual decrease in contrast

sensitivity [12, 29, 30] and spatial frequency sensitivity [16] with angular distance from the

cardinal meridians. Measurements of cone density and retinal ganglion cell density have

emphasized the meridians, so there is less information regarding how the asymmetries vary

with angular distance from the meridians.

Third, there is good correspondence between cortical properties and behavior in the

domain of spatial frequency and contrast sensitivity. Polar angle asymmetries in spatial

frequency sensitivity observed by Barbot *et al.* [16] parallel spatial frequency tuning in V1 cortex.

Specifically, fMRI measurements show that in V1, in behavior spatial frequency thresholds are

higher on the horizontal than vertical visual meridian [16] and the preferred spatial frequency

tuning is higher along the horizontal meridian than vertical visual meridian [133]. Additionally,

polar angle asymmetries in contrast sensitivity covary with polar angle asymmetries in V1

cortical magnification [103]: Observers with larger horizontal-vertical asymmetries in contrast

sensitivity (*i.e.,* better performance on the horizontal vs vertical visual meridian at matched

eccentricities), tend to have larger horizontal-vertical asymmetries in V1 cortical magnification at

corresponding locations in the visual field.

Fourth, polar angle asymmetries in behavior are maintained when tested monocularly

[12, 16], but thresholds are slightly higher compared to binocular testing (at least for spatial

frequency sensitivity [16]). Higher thresholds (*i.e.,* poorer performance) show that performance

benefits from combining information of the two eyes, as twice the amount of information

increases the signal-to-noise ratio [134]. This summation is likely to arise in early visual cortex,

as V1 is the first stage in the visual processing pathways where information of the left and right

visual field merges [135-137].

705   Conclusion

Overall, we have shown that the well documented polar angle asymmetries in visual

performance are associated with differences in the structural organization of cells throughout

the early visual pathway. Polar angle asymmetries in cone density are amplified in downstream

processing, from cones to RGCs and again from RGCs to early visual cortex. Further, we have

extended our computational observer model to include temporal filtering when converting cone

absorptions to photocurrent and spatial filtering of mRGCs, and found that both contributions,

although larger than those of cones, are far from explaining behavior. In future research, we will

aim to integrate cortical data within the computational observer model to explain whether a

significant amount of the polar angle asymmetries can be accounted for by the organization of

cortical space in early visual cortex.

Methods

Reproducible computation and code sharing

All analyses were conducted in MATLAB (MathWorks, MA, USA). Data and code for our previously published and extended computational observer model, including density computations and figure scripts, are made publicly available via the Open Science Framework at the URL: https://osf.io/mygvu/ (previously published) and https://osf.io/ywu5v/ (this study).

Data sources

Data on cone density, midget RGC density, and V1 cortical surface area previously published or from publicly available analysis toolboxes. Both cone and mRGC densities were computed as cells/deg^2 for 0–40° eccentricities (step size 0.05°), at the cardinal meridians (0°, 90°, 180°, and 270° polar angle, corresponding to nasal, superior, temporal, and inferior retina of the left eye. **Fig 1** contains averaged cone and mRGC densities across all meridians as a function of eccentricity. **Fig 2** contains cone and mRGC densities converted to visual field coordinates, where the horizontal visual field meridian is the average of nasal and temporal retina, upper visual field meridian corresponds to the inferior retina and lower visual field meridian to the superior retina.

Cone density

Cone density data for the main results were extracted from post-mortem retinal tissue of 8 human retina's published by Curcio *et al.* [9] using the analysis toolbox ISETBIO [65-67], publicly available via GitHub (https://github.com/isetbio/isetbio).

Cone density in **Supplemental Fig 1** shows two datasets computed by two analysis toolboxes. To extract post-mortem data from Curcio *et al.* [9], we either use ISETBIO or the rgcDisplacementMap toolbox [76], publicly available at GitHub (https://github.com/gkaguirrelab/rgcDisplacementMap). A second cone density dataset comes from an adaptive optics study published by Song *et al.* [10]. From this work, we use "Group 1" (young individuals, 22-35 years old) implemented in ISETBIO.

Midget retinal ganglion cell receptive field density

Midget RGC density for the main results were computed with the quantitative model by Watson [64] implemented in ISETBIO. This model combines cone density data from Curcio *et al.* [9], mRGC cell body data from Curcio and Allen [53] and the displacement model by Drasdo *et al.* [57], to predict the midget RGC receptive fields (RFs).

Midget RGC data in **Supplemental Fig 1** computes mRGC density with two computational models: Watson [64] from ISETBIO and the displacement model by Barnett and Aguirre [76] implemented in the rgcDisplacementMap toolbox.

Cortical magnification factor in early visual cortex

To quantify the fovea-to-periphery gradient in the V1 cortical magnification factor (CMF), we used the areal CMF function published in Horton and Hoyt [68] for 0–40° eccentricity (Fig 1). Because this function does not make separate predictions for the cardinal meridians (Fig 2), we used data from the Human Connectome Project (HCP) 7 Tesla retinotopy dataset (*n*=163), which were first published by Ugurbil, van Essen, and colleagues [138, 139] and analyzed with population receptive field models by Benson *et al.* [79]). V1 CMF surface area data are from Benson *et al.* [78] segmented into bins using hand-drawn ROIs from Benson *et al.* [140] and computed as follows.

To compute V1 CMF from retinotopy data, we used the extracted surface area for ±10° and ±20° wedge ROIs centered on the cardinal meridians in each individual's hemisphere. The wedges on the horizontal, dorsal, and ventral locations represented the horizontal, lower, and upper visual field meridians respectively. Wedge ROIs were computed in the following steps: First, area V1 and V2 were manually labeled with iso-eccentricity and iso-polar angle contour lines using the measured retinotopic maps of each hemisphere [140]. Second, for each cardinal meridian and each 1°-eccentricity bin, we calculated the mean distance along the cortex to reach a 10° or 20° polar angle. All vertices that fell within the eccentricity bin and polar angle distance were included in the particular ROI. We computed wedge strips, rather than an entire wedge or line, to avoid localization errors in defining the exact boundaries.

The wedges were separated into 5 eccentricity bins between 1–6° (1° step size) using the hand-drawn ROIs from Benson *et al.* [140], marking eccentricity lines at 1°, 2°, 4°, and 7°. The 3°, 5° and 6° eccentricity lines were deduced from the 2°, 4° and 7° lines using isotropic interpolation (independently for ±10° and ±20° wedge ROIs, for more details see Benson *et al.*

[78]), and hence are likely to be less accurate than the data points at the exact hand-drawn

eccentricity lines. The cortical surface area (mm^2) was summed across hemispheres within each

subject and divided by the visual field area (deg^2). For each eccentricity bin and cardinal

meridian, mean and standard error V1 CMF were computed from bootstrapped data across

subjects (1,000 iterations). Mean data for each cardinal meridian were fit with a linear function in

log-log space (*i.e.*, power law function in linear coordinates) for 1–6° eccentricity.

The initial ROIs used for the upper and lower vertical meridian included both V1 and V2

sections of the vertical meridian, and therefore contain twice as much visual area as the

horizontal ROI. To have a fair comparison between the horizontal and upper and lower visual

field ROIs, we corrected the upper and lower ROIs as follows. For each subject and eccentricity

bin, we computed a vertical surface area ROI (with both upper and lower visual fields) that

excluded V2 sections of the vertical meridian. When summed over both hemispheres, this

vertical ROI has a size comparable to the horizontal ROI. We then calculated a scale factor for

each subject and eccentricity, by dividing the vertical ROI by the sum of upper and lower

surface area ROIs. This scale factor was on average ~0.5. To get the corrected V1 CMF, we

multiplied the scale factor to corresponding ventral and dorsal surface areas and divided by the

corresponding visual field area. By scaling dorsal and ventral ROIs to only include the V1-side,

we made the assumption that V2 is approximately the same size as V1. These vertical ROIs

may be slightly less precise than the horizontal meridian ROI and affect the horizontal-vertical

asymmetry (HVA). We did not compare differences in pRF sizes for the cardinal meridians.

Although the narrower ±10° wedge ROIs are in closer correspondence to the single line

estimations of cone and mRGC density, we use ±20° wedge ROIs in **Fig 2** as those data are

more robust. This is because narrow wedge ROIs are prone to overestimation of the vertical

meridian surface, caused by ipsilateral representations near the boundaries. Such ipsilateral

representations are sometimes incorrectly counted as part of the ±20° ROI for the ipsilateral

hemisphere, instead of as part of the ±10° ROI for the contralateral hemisphere, and this effect

is exacerbated for smaller wedges. We visualize V1 asymmetries for both ±10° and ±20° wedge

ROI **Supplementary Fig 1**.

Convergence ratios

The cone:mRGC ratio was computed by dividing mRGC density (cells/deg^2) by cone density

(cells/deg^2) for 0–40° eccentricity, in 0.05° bins. The mRGC:CMF ratio was computed in

cells/mm^2. When comparing mRGC density to Horton and Hoyt's CMF prediction, mRGC

density (cells/deg^2) was divided by V1 CMF (deg^2/mm^2) for 0–40° eccentricity, in 0.05° bins.

When comparing HCP's retinotopy CMF to mRGC density, mRGC density was restricted to 1–

6° eccentricity, and divided by the power law functions fitted to the V1 CMF. To compute the

transformation ratios relative to horizontal visual field meridian for cone:mRGC or mRGC:V1

CMF ratios in **Supplementary Fig 2**, we divide the lower and upper visual field transformation

ratio separately by the horizontal visual field transformation ratio.

Asymmetry computation

Polar angle asymmetries between meridians for cone density and mRGC density were

calculated as percent change in retinal coordinates as in Equation 1 and 2:

$$Horizontal\ Vertical\ Asymmetry = 100 \cdot \frac{mean(nasal, temporal) - mean(superior, inferior)}{mean(nasal, temporal, superior, inferior)} \quad \text{(Eq 1)}$$

$$Vertical\ Meridian\ Asymmetry = 100 \cdot \frac{superior - inferior}{mean(superior, inferior)} \quad \text{(Eq 2)}$$

Polar angle asymmetries in V1 CMF and behavior were computed with the same equations, but

for visual field coordinates (*i.e.*, nasal and temporal retina are left and right visual field

meridians, and superior and inferior retina are lower and upper visual field meridians).

Computational observer model

The computational observer uses and extends a published model [50]. The extensions include

(1) a phototransduction stage in the cone outer segment (transforming absorptions to

photocurrent) and (2) a midget RGC layer (transforming photocurrent to mRGC responses)

between the cone isomerization stage and the behavioral inference stage. To compensate for

the increase in computational load and to keep the model tractable, we also made two

simplifications: We used an L-cone only mosaic (instead of L-, M-, S-cone mosaic), and

removed any stimulus location uncertainty by omitting fixational eye movements and stimulus

phase shifts within a single stimulus orientation. With our extended model, we generated new

cone absorption and photocurrent data using a fixed random number generator.

Given that several stages of the model are identical to those to the previous study, we

refer to those methods on *Scene radiance*, *Retinal irradiance*, and *Cone mosaic and*

absorptions. Unlike in our previous study [50], we did not vary the level of defocus in the *Retinal*

irradiance stage nor the ratio of different cone types within a cone mosaic.

Stimulus parameters

The computational model simulates a 2-AFC orientation discrimination task while varying

stimulus contrast. The stimulus parameters are chosen to match the baseline condition of the

psychophysical study by Himmelberg *et al.* [15], whose results have replicated the

psychophysical study used for comparison in our previous computational observer model [13].

The recent psychophysics experiment used achromatic oriented Gabor patches, ±15° oriented

from vertical, with a spatial frequency of 4 cycles per degree. Stimuli were presented at 4.5° iso-

eccentric locations on the cardinal meridians, with a size of 3x3° visual angle (σ = 0.43°) and

duration of 120 ms. These stimulus parameters were identical to those the model, except for

size, duration, and phase randomization of the Gabor. The simulated stimulus by the model was

smaller (2x2° visual angle (σ = 0.25°), shorter (54-ms on, 2-ms sampling) followed by a 164-ms

blank period (mean luminance). We simulated these additional time points without a stimulus

because photocurrent data are temporally delayed (see next section on *Photocurrent*). There

was no stimulus onset period, and the phase of the Gabor patches were identical across all

trials (90°). Instead of simulating 5 experiments with 200 trials per stimulus orientation as in our

previous paper, we simulated one experiment with 5x more trials (*i.e.*, 1,000 trials per stimulus

orientation, 2,000 trials in total) to ensure that our behavioral inference stage had sufficient

number of trials to successfully learn and classify stimulus orientation. To assure psychometric

functions with lower and upper asymptotes, stimulus contrasts ranged from 0.05-100%*.*

Photocurrent

After the cone isomerization stage, we applied ISETBIO's build-in *osLinear* photocurrent

functionality implemented by Cottaris *et al*. [51] to our cone absorption data (separate for each

simulation varying in cone density). This photocurrent stage converts cone excitations into

photocurrent in pA in a linear manner (in contrast to the *osBiophys* functionality in ISETBIO

which contains a more complex and computationally intensive biophysical model to calculate

cone current).

The phototransduction stage takes the cone absorptions and applies three

computations. First, it convolves cone absorptions trials with a linear temporal impulse response

specific to L-cones (see **Fig 3**, panel in between absorptions and photocurrent stage). This

temporal filter delays and blurs the cone photocurrent in time. Second, photocurrent gain is

downregulated by light input, for instance due to increased luminance levels or larger cone

apertures. Third, photocurrents are subject to an additional source of white Gaussian noise,

which are determined by photocurrent measurement by [80] (for more details, see Cottaris *et al*.

[51]). This resulted in a 4D array with m rows by n columns by 109 2-ms time points by 2,000

trials.

Because our simulated experiments do not contain any uncertainty about the stimulus

location (no fixational eye movements or stimulus phase randomization), we were able to

average both cone absorptions and photocurrent data across stimulus time points. We

computed mean cone absorption data by taking the average across the first 54 ms (ignoring the

time points without stimulus). For mean cone photocurrent data, we took a weighted mean

across all 218 ms time points using a temporally delayed stimulus time course. This time course

was constructed by convolving the stimulus on-off boxcar with the temporal photocurrent filter.

This resulted in a 3D array with time-averaged cone photocurrent m rows by n columns by

2,000 trials.

Midget RGC layer

Prior to the mRGC layer, Gabor stimuli were simulated as spectral scene radiance from a visual

display, passed through the simulated human optics, subject to isomerization and

phototransduction by the cones in a rectangular mosaic (2x2° field-of-view) and saved as

separate files for each stimulus contrast. The mRGC layer loaded the simulated 2D cone

absorptions and photocurrent data.

The mRGC layer was built as a rectangular array, with the identical size mosaic as the

cone mosaic (2x2°). Spatial summation by RGC RFs was implemented as 2D Difference of

Gaussians (DoG) filters [81, 82]. The DoG RF was defined on a support of 31 rows by 31

columns. The DoG size was based on Croner and Kaplan [83]: the standard deviation of the

center Gaussian (σ_c) was 1/3 times the cone spacing and the standard deviation of the surround

Gaussian (σ_s) was 6 times the center standard deviation. The center/surround weights were

0.64:0.36, hence unbalanced. These parameters create neighboring DoG RFs that overlap at

1.3 standard deviation from their centers, approximating RGC tiling in human retina based on

overlap of dendrites fields [55]. The support of the DoG filter did not change size, however,

because the mRGC array is matched to the cone array and cone density affects cone spacing

(*i.e.*, a lower cone density results in a sparser array), the width of the DoG varies with cone

density and can be expressed in units of degree visual angle (*i.e.*, scaling with the number of

cones per degree within the cone array).

In the primate fovea, there is one ON and one OFF mRGC cell per cone, for a ratio of 2

mRGCs per cone. Unlike in the eye, our model mRGCs are not rectified, hence one of our

mRGCs can signal either increments or decrements. For comparison to the literature, we

multiply our mRGC counts by 2. We do not model on- and off-center mRGCs separately, but

rather consider one linear mRGC (no rectification) as a pair of rectified on- and off-centers. For

example, we consider an mRGC layer with no subsampling as having an mRGC:cone ratio of

2:1 (2 mRGCs per cone). The mRGC:cone ratios, counted in this way, were 2:1, 0.5:1, 0.22:1,

0.125:1, 0.08:1. The highest ratio (2:1) is similar to the observed in the fovea and the lowest

ratio (0.08:1) is similar to the observed at ~40° eccentricity [64]. We tested a wide range of

ratios because the purpose of the modeling was to assess how variation in mRGC density

affects performance. The relationships between cone density and performance, or between

mRGC:cone ratio and performance, are more robustly assessed by testing a wide range of

parameters.

The spatial computations of the mRGC layer were implemented in three stages. In the

first stage, the 2D DoG filter was convolved with each time-averaged 2D cone photocurrent

frame separately for each trial. The photocurrent images were padded to avoid border artifacts.

We padded the array with the mean of the photocurrent cone array, where the padding doubled

the width and height of the array. The post-convolution array maintained the same size as the

cone array without padding.

In the second stage, white Gaussian noise was added to each time point of the filtered

cone photocurrent response, sampled from a distribution with a standard deviation of 1. This

noise level was determined after testing a range of values showed that doubling or halving the

width of the Gaussian only scaled the absolute performance levels, not the effect as a function

of cone density or mRGC:cone ratios (for results using a standard deviation of 0.5 and 2, see

Supplementary Fig 4). We added noise to our mRGC responses at this stage, because our

mRGC layer without noise would perform a linear transform of the photocurrent responses

(linear filtering and linear subsampling). A transform that a linear support vector machine

classifier should be able to learn the optimal hyperplane with enough training trials to "untangle"

the two stimulus classes. This would mean that our model would not predict any loss of

information introduced by the mRGC layer, the effect we are most interested in. Had we used a

limited number of trials instead, our model would have performed suboptimally and showed

differences in classification accuracy. In such case, it would be difficult to distinguish the extent

to which these performance differences are caused by spatial variations in mRGCs on visual

performance versus the general ability of the SVM algorithm.

In the third stage, the filtered cone responses were linearly subsampled. This was

implemented by resampling each row and column of the filtered cone responses with a sample

rate equal to the mRGC:cone ratio. For instance, an array with an mRGC:cone ratio of 0.5:1

samples from every other cone. The mRGCs are centered on the cones, limiting the resampling

of filtered cone responses to integer numbers of cones. These spatially filtered and subsampled

responses are the mRGC responses in arbitrary units, as we added an arbitrary level of

Gaussian white noise on the filtered photocurrent responses and did not implement spiking non-

linearity in this transformation.

Simulated experiments

A single simulated experiment had a total of 64,000 trials: 2,000 trials per contrast level, 1,000

clockwise and 1,000 counter-clockwise. Stimulus contrast was systematically varied from 0 to

100% Michelson contrast, using 32 contrast levels. The cone mosaic was identical across

contrast levels, only including L-cones, cone density and cone spacing. There were no eye

movements. Cone absorptions and photocurrent simulations used a fixed random number

generator seed. Data from a single contrast level were represented as a 4D array (m rows by n

columns by 218 time points by 2,000 trials). The size of the m by n frame depended on the

defined subsampling ratio used for the mRGC layer.

This single experiment was repeated for 17 different cone mosaics, which varied

systematically in cone density and spacing. The cone density variation was implemented by

simulating cone mosaics at different eccentricities, ranging from a density as high as at the 1°

(4.9 x10^3 cells/deg^2) to as low as at 40° eccentricity on the horizontal meridian (0.047 x10^4

cells/deg^2). This resulted in a total of 1,088,000 simulated trials (64,000 trials x 17 cone

densities).

Simulated experiments for each of the 17 different cone densities were averaged across

time, resulting in a 3D array (m rows by n columns by 2,000 trials). In the mRGC layer, each 3D

array was spatially subsampled by 5 different mRGC:cone ratios. This resulted in a total of

5,440,000 simulated trials (64,000 trials x 17 cone densities x 5 ratios).

Inference engines

The simulated trials were fed into an inference engine. The task of the inference engine was to
classify if a trial contained a clockwise or counter-clockwise oriented Gabor stimulus given the
mRGC responses. Classification was performed separately for every 2,000 trials, *i.e.*,
separately for each contrast level, cone density, and mRGC:cone ratio.

We used a linear SVM classifier as implemented in MATLAB's *fitcsvm* with 10-fold
cross-validation and built-in z-scoring. This procedure is identical to our previously published
model [50]. In contrast to our previous model implementation, we did not transform each 2D
frame of mRGC responses to the Fourier domain and did not discard phase information prior to
classification, because the stimulus was static and did not contain any uncertainty about
stimulus location nor simulated fixational eye movements. The mRGC responses were
concatenated across space, resulting in a matrix of 2,000 trials by mRGC responses. The order
of the trials within this vector was randomized and fed into the linear SVM classifier with a set of
stimulus labels. The classifier trained its weights on 90% of the trials, and tested on the 10%
left-out trials. This resulted in accuracy (percent correct) for each given contrast level, cone
density and ratio.

Accuracy data for a single simulated experiment were fitted with a Weibull function to
extract the contrast threshold. The threshold was defined as the power of 1 over the slope of the
Weibull function, which comes out approximately ~80% correct, given that chance is 50% for a
2-AFC task and our slope was defined as $\beta = 3$.

Comparing model performance to behavior

To quantify the contribution of the spatial filtering by mRGCs, we compared the model
performance to behavior reported by Himmelberg *et al.* [15]. To do so, we extracted the mean
contrast thresholds across all simulated cone densities and mRGC:cone ratios. This resulted in
a matrix of 17 cone densities x 5 mRGC:cone ratios. We placed these data points in a 3D
coordinate space: log cone density (x-dimension) by log mRGC:cone ratio (y-dimension) by log
contrast thresholds (z-dimension). We fitted a 3D mesh using a regression with locally weighted
scatterplot smoothing with MATLAB's *fit.m* (using a LOWESS fit type with a span = 0.2, build-in
normalization and the 'bisquare' robust fitting options). This 3D mesh fit is used to visualize the
effect of cone density at a single mRGC:cone ratio by extracting a single curve from the mesh at
that particular ratio (**Fig 6A**). We then used the 3D mesh fit to predict contrast thresholds for the

four cardinal meridians at 4.5° eccentricity, evaluating the model at the four observed [cone,

mRGC:cone ratio]-density coordinates reported by Curcio *et al.* [9] and Watson [64].

Predicted thresholds for the model up to cone isomerizations and photocurrent were

computed using contrast thresholds for each cone density. These data were fitted separately

per model stage, with the same 3D mesh fit as mRGC responses using a dummy variable for

the mRGC:cone ratio. This fit was used to predict thresholds for each model stage given the

observed cone densities at the four cardinal meridians at 4.5° eccentricity.

Contrast thresholds were converted into contrast sensitivity by taking the reciprocal.

Nasal and temporal retina were averaged to represent the horizontal meridian. Because cone

density can vary dramatically across observers [141, 142], we computed error bars that

represent the amount of variability in predicted sensitivity based on a difference in underlying

cone density.

The upper/lower bound of the error bars in cone and mRGC model predictions were

defined by assuming that our estimates of cone density on the meridians are imperfect.

Specifically, we assumed that the measured asymmetries might be off by as much as a factor of

2. So, for example, if the reported density for the horizontal meridian is 20% above the mean,

and for the vertical meridian is 20% below the mean, we considered the possibility that they

were in fact 40% above or below the mean, or 10% above or below the mean.

References

1.   Wertheim T. Über die indirekte Sehschärfe. Z Psychol, Physiol. 1894;7:172-83.

2.   Ludvigh E. Extrafoveal acuity as measured with Snellen test-letters. American Journal of
Ophtalmology. 1941;24:225-33.

3.   Polyak SL. The retina. Press UoC, editor. Chicago1941.

4.   Strasburger H, Rentschler I, Juttner M. Peripheral vision and pattern recognition: a
review. J Vis. 2011;11(5):13. doi: 10.1167/11.5.13. PubMed PMID: 22207654.

5.   Hering E. Beiträge zur Physiologie [Contributions to physiology]. Leipzig: Wilhelm
Engelmann; 1861.

6.   von Helmholtz H. Handbuch der physiologischen Optik. Hamburg: Leopold Voss; 1896.

7.   Østerberg GA. Topography of the layer of rods and cones in the human retina. Acta
Ophthalmologica. 1935;13(6):1-102.

8.   Curcio CA, Sloan KR, Jr., Packer O, Hendrickson AE, Kalina RE. Distribution of cones in
human and monkey retina: individual variability and radial asymmetry. Science.
1987;236(4801):579-82. doi: 10.1126/science.3576186. PubMed PMID: 3576186.

9.   Curcio CA, Sloan KR, Kalina RE, Hendrickson AE. Human photoreceptor topography. J
Comp Neurol. 1990;292(4):497-523. doi: 10.1002/cne.902920402. PubMed PMID: 2324310.

10.   Song H, Chui TY, Zhong Z, Elsner AE, Burns SA. Variation of cone photoreceptor
packing density with retinal eccentricity and age. Invest Ophthalmol Vis Sci. 2011;52(10):7376-
84. doi: 10.1167/iovs.11-7199. PubMed PMID: 21724911; PubMed Central PMCID:
PMCPMC3183974.

11.   Mackeben M. Sustained focal attention and peripheral letter recognition. Spat Vis.
1999;12(1):51-72. PubMed PMID: 10195388.

12.   Carrasco M, Talgar CP, Cameron EL. Characterizing visual performance fields: effects
of transient covert attention, spatial frequency, eccentricity, task and set size. Spat Vis.
2001;15(1):61-75. Epub 2002/03/15. PubMed PMID: 11893125; PubMed Central PMCID:
PMCPMC4332623.

13.   Cameron EL, Tai JC, Carrasco M. Covert attention affects the psychometric function of
contrast sensitivity. Vision Res. 2002;42(8):949-67. Epub 2002/04/06. PubMed PMID:
11934448.

14.   Corbett JE, Carrasco M. Visual performance fields: frames of reference. PLoS One.
2011;6(9):e24470. doi: 10.1371/journal.pone.0024470. PubMed PMID: 21931727; PubMed
Central PMCID: PMCPMC3169603.

15.   Himmelberg MM, Winawer J, Carrasco M. Stimulus-dependent contrast sensitivity
asymmetries around the visual field. J Vis. 2020;20(9):18. Epub 2020/09/29. doi:
10.1167/jov.20.9.18. PubMed PMID: 32986805; PubMed Central PMCID: PMCPMC7533736.

16.   Barbot A, Xue S, Carrasco M. Asymmetries in visual acuity around the visual field. J Vis.
2021;21(1):2. Epub 2021/01/05. doi: 10.1167/jov.21.1.2. PubMed PMID: 33393963; PubMed
Central PMCID: PMCPMC7794272.

17.   Talgar CP, Carrasco M. Vertical meridian asymmetry in spatial resolution: visual and
attentional factors. Psychon Bull Rev. 2002;9(4):714-22. doi: 10.1093/mnras/94.5.377. PubMed
PMID: 12613674.

18.   Fuller S, Carrasco M. Perceptual consequences of visual performance fields: the case of
the line motion illusion. J Vis. 2009;9(4):13 1-7. doi: 10.1167/9.4.13. PubMed PMID: 19757922;
PubMed Central PMCID: PMCPMC3703960.

19.   Rovamo J, Virsu V. An estimation and application of the human cortical magnification
factor. Exp Brain Res. 1979;37(3):495-510. doi: 10.1007/BF00236819. PubMed PMID: 520439.

20.   Virsu V, Rovamo J. Visual resolution, contrast sensitivity, and the cortical magnification
factor. Exp Brain Res. 1979;37(3):475-94. PubMed PMID: 520438.

21.   Kroon JN, Rijsdijk JP, van der Wildt GJ. Peripheral contrast sensitivity for sine-wave
gratings and single periods. Vision Res. 1980;20(3):243-52. doi: 10.1016/0042-6989(80)90109-
1. PubMed PMID: 7385598.

22.   Rijsdijk JP, Kroon JN, van der Wildt GJ. Contrast sensitivity as a function of position on
the retina. Vision Res. 1980;20(3):235-41. doi: 10.1016/0042-6989(80)90108-X. PubMed PMID:
7385597.

23.   Robson JG, Graham N. Probability summation and regional variation in contrast
sensitivity across the visual field. Vision Res. 1981;21(3):409-18. doi: 10.1016/0042-
6989(81)90169-3. PubMed PMID: 7269319.

24.   Lundh BL, Lennerstrand G, Derefeldt G. Central and peripheral normal contrast
sensitivity for static and dynamic sinusoidal gratings. Acta Ophthalmol (Copenh).
1983;61(2):171-82. Epub 1983/04/01. doi: 10.1111/j.1755-3768.1983.tb01410.x. PubMed PMID:
6880630.

25.   Skrandies W. Human contrast sensitivity: regional retinal differences. Hum Neurobiol.
1985;4(2):97-9. PubMed PMID: 4030428.

26.   Seiple W, Holopigian K, Szlyk JP, Wu C. Multidimensional visual field maps:
relationships among local psychophysical and local electrophysiological measures. J Rehabil
Res Dev. 2004;41(3A):359-72. PubMed PMID: 15543452.

27.   Silva MF, Maia-Lopes S, Mateus C, Guerreiro M, Sampaio J, Faria P, et al. Retinal and
cortical patterns of spatial anisotropy in contrast sensitivity tasks. Vision Res. 2008;48(1):127-
35. doi: 10.1016/j.visres.2007.10.018. PubMed PMID: 18067943.

28.   Silva MF, Mateus C, Reis A, Nunes S, Fonseca P, Castelo-Branco M. Asymmetry of
visual sensory mechanisms: electrophysiological, structural, and psychophysical evidences. J
Vis. 2010;10(6):26. doi: 10.1167/10.6.26. PubMed PMID: 20884575.

29.   Abrams J, Nizam A, Carrasco M. Isoeccentric locations are not equivalent: the extent of
the vertical meridian asymmetry. Vision Res. 2012;52(1):70-8. Epub 2011/11/17. doi:
10.1016/j.visres.2011.10.016. PubMed PMID: 22086075; PubMed Central PMCID:
PMCPMC3345502.

30.   Baldwin AS, Meese TS, Baker DH. The attenuation surface for contrast sensitivity has
the form of a witch's hat within the central visual field. J Vis. 2012;12(11). Epub 2012/10/30. doi:
10.1167/12.11.23. PubMed PMID: 23104816.

31.   Silva MF, d'Almeida OC, Oliveiros B, Mateus C, Castelo-Branco M. Development and
aging of visual hemifield asymmetries in contrast sensitivity. J Vis. 2014;14(12). doi:
10.1167/14.12.19. PubMed PMID: 25326605.

32. Altpeter E, Mackeben M, Trauzettel-Klosinski S. The importance of sustained attention
for patients with maculopathies. Vision Res. 2000;40(10-12):1539-47. doi: 10.1016/S0042-
6989(00)00059-6. PubMed PMID: 10788657.

33. Carrasco M, Williams PE, Yeshurun Y. Covert attention increases spatial resolution with
or without masks: support for signal enhancement. J Vis. 2002;2(6):467-79. doi: 10.1167/2.6.4.
PubMed PMID: 12678645.

34. Montaser-Kouhsari L, Carrasco M. Perceptual asymmetries are preserved in short-term
memory tasks. Atten Percept Psychophys. 2009;71(8):1782-92. doi: 10.3758/APP.71.8.1782.
PubMed PMID: 19933562; PubMed Central PMCID: PMCPMC3697833.

35. Fuller S, Rodriguez RZ, Carrasco M. Apparent contrast differs across the vertical
meridian: visual and attentional factors. J Vis. 2008;8(1):16 1-. doi: 10.1167/8.1.16. PubMed
PMID: 18318619; PubMed Central PMCID: PMCPMC2789458.

36. Chaikin JD, Corbin HH, Volkmann J. Mapping a field of short-time visual search.
Science. 1962;138(3547):1327-8. doi: 10.1126/science.138.3547.1327. PubMed PMID:
14019856.

37. Krose BJ, Julesz B. The control and speed of shifts of attention. Vision Res.
1989;29(11):1607-19. PubMed PMID: 2635484.

38. Carrasco M, Giordano AM, McElree B. Temporal performance fields: visual and
attentional factors. Vision Res. 2004;44(12):1351-65. doi: 10.1016/j.visres.2003.11.026.
PubMed PMID: 15066395.

39. Rezec AA, Dobkins KR. Attentional weighting: a possible account of visual field
asymmetries in visual search? Spat Vis. 2004;17(4-5):269-93. PubMed PMID: 15559106.

40. Pretorius LL, Hanekom JJ. An accurate method for determining the conspicuity area
associated with visual targets. Hum Factors. 2006;48(4):774-84. doi:
10.1518/001872006779166370. PubMed PMID: 17240724.

41. Kristjansson A, Sigurdardottir HM. On the benefits of transient attention across the visual
field. Perception. 2008;37(5):747-64. doi: 10.1068/p5922. PubMed PMID: 18605148.

42. Najemnik J, Geisler WS. Eye movement statistics in humans are consistent with an
optimal search strategy. J Vis. 2008;8(3):4 1-14. doi: 10.1167/8.3.4. PubMed PMID: 18484810;
PubMed Central PMCID: PMCPMC2868380.

43. Najemnik J, Geisler WS. Simple summation rule for optimal fixation selection in visual
search. Vision Res. 2009;49(10):1286-94. doi: 10.1016/j.visres.2008.12.005. PubMed PMID:
19138697.

44. Fortenbaugh FC, Silver MA, Robertson LC. Individual differences in visual field shape
modulate the effects of attention on the lower visual field advantage in crowding. J Vis.
2015;15(2). doi: 10.1167/15.2.19. PubMed PMID: 25761337; PubMed Central PMCID:
PMCPMC4327314.

45. Toet A, Levi DM. The two-dimensional shape of spatial interaction zones in the
parafovea. Vision Res. 1992;32(7):1349-57. Epub 1992/07/01. doi: 10.1016/0042-
6989(92)90227-a. PubMed PMID: 1455707.

46. He S, Cavanagh P, Intriligator J. Attentional resolution and the locus of visual
awareness. Nature. 1996;383(6598):334-7. doi: 10.1038/383334a0. PubMed PMID: 8848045.

47.   Greenwood JA, Szinte M, Sayim B, Cavanagh P. Variations in crowding, saccadic
precision, and spatial localization reveal the shared topology of spatial vision. Proc Natl Acad
Sci U S A. 2017;114(17):E3573-E82. Epub 2017/04/12. doi: 10.1073/pnas.1615504114.
PubMed PMID: 28396415; PubMed Central PMCID: PMCPMC5410794.

48.   Roberts M, Cymerman R, Smith RT, Kiorpes L, Carrasco M. Covert spatial attention is
functionally intact in amblyopic human adults. J Vis. 2016;16(15):30. Epub 2016/12/30. doi:
10.1167/16.15.30. PubMed PMID: 28033433; PubMed Central PMCID: PMCPMC5215291.

49.   Purokayastha S, Roberts M, Carrasco M. Voluntary attention improves performance
similarly around the visual field. Atten Percept Psychophys. 2021;83(7):2784-94. Epub
2021/05/27. doi: 10.3758/s13414-021-02316-y. PubMed PMID: 34036535; PubMed Central
PMCID: PMCPMC8514247.

50.   Kupers ER, Carrasco M, Winawer J. Modeling visual performance differences 'around'
the visual field: A computational observer approach. PLoS Comput Biol. 2019;15(5):e1007063.
Epub 2019/05/28. doi: 10.1371/journal.pcbi.1007063. PubMed PMID: 31125331; PubMed
Central PMCID: PMCPMC6553792.

51.   Cottaris NP, Wandell BA, Rieke F, Brainard DH. A computational observer model of
spatial contrast sensitivity: Effects of photocurrent encoding, fixational eye movements, and
inference engine. J Vis. 2020;20(7):17. Epub 2020/07/22. doi: 10.1167/jov.20.7.17. PubMed
PMID: 32692826; PubMed Central PMCID: PMCPMC7424933.

52.   Drasdo N. Receptive field densities of the ganglion cells of the human retina. Vision Res.
1989;29(8):985-8. Epub 1989/01/01. doi: 10.1016/0042-6989(89)90113-2. PubMed PMID:
2629213.

53.   Curcio CA, Allen KA. Topography of ganglion cells in human retina. J Comp Neurol.
1990;300(1):5-25. Epub 1990/10/01. doi: 10.1002/cne.903000103. PubMed PMID: 2229487.

54.   Dacey DM, Petersen MR. Dendritic field size and morphology of midget and parasol
ganglion cells of the human retina. Proc Natl Acad Sci U S A. 1992;89(20):9666-70. doi:
10.1073/pnas.89.20.9666. PubMed PMID: 1409680; PubMed Central PMCID: PMCPMC50193.

55.   Dacey DM. The mosaic of midget ganglion cells in the human retina. J Neurosci.
1993;13(12):5334-55. PubMed PMID: 8254378.

56.   Sjostrand J, Popovic Z, Conradi N, Marshall J. Morphometric study of the displacement
of retinal ganglion cells subserving cones within the human fovea. Graefes Arch Clin Exp
Ophthalmol. 1999;237(12):1014-23. Epub 2000/02/02. doi: 10.1007/s004170050338. PubMed
PMID: 10654171.

57.   Drasdo N, Millican CL, Katholi CR, Curcio CA. The length of Henle fibers in the human
retina and a model of ganglion receptive field density in the visual field. Vision Res.
2007;47(22):2901-11. Epub 2007/02/27. doi: 10.1016/j.visres.2007.01.007. PubMed PMID:
17320143; PubMed Central PMCID: PMCPMC2077907.

58.   Liu Z, Kurokawa K, Zhang F, Lee JJ, Miller DT. Imaging and quantifying ganglion cells
and other transparent neurons in the living human retina. Proc Natl Acad Sci U S A.
2017;114(48):12803-8. Epub 2017/11/16. doi: 10.1073/pnas.1711734114. PubMed PMID:
29138314; PubMed Central PMCID: PMCPMC5715765.

59.   Webb SV, Kaas JH. The sizes and distribution of ganglion cells in the retina of the owl
monkey. Aotus trivirgatus. Vision Res. 1976;16(11):1247-54. Epub 1976/01/01. doi:
10.1016/0042-6989(76)90049-3. PubMed PMID: 827113.

60.   Perry VH, Oehler R, Cowey A. Retinal ganglion cells that project to the dorsal lateral
geniculate nucleus in the macaque monkey. Neuroscience. 1984;12(4):1101-23. PubMed PMID:
6483193.

61.   Perry VH, Cowey A. The ganglion cell and cone distributions in the monkey's retina:
Implications for central magnification factors. Vision Res. 1985;25(12):1795-810.

62.   Wassle H, Grunert U, Rohrenbeck J, Boycott BB. Cortical magnification factor and the
ganglion cell density of the primate retina. Nature. 1989;341(6243):643-6. Epub 1989/10/19. doi:
10.1038/341643a0. PubMed PMID: 2797190.

63.   Leventhal AG, Rodieck RW, Dreher B. Retinal ganglion cell classes in the Old World
monkey: morphology and central projections. Science. 1981;213(4512):1139-42. Epub
1981/09/04. doi: 10.1126/science.7268423. PubMed PMID: 7268423.

64.   Watson AB. A formula for human retinal ganglion cell receptive field density as a
function of visual field location. J Vis. 2014;14(7). doi: 10.1167/14.7.15. PubMed PMID:
24982468.

65.   Farrell JE, Winawer J, Brainard DH, Wandell B. Modeling visible differences: The
computational observer model. SID Symposium Digest of Technical Papers2014. p. 352-6.

66.   Brainard DH, Jiang H, Cottaris NP, Rieke F, Chichilnisky EJ, Farrell JE, et al. ISETBIO:
Computational tools for modeling early human vision. Imaging and Applied Optics 2015;
2015/06/07; Arlington, Virginia: Optical Society of America; 2015.

67.   Cottaris NP, Jiang H, Ding X, Wandell BA, Brainard DH. A computational-observer
model of spatial contrast sensitivity: Effects of wave-front-based optics, cone-mosaic structure,
and inference engine. J Vis. 2019;19(4):8. doi: 10.1167/19.4.8. PubMed PMID: 30943530.

68.   Horton JC, Hoyt WF. The representation of the visual field in human striate cortex. A
revision of the classic Holmes map. Archives of Ophthalmology. 1991;109(6):816-24. PubMed
PMID: 2043069.

69.   Polyak S. The main afferent fiber systems of the cerebral cortex in primates. Berkeley:
University of California; 1932.

70.   Polyak S. A contribution of the cerebral representation of the retina. The Journal of
Comparative Neurology. 1933;57:541-617.

71.   Pointer JS. The cortical magnification factor and photopic vision. Biol Rev Camb Philos
Soc. 1986;61(2):97-119. Epub 1986/05/01. doi: 10.1111/j.1469-185x.1986.tb00463.x. PubMed
PMID: 3527286.

72.   Tolhurst DJ, Ling L. Magnification factors and the organization of the human striate
cortex. Hum Neurobiol. 1988;6(4):247-54. Epub 1988/01/01. PubMed PMID: 2832355.

73.   Adams DL, Horton JC. A precise retinotopic map of primate striate cortex generated
from the representation of angioscotomas. J Neurosci. 2003;23(9):3771-89. Epub 2003/05/09.
PubMed PMID: 12736348.

74.   Myerson J, Manis PB, Miezin FM, Allman JM. Magnification in striate cortex and retinal
ganglion cell layer of owl monkey: a quantitative comparison. Science. 1977;198(4319):855-7.
Epub 1977/11/25. doi: 10.1126/science.411172. PubMed PMID: 411172.

75.   Van Essen DC, Newsome WT, Maunsell JH. The visual field representation in striate
cortex of the macaque monkey: asymmetries, anisotropies, and individual variability. Vision

Res. 1984;24(5):429-48. Epub 1984/01/01. doi: 10.1016/0042-6989(84)90041-5. PubMed
PMID: 6740964.

76. Barnett M, Aguirre GK. A spatial model of human retinal cell densities and solution for
retinal ganglion cell displacement. Vision Sciences Society Annual Meeting; St. Pete Beach,
FL, USA: Journal of Vision; 2018. p. 23.

77. Fahle M, Schmid M. Naso-temporal asymmetry of visual perception and of the visual
cortex. Vision Res. 1988;28(2):293-300. Epub 1988/01/01. doi: 10.1016/0042-6989(88)90157-5.
PubMed PMID: 3414016.

78. Benson NC, Kupers ER, Barbot A, Carrasco M, Winawer J. Cortical magnification in
human visual cortex parallels task performance around the visual field. Elife. 2021;10. Epub
2021/08/04. doi: 10.7554/eLife.67685. PubMed PMID: 34342581; PubMed Central PMCID:
PMCPMC8378846.

79. Benson NC, Jamison KW, Arcaro MJ, Vu AT, Glasser MF, Coalson TS, et al. The
Human Connectome Project 7 Tesla retinotopy dataset: Description and population receptive
field analysis. J Vis. 2018;18(13):23. Epub 2018/12/29. doi: 10.1167/18.13.23. PubMed PMID:
30593068; PubMed Central PMCID: PMCPMC6314247.

80. Angueyra JM, Rieke F. Origin and effect of phototransduction noise in primate cone
photoreceptors. Nature neuroscience. 2013;16(11):1692-700. Epub 2013/10/08. doi:
10.1038/nn.3534. PubMed PMID: 24097042; PubMed Central PMCID: PMCPMC3815624.

81. Rodieck RW. Quantitative analysis of cat retinal ganglion cell response to visual stimuli.
Vision Res. 1965;5(11):583-601. Epub 1965/12/01. doi: 10.1016/0042-6989(65)90033-7.
PubMed PMID: 5862581.

82. Enroth-Cugell C, Robson JG. The contrast sensitivity of retinal ganglion cells of the cat.
J Physiol. 1966;187(3):517-52. Epub 1966/12/01. doi: 10.1113/jphysiol.1966.sp008107.
PubMed PMID: 16783910; PubMed Central PMCID: PMCPMC1395960.

83. Croner LJ, Kaplan E. Receptive fields of P and M ganglion cells across the primate
retina. Vision Res. 1995;35(1):7-24. doi: 10.1016/0042-6989(94)E0066-T. PubMed PMID:
7839612.

84. Green DG. Regional variations in the visual acuity for interference fringes on the retina. J
Physiol. 1970;207(2):351-6. Epub 1970/04/01. doi: 10.1113/jphysiol.1970.sp009065. PubMed
PMID: 5499023; PubMed Central PMCID: PMCPMC1348710.

85. Williams DR. Aliasing in human foveal vision. Vision Res. 1985;25(2):195-205. Epub
1985/01/01. doi: 10.1016/0042-6989(85)90113-0. PubMed PMID: 4013088.

86. Coletta NJ, Williams DR. Psychophysical estimate of extrafoveal cone spacing. J Opt
Soc Am A. 1987;4(8):1503-13. Epub 1987/08/01. PubMed PMID: 3625330.

87. Williams DR, Coletta NJ. Cone spacing and the visual resolution limit. J Opt Soc Am A.
1987;4(8):1514-23. Epub 1987/08/01. PubMed PMID: 3625331.

88. Anderson SJ, Hess RF. Post-receptoral undersampling in normal human peripheral
vision. Vision Res. 1990;30(10):1507-15. Epub 1990/01/01. doi: 10.1016/0042-6989(90)90031-f.
PubMed PMID: 2247960.

89. Anderson SJ, Mullen KT, Hess RF. Human peripheral spatial resolution for achromatic
and chromatic stimuli: limits imposed by optical and retinal factors. J Physiol. 1991;442:47-64.
Epub 1991/10/01. PubMed PMID: 1798037; PubMed Central PMCID: PMCPMC1179877.

90.   Banks MS, Sekuler AB, Anderson SJ. Peripheral spatial vision: limits imposed by optics,
photoreceptors, and receptor pooling. J Opt Soc Am A. 1991;8(11):1775-87. Epub 1991/11/01.
PubMed PMID: 1744774.

91.   Sjostrand J, Olsson V, Popovic Z, Conradi N. Quantitative estimations of foveal and
extra-foveal retinal circuitry in humans. Vision Res. 1999;39(18):2987-98. Epub 2000/02/09. doi:
10.1016/s0042-6989(99)00030-9. PubMed PMID: 10664798.

92.   Popovic Z, Sjostrand J. Resolution, separation of retinal ganglion cells, and cortical
magnification in humans. Vision Res. 2001;41(10-11):1313-9. Epub 2001/04/27. doi:
10.1016/s0042-6989(00)00290-x. PubMed PMID: 11322976.

93.   Popovic Z, Sjostrand J. The relation between resolution measurements and numbers of
retinal ganglion cells in the same human subjects. Vision Res. 2005;45(17):2331-8. Epub
2005/06/01. doi: 10.1016/j.visres.2005.02.013. PubMed PMID: 15924946.

94.   Wilkinson MO, Anderson RS, Bradley A, Thibos LN. Neural bandwidth of veridical
perception across the visual field. J Vis. 2016;16(2):1. Epub 2016/01/30. doi: 10.1167/16.2.1.
PubMed PMID: 26824638; PubMed Central PMCID: PMCPMC5833322.

95.   Anstis SM. Letter: A chart demonstrating variations in acuity with retinal position. Vision
Res. 1974;14(7):589-92. Epub 1974/07/01. doi: 10.1016/0042-6989(74)90049-2. PubMed
PMID: 4419807.

96.   Duncan RO, Boynton GM. Cortical magnification within human primary visual cortex
correlates with acuity thresholds. Neuron. 2003;38(4):659-71. PubMed PMID: 12765616.

97.   Carrasco M, Frieder KS. Cortical magnification neutralizes the eccentricity effect in
visual search. Vision Res. 1997;37(1):63-82. Epub 1997/01/01. doi: 10.1016/s0042-
6989(96)00102-2. PubMed PMID: 9068831.

98.   Carrasco M, McLean TL, Katz SM, Frieder KS. Feature asymmetries in visual search:
effects of display duration, target eccentricity, orientation and spatial frequency. Vision Res.
1998;38(3):347-74. Epub 1998/04/16. doi: 10.1016/s0042-6989(97)00152-1. PubMed PMID:
9536360.

99.   Schwarzkopf DS, Song C, Rees G. The surface area of human V1 predicts the
subjective experience of object size. Nature neuroscience. 2011;14(1):28-30. doi:
10.1038/nn.2706. PubMed PMID: 21131954; PubMed Central PMCID: PMC3012031.

100.   Liu T, Heeger DJ, Carrasco M. Neural correlates of the visual vertical meridian
asymmetry. J Vis. 2006;6(11):1294-306. Epub 2007/01/11. doi: 10.1167/6.11.12. PubMed
PMID: 17209736; PubMed Central PMCID: PMCPMC1864963.

101.   Benson NC, Butt OH, Datta R, Radoeva PD, Brainard DH, Aguirre GK. The retinotopic
organization of striate cortex is well predicted by surface topology. Current biology : CB.
2012;22(21):2081-5. doi: 10.1016/j.cub.2012.09.014. PubMed PMID: 23041195; PubMed
Central PMCID: PMC3494819.

102.   Silva MF, Brascamp JW, Ferreira S, Castelo-Branco M, Dumoulin SO, Harvey BM.
Radial asymmetries in population receptive field size and cortical magnification factor in early
visual cortex. Neuroimage. 2018;167:41-52. doi: 10.1016/j.neuroimage.2017.11.021. PubMed
PMID: 29155078.

103.   Himmelberg MM, Winawer J, Carrasco M. Linking contrast sensitivity to cortical
magnification in human primary visual cortex. bioRxiv. 2021;10.04.463138. doi:
https://doi.org/10.1101/2021.10.04.463138.

104.   Daniel PM, Whitteridge D. The representation of the visual field on the cerebral cortex in
monkeys. J Physiol. 1961;159:203-21. Epub 1961/12/01. doi: 10.1113/jphysiol.1961.sp006803.
PubMed PMID: 13883391; PubMed Central PMCID: PMCPMC1359500.

105.   Rolls ET, Cowey A. Topography of the retina and striate cortex and its relationship to
visual acuity in rhesus monkeys and squirrel monkeys. Exp Brain Res. 1970;10(3):298-310.
Epub 1970/01/01. doi: 10.1007/BF00235053. PubMed PMID: 4986000.

106.   Braccini C, Gambardella G, Sandini G, Tagliasco V. A model of the early stages of the
human visual system: functional and topological transformations performed in the peripheral
visual field. Biol Cybern. 1982;44(1):47-58. Epub 1982/01/01. doi: 10.1007/BF00353955.
PubMed PMID: 7093369.

107.   Schutt HH, Wichmann FA. An image-computable psychophysical spatial vision model. J
Vis. 2017;17(12):12. Epub 2017/10/21. doi: 10.1167/17.12.12. PubMed PMID: 29053781.

108.   Bradley C, Abrams J, Geisler WS. Retina-V1 model of detectability across the visual
field. J Vis. 2014;14(12). Epub 2014/10/23. doi: 10.1167/14.12.22. PubMed PMID: 25336179;
PubMed Central PMCID: PMCPMC4204678.

109.   Watson AB, Ahumada AJ. Letter identification and the neural image classifier. J Vis.
2015;15(2). Epub 2015/03/12. doi: 10.1167/15.2.15. PubMed PMID: 25761333.

110.   Kwon M, Liu R. Linkage between retinal ganglion cell density and the nonuniform spatial
integration across the visual field. Proc Natl Acad Sci U S A. 2019;116(9):3827-36. Epub
2019/02/10. doi: 10.1073/pnas.1817076116. PubMed PMID: 30737290; PubMed Central
PMCID: PMCPMC6397585.

111.   Himmelberg MM, Kurzawski JW, Benson NC, Pelli DG, Carrasco M, Winawer J. Cross-
dataset reproducibility of human retinotopic maps. Neuroimage. 2021;244:118609. Epub
2021/09/29. doi: 10.1016/j.neuroimage.2021.118609. PubMed PMID: 34582948; PubMed
Central PMCID: PMCPMC8560578.

112.   Perry VH, Cowey A. The lengths of the fibres of Henle in the retina of macaque
monkeys: implications for vision. Neuroscience. 1988;25(1):225-36. Epub 1988/04/01. doi:
10.1016/0306-4522(88)90021-8. PubMed PMID: 3393279.

113.   Tootell RB, Switkes E, Silverman MS, Hamilton SL. Functional anatomy of macaque
striate cortex. II. Retinotopic organization. The Journal of neuroscience : the official journal of
the Society for Neuroscience. 1988;8(5):1531-68. Epub 1988/05/01. PubMed PMID: 3367210;
PubMed Central PMCID: PMCPMC6569212.

114.   Andrews TJ, Halpern SD, Purves D. Correlated size variations in human visual cortex,
lateral geniculate nucleus, and optic tract. The Journal of neuroscience : the official journal of
the Society for Neuroscience. 1997;17(8):2859-68. PubMed PMID: 9092607.

115.   Kastner S, Schneider KA, Wunderlich K. Beyond a relay nucleus: neuroimaging views
on the human LGN. Progress in brain research. 2006;155:125-43. doi: 10.1016/S0079-
6123(06)55008-3. PubMed PMID: 17027384.

116.   Malpeli JG, Baker FH. The representation of the visual field in the lateral geniculate
nucleus of Macaca mulatta. J Comp Neurol. 1975;161(4):569-94. Epub 1975/06/15. doi:
10.1002/cne.901610407. PubMed PMID: 1133232.

117.   Connolly M, Van Essen D. The representation of the visual field in parvicellular and
magnocellular layers of the lateral geniculate nucleus in the macaque monkey. J Comp Neurol.
1984;226(4):544-64. Epub 1984/07/10. doi: 10.1002/cne.902260408. PubMed PMID: 6747034.

118.   Schein SJ, de Monasterio FM. Mapping of retinal and geniculate neurons onto striate
cortex of macaque. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 1987;7(4):996-1009. Epub 1987/04/01. PubMed PMID: 3033167; PubMed
Central PMCID: PMCPMC6568992.

119.   Azzopardi P, Cowey A. Preferential representation of the fovea in the primary visual
cortex. Nature. 1993;361(6414):719-21. Epub 1993/02/25. doi: 10.1038/361719a0. PubMed
PMID: 7680108.

120.   Malpeli JG, Lee D, Baker FH. Laminar and retinotopic organization of the macaque
lateral geniculate nucleus: magnocellular and parvocellular magnification functions. J Comp
Neurol. 1996;375(3):363-77. Epub 1996/11/18. doi: 10.1002/(SICI)1096-
9861(19961118)375:3<363::AID-CNE2>3.0.CO;2-0. PubMed PMID: 8915836.

121.   Shriki O, Kohn A, Shamir M. Fast coding of orientation in primary visual cortex. PLoS
Comput Biol. 2012;8(6):e1002536. Epub 2012/06/22. doi: 10.1371/journal.pcbi.1002536.
PubMed PMID: 22719237; PubMed Central PMCID: PMCPMC3375217.

122.   Barlow HB. Single units and sensation: a neuron doctrine for perceptual psychology?
Perception. 1972;1(4):371-94. Epub 1972/01/01. doi: 10.1068/p010371. PubMed PMID:
4377168.

123.   Hochstein S, Shapley RM. Linear and nonlinear spatial subunits in Y cat retinal ganglion
cells. J Physiol. 1976;262(2):265-84. Epub 1976/11/01. doi: 10.1113/jphysiol.1976.sp011595.
PubMed PMID: 994040; PubMed Central PMCID: PMCPMC1307643.

124.   Demb JB, Haarsma L, Freed MA, Sterling P. Functional circuitry of the retinal ganglion
cell's nonlinear receptive field. The Journal of neuroscience : the official journal of the Society
for Neuroscience. 1999;19(22):9756-67. Epub 1999/11/13. PubMed PMID: 10559385; PubMed
Central PMCID: PMCPMC6782950.

125.   Schwartz GW, Okawa H, Dunn FA, Morgan JL, Kerschensteiner D, Wong RO, et al. The
spatial structure of a nonlinear receptive field. Nature neuroscience. 2012;15(11):1572-80. Epub
2012/09/25. doi: 10.1038/nn.3225. PubMed PMID: 23001060; PubMed Central PMCID:
PMCPMC3517818.

126.   Shah NP, Brackbill N, Rhoades C, Kling A, Goetz G, Litke AM, et al. Inference of
nonlinear receptive field subunits with spike-triggered clustering. Elife. 2020;9. Epub
2020/03/10. doi: 10.7554/eLife.45743. PubMed PMID: 32149600; PubMed Central PMCID:
PMCPMC7062463.

127.   Schwartz G, Rieke F. Perspectives on: information and coding in mammalian sensory
physiology: nonlinear spatial encoding by retinal ganglion cells: when 1 + 1 not equal 2. J Gen
Physiol. 2011;138(3):283-90. Epub 2011/08/31. doi: 10.1085/jgp.201110629. PubMed PMID:
21875977; PubMed Central PMCID: PMCPMC3171084.

128.   Pillow JW, Shlens J, Paninski L, Sher A, Litke AM, Chichilnisky EJ, et al. Spatio-
temporal correlations and visual signalling in a complete neuronal population. Nature.
2008;454(7207):995-9. Epub 2008/07/25. doi: 10.1038/nature07140. PubMed PMID: 18650810;
PubMed Central PMCID: PMCPMC2684455.

129.   Ala-Laurila P, Greschner M, Chichilnisky EJ, Rieke F. Cone photoreceptor contributions
to noise and correlations in the retinal output. Nature neuroscience. 2011;14(10):1309-16. Epub
2011/09/20. doi: 10.1038/nn.2927. PubMed PMID: 21926983; PubMed Central PMCID:
PMCPMC3183110.

130.   Shapley R, Perry VH. Cat and monkey retinal ganglion cells and their visual functional
roles. Trends in Neurosciences. 1986;9:229-35.

131.   Rauschecker AM, Bowen RF, Parvizi J, Wandell BA. Position sensitivity in the visual
word form area. Proc Natl Acad Sci U S A. 2012;109(24):E1568-77. doi:
10.1073/pnas.1121304109. PubMed PMID: 22570498; PubMed Central PMCID: PMC3386120.

132.   Gibaldi A, Benson NC, Banks MS. Crossed-uncrossed projections from primate retina
are adapted to disparities of natural scenes. Proc Natl Acad Sci U S A. 2021;118(7). Epub
2021/02/13. doi: 10.1073/pnas.2015651118. PubMed PMID: 33574061; PubMed Central
PMCID: PMCPMC7896330.

133.   Aghajari S, Vinke LN, Ling S. Population spatial frequency tuning in human early visual
cortex. J Neurophysiol. 2020;123(2):773-85. Epub 2020/01/16. doi: 10.1152/jn.00291.2019.
PubMed PMID: 31940228; PubMed Central PMCID: PMCPMC7052645.

134.   Campbell FW, Green DG. Monocular versus binocular visual acuity. Nature.
1965;208(5006):191-2. Epub 1965/10/09. doi: 10.1038/208191a0. PubMed PMID: 5884255.

135.   Hubel DH, Wiesel TN. Receptive fields, binocular interaction and functional architecture
in the cat's visual cortex. J Physiol. 1962;160:106-54. PubMed PMID: 14449617; PubMed
Central PMCID: PMCPMC1359523.

136.   Banton T, Levi DM. Binocular summation in vernier acuity. J Opt Soc Am A.
1991;8(4):673-80. PubMed PMID: 2045969.

137.   Dougherty K, Cox MA, Westerberg JA, Maier A. Binocular Modulation of Monocular V1
Neurons. Current biology : CB. 2019;29(3):381-91 e4. doi: 10.1016/j.cub.2018.12.004. PubMed
PMID: 30661798; PubMed Central PMCID: PMCPMC6363852.

138.   Ugurbil K, Xu J, Auerbach EJ, Moeller S, Vu AT, Duarte-Carvajalino JM, et al. Pushing
spatial and temporal resolution for functional and diffusion MRI in the Human Connectome
Project. Neuroimage. 2013;80:80-104. Epub 2013/05/25. doi:
10.1016/j.neuroimage.2013.05.012. PubMed PMID: 23702417; PubMed Central PMCID:
PMCPMC3740184.

139.   Van Essen DC, Smith SM, Barch DM, Behrens TE, Yacoub E, Ugurbil K, et al. The WU-
Minn Human Connectome Project: an overview. Neuroimage. 2013;80:62-79. doi:
10.1016/j.neuroimage.2013.05.041. PubMed PMID: 23684880; PubMed Central PMCID:
PMC3724347.

140.   Benson NC, Yoon JMD, Forenzo D, Engel SA, Kay KN, Winawer J. Variability of the
Surface Area of the V1, V2, and V3 Maps in a Large Sample of Human Observers. BioRXiv.
2021. doi: https://doi.org/10.1101/2020.12.30.424856.

141.   Roorda A, Williams DR. The arrangement of the three cone classes in the living human
eye. Nature. 1999;397(6719):520-2. Epub 1999/02/24. doi: 10.1038/17383. PubMed PMID:
10028967.

142.   Hofer H, Carroll J, Neitz J, Neitz M, Williams DR. Organization of the human trichromatic
cone mosaic. The Journal of neuroscience : the official journal of the Society for Neuroscience.
2005;25(42):9669-79. Epub 2005/10/21. doi: 10.1523/JNEUROSCI.2414-05.2005. PubMed
PMID: 16237171.

1433

1434

Supplementary Material

1436

Supplementary Fig 1. Polar angle asymmetries for cone density, mRGC density and V1-V2 surface area computed from different publicly available datasets. Asymmetries are in percent change, calculated as the difference between horizontal and vertical meridians divided by their mean (left column), the difference between upper and lower vertical meridians divided by their means (right column). Positive asymmetries would positively correlate with observed differences in behavior. (Top row) Cone data are from either Curcio *et al.* [9] (black lines) or Song *et al.* [10] (orange line) computed with either ISETBIO (solid lines) or rgcDisplacementMap toolbox (dotted lines). (Middle row) Midget RGC RF data are computed using the computational model by Watson (2014) implemented in the ISETBIO toolbox (solid black line) or Barnett and Aguirre [76] implemented in the rgcDisplacementMap toolbox (dotted black line). (Bottom row) V1-V2 surface is computed from the Human Connectome Project 7T retinotopy dataset (*n*=163), using the analyzed dataset by Benson *et al.* [78, 79]. Surface areas are defined as ±10° (black) and ±20° (red) wedge ROIs from 1-6° eccentricity around the meridians, avoiding the central one degree and stimulus border (7-8°) as those data can be noisy. Note that the x-axis is truncated as cortical measurements are limited by the field-of-view in the fMRI experiment. Data are fit with a 2nd degree polynomial, $R^2 = 0.48$ (±10°) and $R^2 = 0.89$ (±20°) for horizontal-vertical and $R^2 = 0.94$ (±10°) and $R^2 = 0.72$ (±20°) for vertical-meridian asymmetries).

1452

Supplementary Fig 2. Transformation ratios relative to horizontal visual field meridian. Relative ratio is
computed taking the lower or upper visual field transformation ratio and horizontal visual field transformation ratio
from panel B, and divide the two for cone:mRGC ratios (left panel) and mRGC:V1 CMF ratios (right panel).

1456

1457

1458 **Supplementary Fig 3. Classifier performance varying with cone density, separately for each mRGC:cone**
1459 **ratio.** Linear SVM classifier accuracy is computed for each contrast level in a simulated experiment with 1,000
1460 clockwise and 1,000 counter-clockwise trials. Average accuracy data are fitted with a Weibull function.

1461
Supplementary Fig 4. The effect of noise in mRGC layer on contrast thresholds as a function of cone density,
separately for each mRGC:cone ratio. (A) Contrast thresholds as a function of cone density when adding white
noise following a Gaussian distribution with a standard deviation of 0.5 (left panel), 1 (middle panel), 2 (right panel).
Data are fit with a locally weighted regression using the same procedure as the fit shown in Fig 6. Middle panel (1
std) is identical to Fig 6A. (B) Same data as panel A, visualizing the three mRGC noise levels separately per
mRGC:cone ratio. Decreasing opacity of fits and data correspond to decreasing levels of noise.