Title:

Cell-Type-Specific Decrease of the Intrinsic Excitability of Motor Cortical Pyramidal

Neurons in Parkinsonian Mice

Abbreviated title:

Intrinsic Adaptations of M1 Pyramidal Cells in PD

Authors: Liqiang Chen, Yerim Kim, Hong-Yuan Chu

Affiliation: Center for Neurodegenerative Science, Van Andel Institute

Address: 333 Bostwick Ave, N.E., Grand Rapids, MI 49503

Correspondence: Hong-Yuan Chu, Ph.D., email: hongyuan.chu@vai.org

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by fund from Van Andel Institute

(H.Y.C). The authors thank Mr. David Nadziejka for technical editing.

Abstract

The hypokinetic motor symptoms of Parkinson's disease (PD) are closely linked with a

decreased motor cortical output as a consequence of elevated basal ganglia inhibition.

However, whether and how the loss of dopamine alters the cellular properties of motor

cortical neurons in PD remains undefined. We induced experimental parkinsonism in

adult C57BL6 mice of both sexes by injecting neurotoxin, 6-hydroxydopamine, into the

medial forebrain bundle. By using *ex vivo* patch-clamp recording and retrograde tracing

approach, we found that the intrinsic excitability of pyramidal tract neurons (PTNs) in the

motor cortical layer 5b was greatly decreased following the degeneration of midbrain

dopaminergic neurons; but the intratelencephalic neurons (ITNs) were not affected. The

cell-type-specific intrinsic adaptations were associated with a significant broadening of

the action potentials in PTNs but not in ITNs. Moreover, the loss of midbrain

dopaminergic neurons impaired the capability of M1 PTNs to sustain high-frequency

firing, which could underlie their abnormal pattern of activity in the parkinsonian state.

We also showed that the decreased excitability and broadened action potentials were

largely caused by a disrupted function of the large conductance, Ca^{2+}-activated K^+

channels. The restoration of dopaminergic neuromodulation failed to rescue the

impaired intrinsic excitability of M1 PTNs in parkinsonian mice. Altogether, our data

show cell-type-specific decreases of the excitability of M1 pyramidal neurons following

the loss of midbrain dopaminergic neurons. Thus, intrinsic adaptations in the motor

cortex, together with pathological basal ganglia inhibition, underlie the decreased motor

cortical output in parkinsonian state and exacerbate parkinsonian motor deficits.

Significance statement

The degeneration of midbrain dopaminergic neurons in Parkinson's disease remodels

the connectivity and function of cortico–basal ganglia–thalamocortical network.

However, whether and how the loss of dopamine and aberrant basal ganglia activity

alter motor cortical circuitry remain undefined. We found that pyramidal neurons in the

layer 5b of the primary motor cortex (M1) exhibit distinct adaptations in response to the

loss of midbrain dopaminergic neurons, depending on their long-range projections.

Besides the decreased thalamocortical synaptic excitation as proposed by the classical

model of Parkinson's pathophysiology, these results, for the first time, show novel

cellular and molecular mechanisms underlying the abnormal motor cortical output in

parkinsonian state.

Introduction

The degeneration of dopamine (DA) neurons in the substantia nigra (SN) alters the

connection and computation of cortico–basal ganglia–thalamocortical network, which

underlies the devastating motor symptoms in Parkinson's disease (PD), including

akinesia, bradykinesia, and rigidity (Albin et al., 1989; Galvan and Wichmann, 2008;

McGregor and Nelson, 2019). Specially, the loss of SN DA neurons increases and

decreases the activities of indirect and direct pathways, respectively, which disrupts the

balanced activity between striatal direct and indirect pathways, leading to the

hypokinetic symptoms in PD (Albin et al., 1989; DeLong, 1990). The pathway-specific

alterations in the striatum following the loss of SN DA neurons induce numerous cellular

and synaptic changes in the basal ganglia nuclei and extended brain regions, which

further drive the abnormal neural activity throughout the cortico–basal ganglia–

thalamocortical network in experimental parkinsonism (Gittis et al., 2011; Kita and Kita,

2011; Fieblinger et al., 2014; Chu et al., 2015, 2017; Mathai et al., 2015; Shen et al.,

2015; Parker et al., 2016, 2018; Sharott et al., 2017; McIver et al., 2019; Willard et al.,

2019).

The primary motor cortex (M1) plays essential and complex roles in motor control

and motor learning and is a key node in the cortico–basal ganglia–thalamocortical

network (Shepherd, 2013; Ebbesen and Brecht, 2017; Ebbesen et al., 2018). M1 is a

laminar structure and contains a heterogeneous group of neurons that differ in gene

expression, morphology, connectivity, and electrophysiological properties (Sheets et al.,

2011; Oswald et al., 2013; Shepherd, 2013; Suter et al., 2013; Economo et al., 2018).

These neurons can be classified into the intratelencephalic neurons (ITNs) distributed

across the layers (L) 2-6 and the pyramidal tract neurons (PTNs) that mainly locate

within the L5b (Oswald et al., 2013; Shepherd, 2013). Both PTNs and ITNs project to

the striatum and receive basal ganglia feedbacks through the transition of the motor

thalamus (Bodor et al., 2008; Kita and Kita, 2011; Kress et al., 2013; Lee et al., 2020).

In physiological state, M1 network dynamics plays an essential role in the execution and

coordination of complex movements and the acquisition of motor skills (Guo et al.,

2015a; Kawai et al., 2015; Sreenivasan et al., 2016; Barthas and Kwan, 2017; Ebbesen

et al., 2017; Wang et al., 2017; Economo et al., 2018; Sauerbrei et al., 2020). In

parkinsonian state, M1 exhibits an aberrant oscillation and bursting pattern of activity at

both individual neuron and population levels (Goldberg et al., 2002; Mallet et al., 2008;

Pasquereau and Turner, 2011; Shimamoto et al., 2013; Hemptinne et al., 2015;

Pasquereau et al., 2016), which disrupt its normal function in motor control and motor

learning.

The abnormal neuronal activity in the M1 has been hypothesized to a

consequence of pathological basal ganglia output (Hosp et al., 2011, 2015; Pasquereau

and Turner, 2011; Guo et al., 2015b; Pasquereau et al., 2016), but the cellular

adaptations could also play a key role in such abnormal activities. Moreover, compelling

evidence suggests that PTNs and ITNs may adapt differently to the degeneration of

midbrain DA neurons (Pasquereau and Turner, 2011; Pasquereau et al., 2016).Thus,

we hypothesized that loss of SN DA neurons alters the intrinsic properties of M1

pyramidal neurons in a cell-type-specific manner, and that the intrinsic adaptations

contribute to the abnormal neuronal activity of M1 in parkinsonian state. We addressed

these hypotheses using electrophysiology and retrograde tracing approach in mice with

6-hydroxydopmaine (6-OHDA) lesion, an established model of parkinsonism.

Material and Methods

Animals

Mice were housed up to four animals per cage under a 12-h light/12-h dark cycle with

access to food and water *ad libitum* in accordance with Van Andel Institute IACUC and

NIH guidelines for care and use of animals.

Stereotaxic surgery

Mice were deeply anesthetized under 2% isoflurane, were placed in a stereotaxic frame

(Kopf), and were supported by a thermostatic heating pad. To induce degeneration of

SN DA neurons, 6-OHDA (3-4 mg/mL, 1.0 µL) was injected unilaterally into the medial

forebrain bundle (MFB, from bregma (in mm): AP, -0.7, ML, +1.2, DV, -4.7) over 10 min

using a 10-µl syringe (Hamilton) and a motorized microinjector (Stoelting, Wood Dale,

IL, USA) (Chu et al., 2017). Controls were injected in the same location with vehicle.

Desipramine (25 mg/kg) and pargyline (50 mg/kg) were subcutaneously injected 30-40

min prior to 6-OHDA injection, to enhance the sensitivity and toxicity of 6-OHDA (Chu et

al., 2015, 2017). To label and identify PTNs and ITNs in the M1, animals also received

red or green Retrobeads (0.3 µL, Lumafluor, Inc) injected into the ipsilateral pontine

nuclei (from bregma (in mm): AP, -5.0; ML, +0.6; DV, -5.0; 0.3 µl per injection) or the

contralateral dorsolateral striatum (from bregma (in mm): AP, +0.4; ML, -2.0; DV, -2.8).

The surgical wound was closed using sutures and mice were allowed to recover after

surgery in a heated cage with access to food and water at the cage-floor level. Motor

cortex slices containing the forelimb regions from 6-OHDA- or vehicle-injected mice

were prepared 3-4 weeks after surgery for *ex vivo* electrophysiology (days post-surgery:

control = 27 [22, 28] days, n = 19 mice; 6-OHDA = 24 [22, 27] days, n = 19 mice).

Slice preparation

Mice were deeply anesthetized with intraperitoneal avertin (250 mg/kg) and then were

perfused transcardially with ice-cold, sucrose-based, artificial cerebrospinal fluid (aCSF)

containing (in mM) 230 sucrose, 26 $NaHCO_3$, 10 glucose, 10 $MgSO_4$, 2.5 KCl, 1.25

NaH_2PO_4, and 0.5 $CaCl_2$. Next, coronal brain slices (250 µm) containing M1 forelimb

regions were prepared in the same slicing solution using a vibratome (VT1200S; Leica

Microsystems Inc., Buffalo Grove, IL, USA). Brain slices were kept in normal aCSF (in

mM: 126 NaCl, 26 $NaHCO_3$, 10 glucose, 2.5 KCl, 2 $CaCl_2$, 2 $MgSO_4$, 1.25 NaH_2PO_4, 1

sodium pyruvate and 0.005 L-glutathione) equilibrated with 95% O_2 and 5% CO_2 for 30

min at 35 °C and then held at room temperature until electrophysiological recording.

Ex vivo **electrophysiology recording**

Brain slices were transferred into a recording chamber perfused at a rate of 4 mL/min

with synthetic interstitial fluid (in mM: 126 NaCl, 26 $NaHCO_3$, 10 glucose, 3 KCl, 1.6

$CaCl_2$, 1.5 $MgSO_4$, 1.25 NaH_2PO_4) equilibrated with 95% O_2 and 5% CO_2 at 35 °C via a

feedback-controlled in-line heater (TC-324C, Warner Instruments). DNQX (20 µM), D-

APV (50 µM), and SR-95531 (10 µM) were routinely added to block synaptic

transmission mediated by ionotropic glutamatergic and GABAergic receptors. Neurons

were visualized and recorded under gradient contrast SliceScope 6000 (Scientifica, UK)

with infrared illumination using a CCD camera (SciCam Pro, Scientifica, UK) and

motorized micromanipulators (Scientifica, UK). Individual neurons labeled with

Retrobeads in the M1 L5b were identified using a 60X water immersion objective lens

(Olympus, Japan) and targeted for whole-cell patch-clamp recording, using a

MultiClamp 700B amplifier and a Digidata 1500B digitizer under the control of

pClamp11 (Molecular Devices, San Jose, USA). Data were collected at a sampling rate

of 20-50 KHz. Borosilicate glass pipettes for patch clamp recordings (4-7 MΩ) were

pulled using a micropipette puller (P1000, Sutter Instruments). Pipette capacitance was

compensated for before the formation of the whole-cell configuration. Series resistance

(Rs) was regularly monitored and compensated for with the bridge balance circuit in

current clamp mode. Liquid junction potential (about 11 mV) was corrected.

Retrobeads-labeled neurons in the L5b of M1 were recorded in whole-cell

current-clamp mode to study their intrinsic properties. In these experiments, glass

pipettes were filled with a potassium gluconate–based internal solution of (in mM) 140

K-gluconate, 3.8 NaCl, 1 $MgCl_2$, 10 HEPES, 0.1 Na_4-EGTA, 2 ATP-Mg, and 0.1 GTP-

Na, pH 7.3, osmolarity 290 mOsm. The resting membrane potential (V_m) was recorded

once the whole-cell configuration was obtained. The intrinsic properties and excitability

of M1 pyramidal neuron subtypes were studied by injecting a family of current steps

ranging from -80 pA to 720 pA in 40-pA increments and with a duration of 1 s. Current

injections were from V_m. Input resistance was determined by measuring the steady-

state voltage responses to a series of 1-s hyperpolarizing currents.

The rheobase, as an indicator of neuronal excitability, was defined as the current

intensity to elicit an action potential (AP) during a 1-s current injection. AP waveforms at

the rheobase were systematically analyzed and quantified. Specifically, the threshold of

AP was determined as the voltage level at which dV/dt exceeded 20 mV/ms. AP

amplitude was defined as the voltage difference between the threshold and the peak

voltage. AP half-width was measured as the time difference at 50% of AP amplitude.

Fast and medium afterhyperpolarizations (fAHPs and mAHPs) were measured as the

negative voltage peaks relative to the threshold within 2-5 ms and 20-50 ms from AP

threshold, respectively (Villalobos et al., 2004; Bean, 2007). To quantify spike trains,

spike frequency adaptation was measured as the ratio between the last inter-spike-

interval and the average of the first two inter-spike-intervals. The gain of the spike-

current curve was defined as the slope of the instantaneous spike-current curve.

Immunohistochemistry

Striatal immunoreactivity of tyrosine hydroxylase (TH) was assessed to validate the

degeneration of the nigrostriatal DA projections in vehicle- and 6-OHDA-injected mice.

Briefly, brain tissues were first fixed in 4% paraformaldehyde in 0.1 M phosphate buffer,

pH7.4, at 4 °C for 12 hours before rinsing in phosphate-buffered saline (PBS, 0.05 M;

pH 7.4). Tissues was re-sectioned at 70 μm using a vibratome (Leica VT1000S;

Microsystems Inc.). Immunochemical detection of TH was conducted in PBS containing

0.2% Triton X-100 (Fisher Scientific) and 2% normal donkey serum (Sigma-Aldrich, St.

Louis, MO). Brain sections were incubated in primary antibody (mouse anti-TH, 1:1000,

cat#: MAB318, Sigma-Aldrich, St. Louis, MO, USA) for 48 h at 4 °C or overnight at room

temperature (RT), washed in PBS, and then incubated in the secondary antibody

(donkey anti-mouse Alexa Fluor 488, cat#: 715-545-150 or donkey anti-mouse Alexa

Fluor 594, cat#715-585-150; 1: 500, Jackson ImmunoReseach) for 90 minutes at RT

before washing with PBS. Brain sections were mounted on glass slides using

VECTASHIELD antifade mounting medium (cat#: H-1000, Vector Laboratories,

Burlingame, CA, USA) and were cover-slipped. TH immunoreactivity was imaged using

an Olympus BX63F microscope equipped with an Olympus DP80 camera or a confocal

laser scanning microscope (A1R; Nikon, Melville, USA). The intensity of TH

immunofluorescence was quantified using ImageJ (NIH). All the mice with 6-OHDA

injection showed > 80% TH immunoreactivity loss in the ipsilateral dorsal striatum

versus contralateral dorsal striatum.

Experimental design and statistical analysis

Nineteen male and 19 female adult (4-5 months old) C57BL/6 mice were used in the

study. Data were analyzed in Clampfit (Molecular Devices) and ImageJ (NIH). Statistics

were done using Prism8 (GraphPad Software, San Diego, CA, USA). To minimize the

assumption of the data normality, we used non-parametric, distribution-independent

Mann-Whiney U (MWU) or Wilcoxon signed rank (WSR) tests for paired or non-paired

data comparisons, respectively. The Holm-Bonferroni correction was applied for multiple

comparisons. All tests were two-tailed, and an α-level of 0.05 was used to determine

statistical significance. Results are reported as median and interquartile range. Boxplots

illustrate the median (central line), interquartile range (box), and 10-90% range

(whiskers) of data. Data are available upon request.

Results

Cell-type-specific decrease in the intrinsic excitability of M1 L5b pyramidal

neurons following the loss of midbrain DA neurons.

We compared the intrinsic excitability of projection-defined pyramidal neurons in the L5b

of M1 from vehicle-injected controls (hereinafter, "controls") and from mice with 6-OHDA

lesion (hereinafter, "6-OHDA mice"). To distinguish M1 pyramidal neurons based on

long-range projections, Retrobeads were stereotaxically injected into the ipsilateral

pontine nuclei (Fig. 1A) and the contralateral striatum (Fig. 2A) to retrogradely label

PTNs (Fig. 1B) and ITNs (Fig. 2B), respectively. The intrinsic electrophysiological

properties of M1 PTNs and ITNs were assessed using whole-cell current-clamp

recording in the presence of antagonists of ionotropic glutamatergic and GABAergic

receptors, including DNQX (20 µM), D-APV (50 µM), and SR95531 (10 µM). All

recordings were conducted between 3 and 4 weeks post-surgery, when the level of

midbrain DA degeneration and functional adaptations in the brain have reached

maximum and been stabilized (Vila et al., 2000; Viaro et al., 2011). The data are

reported as median and interquartile range in the text and figures.

The intrinsic excitability of M1 pyramidal neuron subtypes was assessed by

somatic injection of a range of depolarizing currents in controls and 6-OHDA mice

(Figures 1C, D). We found that the gain of spike-current curve (control = 78.2 [67.4,

91.25] spikes/nA, n = 25 neurons/5 mice; 6-OHDA = 52.0 [44.7, 72.2] spikes/nA, n = 25

neurons/4 mice; p = 0.0015, MWU; Figures 1E) was significantly decreased in M1 PTNs

from 6-OHDA mice relative to those from controls. The same reduction was also

detected for the spike frequency (e.g. frequency at 720 pA current injection, control = 55

[47, 59] Hz, n = 25 neurons/5 mice; 6-OHDA = 38 [29, 53] Hz, n = 25 neurons/4 mice; p

= 0.0016, MWU; Figure 1F). M1 PTNs could sustain persistent firing with minimal spike

adaptation during repetitive firing in the physiological state (Figure 1C) (Oswald et al.,

2013; Suter et al., 2013). However, following the loss of midbrain DA neurons, PTNs

showed significantly enhanced spike adaptation. For instance, the spike adaptation of a

10- to15-Hz spike train, control = 1.06 [0.97, 1.26], n = 25 neurons/5 mice; 6-OHDA =

1.38 [1.17, 1.60], n = 25 neurons/4 mice; p = 0.0001, MWU (Figures 1C, G). These

results suggest a significant reduction in the intrinsic excitability of M1 PTNs following

the loss of midbrain DA neurons.

Figure 1. Decreased intrinsic excitability of PTNs in the L5b of M1 following the loss of midbrain DA neurons. A and B) Representative graphs showing the injection site of red Retrobeads in the pontine nuclei (**A**) and a retrogradely labeled PTN in L5b of M1 (**B**). **C and D**) Representative spike trains of PTNs evoked by somatic current injections from controls and 6-OHDA mice (**C**), and the frequency-current relationship of the PTNs from controls and 6-OHDA mice (**D**). **E-G**) Summarized results showing a reduced gain of spike-current curve (**E**), a decreased firing frequency in response to maximal current injections (i.e., 720 pA) (**F**), and an enhanced spike adaptations (**G**) in PTNs from 6-OHDA mice relative to those from controls. *, p < 0.05, MWU.

In contrast, M1 ITNs showed comparable spike-current relationships between the

controls and 6-OHDA mice. This was reflected by three measures, namely, a) the

unaltered gain of spike-current curve (control = 78 [54, 93] spikes/nA, n = 38 neurons/6

mice; 6-OHDA = 77 [67, 88] spikes/nA, n= 35 neurons/6 mice; p = 0.94, MWU; Figure

2D, E); b) the spike frequency at maximal (720 pA) current injection (control = 53 [38,

64] Hz, n = 38 neurons/6 mice; 6-OHDA = 50 [40, 58] Hz, n = 35 neurons/6 mice; p =

0.4, MWU; Figure 2D, F); and c) spike adaptations (control = 1.36 [1.08, 1.67], n= 38

neurons/6 mice; 6-OHDA = 1.20 [1.02, 1.61], n = 35 neurons/6 mice; p = 0.24, MWU;

Figure 2C, G). Altogether, our results thus suggest that the intrinsic excitability of PTNs,

but not ITNs, was significantly decreased following the loss of midbrain DA neurons,

that is, the M1 L5b pyramidal neurons exhibited cell-type-specific adaptations in their

intrinsic excitability in the parkinsonian state.

Figure 2. Intact intrinsic excitability of ITNs in the M1 L5b following the loss of midbrain DA neurons. A and B) Representative graphs showing the injection site of green Retrobeads in the dorsal striatum (**A**) and retrogradely labeled M1 L5b ITNs (**B**). **C and D)** Representative spikes trains evoked by somatic current injections into ITNs from control and 6-OHDA mice (**C**), and the frequency-current relationship of ITNs from controls and 6-OHDA mice (**D**). **E-G)** Boxplots showing an unaltered gain of spike-current curve (**E**), firing frequency in response to maximal current injections (i.e., 720 pA) (**F**), and spike adaptation (**G**) between ITNs from 6-OHDA mice and controls. ns, not significant, MWU.

256 **Cell-type-specific changes in the passive membrane properties of M1 L5b**

257 **pyramidal neurons following the loss of midbrain DA neurons.**

258 In the absence of significant synaptic inputs, PTNs in 6-OHDA mice and controls had

259 similar resting membrane potentials (V_m, control = -79.3 [-81.6, -77.9] mV, n = 25

260 neurons/5 mice; 6-OHDA = -79.7 [-81.5, -78.4] mV, n=25 neurons/4 mice; p = 0.85,

261 MWU, Figure 3A). However, PTNs from 6-OHDA mice had an increased input

262 resistance (R_m, control = 69.3 [60.1, 81.5] MΩ, n = 25 neurons/5 mice; 6-OHDA = 109.6

263 [73.0, 136.4] MΩ, n= 25 neurons/4 mice; p = 0.0028, MWU; Figures 3A, B). In contrast,

264 ITNs showed a hyperpolarized V_m in 6-OHDA mice (control = -77.7 [-80.2, -75.0] mV,

265 n=38 neurons/6 mice; 6-OHDA = -79.6 [-82.1, -77.2] mV, n = 35 neurons/6 mice; p =

266 0.012, MWU; Figure 3D). ITNs in 6-OHDA mice also had an unaltered R_m (control =

267 89.7 [60.4, 121.3] MΩ, n = 38 neurons/6 mice; 6-OHDA = 70.9 [54.7, 93] MΩ, n = 35

268 neurons/6 mice; p = 0.067, MWU; Figure 3E). Changes in V_m and R_m were associated

269 with a significant increase in rheobase in ITNs (control= 80.8 [51.3, 125] pA, n = 36

270 neurons/6 mice; 6-OHDA = 125 [86, 191] pA, n = 35 neurons/6 mice; p = 0.002, MWU;

271 Figure 3F). PTNs had unaltered rheobase following the loss of midbrain DA neurons

272 (control = 82 [47, 163] pA, n= 25 neurons/5 mice; 6-OHDA = 79 [56, 120] pA, n = 25

273 neurons/4 mice; p = 0.645, MWU; Figure 3C). Thus, the loss of midbrain DA neurons

274 induced distinct adaptations in passive membrane properties of PTNs and ITNs in M1.

Figure 3. Distinct changes in the passive membrane properties of PTNs and ITNs in the M1 L5b following the loss of midbrain DA neurons. A-C) Boxplots showing changes in the resting membrane potential (**A**), input resistance (**B**), and rheobase (**C**) of PTNs from 6-OHDA mice relative to those from controls. **D-F)** Boxplots showing changes in in the resting membrane potential (**D**), input resistance (**E**), and rheobase (**F**) of ITNs from 6-OHDA mice relative to those from controls. ns, not significant; *, p < 0.05, MWU.

Cell-type-specific changes in action potential waveforms of M1 L5b pyramidal neurons following the loss of midbrain DA neurons.

The shape of the action potential is determined by the interaction of numerous voltage-gated ion channels and has profound effects on firing rate and pattern of neurons (Bean, 2007). We therefore analyzed the morphology of AP waveforms of PTNs (Figure 4A) and ITNs (Figures 4G) from controls and 6-OHDA mice. Following the loss of the midbrain DA neurons, M1 PTNs showed a striking AP broadening (AP half-width, controls = 0.69 [0.62, 0.75] ms, n = 25 neurons/5 mice; 6-OHDA = 0.85 [0.68, 1.23] ms, n = 25 neurons/4 mice; p = 0.005, MWU; Figure 4B). The broadened AP of PTNs was associated with a significantly prolonged AP rise time (controls = 0.22 [0.19, 0.24] ms, n = 25 neurons/5 mice; 6-OHDA = 0.25 [0.22, 0.32] ms, n = 25 neurons/4 mice; p < 0.002, MWU; Figures 4C) and also a prolonged decay time (controls = 0.60 [0.53, 0.68] ms, n = 25 neurons/5 mice, 6-OHDA = 0.80 [0.58, 1.3] ms, 25 neurons/4 mice; p = 0.0035, MWU; Figure 4D). Further, the maximal rates of AP depolarization and repolarization of M1 PTNs also greatly decreased. The maximal AP rise rate in controls = 358 [306, 437]

mV/ms, n = 25 neurons/5 mice, while the maximal AP rise rate in 6-OHDA mice = 295

[245, 339] mV/ms, n= 25 neurons/4 mice; p = 0.0008, MWU (Figure 4E). The maximal

AP decay rate, controls = -112 [-104, -130] mV/ms, n = 25 neurons/5 mice, and 6-

OHDA = -84 [-64, -106] mV/ms, n = 25 neurons/4 mice; p = 0.0001, MWU (Figure 4F).

Thus, the decreased excitability of PTNs is largely caused by an abnormal AP

broadening, involving impaired AP depolarization and repolarization.

Figure 4. Distinct changes in AP waveforms of PTNs and ITNs in the M1 L5b following loss of midbrain DA neurons. A) Top, representative AP waveforms of PTNs from controls and 6-OHDA mice. APs were aligned at threshold and overlaid for comparison. Bottom, the corresponding dV/dt traces plotted against time, showing depolarizing and repolarizing rates of APs. B) Boxplot showing broadened AP width in PTNs from 6-OHDA mice relative to those from controls. C and D), Boxplots showing prolonged AP rise time (C) and decay time (D) in PTNs from 6-OHDA mice relative to those from controls. E and F), Boxplots showing decreased AP rise rate (E) and decay rate (F) in PTNs from 6-OHDA mice relative to those from controls. G) Top, representative AP waveforms of ITNs from controls and 6-OHDA mice. APs were aligned at threshold and overlaid for comparison. Bottom, the corresponding dV/dt traces plotted against time, showing depolarizing and repolarizing rates of APs. H) Boxplot showing similar AP width in ITNs from 6-OHDA mice relative to those from controls. I-J) Boxplots showing a shorter AP rise time (I) but similar decay time (J) in ITNs from 6-OHDA mice relative to those from controls. K-L) Boxplots showing an increased AP rise rate (K) but similar decay rate (L) in ITNs from 6-OHDA mice relative to those from controls. ns, not significant; *, p < 0.05, MWU.

In contrast, ITNs from controls and 6-OHDA mice showed similar AP width

(control = 0.73 [0.64, 0.81] ms; n = 38 neurons/6 mice, 6-OHDA = 0.70 [0.61, 0.75] ms,

n = 36 neurons/6 mice; p = 0.2, MWU; Figure 4H). The similar AP width of ITNs was

associated with an unaltered decay time and decay rate. The decay time for controls =

0.67 [0.58, 0.77] ms, n = 38 neurons/6 mice, while for 6-OHDA = 0.63 [0.57, 0.70] ms, n

= 35 neurons/6 mice; p = 0.15, MWU (Figure 4J). The decay rate values were, for

controls = -102 [-82, -122] mV/ms, n = 38 neurons/6 mice, and for 6-OHDA = -112 [-

100, -122] mV/ms, n = 35 neurons/6 mice; p = 0.1, MWU (Figure 4L). Moreover, ITNs

from 6-OHDA mice exhibited a faster AP depolarization relative to those from controls,

as reflected by a shorter rise time (controls = 0.24 [0.21, 0.26) ms, n = 38 neurons/6

mice; 6-OHDA = 0.21 [0.19, 0.24] ms, n = 35 neurons/6 mice; p = 0.03, MWU; Figure 4I]

and also a faster rise rate (control = 314 [276, 383] mV/ms, n = 38 neurons/6 mice; 6-

OHDA = 377 [329, 431] mV/ms, n = 35 neurons/6 mice; p = 0.009, MWU; Figure 4K).

Thus, these results suggest that ITNs of 6-OHDA mice had subtler changes in AP

waveforms and that efficient AP depolarization is likely a compensatory mechanism for

the hyperpolarized V_m and reduced R_m (Figures 3D, E), leading to an unaltered

superathreshold excitability (Figures 2C, D).

Progressive alterations in AP waveforms of PTNs during repetitive firing

following the loss of SN DA neurons.

In the physiological state, the burst pattern of activity of M1 PTNs is closely linked to

movement parameters, but this specificity of movement encoding is disrupted in PD

(Pasquereau et al., 2016; Economo et al., 2018; Sauerbrei et al., 2020). In addition,

PTNs show a decreased intraburst firing rate in MPTP-treated animals (Pasquereau

and Turner, 2011). We predicted that the altered intrinsic properties of PTNs contribute

to these *in vivo* observations in the parkinsonian state. Thus, spike trains containing 10-

15 spikes were elicited from M1 PTNs in controls and 6-OHDA mice, and the waveform

of the first and the last APs within a spike train were analyzed to study progressive

changes during repetitive firing.

Figure 5. Progressive changes in AP waveforms of M1 PTNs during repetitive firing. **A-B**) APs from a 10- to15-Hz spike train were superimposed for PTNs from control (**A**) or 6-OHDA mice (**B**). APs were aligned at threshold, and the solid and dashed lines indicated the first and the last APs from the spike train, respectively. All the other APs were shown as transparent lines. **C-D**) Summarized graphs showing the alterations of AP width between the first and the last APs. Each line in (**C**) connects the AP width of the first and the last APs from the same spike train. Filled and open circles indicate the median plus interquartile range of the first and the last AP width of PTNs, respectively. (**D**) Boxplots showing the AP width difference between the first and the last APs was greater in PTNs from 6-OHDA mice relative to those from controls. **E-F**) Summarized graphs showing alterations of AP decay rate between the first and the last APs (**E**) and greater changes of AP decay rate in PTNs from 6-OHDA mice relative to those from controls (**F**). **G-H**) Summarized graphs showing alterations of AP decay time between the first and the last APs (**G**), and greater changes of AP decay time in PTNs from 6-OHDA mice relative to those from controls (**H**). **I-J**) Summarized graphs showing alterations of AP threshold between the first and the last APs (**I**), and greater changes of AP threshold in PTNs from 6-OHDA mice relative to those from controls (**J**). ns, not significant; *, p < 0.05, WSR.

The literature suggests that M1 PTNs have the capacity for high-frequency AP

firing (300-450 Hz) (Suter et al., 2013). We also noted constant AP waveforms during

repetitive firing in the control mice (Fig. 5A). Specifically, the PTNs from control mice

showed subtle AP broadening during repetitive firing (AP width in controls, the first AP =

0.68 [0.60, 0.72] ms, the last AP = 0.69 [0.60, 0.75] ms, n = 25 neurons/5 mice; p =

0.013, WSR; Figure 5C). They also showed an unaltered decay rate (the first AP = -114

[-110, -131] mV/ms, the last AP = -115 [-105, -132] mV/ms, n = 25 neurons/5 mice; p =

0.57, WSR; Figure 5E) and an unaltered decay time (the first AP = 0.59 [0.53, 0.68] ms,

the last AP = 0.57 [0.50, 0.68] ms, n = 25 neurons/5 mice; p = 0.26, WSR; Figure 5G)

between the first and the last APs during repetitive firing. In addition, the AP threshold of

M1 PTNs from controls was also constant between the first and the last AP (threshold,

the first AP = -53.5 [-57.2, -51.4] mV, and the last AP = -52.3 [-54.9, -50.9] mV; n = 25

neurons/5 mice; p = 0.26, WSR; Figure 5I). The constant AP parameters of PTNs during

repetitive firings are key to delivering motor commands to subcortical premotor centers

with temporal precision during movement initiation and execution.

In contrast, PTNs from 6-OHDA mice showed progressive AP broadening during

repetitive firing (the first AP width = 0.78 [0.67, 0.97] ms, the last AP = 0.88 [0.70, 1.08]

ms, n = 25 neurons/4 mice; p = 0.0001, WSR; Figure 5C). This AP broadening was

associated with a significantly decreased AP decay rate (the first AP = -91.6 [-69.3, -

114.7] mV/ms, the last AP = -78.7 [-63.2, -104.4] mV/ms, n = 25 neurons/4 mice; p =

0.0001, WSR; Figure 5E) and decay time (the first AP = 0.72 [0.61, 1.0] ms, the last AP

= 0.77 [0.61, 1.20] ms, n = 25 neurons/4 mice; p = 0.016, WSR; Figure 5G). Moreover,

PTNs in 6-OHDA mice had a progressive increase in AP threshold during repetitive

firing (the first AP = -53.7 [-59.1, -50.8] mV, the last AP = -51.1 [-56.6, -44.8] mV, n = 25

neurons/4 mice; p = 0.0001, WSR; Figure 5I).

When the difference between the first and last APs was analyzed, we found

greater differences in the above parameters in PTNs from 6-OHDA mice than those

from controls. The AP width$_{(last-first)}$ for controls was 0.006 (-0.004, 0.04) ms, n = 25

neurons/5 mice, while for 6-OHDA mice was 0.05 (0.01, 0.10) ms, n = 25 neurons/4

mice; p = 0.016, MWU (Figure 5D). The AP decay rate$_{(last-first)}$ for the controls was 2.44 [-

3.66, 5.56] mV/ms, n = 25 neurons/5 mice, while the value for 6-OHDA mice was 7.94

[4.58, 11.90] mV/ms, n = 25 neurons/4 mice; p = 0.0004, MWU (Figure 5F). The AP

decay time$_{(last-first)}$ for controls was -0.015 (-0.045, 0.016) ms, n = 25 neurons/5 mice,

while the value for 6-OHDA mice was 0.059 (-0.03, 0.12) ms, n = 25 neurons/4 mice; p

= 0.018, MWU (Figure 5H). Finally, the AP threshold$_{(last-first)}$ value for controls was 0.40 [-

1.65, 2.78] mV, n = 25 neurons/5 mice, while the value for 6-OHDA mice was 2.96

[1.05, 4.07] mV, n = 25 neurons/4 mice; p < 0.006, MWU (Figure 5J). These results

suggest that following the loss of midbrain DA neurons the altered intrinsic properties of

M1 PTNs impair their capability to sustain constant AP waveforms during repetitive

firing, which likely compromises their capacity of fast firing and the temporal precision of

motor commands for motor control (Suter et al., 2013; Sauerbrei et al., 2020).

Impaired activity of BK channels underlies the decreased intrinsic excitability of

M1 PTNs following the loss of midbrain DA neurons.

AP width is determined by the time and velocity of depolarization and repolarization

(Bean, 2007). Our results above indicated that broadened APs of PTNs following the

loss of DA neurons are largely attributed to prolonged AP repolarization. Furthermore,

following the loss of midbrain DA neurons, the amplitude of the fast

afterhyperpolarization (fAHP) in PTNs diminished (controls = 12.6 [9.8, 14.5] mV, n = 25

neurons/5 mice; 6-OHDA = 9.1 [6.0, 12.3] mV, n = 25 neurons/5 mice; p = 0.003, MWU;

Figures 6A, B), but the amplitude of median afterhyperpolarization (mAHP) was similar

to that of controls (controls = 21.7 [16.6, 24.4] mV, n = 25 neurons/5 mice; 6-OHDA =

23.0 [20.1, 26.4] mV, n = 25 neurons/5 mice; p = 0.63, MWU; Figure 6C).

Figure 6. Decreased BK channel activity underlies AP broadening and impaired intrinsic excitability of PTNs following loss of midbrain DA neurons. A) Representative AP waveforms showing diminished fAHPs (arrow) in PTNs from 6-OHDA mice, relative to those from controls. **B-C**) Boxplot showing decreased amplitude of fAHPs (**B**) and unaltered mAHPs (**C**) in PTNs from 6-OHDA mice relative to those from controls. **D**) Representative APs waveforms of PTNs from controls and 6-OHDA mice in the presence of BAPTA. APs were aligned at the threshold and overlaid for comparison. **E**) Boxplot showing reversal of broadened AP of PTNs in 6-OHDA mice relative to controls after intracellular BAPTA dialysis. **F**) Summarized graph showing unaltered excitability of PTNs from 6-OHDA mice relative to those from controls in the presence of BAPTA. The number of APs elicited by 400-, 600-, and 800-pA somatic current injections were compared. **G and H**) Representative APs waveforms before and after paxillin application in PTNs from controls (**G**) and 6-OHDA mice (**H**). APs were aligned at the threshold and overlaid for comparison. **I**) Summarized graph showing paxillin broadened APs of PTNs from both controls and 6-OHDA mice. **J**) Summarized graph showing that paxillin significantly decreased the number of spikes (evoked by injecting 720-pA current) of PTNs from controls but did not affect those from 6-OHDA mice. Application of paxillin occluded the decreased number of spikes of PTNs from 6-OHDA mice relative to those from controls. Filled and open symbols in (**I and J**) indicate the values of before and after paxillin application, respectively. ns, not significant; *, p < 0.05, WSR or MWU.

BK currents contribute to fAHPs, AP repolarization and/or AP broadening during

repetitive neuronal firing in multiple brain regions (Faber and Sah, 2003; Bean, 2007;

Contet et al., 2016). The reduced fAHP amplitude in PTNs from 6-OHDA mice indicates

that altered BK channel activity probably underlies broadened AP of PTNs in 6-OHDA

mice.

BK channel activation requires both membrane depolarization and Ca^{2+} influx

during APs. Therefore, we decreased the activation of BK channels by intracellularly

dialyzing a high-affinity Ca^{2+} chelator, BAPTA (10 mM), and determined its effect on the

AP width of PTNs from both controls and 6-OHDA mice. In the presence of BAPTA, the

AP of PTNs from 6-OHDA mice was slightly and statistically significant narrower (half-

width: controls = 0.99 [0.92, 1.17] ms, n = 40 neurons/3 mice; 6-OHDA = 0.95 [0.83,

1.03] ms, n = 39 neurons/3 mice; p = 0.029, MWU; Figures 6D, E). In addition, the

PTNs from both groups also showed similar intrinsic excitability in the presence of

BAPTA (Figure 6F). For example, the number of spikes in response to maximal current

injection at 800 pA for controls = 42 [28, 55] spikes, n = 40 neurons/3 mice, while the

value for 6-OHDA = 40 [31, 49] spikes, n = 39 neurons/3 mice; p = 0.84, MWU (Figure

6F).

Furthermore, we blocked BK channels activity by a selective BK channel blocker,

paxillin (10 µM), through bath application and we found that it significantly prolonged the

APs of PTNs in both groups (Figure 6G, H). The half-width of controls before and after

paxillin were 0.77 [0.64, 0.81] ms, and 0.84 [0.8, 0.96] ms, respectively (n = 15

neurons/3 mice; p = 0.0001, WSR; Figures 6I). For PTNs from 6-OHDA mice, the half-

width before and after paxillin were 0.82 [0.75, 0.87] ms and 0.89 [0.78, 0.98] ms,

respectively (n= 14 neurons/3 mice; p = 0.0009, WSR; Figures 6I). With the blockade of

BK channels by paxillin, the AP width was not statistically different between PTNs from

controls and 6-OHDA mice (MWU, p = 0.62; Figure 6G, I). In addition, paxillin

significantly decreased the intrinsic excitability of PTNs from controls (the number of

spikes in response to 480-pA current injection, baseline = 36 [32, 48] Hz, paxillin = 31

[17, 43] Hz; n = 15 neurons/ 3 mice; p = 0.0002, WSR; Figure 6J). In contrast, paxillin

did not affect the number of spikes in PTNs of 6-OHDA mice (spike frequency at 480-pA

current injection, baseline = 32 [27, 39] Hz, paxillin = 29 [28, 39] Hz; n = 14 neurons/ 3

mice; p = 0.71, WSR; Figure 6J). In the presence of paxillin, there is no difference in the

intrinsic excitability of PTNs between 6-OHDA mice and controls (MWU, p = 0.7; Figure

6J). Together, these results suggested that impaired BK channel function underlies the

broadened APs and decreased intrinsic excitability of PTNs following the loss of

midbrain DA neurons.

Intrinsic adaptations of M1 pyramidal neurons were not due to the decreased DA

neuromodulation in M1

M1 receives dopaminergic inputs from the ventral tegmental area and these neurons

are partially degenerate in 6-OHDA mice (Hosp et al., 2011, 2015; Hosp and Luft,

2013). Thus, we predicted that loss of direct mesocortical DA neuromodulation

contributes to the intrinsic adaptations of PTNs and ITNs in M1. To test this prediction,

we activated DA receptors on PTNs and ITNs through bath perfusion of DA (10 µM) and

determined its impact on the intrinsic properties of M1 PTNs and ITNs from both

controls and 6-OHDA mice. The activation of DA receptors by DA produced negligible

effects to the intrinsic excitability of M1 PTNs in controls (e.g., the firing frequency at a

360-pA current injection, baseline = 37 [30, 47] Hz; DA = 34 [20, 39] Hz; n = 10

neurons/3 mice; p = 0.25, WSR; Figures. 7A, B). In addition, the application of DA (10

µM) did not produce significant effects to the intrinsic excitability of M1 PTNs in 6-OHDA

mice either (e.g., the firing frequency at a 360-pA current injection, baseline = 26 [20,

33] Hz, DA = 27 [19, 36] Hz, n = 14 neurons/3 mice; p = 0.72, WSR; Figures. 7C, D).

Figure 7. DA application could not rescue intrinsic adaptations in PTNs and ITNs of M1 following loss of midbrain DA neurons. **A-B)** Representative AP traces showing the effect of DA application (10 µM) on the number of APs of PTNs from controls (**A**) and the summarized results (**B**). **C-D**) Representative AP traces showing the effect of DA application on the number of APs of PTNs from 6-OHDA mice (**C**) and the summarized results (**D**). **E-G)** Summarized graphs showing the impact of DA (10 µM) on intrinsic properties of ITNs from controls and 6-OHDA mice, including the V_m (**E**), R_m (**F**), rheobase (**G**) of ITNs from controls and 6-OHDA mice. Filled and open circles in (**E-G**) indicate the values of before and after DA application, respectively Each line connected the rheobase before (filled circle) and after (open circle) DA application. *, p < 0.05, WSR.

Furthermore, DA (10 µM) did not alter the AP width of PTNs in both group either

(controls: baseline = 0.72 [0.56, 0.8] ms, DA = 0.71 [0.55, 0.84] ms, n = 10 neurons/3

mice, p = 0.32, WSR; 6-OHDA: baseline = 0.74 [0.62, 0.85] ms, 6-OHDA DA = 0.74

[0.57, 0.88] ms, n = 14 neurons/3 mice, p = 0.24, WSR). These data suggested that

restoration of local DA neuromodulation cannot rescue the impaired intrinsic excitability

and AP broadening of M1 PTNs following the loss of midbrain DA neurons.

Next, we tested whether the altered passive membrane properties of M1 ITNs of

6-OHDA mice (Figure 3) are caused by the loss of local DA neuromodulation. In control

mice, the activation of DA receptors by DA (10 µM) did not produce a significant effect

on the V_m (baseline = -78 [-81, -74] mV, DA = -78 [-80, -75] mV, n = 10 neurons/2 mice;

p > 0.99, WSR; Figure 7E) or on the R_m of ITNs (baseline = 88 [51, 109] MΩ, DA = 86

[50, 108] MΩ, n = 10 neurons/2 mice; p = 0.16, WSR; Figure 7F). However, the

rheobase of ITNs was significantly increased (baseline = 66 [43, 112] pA, DA = 116 [77,

212] pA, n = 10 neurons/ 2 mice; p = 0.027, WSR; Figure 7G). On the other hand, in

ITNs of 6-OHDA mice, the application of DA (10 μM) did not alter the V_m (baseline = -79

[-82, -76] mV, DA = -77 [-80, -75] mV, n = 14 neurons/3 mice; p = 0.22, WSR; Figure

7E); the R_m (baseline = 65 [53, 88] MΩ, DA = 64 [53, 75] MΩ, n = 14 neurons/3 mice; p

= 0.06, WSR; Figure 7F); or the rheobase (baseline = 95 [80, 145] pA, DA = 90 [34,

161] pA, n = 14 neurons/3 mice; p = 0.33, WSR; Figure 7G). Thus, the loss of DA

neuromodulation did not play a major role in the cell-type-specific intrinsic adaptations

of M1 pyramidal neurons in the parkinsonian state.

Discussion

We systematically studied adaptative changes in the intrinsic properties of M1

pyramidal neurons in experimental parkinsonism. We found cell-type-specific decrease

of the intrinsic excitability PTNs and adaptive changes in the passive membrane

properties of ITNs in M1. We also found that prolonged AP repolarization led to AP

broadening and resulted in the decreased intrinsic excitability of M1 PTNs. Further, we

revealed that impaired BK channel function was the main ionic mechanism underlying

the altered AP waveforms and intrinsic excitability of M1 PTNs. Last, we showed that

the intrinsic adaptations in M1 PTNs and ITNs were not caused by a loss of

mesocortical dopaminergic neuromodulation. Our findings strongly suggest that the loss

of midbrain DA neurons not only alters basal ganglia activity, but also triggers intrinsic

adaptations in M1 circuitry, which contribute to the reduced motor cortical output and

motor deficits in the parkinsonian state.

Compelling evidence supports an abnormal rate and pattern of activity in M1 in

parkinsonian state, which could be caused by loss of direct mesocortical DA projection,

the pathological basal ganglia output associated with nigrostriatal degeneration, or both.

(Goldberg et al., 2002; Pasquereau and Turner, 2011; Pasquereau et al., 2016). To

distinguish between these possibilities, we employed *ex vivo* electrophysiology to study

the intrinsic properties of projection-defined M1 pyramidal neurons. Our findings that

PTNs, but not the ITNs, in the L5b of M1 had decreased intrinsic excitability in 6-OHDA

mice are consistent with the earlier findings in MPTP-treated primates, showing a cell-

type-specific reduction of the firing frequency of M1 pyramidal neurons (Pasquereau

and Turner, 2011). Given the innervation of subcortical motor centers by M1 PTNs,

these results suggested that intrinsic adaptations of M1 pyramidal neurons contribute to

an insufficient motor cortical outputs following the loss of midbrain DA neurons, in

addition to an elevated basal ganglia inhibition as proposed by the classical model

(Albin et al., 1989; DeLong, 1990; Galvan and Wichmann, 2008). We postulate that

these adaptations exacerbate the hypokinetic parkinsonian motor deficits.

Distinct changes in the intrinsic excitability of M1 pyramidal neurons cannot be

explained by alterations in their passive membrane properties. For example, we found

an increased R_m of PTNs in 6-OHDA mice, which might be a homeostatic compensation

for, rather than a cause of, their decreased intrinsic excitability. It is possible that the

increased R_m makes M1 PTNs more sensitive to and more easily to be entrained by

synaptic inputs in the parkinsonian state, including the abnormal basal ganglia signals

through the thalamocortical projections. The increased sensitivity of PTNs to synaptic

inputs may also disrupts their selectivity in encoding movements. In contrast, ITNs were

hyperpolarized and showed a trend of decreased input resistance following the loss of

midbrain DA neurons, but these changes were compensated for by other adaptions

(e.g., an increased Na_v availability for AP generation), so they did not significantly alter

the superathreshold excitability. Together, our data suggested that, intrinsic adaptations

play an important role in the cell-type-specific alterations of the activity of M1 pyramidal

neurons following the loss of midbrain DA neurons. The molecular mechanisms

underlying cell-type specific adaptations in M1 in parkinsonian state should be

determined in future studies.

APs of PTNs were significantly broadened following the loss of DA neurons and it

was largely due to a prolonged repolarization (Figures 4A-E). AP width has significant

impact on Ca^{2+} influx during APs, which further controls the quantity and kinetics of

neurotransmitter release and synaptic plasticity at axon terminal (Bean, 2007). Thus, we

suggest that AP broadening is a compensatory mechanism for the decreased intrinsic

excitability of M1 PTNs in parkinsonian state in order to maintain stable motor cortical

outputs, though this possibility should be further studied. We further showed that the AP

broadening in PTNs from 6-OHDA mice was occluded in the presence of BAPTA and

paxillin, indicating a key role of BK channels in the intrinsic adaptations of M1 PTNs in

parkinsonian state (Shao et al., 1999; Faber and Sah, 2003; Bean, 2007). Indeed, it has

been proposed that impaired function or inactivation of BK channels that underlies AP

repolarization and/or fAHPs could slow the de-inactivation of Na_v channels during AP

and reduce Na_v availability for the subsequent AP generation, leading to a decreased

excitability (Bean, 2007; Jaffe et al., 2011). A progressive and accumulative Na_v

inactivation is consisted with our observations that PTNs in 6-OHDA mice showed

stronger spike adaptation during repetitive firing relative to those from controls

(Fleidervish et al., 1996), though the contribution of the other mechanisms such as the

M-current can't be discounted (Buchta et al., 2017; Parrilla-Carrero et al., 2018). It is

intriguing that in the presence of BAPTA PTNs from 6-OHDA mice exhibited narrower

APs than those from controls (Figures 6D, E). It indicates the involvement of other ionic

mechanisms in controlling of AP repolarizationof PTNs, e.g., K_v conductance (e.g., K_v3

and K_v4) (Rudy and McBain, 2001; Pathak et al., 2016; Soares et al., 2017).

An acute application of DA did not rescue altered intrinsic excitability of PTNs

and had minimal effects to intrinsic properties of ITNs in 6-OHDA mice. This indicated

that loss of dopaminergic neuromodulation does not play a major role in the intrinsic

adaptations of M1 pyramidal neurons in parkinsonian state, which is consistent with

recent findings that dopaminergic modulation of pyramidal neurons is mainly mediated

indirectly by parvalbumin-expressing interneurons in M1 (Vitrac et al., 2014; Cousineau

et al., 2020). Thus, we propose that circuitry mechanisms underlie cellular adaptions in

M1 in parkinsonian state. For example, recent work highlighted the critical role indirect

pathway SPN hyperactivity in driving cellular and synaptic adaptations in basal ganglia

nuclei, such as the subthalamic nucleus (Chu et al., 2017; Sharott et al., 2017; Ryan et

al., 2018; McIver et al., 2019). Thus, it is likely that the pathological basal ganglia

outputs following loss of DA that drive adaptive changes in the intrinsic properties of M1

pyramidal neurons, which are mediated by thalamocortical projections.

The exaggerated rhythmic activity and aberrant oscillations in the basal ganglia

circuitry are closely related to the expression of motor symptoms in PD (Bevan et al.,

2002; Little and Brown, 2014; Hemptinne et al., 2015). Indeed, the motor cortex exhibits

abnormal neural activity in parkinsonian state at both the single-cell and population

levels (Goldberg et al., 2002; Pasquereau and Turner, 2011; Hemptinne et al., 2015;

Pasquereau et al., 2016). Moreover, both experimental and computation studies have

indicated that cerebral cortical inputs play a crucial role in orchestrating pathological

activity in basal ganglia nuclei (e.g., the external global pallidus and the subthalamic

nucleus) (Bevan et al., 2002; Magill et al., 2004; Kita and Kita, 2011; Corbit et al., 2016;

Chu et al., 2017; Sharott et al., 2017). Recent studies further showed that electrical or

optogenetic stimulation of the motor cortical pyramidal neurons and/or their projections

produced therapeutic effects in parkinsonian animals (Gradinaru et al., 2009; Li et al.,

2012; Hemptinne et al., 2015; Sanders and Jaeger, 2016; Magno et al., 2019). Thus, an

in-depth understanding of motor cortical adaptations in PD at the cellular and circuitry

levels will help to design better therapeutic approaches with fewer off-target effects.

Figure legends

Figure 1. **Decreased intrinsic excitability of PTNs in the L5b of M1 following the loss of midbrain DA neurons. A and B**) Representative graphs showing the injection site of red Retrobeads in the pontine nuclei (**A**) and a retrogradely labeled PTN in L5b of M1 (**B**). **C and D**) Representative spike trains of PTNs evoked by somatic current injections from controls and 6-OHDA mice (**C**), and the frequency-current relationship of the PTNs from controls and 6-OHDA mice (**D**). **E-G**) Boxplots showing a reduced gain of spike-current curve (**E**), a decreased firing frequency in response to maximal current injections (i.e., 720 pA) (**F**), and an enhanced spike adaptations (**G**) in PTNs from 6-OHDA mice relative to those from controls. *, $p < 0.05$, MWU.

Figure 2. Intact intrinsic excitability of ITNs in the M1 L5b following the loss of midbrain DA neurons. A and B) Representative graphs showing the injection site of green Retrobeads in the dorsal striatum (**A**) and retrogradely labeled ITNs (**B**). **C and D)** Representative spikes trains evoked by somatic current injections into ITNs from control and 6-OHDA mice (**C**), and the frequency-current relationship of ITNs from controls and 6-OHDA mice (**D**). **E-G**) Boxplots showing an unaltered gain of spike-current curve (**E**), firing frequency in response to maximal current injections (i.e., 720 pA) (**F**), and spike adaptation (**G**) between ITNs from 6-OHDA mice and controls. ns, not significant, MWU.

Figure 3. **Distinct changes in the passive membrane properties of PTNs and ITNs in the M1 L5b following loss of midbrain DA neurons**. **A-C**) Boxplots showing

changes in the resting membrane potential (**A**), input resistance (**B**), and rheobase (**C**)

of PTNs from 6-OHDA mice relative to those from controls. **D-F**) Boxplots showing

changes in in the resting membrane potential (**D**), input resistance (**E**), and rheobase

(**F**) of ITNs from 6-OHDA mice relative to those from controls. ns, not significant; *, p <

0.05, MWU.

Figure 4. Distinct changes in AP waveforms of PTNs and ITNs in the M1 L5b

following loss of midbrain DA neurons. A) **Top**, representative AP waveforms of

PTNs from controls and 6-OHDA mice. APs were aligned at threshold and overlaid for

comparison. **Bottom**, the corresponding dV/dt traces plotted against time, showing

depolarizing and repolarizing rates of APs. **B**) Boxplot showing broadened AP width in

PTNs from 6-OHDA mice relative to those from controls. **C and D**), Boxplots showing

prolonged AP rise time (**C**) and decay time (**D**) in PTNs from 6-OHDA mice relative to

those from controls. **E and F**), Boxplots showing decreased AP rise rate (**E**) and decay

rate (**F**) in PTNs from 6-OHDA mice relative to those from controls. **G**) **Top**,

representative AP waveforms of ITNs from controls and 6-OHDA mice. APs were

aligned at threshold and overlaid for comparison. **Bottom**, the corresponding dV/dt

traces plotted against time, showing depolarizing and repolarizing rates of APs. **H**)

Boxplot showing similar AP width in ITNs from 6-OHDA mice relative to those from

controls. **I-J**) Boxplots showing a shorter AP rise time (**I**) but similar decay time (**J**) in

ITNs from 6-OHDA mice relative to those from controls. **K-L**) Boxplots showing an

increased AP rise rate (**K**) but similar decay rate (**L**) in ITNs from 6-OHDA mice relative

to those from controls. ns, not significant; *, p < 0.05, MWU.

Figure 5. **Progressive changes in AP waveforms of M1 PTNs during repetitive**

firing. A-B) APs from a 10- to15-Hz spike train were superimposed for PTN from

control (**A**) or 6-OHDA mice (**B**). APs were aligned at threshold, and the solid and

dashed lines indicated the first and the last APs from the spike train, respectively. All the

other APs were shown as transparent lines. **C-D**) Summarized graphs showing the

alterations of AP width between the first and the last APs. Each line in (**C**) connects the

AP width of the first and the last APs from the same spike train. Filled and open circles

indicate the median plus interquartile range of the first and the last AP width of PTNs,

respectively. (**D**) Boxplots showing the difference between the AP width of the first and

the last APs was greater in PTNs from 6-OHDA mice relative to those from controls. **E-**

F) Summarized graphs showing alterations of AP decay rate between the first and the

last APs (**E**) and greater changes of AP decay rate in PTNs from 6-OHDA mice relative

to those from controls (**F**). **G-H**) Summarized graphs showing alterations of AP decay

time between the first and the last APs (**G**), and greater changes of AP decay time in

PTNs from 6-OHDA mice relative to those from controls (**H**). **I-J**) Summarized graphs

showing alterations of AP threshold between the first and the last APs (**I**), and greater

changes of AP threshold in PTNs from 6-OHDA mice relative to those from controls (**J**).

ns, not significant; *, $p < 0.05$, WSR.

Figure 6. Decreased BK channel activity underlies AP broadening and impaired

intrinsic excitability of PTNs following loss of midbrain DA neurons. A)

Representative AP waveforms showing diminished fAHPs (arrow) in PTNs from 6-OHDA

mice, relative to those from controls. **B-C**) Boxplot showing decreased amplitude of fAHPs

(**B**) and unaltered mAHPs (**C**) in PTNs from 6-OHDA mice relative to those from controls.

D) Representative APs waveforms of PTNs from controls and 6-OHDA mice in the

presence of BAPTA. APs were aligned at the threshold and overlaid for comparison. **E**)

Boxplot showing reversal of broadened AP of PTNs in 6-OHDA mice relative to controls

after intracellular BAPTA dialysis. **F**) Summarized graph showing unaltered excitability of

PTNs from 6-OHDA mice relative to those from controls in the presence of BAPTA. The

number of APs elicited by 400-, 600-, and 800-pA somatic current injections were

compared. **G and H**) Representative APs waveforms before and after paxillin application

in PTNs from controls (**G**) and 6-OHDA mice (**H**). APs were aligned at the threshold and

overlaid for comparison. **I**) Summarized graph showing paxillin broadened APs of PTNs

from both controls and 6-OHDA mice. **J**) Summarized graph showing that paxillin

significantly decreased the number of spikes (evoked by injecting 720-pA current) of

PTNs from controls but did not affect those from 6-OHDA mice. Application of paxillin

occluded the decreased number of spikes of PTNs from 6-OHDA mice relative to those

from controls. Filled and open symbols in (**I and J**) indicate the values of before and after

paxillin application, respectively. ns, not significant; *, $p < 0.05$, WSR or MWU.

Figure 7. DA application could not rescue intrinsic adaptations in PTNs and ITNs of M1

following loss of midbrain DA neurons. **A-B**) Representative AP traces showing the

effect of DA application (10 µM) on the number of APs of PTNs from controls (**A**) and

the summarized results (**B**). **C-D**) Representative AP traces showing the effect of DA

application on the number of APs of PTNs from 6-OHDA mice (**C**) and the summarized

results (**D**). **E-G**) Summarized graphs showing the effect of DA (10 µM) on intrinsic

properties of ITNs from controls and 6-OHDA mice, including the V_m (**E**), R_m (**F**),

rheobase (**G**) of ITNs from controls and 6-OHDA mice. Filled and open circles in (**E-G**)

indicate the values of before and after DA application, respectively. Each line connected

the rheobase before (filled circle) and after (open circle) DA application. *, $p < 0.05$,

WSR.

References cited:

Albin RL, Young AB, Penney JB (1989) The functional anatomy of basal ganglia

disorders. Trends Neurosci 12:366–375.

Barthas F, Kwan AC (2017) Secondary motor cortex: where 'sensory' meets 'motor' in

the rodent frontal cortex. Trends Neurosci 40:181–193.

Bean BP (2007) The action potential in mammalian central neurons. Nat Rev Neurosci

8:451–465.

Bevan MD, Magill PJ, Terman D, Bolam JP, Wilson CJ (2002) Move to the rhythm:

oscillations in the subthalamic nucleus–external globus pallidus network. Trends

Neurosci 25:525–531.

Bodor ÁL, Giber K, Rovó Z, Ulbert I, Acsády L (2008) Structural correlates of efficient

GABAergic transmission in the basal ganglia–thalamus pathway. J Neurosci 28:3090–

3102.

Buchta WC, Mahler SV, Harlan B, Aston-Jones GS, Riegel AC (2017) Dopamine

terminals from the ventral tegmental area gate intrinsic inhibition in the prefrontal cortex.

Physiol Rep 5:e13198.

Chu H-Y, Atherton JF, Wokosin D, Surmeier DJ, Bevan MD (2015) Heterosynaptic

regulation of external globus pallidus inputs to the subthalamic nucleus by the motor

cortex. Neuron 85:364–376.

Chu H-Y, McIver EL, Kovaleski RF, Atherton JF, Bevan MD (2017) Loss of hyperdirect

pathway cortico-subthalamic inputs following degeneration of midbrain dopamine

neurons. Neuron 95:1306-1318.e5.

Contet C, Goulding SP, Kuljis DA, Barth AL (2016) BK channels in the central nervous

system. Int Rev Neurobiol 128:281–342.

Corbit VL, Whalen TC, Zitelli KT, Crilly SY, Rubin JE, Gittis AH (2016) Pallidostriatal

projections promote β oscillations in a dopamine-depleted biophysical network model. J

Neurosci 36:5556–5571.

Cousineau J, Lescouzères L, Taupignon A, Delgado-Zabalza L, Valjent E, Baufreton J,

Bon-Jégo ML (2020) Dopamine D2-like receptors modulate intrinsic properties and

synaptic transmission of parvalbumin interneurons in the mouse primary motor cortex.

Eneuro:ENEURO.0081-20.2020.

DeLong MR (1990) Primate models of movement disorders of basal ganglia origin.

Trends Neurosci 13:281–285.

Ebbesen CL, Brecht M (2017) Motor cortex — to act or not to act? Nat Rev Neurosci

18:694–705.

Ebbesen CL, Doron G, Lenschow C, Brecht M (2017) Vibrissa motor cortex activity

suppresses contralateral whisking behavior. Nat Neurosci 20:82–89.

Ebbesen CL, Insanally MN, Kopec CD, Murakami M, Saiki A, Erlich JC (2018) More

than just a "motor": recent surprises from the frontal cortex. J Neurosci 38:9402–9413.

Economo MN, Viswanathan S, Tasic B, Bas E, Winnubst J, Menon V, Graybuck LT,

Nguyen TN, Smith KA, Yao Z, Wang L, Gerfen CR, Chandrashekar J, Zeng H, Looger

LL, Svoboda K (2018) Distinct descending motor cortex pathways and their roles in

movement. Nature 563:79–84.

Faber ESL, Sah P (2003) Ca2+-activated K+ (BK) channel inactivation contributes to

spike broadening during repetitive firing in the rat lateral amygdala. J Physiol 552:483–

497.

Fieblinger T, Graves SM, Sebel LE, Alcacer C, Plotkin JL, Gertler TS, Chan CS,

Heiman M, Greengard P, Cenci MA, Surmeier DJ (2014) Cell type-specific plasticity of

striatal projection neurons in parkinsonism and L-DOPA-induced dyskinesia. Nat

Commun 5:5316.

Fleidervish IA, Friedman A, Gutnick MJ (1996) Slow inactivation of Na+ current and

slow cumulative spike adaptation in mouse and guinea-pig neocortical neurones in

slices. J Physiol 493:83–97.

Galvan A, Wichmann T (2008) Pathophysiology of parkinsonism. Clin Neurophysiology

Official J Int Fed Clin Neurophysiology 119:1459–1474.

Gittis AH, Hang GB, LaDow ES, Shoenfeld LR, Atallah BV, Finkbeiner S, Kreitzer AC

(2011) Rapid target-specific remodeling of fast–spiking inhibitory circuits after loss of

dopamine. Neuron 71:858–868.

Goldberg JA, Boraud T, Maraton S, Haber SN, Vaadia E, Bergman H (2002) Enhanced

synchrony among primary motor cortex neurons in the 1-Methyl-4-Phenyl-1,2,3,6-

Tetrahydropyridine primate model of Parkinson's disease. J Neurosci 22:4639–4653.

Gradinaru V, Mogri M, Thompson KR, Henderson JM, Deisseroth K (2009) Optical

deconstruction of parkinsonian neural circuitry. Science 324:354–359.

Guo J-Z, Graves AR, Guo WW, Zheng J, Lee A, Rodríguez-González J, Li N, Macklin

JJ, Phillips JW, Mensh BD, Branson K, Hantman AW (2015a) Cortex commands the

performance of skilled movement. Elife 4:e10774.

Guo L, Xiong H, Kim J-I, Wu Y-W, Lalchandani RR, Cui Y, Shu Y, Xu T, Ding JB

(2015b) Dynamic rewiring of neural circuits in the motor cortex in mouse models of

Parkinson's disease. Nat Neurosci 18:1299–1309.

Hemptinne C de, Swann NC, Ostrem JL, Ryapolova-Webb ES, Luciano MS,

Galifianakis NB, Starr PA (2015) Therapeutic deep brain stimulation reduces cortical

phase-amplitude coupling in Parkinson's disease. Nat Neurosci 18:779–786.

Hosp JA, Luft AR (2013) Dopaminergic meso-cortical projections to M1: role in motor

learning and motor cortex plasticity. Front Neurol 4:145.

Hosp JA, Nolan HE, Luft AR (2015) Topography and collateralization of dopaminergic

projections to primary motor cortex in rats. Exp Brain Res 233:1365–1375.

Hosp JA, Pekanovic A, Rioult-Pedotti MS, Luft AR (2011) Dopaminergic projections

from midbrain to primary motor cortex mediate motor skill learning. J Neurosci 31:2481–

2487.

Jaffe DB, Wang B, Brenner R (2011) Shaping of action potentials by type I and type II

large-conductance Ca2+-activated K+ channels. Neuroscience 192:205–218.

Kawai R, Markman T, Poddar R, Ko R, Fantana AL, Dhawale AK, Kampff AR, Ölveczky

BP (2015) Motor cortex is required for learning but not for executing a motor skill.

Neuron 86:800–812.

Kita H, Kita T (2011) Cortical stimulation evokes abnormal responses in the dopamine-

depleted rat basal ganglia. J Neurosci 31:10311–10322.

Kress GJ, Yamawaki N, Wokosin DL, Wickersham IR, Shepherd GMG, Surmeier DJ

(2013) Convergent cortical innervation of striatal projection neurons. Nat Neurosci

16:665–667.

Lee J, Wang W, Sabatini BL (2020) Anatomically segregated basal ganglia pathways

allow parallel behavioral modulation. Nat Neurosci. In press.

Li Q, Ke Y, Chan DCW, Qian Z-M, Yung KKL, Ko H, Arbuthnott GW, Yung W-H (2012)

Therapeutic deep brain stimulation in parkinsonian rats directly influences motor cortex.

Neuron 76:1030–1041.

Little S, Brown P (2014) The functional role of beta oscillations in Parkinson's disease.

Parkinsonism Relat D 20 Suppl 1:S44-8.

Magill PJ, Sharott A, Bevan MD, Brown P, Bolam JP (2004) Synchronous unit activity

and local field potentials evoked in the subthalamic nucleus by cortical stimulation. J

Neurophysiol 92:700–714.

Magno LAV, Tenza-Ferrer H, Collodetti M, Aguiar MFG, Rodrigues APC, Silva RS da,

Silva J do P, Nicolau NF, Rosa DVF, Birbrair A, Miranda DM, Romano-Silva MA (2019)

Optogenetic stimulation of the M2 cortex reverts motor dysfunction in a mouse model of

Parkinson's disease. J Neurosci 39:3234–3248.

Mallet N, Pogosyan A, Sharott A, Csicsvari J, Bolam JP, Brown P, Magill PJ (2008)

Disrupted dopamine transmission and the emergence of exaggerated beta oscillations

in subthalamic nucleus and cerebral cortex. J Neurosci 28:4795–4806.

Mathai A, Ma Y, Paré J-F, Villalba RM, Wichmann T, Smith Y (2015) Reduced cortical

innervation of the subthalamic nucleus in MPTP-treated parkinsonian monkeys. Brain

138:946–962.

McGregor MM, Nelson AB (2019) Circuit mechanisms of Parkinson's disease. Neuron

101:1042–1056.

McIver EL, Atherton JF, Chu H-Y, Cosgrove KE, Kondapalli J, Wokosin D, Surmeier DJ,

Bevan MD (2019) Maladaptive downregulation of autonomous subthalamic nucleus

activity following the loss of midbrain dopamine neurons. Cell Reports 28:992-1002.e4.

Oswald MJ, Tantirigama MLS, Sonntag I, Hughes SM, Empson RM (2013) Diversity of

layer 5 projection neurons in the mouse motor cortex. Front Cell Neurosci 7:174.

Parker JG, Marshall JD, Ahanonu B, Wu Y-W, Kim TH, Grewe BF, Zhang Y, Li JZ, Ding
JB, Ehlers MD, Schnitzer MJ (2018) Diametric neural ensemble dynamics in
parkinsonian and dyskinetic states. Nature 557:177–182.

Parker PRL, Lalive AL, Kreitzer AC (2016) Pathway-specific remodeling of
thalamostriatal synapses in parkinsonian mice. Neuron 89:734–740.

Parrilla-Carrero J, Buchta WC, Goswamee P, Culver O, McKendrick G, Harlan B,
Moutal A, Penrod R, Lauer A, Ramakrishnan V, Khanna R, Kalivas P, Riegel AC (2018)
Restoration of Kv7 channel-mediated inhibition reduces cued-reinstatement of cocaine
seeking. J Neurosci 38:4212–4229.

Pasquereau B, DeLong MR, Turner RS (2016) Primary motor cortex of the parkinsonian
monkey: altered encoding of active movement. Brain 139:127–143.

Pasquereau B, Turner RS (2011) Primary motor cortex of the parkinsonian monkey:
differential effects on the spontaneous activity of pyramidal tract-type neurons. Cereb
Cortex 21:1362–1378.

Pathak D, Guan D, Foehring RC (2016) Roles of specific Kv channel types in
repolarization of the action potential in genetically identified subclasses of pyramidal
neurons in mouse neocortex. J Neurophysiol 115:2317–2329.

Rudy B, McBain CJ (2001) Kv3 channels: voltage-gated K+ channels designed for high-
frequency repetitive firing. Trends Neurosci 24:517–526.

Ryan MB, Bair-Marshall C, Nelson AB (2018) Aberrant striatal activity in parkinsonism

and levodopa-induced dyskinesia. Cell Reports 23:3438-3446.e5.

Sanders TH, Jaeger D (2016) Optogenetic stimulation of cortico-subthalamic projections

is sufficient to ameliorate bradykinesia in 6-OHDA lesioned mice. Neurobiol Dis 95:225–

237.

Sauerbrei BA, Guo J-Z, Cohen JD, Mischiati M, Guo W, Kabra M, Verma N, Mensh B,

Branson K, Hantman AW (2020) Cortical pattern generation during dexterous

movement is input-driven. Nature 577:386–391.

Shao L, Halvorsrud R, Borg-Graham L, Storm JF (1999) The role of BK-type Ca2+-

dependent K+ channels in spike broadening during repetitive firing in rat hippocampal

pyramidal cells. J Physiol 521:135–146.

Sharott A, Vinciati F, Nakamura KC, Magill PJ (2017) A population of indirect pathway

striatal projection neurons is selectively entrained to parkinsonian beta oscillations. J

Neurosci 37:9977–9998.

Sheets PL, Suter BA, Kiritani T, Chan CS, Surmeier DJ, Shepherd GMG (2011)

Corticospinal-specific HCN expression in mouse motor cortex: Ih -dependent synaptic

integration as a candidate microcircuit mechanism involved in motor control. J

Neurophysiol 106:2216–2231.

Shen W, Plotkin JL, Francardo V, Ko WKD, Xie Z, Li Q, Fieblinger T, Wess J, Neubig

RR, Lindsley CW, Conn PJ, Greengard P, Bezard E, Cenci MA, Surmeier DJ (2015) M4

muscarinic receptor signaling ameliorates striatal plasticity deficits in models of L-

DOPA-induced dyskinesia. Neuron 88:762–773.

Shepherd GMG (2013) Corticostriatal connectivity and its role in disease. Nat Rev

Neurosci 14:278–291.

Shimamoto SA, Ryapolova-Webb ES, Ostrem JL, Galifianakis NB, Miller KJ, Starr PA

(2013) Subthalamic nucleus neurons are synchronized to primary motor cortex local

field potentials in Parkinson's disease. J Neurosci 33:7220–7233.

Soares D, Goldrick I, Lemon RN, Kraskov A, Greensmith L, Kalmar B (2017)

Expression of Kv3.1b potassium channel is widespread in macaque motor cortex

pyramidal cells: A histological comparison between rat and macaque. J Comp Neurol

525:2164–2174.

Sreenivasan V, Esmaeili V, Kiritani T, Galan K, Crochet S, Petersen CCH (2016)

Movement initiation signals in mouse whisker motor cortex. Neuron 92:1368–1382.

Suter BA, Migliore M, Shepherd GMG (2013) Intrinsic electrophysiology of mouse

corticospinal neurons: a class-specific triad of spike-related properties. Cereb Cortex

23:1965–1977.

Viaro R, Morari M, Franchi G (2011) Progressive motor cortex functional reorganization

following 6-hydroxydopamine lesioning in rats. J Neurosci 31:4544–4554.

Vila M, Périer C, Féger J, Yelnik J, Faucheux B, Ruberg M, Raisman-Vozari R, Agid Y,

Hirsch EC (2000) Evolution of changes in neuronal activity in the subthalamic nucleus of

rats with unilateral lesion of the substantia nigra assessed by metabolic and

electrophysiological measurements. Eur J Neurosci 12:337–344.

Villalobos C, Shakkottai VG, Chandy KG, Michelhaugh SK, Andrade R (2004) SKCa

channels mediate the medium but not the slow calcium-activated afterhyperpolarization

in cortical neurons. J Neurosci 24:3537–3542.

Vitrac C, Péron S, Frappé I, Fernagut P-O, Jaber M, Gaillard A, Benoit-Marand M

(2014) Dopamine control of pyramidal neuron activity in the primary motor cortex via D2

receptors. Front Neural Circuit 8:13.

Wang X, Liu Y, Li X, Zhang Z, Yang H, Zhang Y, Williams PR, Alwahab NSA, Kapur K,

Yu B, Zhang Y, Chen M, Ding H, Gerfen CR, Wang KH, He Z (2017) Deconstruction of

corticospinal circuits for goal-directed motor skills. Cell 171:440-455.e14.

Willard AM, Isett BR, Whalen TC, Mastro KJ, Ki CS, Mao X, Gittis AH (2019) State

transitions in the substantia nigra reticulata predict the onset of motor deficits in models

of progressive dopamine depletion in mice. Elife 8:e42746.

846