Auditory Decisions in the Supplementary Motor Area

Isaac Morán,[1][§] Javier Perez-Orive,[2][§] Jonathan Melchor,[1] Tonatiuh Figueroa,[1] and Luis Lemus[1]*

[1] Department of Cognitive Neuroscience, Institute of Cell Physiology, Universidad Nacional Autónoma de México (UNAM). 04510. Mexico City, Mexico.

[2] Instituto Nacional de Rehabilitacion "Luis Guillermo Ibarra Ibarra". Mexico City, Mexico.

§ These authors contributed equally

* Corresponding author

lemus@ifc.unam.mx
Telephone: (+52) 55 5622 5675

Keywords

Word recognition, choices, premotor cortex, macaque, psychophysics

Abstract

In human speech and communication across various species, recognizing and categorizing

sounds is fundamental for the selection of appropriate behaviors. But how does the brain decide

which action to perform based on sounds? We explored whether the premotor supplementary

motor area (SMA), responsible for linking sensory information to motor programs, also accounts

for auditory-driven decision making. To this end, we trained two rhesus monkeys to discriminate

between numerous naturalistic sounds and words learned as target (T) or non-target (nT)

categories. We demonstrated that the neural population is organized differently during the

auditory and the movement periods of the task, implying that it is performing different

computations in each period. We found that SMA neurons perform acoustic-decision-related

computations that transition from auditory to movement representations in this task. Our results

suggest that the SMA integrates sensory information while listening to auditory stimuli in order

to form categorical signals that drive behavior.

Introduction

Recognition and categorization of sensory information are essential for deciding which action to

take (Parker & Newsome, 1998; Freedman et al., 2001; Romo et al., 2003; Seger & Miller,

2010). In particular, categorizing dynamic sounds into classes is necessary for social

communication in various species (Prather et al., 2009; Miller et al., 2003), including monkeys

(May, Moody & Stebbins, 1989) and humans, where grouping of discrete sounds into words is

fundamental for speech recognition (Repp, 1984; Leonard & Chang, 2014). However, it is not

clear how the brain links the recognition of complex sounds to behavior. In primates, auditory

information is carried from the auditory cortex to the prefrontal and premotor cortices via ventral

and dorsal auditory streams (Romanski et al., 1999; Kusmierek & Rauschecker, 2014). Recent

studies with macaques have shown that premotor regions may represent an embodiment of

acoustic recognitions (Archakov et al., 2020). The premotor supplementary motor area (SMA)

has been linked to voluntary action and cognitive control of movement (Nachev et al., 2008;

Lara et al., 2018; Shima & Tanji, 2000). The SMA also participates in working memory and

decision making during tactile discrimination (Romo et al., 1997; Hernandez et al., 2002; Lemus

et al., 2007), imagery, and it links auditory information with motor programs (Lima et al., 2016;

Vergara et al., 2016). Similarly, in songbirds, the premotor HVC nucleus is involved in coding

complex sounds and orchestrates singing (Prather et al., 2009).

Nevertheless, few studies have been published on naturalistic sound coding

(Rauschecker, 1998; Gentner & Margoliash, 2003; Tsunada et al., 2011; Town et al., 2018), and

even fewer in behaving non-human primates (Ng et al., 2009; Scott et al., 2012; Chandrasekaran

et al., 2013; Melchor et al., 2020). This could be because there is still plenty to learn from

experiments using simple, non-naturalistic sounds that do not require behavior (e.g. pure tones,

amplitude-modulated noise; Liang et al., 2002; Selezneva et al., 2006; Bendor & Wang, 2008;

Yin et al., 2020). Additionally, although macaques have brain organizations that are similar to

those of humans and although they, too, use vocalizations to communicate (Chang et al., 2010;

Daube et al., 2019), some consider that non-human primates are challenging for training and

studying auditory behavior (Ng et al., 2009; Scott et al., 2012). However, Melchor et al. (2020)

recently showed that macaques attend to and memorize numerous naturalistic sounds. Therefore,

in macaques, the SMA is amenable for studying the neural correlates of decisions within an

auditory categorization task.

Here we present data from extracellular recordings of SMA neurons from two rhesus

monkeys trained to discriminate between two groups, or categories, of naturalistic sounds and

then choose an action. We also tested acoustic morphings to reveal the underlying responses to

linear physical changes rather than discrete categorizations. We found robust categorical

responses at both the single neuron and population levels. A small number of neurons were also

found to carry information about individual sensory stimuli during the auditory period. By

mapping the neural population activity during different periods of the task, we found that it

organizes into orthogonal subspaces during the task's auditory and movement periods. Finally,

the linear morphings of one stimulus into another indicate that the coding in SMA changes from

sensory, during the auditory period, to categorical during the movement period. Taken together,

our results support the hypothesis that SMA integrates auditory information necessary for a

categorization that provides action signals for behavioral choices.

Results

Monkeys discriminate between learned naturalistic sounds

We trained two monkeys (*Macaca mulatta*) to perform an auditory recognition task. During the task, the monkeys obtained a drop of liquid as a reward for releasing the lever only after identifying a target sound (T). A T followed 0, 1, or 2 non-target sounds (nT). In other words, the T could appear in three different temporal positions (P1, P2, and P3; Figs. 1a,c, see Methods). It is important to note that we incorporated a delay and a green visual cue (VC) after each sound to disentangle auditory and movement signals. Thus, after a T, the monkeys waited for a 0.5 s delay before a 0.5 s VC-release period (Figs. 1a,c). The monkeys learned, by trial and error, to recognize up to fourteen sounds (half of them T, half nT) belonging to four different classes: conspecific vocalizations, interspecific vocalizations, words, and artificial sounds (Figs. 1b,d; see Methods). Once the monkeys performed above 85% correct responses (Fig. 1d), we recorded single neurons (1341 from monkey 1; 524 from monkey 2) from SMA (Fig. 1c (inset) during task execution).

Single units and population activity represent naturalistic sounds

Rasters of a characteristic SMA neuron (Fig. 2a) show a substantial firing rate change during a T regardless of its position (i.e. P1, P2, or P3). In other words, the responses show a clear distinction between T and nT, but with small or no differences within each category (Figs. 2b and c). The majority of neurons recorded in both monkeys (68.5% and 85.9%, respectively) exhibited this type of categorical coding (Fig.2d), as measured with the f-statistic and bootstrapping (see Methods). Interestingly, a small percentage of neurons displayed significant coding for specific sensory stimuli within the T and nT categories (2.4% and 1.1% for T; 1.1% and 2.1% for nT in each monkey, respectively; Fig. 2d). As shown in Figures 2b, 2c, and 2e, neurons with this particular sensory coding did this during the auditory periods when sounds were presented (0.0-0.5 s). In contrast, the f-statistic shows that the categorical coding starting at the end of auditory periods rises further during the delays (0.5-1.0 s) and movement periods (1.0-1.5 s; Fig. 2e).

To study the neuronal population dynamics, we used principal component analysis (PCA), a

dimensionality reduction technique. Similarly to the observed results of individual neurons, the

population dynamics differentiated between T and nT, independently of the position in which T

appears (Fig. 3a). The component explaining most of the variance shows a small increase at the

beginning of the T stimulus, followed by a larger increase during the movement period, as

expected for this premotor area. The second component shows acoustic-anticipatory activity that

continues to rise with Ts but decreases for nTs, providing a clear auditory-based signal for

premotor activity. The third principal component shows a large increase at the beginning of the

trial and an additional increase before the lever release indicating the end of the trial.

We also used demixed PCA (dPCA), a dimensionality reduction tool that can separate the

effects on the population response of different task parameters (Kobak et al., 2016), to separate

the effect of the auditory stimulus coding from the condition-independent activity (Fig.3b). Since

the previous results from single units showed strong categorical codification, we computed

dPCA for the T and nT categories. The demixed principal components (dPCs) with the most

weight explaining the variance of the population activity are those for the categorical coding

(first two categorical coding dPCs explain 57.7% and 65.3% of the total variance for each

monkey, respectively, while the first two condition-independent dPCs explain 29.6% and 29.5%

of the variance). For both monkeys, the dPCs separated the T and nT trajectories during the

auditory periods. For dPC1, this separation persisted during the delay and reached its maximum

during the movement period (Fig.3b, top). Here, dPC2 and dPC3 shift their signs in the delay or

during the movement period. The neural activity projections over the condition-independent

dPCs are associated mainly with the task's temporal dynamics.

Auditory and movement computations occupy different subspaces

After observing the differences between auditory and movement period responses of single

neurons and the SMA population, we set out to determine whether such representations

proceeded from distinct population organizations, each coming together at different time periods

to perform different computations (Elsayed et al., 2016). Figure 3c highlights the correlation

matrices for the full neural population, organized so the structure during the auditory period is

highlighted, while keeping this same ordering for the movement period. The correlation matrices

show two large groups of neurons for both monkeys, with high correlations during the auditory

period, but not during the movement period (Fig.3c). To evaluate whether this difference was

due to two different populations, we calculated a simple epoch-preference index (Elsayed et al.,

2016; see Methods). If the preference index distribution for all neurons is bimodal, this implies

two pools of neurons. However, we found evidence that both monkeys' data came from a

unimodal distribution (Hartigan's dip test p= 0.98 and p = 0.96 for monkeys 1 and 2,

respectively), suggesting that the auditory and movement activities did not come from two

different populations.

We used PCA to investigate whether these population responses during the auditory and

movement epochs occupy partially overlapping or orthogonal subspaces within the neural

activity space. By definition, the top five PCs during the auditory period capture a large amount

of variance from this period. However, we found that they captured minimal variance from the

movement period (monkey 1: 76.9% vs. 3.3%; monkey 2: 88.8% vs. 10.9%, auditory vs.

movement period variances, respectively, for PC1 to 5 combined; Fig.3d). Similarly, the PCs

calculated for the movement period captured a large proportion of movement period variance,

but a minimum of auditory period variance (monkey 1: 4.8% vs. 86.5%; monkey 2: 12.0% vs.

92.8%, auditory- vs. movement-period variances, respectively, for PC1 to 5; Fig.3d). This result

supports the hypothesis that the auditory and movement periods use close to orthogonal

subspaces to perform their computations.

We also calculated an alignment index to quantify the subspace overlap between the

activities of both signals. Here, an index of zero means no alignment and one full alignment. The

alignment index was 0.08 and 0.23 for monkeys 1 and 2, respectively, significantly lower than an

alignment index calculated in a random neural sampling distribution ($p < 0.05$ for both

monkeys). This result indicates that the population rearranges its activity for auditory and

movement periods in a nearly orthogonal manner. Therefore, since the neuronal population's

activity behaves in an almost orthogonal fashion in the auditory and movement periods, it is

possible to implement a linear decoder to generate two orthogonal subspaces in order to explain

the neuronal variability population in both periods. Thus, we applied an algorithm (Elsayed et al.,
2016) to find two sets of bases that identify principal components of each period in an orthogonal
manner (Figs. 3e and f). The top four PCs from the auditory subspace captured 63.7% and 75.8%
of the auditory variance in monkeys 1 and 2, respectively. The top four PCs from the movement
subspace captured 76.7% and 82.0% of the movement variance for monkeys 1 and 2,
respectively. Plotting one against the other revealed that the T and nT categories separate and
move in opposite directions during the auditory period. Meanwhile, T elicits a broad trajectory
during the movement period while nT remains close to the origin (Figs. 3g and h). There is a
striking similarity between the monkeys' results, indicating that the appearance of these different
population organizations at different times in the task is not monkey-dependent.
164
165   *Linear stimulus changes modulate behavior and neural activity*
166
Gradually morphing sounds into one another (Fig.4a) may reveal the degree to which the
neuronal activity changes as a function of the physical dimensions of the stimuli or reflect the
behavioral outcome computations. By fitting sigmoidal functions to the monkeys' behavior (Fig.
4b; Table 1) and the neuronal responses to the morphings (Figs. 4c-d) it is possible to observe the
relationship between the neuronal activity and behavior. The neuronal responses of an example
neuron (Fig. 3c) identified T at morphing values between 60% and 100%, and nT at morphing
values between 0% and 40%. Remarkably, a stimulus at 50% produced a neuronal response
between the two groups. However, there was a gradual change during the auditory period,
whereas the responses transition more abruptly in the movement period. Figure 4d shows this
more precisely. The neurometric and psychometric functions are compared (gray and black lines,
respectively) during the auditory, delay, and movement periods. During the latter periods of the
task, the neurometric curves of all recorded neurons became more similar to the psychometric
curve, i.e. with steeper slopes around the 50% midpoint (Fig.4e). The overall ratios of the
neurometric/psychometric midpoint slopes increased progressively from the auditory, to the
delay, to the movement periods: 0.47, 0.60, and 0.93 for monkey 1, and 0.38, 0.69, and 0.92 for
monkey 2. Furthermore, figure 4f shows increasing Spearman correlations between neurometric
and psychometric curves for all recorded neurons as the task progresses. These results imply that
SMA neurons have more information about sensory stimuli during the auditory period and more

categorical information (thus becoming more similar to the monkeys' behavior) during the

movement period.

Demixed PCA revealed that, in response to morphing stimuli, the population activity

followed a pattern similar to that of the single neurons, distinguishing between T and nT and

with an in-between response at 50% morphing (Fig.4g). Divergence started during the auditory

period, followed by a separation to two distinct attractors during the movement period (Figs. 4g-

h). The two-dimensional dPC projections of the population activity (Fig.4H) show how neuronal

responses to T and nT sounds form two groups and move towards one of two attractors. The

transition from one attractor to the other occurred abruptly, further evidencing the SMA's role in

the categorical coding of naturalistic sounds.

The non-alert condition and errors also point to a role played by SMA in acoustic decisions

To further distinguish between sensory and categorical coding, we projected the neural activity

during the errors over the categorical dPCs calculated in correct responses. We found an

inversion of activity resembling that of the correct trials (Fig. 5a). That is, in error trials the

population activity in SMA reflected the behavioral decision made, rather than the sensory

stimulus presented. Moreover, such population activity suggests that when a signal separation

during the auditory period does not occur, monkeys make errors. We also recorded the neural

activity during sets of passive listening, i.e. when the monkeys listened but did not perform the

task because we withdrew the lever and the visual go-cues. Figure 5b shows that during these

sets, SMA ceased coding for auditory or categorical decisions. This result indicates that SMA is

only active when the monkeys engage in the task, further supporting the role of SMA

involvement during acoustic-driven decision making.

Discussion

To study whether the SMA codes for acoustic decisions, we trained two monkeys to categorize

naturalistic sounds. Observing the time courses of the neural responses, analyzed in different

ways, we present here several lines of evidence supporting the concept that SMA integrates

sensory information during the auditory period, in support of categorical representations during

the movement period: A) Demixed PCA, in which the main dPC began to separate in the

auditory period, remained constant during the delay, and then rose sharply during the movement

period. B) The two different organizational states were indicative of two different computations

being performed in each, during the auditory and movement periods. C) Single neuron

recordings during morphing sets showed coding-shifts from sensory to categorical

representations as the task progressed. D) The activity of the neuronal population diverged to

two distinct attractors, i.e. T and nT. E) Trials in which the monkeys made errors suggested that

a separation in the neural population activity seems to be needed during the auditory period;

additionally during the movement period the neuronal population with false alarms, had the same

activity as during hits.

In previous work with monkeys trained to solve a delayed match-to-sample acoustic task,

Scott et al. (2012) reported that the monkeys performed poorly in such tasks. In our study,

although the monkeys' training was more prolonged than in visual or tactile paradigms (Lemus

et al., 2009), the monkeys ended up being highly proficient in the categorization of numerous

naturalistic sounds (see also Melchor et al., 2020). Therefore, our task presents multiple

advantages for studying the neural basis of listening. However, even when the task comprised a

great diversity of complex sounds, the monkeys learned to include them in just two categories,

i.e. T and nT. Therefore, we did not explore each sound's semantics, but probably only

synonyms, i.e. that multiple sounds mean T, while others mean nT. Moreover, since the monkeys

expressed their decisions by releasing a lever or holding it down waiting for a T, one could argue

that the monkeys only had to recognize one of the two categories of sounds. Nevertheless, this

argument is not plausible since SMA neurons, like the monkeys, acknowledged both categories.

As expected for a premotor area (Nachev et al., 2008; Lara et al., 2018; Shima & Tanji,

2000), during the task's movement period, single neurons and population responses coded for the

monkeys' dichotomous decisions in the task. Furthermore, dPCA showed that the task's

categorical component captured most of the variance, as opposed to other commonly observed

results with this method, in which the condition-independent components are often those with the

most weight (Romo et al., 1999; Kobak et al., 2016). Interestingly, the second principal

component (Fig. 3a) rises before each auditory stimulus in a predictable fashion, apparently

consistent with models of accumulation of information (Leon & Shadlen, 1999). Despite the fact

that such a phenomenon, similar to a hazard function of the probability of appearance of a T, was

not previously described for auditory processing in SMA, it is consistent with its role in premotor

processing (Shima & Tanji, 2000; Nachev et al., 2008; Zimnik et al., 2019). Furthermore, this is

the first description of how SMA neuronal activity correlates with decisions based on active

listening to naturalistic sounds.

Interestingly, a small group of neurons coded for sounds during auditory periods, as

predicted by imaging studies that suggested that the premotor cortex is activated by auditory

stimuli (Lima et al., 2016; Vergara et al., 2016; Archakov et al., 2020). Nevertheless, to our

knowledge, our results constitute the first electrophysiological evidence of SMA coding natural

sounds. Previous fMRI studies using acoustic morphings (Jiang et al., 2018) suggested a two-

step category-learning process. First, a perceptual stage in the auditory cortex, and later a

categorization stage in the lateral prefrontal cortex. While our current data do not contradict

those findings, it adds further nuance by providing evidence that there is a separate computation

that processes the physical features of sensory stimuli in premotor cortex. Elsayed et al. (2016)

showed that a population of neurons in motor cortex acts as two coordinated circuits of

preparatory and movement computations. When we applied their analytical method, we, too,

found that the activity of a single neural population in SMA occupies orthogonal subspaces

during the auditory and movement periods, implying different computations in each period.

Therefore, those two periods' computations require no more than a single population of neurons

organizing itself into two different circuits.

We presented the monkeys with sounds that morphed gradually into one another, in order

to determine how SMA responds to linearly changing physical stimuli. Measuring the slope at

the midpoint of the sigmoidal curve of the neurometric and psychometric curves, we found that

during the auditory period, the neurometric curves have smaller slopes than the psychometric

curve, implying that at this moment of the task the neural response is more gradual, and varies

more linearly with the changing sensory stimulus. During the movement period, both the

neurometric and the psychometric curves had similar slopes, indicating that the neural responses

are as categorical as the behavioral choices the animals make. From a population perspective,

dPCA shows that sounds are grouped into two clear categories, with the dPC of the 50%

response in between these groups. Furthermore, as revealed by the first and second principal

components, the population response evolves towards two distinct attractors corresponding to the

T and nT categories. The transition from one of these attractors to the other is abrupt rather than

gradual, which is a hallmark of categorical coding (Wills et al., 2005; Freedman et al., 2001).

Taken together, our results show how the SMA participates in naturalistic sound-driven

decision making by integrating acoustic information to produce decision signals and motor

commands. Future work in other cortical areas that lie upstream from the SMA may provide

additional information with regard to how this categorization occurs.

Methods

Ethics statement

Animals were handled following the Official Mexican Standard for the Care and Use of

Laboratory Animals (NOM-062-ZOO-1999) and the study was approved by the Internal

Committee for the Use and Care of Laboratory Animals of the Institute of Cell Physiology,

UNAM (CICUAL; LLS80-16).

Animals and experimental setup

Two adult rhesus macaques (*Macaca mulatta*; monkey 1: a 10-year-old male weighing 13 kg;

Monkey 2: a 10-year-old female weighing 7 kg) participated in the experiments. Training and

recording sessions took place in a soundproof booth. Each monkey sat on a primate chair with its

head fixed at a distance of 60 cm from a 21 color LCD monitor (1920 x 1080 resolution, 60 Hz

refresh rate). A Yamaha MSP5 speaker (0.05-40 kHz frequency range) was situated fifteen

centimeters above and behind the monitor delivering sounds at ~ 65 dB SPL (measured at the

monkeys' ear level). Additionally, a Logitech® Z120 speaker was placed directly below the

Yamaha speaker to generate background white noise at ~ 55 dB SPL. Finally, a metal lever

placed at the monkeys' waists captured their responses.

Task

We trained two rhesus macaques to perform an acoustic recognition task. During the task, the

monkeys categorized sounds as either Target (T) or non-target (nT). In a trial, a gray circle of 3°

aperture appeared at the center of the screen. Then, a sequence of one to three sounds arose. Each

0.5 s sound continued with a 0.5 s delay and then a 0.5 s green visual cue that replaced the initial

gray circle. The cue indicated that the monkey should release the lever if the sound consisted of a

T. The monkeys received a reward for releasing the lever in a time window of 0.7 s from the start

of the T's cue. However, after an nT, the monkey kept the lever pressed, waiting for a T. If the

monkey released the lever before a T, the trial was classified as a *false alarm*, and a new one

commenced. We required that the monkeys' performance in this task remain above an 80% hit

rate before the electrophysiological recordings were made. The programming task was done in

LabVIEW 2014 (64-bit SP1, National Instruments®).

Auditory stimuli

The sounds were recorded in our laboratory or downloaded from open access online libraries

(*https://freesound.org/*). The monkeys learned more than 37 sounds from 4 different categories

(Melchor el al., 2020): conspecific vocalizations, heterospecific vocalizations, words, and

artificial sounds. However, for statistical repetitions the recordings were made with only 14

sounds (T = 7; nT = 7): conspecific vocalizations (T = 2; nT = 1), interspecific vocalizations (T =

1; nT = 1) and words (T = 2; nT = 3). The sounds' sample rate was 44.1 kHz; their amplitudes

were normalized at -10dB SPL (RMS), and their durations were shortened or lengthened to 0.5 s

(Adobe Audition®).

Morphing sounds

We created sets of morphing sounds that change from an nT to a T in a continuum sequence

(STRAIGHT software; Kawahara et al., 1999). This procedure consists of a linear interpolation

of five spectro-temporal characteristics of a pair of sounds, i.e. fundamental frequencies,

formants, duration, spectro-temporal density, and aperiodicity. The resulting sounds contained

some percentage of stimulus A and a complementary percentage of B. Thus, each set consisted

of seven morphed sounds (i.e., 0 20 40 50 60 80 and 100 % T). The monkeys trained in various

sets; however, monkeys 1 and 2 executed 6 and 4 sets during recordings.

Passive condition

In this condition, the sounds appeared in a continuous clip separated by 0.3 s interlapses, and with no other cue or visual stimulus. The monkeys did not perform since the lever was not present. However, the monkeys received a reward randomly every 15-30 s.

Neural recordings

Extracellular recordings of neurons proceeded from an array of five independently movable microelectrodes (1-3 MΩ; Thomas Recording®) inserted at different SMA locations in each session. We positioned 20 mm diameter recording chambers according to the Paxinos brain atlas coordinates (Paxinos et al., 2000) and structural magnetic resonance imaging of both monkeys (interaural 27 mm, left hemisphere: lateral 6 mm). The sampled extracellular membrane potentials were 40 kHz (Plexon®). Isolation of individual neurons was performed manually using Plexon offline sorter software (Plexon®; monkey 1) and automatically using MountainSort (Chung et al., 2017; monkey 2).

Data analysis

Individual neural coding

To evaluate whether SMA neurons coded for T and nT categories or the identity of particular sounds, we performed F-statistics in a one-way ANOVA of the T vs. nT dependent activity and individual sounds. We aligned the neuronal activity to the start of sounds and calculated the mean firing rate in 200-ms windows sliding in steps of 20 ms. To avoid biases in the F-statistic due to comparing classes of different numbers of trials, we equalized the number of trials using the lowest common number of trials in all classes. We repeated the analysis 1000 times to create an F-statistic distribution. To determine the time bins where the F-statistic was significant, we compared the actual distribution against an F-statistic random distribution obtained by shuffling the stimulus labels 1000 times. A neuron was significant if the 95% confidence intervals of the actual and random distributions did not overlap in at least one time bin.

Population analysis with PCA

We used principal component analysis (PCA) to describe the SMA's neuronal population

dynamics during the task. We calculated the neurons' firing rate in blocks of correct trials of 1, 2,

or 3 stimuli. We calculated the firing rate using a window of 200 ms moving in steps of 20 ms.

Subsequently, we concatenated the firing rate of these three types of trials to obtain a matrix $M \in$

$\mathbb{R}^{N \times ET}$ of N number of neurons, E types of trials (with 1, 2, or 3 stimuli), and T total time points.

Next, we used the demeaned firing rate of matrix M to calculate the principal components (PCs),

considering each row as a variable.

Population analysis with demixed PCA

We performed demixed principal component analysis (dPCA; Kobak et al., 2016) in order to

observe the effects of the task parameters on the population responses. During the supervised

part, dPCA decomposes the neural activity for each variable using the covariance matrices of different

marginalizations. The unsupervised analysis of dPCA was similar to that of the PCA, for each

marginalization matrix. We marginalized the population activity for T and nT and time. We

studied the neuronal activity within times from the starts of sounds to the monkeys' response, in

200 ms windows sliding every 20 ms. The dPCA algorithm separates the matrices from the

decoder (D) and the encoder (F) for each parameter of the task by minimizing the loss function:

$$L_\phi = \|\underline{X}_\phi - F_\phi D_\phi \underline{X}\|$$

Where ϕ denotes the marginalization for each parameter and $\underline{X}$ is the mean-centered

population matrix (the average of each neuron's activity is 0). Each component can be ordered

based on the amount of variance explained. The axes obtained by the decoder and encoder allow

reduction of the data in a few components that capture the most significant variance of each

parameter of the task.

We implemented a method to determine the time bins with statistically correct decoding

(Kobak et al., 2016). First, we separated the data into training and testing sets. We constructed

the testing set by randomly separating one trial from each category (T and nT). In contrast, the

training set was the average of the rest of the trials in each category. We performed dPCA from

the training set to obtain the decoding axes. Then we projected the testing set on these axes and

classified them according to the closest category mean (T or nT). We repeated this procedure

1000 times and measured the proportion of correct classification. We used 100 shuffles to

calculate the classification accuracies expected by chance. The shuffled distribution was

computed by choosing random trials between categories for each neuron. We ran the same

method described above and calculated the proportion of correct classifications for each

iteration. Then we compared the proportion of correct classifications between the real data vs.

the shuffled data. The time bins where the real classification exceeded the shuffled classification

distribution were considered statistically different from chance.

Identifying different subspaces in auditory and movement periods

To compare the population states during the auditory and movement periods, we performed four

procedures to describe the transition between one to other state and a new algorithm developed

by Elsayed et al. (2016) to calculate optimally two orthogonal subspaces, where each one

captures auditory and movement activity. 1) To compare the population covariation between

auditory and movement periods, we computed the correlation between all neurons in each

period. We created two matrices for each period: $A \in R^{N \times CT}$ and $M \in R^{N \times CT}$ where N is the

number of neurons, C is the number of stimuli and T is the time. The correlation matrix of each

period was obtained computing the Pearson correlation between the rows of each matrix (A and

M). 2) To quantify the subspace overlapping between subspaces of both periods, we computed a

PCA over $A \in R^{N \times CT}$ and $M \in R^{N \times CT}$ to obtain the top ten auditory and movement PCs. First,

the firing rate of each period was projected onto the auditory PCs and the percentage of

explained variance relative to the total variance of the auditory period was quantified. Then, the

same procedure was executed for movement PCs. 3) We calculated an alignment index to

measure the amount of auditory period variance captured by the top ten movement PCs

normalized by the auditory period variance captured by the top ten auditory PCs. The index

range varied from 0 to 1, where 0 indicates the orthogonality between both periods and 1 means

a perfect alignment. 4) To rule out that two groups of neurons are activated differentially in the

auditory and movement periods we use an epoch-preference index, which evaluates the relative

weight of each neuron in both periods. We calculated the firing rate range separately in each

period and divided it by the mean of the firing rate of the entire trial. Then the range of each

neuron was normalized by the mean of the firing rate of all neurons. A neuron with a positive

value indicates a preference for the auditory period over the movement period, a negative value
indicates the opposite. A bimodal distribution of the index signifies the existence of two neural
subpopulations, one that is more selective for auditory period and other for movement period.
Finally, we used a method developed by Elsayed et al. (2016) to derive two orthogonal
subspaces, each one capturing auditory and movement activity, respectively. This method
maximizes the sum of the variance of each period activity in each period subspace and was
calculated with the following loss function:

$$[\hat{Q}_{aud}, \hat{Q}_{mov}] = argmax_{[Q_{aud}, Q_{mov}]} \frac{1}{2} \left(\frac{Tr(Q_{aud}^T C_{aud} Q_{aud})}{\sum_{i=1}^{d_{aud}} \sigma_{aud}(i)} + \frac{Tr(Q_{mov}^T C_{mov} Q_{mov})}{\sum_{i=1}^{d_{mov}} \sigma_{mov}(i)} \right)$$

Where C_{aud} and C_{mov} are covariance matrices of the neural activity in auditory and
movement period, respectively. $\sigma_{aud}(i)$y $\sigma_{mov}(i)$ are ith singular value of the C_{aud} and C_{mov},
respectively. Q_{aud}y Q_{mov} are the eigenvectors of auditory and movement subspaces. We
selected four dimensions for Q_{aud} and Q_{mov} due they capture close to 70% variance of each
period. This method identifies simultaneously the auditory and movement subspaces while
constraining them to be orthogonal.

Psychometric and Neurometric Measures
We adjusted hyperbolic tangent functions to the psychophysical and neurometric performance
during morphing sets. This function relates the percentage of times in which the monkey
(psychometric curve) or the neuron (neurometric curve) recognized a sound like a T as a function
of a morphing sound. The function is:

$$f(\%_{morph}) = \beta 0 + \beta 2 * tanh\left(\beta 1 * (\%_{morph} - \theta)\right)$$

θ and β0 represent the midpoint of the function on the horizontal and vertical axis, respectively,
β2 represents the amplitude function from β0, and β1 defines the morphing percentage dependent
of behavior, a high value indicates an acute sigmoid. Additionally, we measured a linear slope at
the sigmoid's midpoint to assess the degree of perceptual and neural modulation as a function of
the morphing rate. We calculated the maximum and minimum values of the sigmoid's second
derivative to find the central part of the sigmoid. This section corresponds to the behavioral

transition in the categorization of morphing stimuli. After obtaining the abscissa's and the

ordinate's coordinates, we used the following equation to determine the slope:

465
$$slope = \frac{y_2 - y_1}{x_2 - x_1}$$

Neurometric function

We derived the neurometric functions from each neuron's correct trials (de Lafuente & Romo,

2005). A neurometric curve represents the probability that an ideal observer reports the stimulus

category (T or nT) based solely on the firing rate. For each neuron and time bin, the neurometric

curve was the percentage of trials in which the maximum firing rate reached or exceeded an

arbitrary threshold. The threshold was selected to maximize the number of correct answers. For

the neurometric curves, we used 300 ms windows, sliding every 50 ms. Psychometric and

neurometric sensitivity were compared based on sigmoidal slopes. We used a Spearman's

correlation between the raw data from the psychometric and neurometric curves.

Acknowledgements

We thank Elizabeth Cabrera for valuable comments. We are also grateful for the financial

support provided by CONACYT, CB-256767, and Programa de Apoyo a Proyectos de

Investigación e Innovación Tecnológica (PAPIIT) IN207919. Special thanks to Gerardo Coello

and Ana Escalante of the computing department of the IFC, to Gabriel Pérez Ruelas for technical

assistance, and to Mr Patrick Weill for proofreading the manuscript. Isaac Morán Martinez is a

doctoral student from Programa de Doctorado en Ciencias Biomédicas, Universidad Nacional

Autónoma de México (UNAM) and received fellowship 288348 from CONACYT, México. Data

in this work are part of his doctoral dissertation.

Author Contributions

IM, JM, TF & LL performed experiments. IM, & JPO analyzed data and prepared figures. LL

designed the paradigm. TF programmed the task. JPO & LL wrote the paper.

Disclosure

The authors declare no conflicts of interest.

References

Archakov, D., DeWitt, I., Kuśmierek, P., Ortiz-Rios, M., Cameron, D., Cui, D., … Rauschecker, J. P.

(2020). Auditory representation of learned sound sequences in motor regions of the

macaque brain. *Proceedings of the National Academy of Sciences of the United States of*

America, *117*(26), 15242–15252. https://doi.org/10.1073/pnas.1915610117

Bendor, D., & Wang, X. (2008). Neural response properties of primary, rostral, and

rostrotemporal core fields in the auditory cortex of marmoset monkeys. *Journal of*

Neurophysiology, *100*(2), 888–906. https://doi.org/10.1152/jn.00884.2007

Chandrasekaran, C., Lemus, L., Ghazanfar, A, A. «Dynamic faces speed up the onset of auditory

cortical spiking responses during vocal detection.» *Proceedings of the National Academy*

of Sciences 110, nº 48 (2013): E4668-E4677.

Chang, E. F., Rieger, J. W., Johnson, K., Berger, M. S., Barbaro, N. M., & Knight, R. T. (2010).

Categorical speech representation in human superior temporal gyrus. *Nature*

Neuroscience, *13*(11), 1428–1432. https://doi.org/10.1038/nn.2641

Chung, J. E., Magland, J. F., Barnett, A. H., Tolosa, V. M., Tooker, A. C., Lee, K. Y., … Greengard, L.

F. (2017). A Fully Automated Approach to Spike Sorting. *Neuron*, *95*(6), 1381-1394.e6.

https://doi.org/10.1016/j.neuron.2017.08.030

Daube, C., Ince, R. A. A., & Gross, J. (2019). Simple Acoustic Features Can Explain Phoneme-

Based Predictions of Cortical Responses to Speech. *Current Biology*, *29*(12), 1924-1937.e9.

https://doi.org/10.1016/j.cub.2019.04.067

De Lafuente, V., & Romo, R. (2005). Neuronal correlates of subjective sensory experience.

Nature Neuroscience, *8*(12), 1698–1703. https://doi.org/10.1038/nn1587

Elsayed, G. F., Lara, A. H., Kaufman, M. T., Churchland, M. M., & Cunningham, J. P. (2016).

Reorganization between preparatory and movement population responses in motor

cortex. *Nature Communications*, *7*(1), 1–15. https://doi.org/10.1038/ncomms13239

Freedman, D. J., Riesenhuber, M., Poggio, T., & Miller, E. K. (2001). Categorical representation

521   of visual stimuli in the primate prefrontal cortex. *Science*, *291*(5502), 312–316.

522   https://doi.org/10.1126/science.291.5502.312

523 Gentner, T. Q., & Margoliash, D. (2003). Neuronal populations and single cells representing

524   learned auditory objects. *Nature*, *424*(6949), 669–674.

525   https://doi.org/10.1038/nature01731

526 Hernández, Adrián and Zainos, Antonio and Romo, R. (2002). Temporal Evolution of a Decision-

527   Making Process in Medial Premotor Cortex. *Neuron*, *33*(6), 959–972. Retrieved from

528   http://www.ncbi.nlm.nih.gov/pubmed/11906701

529 Jiang, X., Chevillet, M. A., Rauschecker, J. P., & Riesenhuber, M. (2018). Training Humans to

530   Categorize Monkey Calls: Auditory Feature- and Category-Selective Neural Tuning

531   Changes. *Neuron*. https://doi.org/10.1016/j.neuron.2018.03.014

532 Kawahara, H., Masuda-Katsuse, I., & De Cheveigné, A. (1999). Restructuring speech

533   representations using a pitch-adaptive time-frequency smoothing and an instantaneous-

534   frequency-based F0 extraction: Possible role of a repetitive structure in sounds. *Speech*

535   *Communication*, *27*(3), 187–207. https://doi.org/10.1016/S0167-6393(98)00085-5

536 Kobak, D., Brendel, W., Constantinidis, C., Feierstein, C. E., Kepecs, A., Mainen, Z. F., …

537   Machens, C. K. (2016). Demixed principal component analysis of neural population data.

538   *ELife*, *5*(APRIL2016), 1–36. https://doi.org/10.7554/eLife.10989

539 Kuśmierek, P., & Rauschecker, J. P. (2014). Selectivity for space and time in early areas of the

540   auditory dorsal stream in the rhesus monkey. *Journal of Neurophysiology*, *111*(8), 1671–

541   1685. https://doi.org/10.1152/jn.00436.2013

542 Lara, A. H., Cunningham, J. P., & Churchland, M. M. (2018). Different population dynamics in the

543   supplementary motor area and motor cortex during reaching. *Nature Communications*,

544   *9*(1). https://doi.org/10.1038/s41467-018-05146-z

545 Lemus, L., Hernández, A., Luna, R., Zainos, A., Nácher, V., & Romo, R. (2007). Neural correlates

546   of a postponed decision report. *Proceedings of the National Academy of Sciences of the*

547   *United States of America*, *104*(43), 17174–17179.

548   https://doi.org/10.1073/pnas.0707961104

549 Lemus, L., Hernández, A., & Romo, R. (2009). Neural codes for perceptual discrimination of

acoustic flutter in the primate auditory cortex. *Proceedings of the National Academy of*

Sciences of the United States of America, *106*(23), 9471–9476.

https://doi.org/10.1073/pnas.0904066106

Leon, M.I., & Shadlen, M.N. (1999). Effect of expected reward magnitude on the response of

neurons in the dorsolateral prefrontal cortex of the macaque. *Neuron*, 24(2), 415-425.

Leonard, M. K., & Chang, E. F. (2014). Dynamic speech representations in the human temporal

lobe. *Trends in Cognitive Sciences*, *18*(9), 472–479.

https://doi.org/10.1016/j.tics.2014.05.001

Liang, L., Lu, T., & Wang, X. (2002). Neural representations of sinusoidal amplitude and

frequency modulations in the primary auditory cortex of awake primates. *Journal of*

Neurophysiology, *87*(5), 2237–2261. https://doi.org/10.1152/jn.2002.87.5.2237

Lima, C. F., Krishnan, S., & Scott, S. K. (2016). Roles of Supplementary Motor Areas in Auditory

Processing and Auditory Imagery. *Trends in Neurosciences*, *39*(8), 527–542.

https://doi.org/10.1016/j.tins.2016.06.003

May, B., Moody, D. B., & Stebbins, W. C. (1989). Categorical Perception Of Conspecific

Communication Sounds By Japanese Macaques. Macaca Fuscata. *Journal of the Acoustical*

Society of America, *85*(2), 837–847. https://doi.org/10.1121/1.397555

Melchor, J., Morán, I., Figueroa, T. and, & Lemus, L. (2019). Perceptual Invariance of Words and

Other Learned Sounds in Non-human Primates. *BioRxiv*, *5*(564), 1–19.

https://doi.org/10.4324/9781315853178

Miller, E. K., Nieder, A., Freedman, D. J., & Wallis, J. D. (2003). Neural correlates of categories

and concepts. *Current Opinion in Neurobiology*, *13*(2), 198–203.

https://doi.org/10.1016/S0959-4388(03)00037-0

Nachev, P., Kennard, C., & Husain, M. (2008). Functional role of the supplementary and pre-

supplementary motor areas. *Nature Reviews Neuroscience*, *9*(11), 856–869.

https://doi.org/10.1038/nrn2478

Ng, C. W., Plakke, B., & Poremba, A. (2009). Primate auditory recognition memory performance

varies with sound type. *Hearing Research*, *256*(1–2), 64–74.

https://doi.org/10.1016/j.heares.2009.06.014

Niessing, J., & Friedrich, R. W. (2010). Olfactory pattern classification by discrete neuronal

network states. *Nature*, *465*(7294), 47–52. https://doi.org/10.1038/nature08961

Parker, A. J., & Newsome, W. T. (1998). Sense and the single neuron: Probing the physiology of

perception. *Annual Review of Neuroscience*, *21*, 227–277.

https://doi.org/10.1146/annurev.neuro.21.1.227

Paxinos, G., Huang, X.-F., & Toga, A. W. (2000). The rhesus monkey brain in stereotaxic

coordinates.

Prather, J. F., Nowicki, S., Anderson, R. C., Peters, S., & Mooney, R. (2009). Neural correlates of

categorical perception in learned vocal communication. *Nature Neuroscience*, *12*(2), 221–

228. https://doi.org/10.1038/nn.2246

Quian Quiroga, R., & Panzeri, S. (2009). Extracting information from neuronal populations:

Information theory and decoding approaches. *Nature Reviews Neuroscience*, *10*(3), 173–

185. https://doi.org/10.1038/nrn2578

Rauschecker, J. P. (1998). Cortical processing of complex sounds. *Current Opinion in*

Neurobiology, *8*(4), 516–521. https://doi.org/10.1016/S0959-4388(98)80040-8

Repp, B. H. (1984). Categorical perception: Issues, methods, findings. *Speech and Language*, *10*,

243–335. https://doi.org/10.1017/CBO9781107415324.004

Romanski, L. M., Tian, B., Fritz, J., Mishkin, M., Goldman-Rakic, P. S., & Rauschecker, J. P. (1999).

Dual streams of auditory afferents target multiple domains in the primate prefrontal

cortex. *Nature Neuroscience*, *2*(12), 1131–1136. https://doi.org/10.1038/16056

Romo, R., Brody, C. D., Hernández, A., & Lemus, L. (1999). Neuronal correlates of parametric

working memory in the prefrontal cortex. *Nature*, *399*(6735), 470–473.

https://doi.org/10.1038/20939

Romo, R., Merchant, H., Zainos, A., & Hernández, A. (1997). Categorical perception of

somesthetic stimuli: Psychophysical measurements correlated with neuronal events in

primate medial premotor cortex. *Cerebral Cortex*, *7*(4), 317–326.

https://doi.org/10.1093/cercor/7.4.317

Romo, R., & Salinas, E. (2003). Cognitive neuroscience: Flutter Discrimination: Neural codes,

perception, memory and decision making. *Nature Reviews Neuroscience*, *4*(3), 203–218.

https://doi.org/10.1038/nrn1058

Scott, B. H., Mishkin, M., & Yin, P. (2012). Monkeys have a limited form of short-term memory

in audition. *Proceedings of the National Academy of Sciences of the United States of*

America, *109*(30), 12237–12241. https://doi.org/10.1073/pnas.1209685109

Seger, C. A., & Miller, E. K. (2010). Category learning in the brain. *Annual Review of*

Neuroscience, *33*, 203–219. https://doi.org/10.1146/annurev.neuro.051508.135546

Selezneva, E., Scheich, H., & Brosch, M. (2006). Dual Time Scales for Categorical Decision

Making in Auditory Cortex. *Current Biology*, *16*(24), 2428–2433.

https://doi.org/10.1016/j.cub.2006.10.027

Shima, K., & Tanji, J. (2000). Neuronal activity in the supplementary and presupplementary

motor areas for temporal organization of multiple movements. *Journal of*

Neurophysiology, *84*(4), 2148–2160. https://doi.org/10.1152/jn.2000.84.4.2148

Town, S. M., Wood, K. C., & Bizley, J. K. (2018). Sound identity is represented robustly in

auditory cortex during perceptual constancy. *Nature Communications*, *9*(1).

https://doi.org/10.1038/s41467-018-07237-3

Tsunada, J., Lee, J. H., & Cohen, Y. E. (2011). Representation of speech categories in the primate

auditory cortex. *Journal of Neurophysiology*, *105*(6), 2634–2646.

https://doi.org/10.1152/jn.00037.2011

Vergara, J., Rivera, N., Rossi-Pool, R., & Romo, R. (2016). A Neural Parametric Code for Storing

Information of More than One Sensory Modality in Working Memory. *Neuron*, *89*(1), 54–

62. https://doi.org/10.1016/j.neuron.2015.11.026

Wills, T. J., Lever, C., Cacucci, F., Burgess, N., & O'Keefe, J. (2005). Attractor dynamics in the

hippocampal representation of the local environment. *Science*, *308*(5723), 873–876.

https://doi.org/10.1126/science.1108905

Yin, P., Strait, D. L., Radtke-Schuller, S., Fritz, J. B., & Shamma, S. A. (2020). Dynamics and

Hierarchical Encoding of Non-compact Acoustic Categories in Auditory and Frontal Cortex.

Current Biology, *30*(9), 1649-1663.e5. https://doi.org/10.1016/j.cub.2020.02.047

Zimnik, A. J., Lara, A. H., & Churchland, M. M. (2019). Perturbation of macaque supplementary

motor area produces context-independent changes in the probability of movement

637   initiation. *Journal of Neuroscience*, *39*(17), 3217–3233.

638   https://doi.org/10.1523/JNEUROSCI.2335-18.2019

640 **Figure Legends**

642 **Figure 1.** *Acoustic recognition task and behavior*

643 **a)** Sequence of events in a trial: The monkey pressed the lever after a gray visual circle indicated

644 the beginning of a trial. Then, a playback of one to three sequential sounds commenced. Each

645 sound lasted 0.5 s followed by a 0.5 s delay and a 0.5 s green visual circle (go cue). The monkey

646 received a reward for releasing the lever within a 0.7 s window after the green cue following a

647 Target sound (T) (red sonograms). False alarms occurred when the monkey released the lever

648 earlier, e.g. during a non-Target sound (nT) (blue sonograms). Finally, not releasing the lever

649 during the T response window counted as a miss trial. **b)** Examples of sonograms and

650 spectrograms of four T (top) and four nT (bottom) sounds. T and nT included four classes:

651 conspecific vocalizations, interspecific vocalizations, words, and artificial sounds. **c)** Types of

652 trials regarding the number of sounds. **Inset,** Brain scheme of MRI of monkey 2 showing the

653 location of the recording chamber over SMA. **d)** Percentage of correct responses for T (red) and

654 nT (blue) for each monkey (monkey 1: circles; monkey 2: triangles). The color code of stimuli

655 will be referenced in the following figures.

657 **Figure 2.** *Individual SMA neurons code behavioral categories*

658 **a)** Rasters and peristimulus time histograms (PSTHs) of a representative neuron from monkey 1.

659 Each row of ticks is one trial aligned with the lever press; each tick is an action potential. Each

660 plot corresponds to the possible number of sounds in a trial. Thick black ticks indicate the

661 beginning of a trial. Red ticks, lever pressings. **b)** Example PSTHs of two neurons for monkey 1

662 aligned with the start of the sounds. The color lines represent the PSTHs for each T and nT

663 stimulus. The lines below (black, categorical; red, T; blue, nT) indicate bins of significant

664 coding. T stimuli are depicted in red and nT in blue palettes, as in Figure 1d. The top neuron is

665 the same as in (a). **c)** Same as (b) but for monkey 2. **d)** Percentage of recorded SMA neurons

666 with statistically significant coding of behavioral categories (Cat) or sensory information (nT and

667 T), and for each monkey. **e)** Average F-statistic for selective SMA neurons as a function of time

<center>23</center>

for both monkeys (monkeys 1: continuous lines; 2: dashed lines); **Inset**, a magnification showing

small F-statistic values for the few neurons with significant sensory coding for particular T or nT

sounds.

Figure 3. *Categorical responses of SMA population project in auditory and movement*

orthogonal subspaces

a) Projections of the neural population responses in the first three principal components. The plot

is grouped by the number of sounds, as in Figure 2a. Notice PC1 codes for lever release, PC2 for

anticipation of sounds together with the likelihood of a T, and PC3 for Ts and movements. **b)**

Left: First three dPCs for categorical coding. Each trace corresponds to T (red) and nT (blue)

projections. Right: First three condition-independent dPCs. The numbers on the top right are the

percentages of explained variance of all dPCs. The first categorical coding components explain a

greater amount of variance compared to the condition-independent components. **c)** Correlation

matrices for auditory (left) and movement (right) periods of all neurons of monkeys 1 (top) and 2

(bottom). Each dot corresponds to the correlation between two neurons. Matrices for both

periods sorted as a function of the auditory period. The groups of correlated neurons that exist in

the auditory period disappear in the movement period. **d)** Percentage of explained variance by

each of the top five acoustic (left) and movement (right) PCs projected onto the population

activity during auditory and movement periods. **e)** Projections of the neural population activity

onto the first PC of the auditory subspace for monkeys 1 (top) and 2 (bottom). **f)** Same as (e), but

for the movement subspace. **g)** Two-dimensional projection of the auditory and movement

orthogonal subspaces for monkey 1. Circles and thick lines show the auditory and movement

periods, respectively. Arrows indicate flow of time. Same color code as in Figure 1d. **h)** Same as

(g), but for monkey 2.

Figure 4. *Neural and behavioral responses to morphing sounds*

a) Spectrogram examples of gradually morphing sounds from grunt (nT) to coo (T). Every

morphing set comprises seven morphed sounds. The percentage of T contribution in each sound

is at the top of each spectrogram. **b)** Psychometric functions of monkeys (1: top; 2: bottom) as a

function of percentage of morphing. Each line is a sigmoidal fit to the behavioral performance.

The black line shows the mean performance for all morphings. **c)** Raster (top) and PSTHs

(bottom) of a representative SMA neuron during the morphing set (grunt-coo) for monkey 2. **d)**

Psychometric (black) and neurometric (gray) comparison in three periods of the neuron in c:

auditory, delay, and movement. Each plot shows the percentage of times that behavior or

neuronal response considered each morph as T. **e)** Slope ratios (neurometric/psychometric) of all

neurons in three periods: auditory, delay, and movement. Monkeys 1: black, 2: gray. Dashed

lines and numbers represent means of the distributions (monkeys 1: black; 2: grey). Note that the

neurometric curves' slopes are smaller in the auditory period and become more similar to the

psychometric curves during the movement period. **f)** Time-dependent Spearman correlations

between neurometric and psychometric data points for both monkeys. **g)** Projection of the first

two stimulus dPCs in both monkeys (1: left; 2: right). Colors correspond to a particular

percentage of morphing (color code in c). **h)** Two-dimensional projection with the first two dPCs

of the population activity for both monkeys (1: left; 2: right)—color code as in g. Circles and

thick lines show auditory and movement periods, respectively. Arrows indicate time flow.

Figure 5. *Error trials and the non-alert condition point to a participation of SMA in acoustic*

decisions

a) The categorical dPCs calculated from correct trials (top) were used to project the population

neural activity in error trials (bottom) in monkey 1. Note that when the dPCs do not separate

during the auditory period, the monkeys make errors. **b)** Projection of first dPC over the neural

population during passive trials shows no modulations.

Table 1. Overall fitting coefficients of psychometric curves.

a

non-Target   non-Target   Target

Hold lever

0.5   0.5   0.5   0.3

Time (s)

Release lever

b

coo   warble   moo   [ko.'mi.ða]

grunt   woof   [si]   ['tʃaŋ.gi]

kHz

0   0.5

Time (s)

Power (dB)
-10
-50

c

Sound position

1   2   3

Sounds per trial

d

Target

% Correct

coo   warble   moo   0.5 kHz   toing   [ko.'mi.ða]   ['pwɛɾ.ta]

non-Target

grunt   woof   miau   10 kHz   ['xaw.ʃa]   [si]   ['tʃaŋ.gi]

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

	non-Target	Target	Number of sessions	Number of neurons	Slope	θ	β0	β1	β2	R^2
Monkey 1										
	grunt	coo	7	24	2.0	35.2	34.9	0.6	41.7	0.80
	grunt	moo	6	36	1.9	60.3	51.3	0.5	51.0	0.91
	grunt	w arble	12	57	2.3	45.6	47.3	0.5	51.7	0.96
	grunt	['pw εɼ.ta]	10	47	1.9	59.6	52.2	0.4	55.9	0.93
	grunt	[ko.'mi.ða]	8	38	1.8	60.2	56.2	0.4	52.5	0.93
	[si]	moo	2	10	1.6	46.7	46.6	0.4	45.5	0.94
Monkey 2										
	grunt	coo	13	91	1.6	52.9	52.0	0.3	59.4	0.96
	grunt	moo	3	19	2.6	45.6	52.3	0.6	51.9	0.96
	grunt	w arble	3	24	1.5	51.6	51.7	0.3	56.6	0.94
	['xaw .la]	['pw εɼ.ta]	8	52	2.3	36.6	46.2	0.5	52.2	0.95

Table 1