**The molecular basis of *Acinetobacter baumannii* cadmium toxicity and**

**resistance**

Saleh F. Alquethamy[1,2#], Felise G. Adams[3#], Ram Maharjan[4#], Natasha N. Delgado[4], Maoge

Zang[3], James C. Paton[1], Karl A. Hassan[5], Ian T. Paulsen[4], Christopher A. McDevitt[1,2], Amy K.

Cain[4], Bart A. Eijkelkamp[3*]

[1] Research Centre for Infectious Diseases, School of Biological Sciences, University of

Adelaide, Adelaide, South Australia, Australia.

[2] Department of Microbiology and Immunology, The Peter Doherty Institute for Infection and

Immunity, University of Melbourne, Melbourne, Victoria, Australia.

[3] College of Science and Engineering, Flinders University, Bedford Park, S.A., 5005, Australia

[4] ARC Centre of Excellence in Synthetic Biology, Department of Molecular Sciences,

Macquarie University, Sydney, New South Wales, Australia.

[5] School of Environmental and Life Sciences, University of Newcastle, Callaghan, New South

Wales, Australia.

[#]Saleh F. Alquethamy, Felise G. Adams and Ram Maharjan contributed equally to this work.

Author order was determined by the time they joined the study.

College of Science and Engineering, Flinders University, Bedford Park, S.A., 5005, Australia.

Phone: +61-8-8201-7779. E-mail: bart.eijkelkamp@flinders.edu.au.

Running Title: *A. baumannii* cadmium stress

**ABSTRACT**

*Acinetobacter* species are ubiquitous Gram-negative bacteria that can be found in water, soil and as commensals of the human skin. The successful inhabitation of *Acinetobacter* species in diverse environments is primarily attributable to the expression of an arsenal of stress resistance determinants, which includes an extensive repertoire of metal ion efflux systems. Although metal ion homeostasis in the hospital pathogen *Acinetobacter baumannii* is known to contribute to pathogenesis, insights into its metal ion transporters for environmental persistence are lacking. Here, we studied the impact of cadmium stress on *A. baumannii*. Our functional genomics and independent mutant analyses revealed a primary role for CzcE, a member of the cation diffusion facilitator (CDF) superfamily, in resisting cadmium stress. Further, we show that the CzcCBA heavy metal efflux system also contributes to cadmium efflux. Analysis of the *A. baumannii* metallome under cadmium stress showed zinc depletion and copper enrichment, which are likely to influence cellular fitness. Overall, this work expands our understanding of the role of membrane transporters in *A. baumannii* metal ion homeostasis.

**IMPORTANCE**

Cadmium toxicity is a widespread problem, yet the interaction of this heavy metal with biological systems is poorly understood. Some microbes have evolved traits to proactively counteract cadmium toxicity, which includes *Acinetobacter baumannii*. Here we show that *A. baumannii* utilises a dedicated cadmium efflux protein in concert with a system that is primarily attuned to zinc efflux, to efficiently overcome cadmium stress. The molecular characterization of *A. baumannii* under cadmium stress revealed how active cadmium efflux plays a key role in preventing the dysregulation of bacterial metal ion homeostasis, which appeared to be the primary means by which cadmium exerts toxicity upon the bacterium.

## INTRODUCTION

The heavy metal cadmium is enriched in industrial settings and mining sites through anthropogenic activity. Although various metalloproteins retain their function when bound to cadmium (1), cadmium is primarily associated with toxicity in biological systems (2). Despite being redox inactive, cadmium stress is commonly connected to the induction of oxidative stress. Recent advances in the field have revealed cadmium-metalloprotein interactions as a primary feature of cadmium toxicity (3, 4). In addition to dysregulation of central carbon metabolism and lipid homeostasis, cadmium stress in *Streptococcus pneumoniae* causes manganese and zinc starvation via interaction with non-cognate metalloproteins, these being a manganese-recruiting protein and a zinc-responsive transcriptional regulator (4). Manganese is the primary co-factor for superoxide dismutase in *S. pneumoniae*, hence, cadmium-induced manganese depletion leads to oxidative stress susceptibility (4, 5). In *Salmonella enterica*, excess cadmium was found to affect zinc and manganese accumulation, but not iron (3). These studies exemplify that dysregulation of metal ion homeostasis and protein mismetallation by cadmium is central to its toxicity in cells.

*Acinetobacter baumannii* is rapidly emerging as one of the world's most antimicrobial resistant bacterial pathogens (6-8). Its highly effective intrinsic and acquired stress resistance features have allowed this human pathogen to survive in the hospital environment, and thrive when colonising immunocompromised individuals (9, 10). Various members of the *Acinetobacter* genus, including some strains of *A. baumannii*, have a strong environmental presence. This has resulted in the *A. baumannii* genome harbouring diverse metabolic and stress resistance features that are typically not linked to pathogenicity (11, 12). Its environmental roots are also reflected in the extensive arsenal of putative heavy metal export mechanisms that may enable persistence under diverse metal ion stresses (13, 14). *A. baumannii* metal ion biology is a

rapidly expanding field of research. In particular its diverse array of metal ion acquisition

systems have been studied intently (15-21). In contrast, our understanding of heavy metal

resistance is limited, with only the roles of zinc and copper toxicity being studied in recent

years (13, 14, 22-24). Metal ion toxicity analyses in *A. baumannii* have revealed widespread

implications on cellular physiology, including dysregulation of metal ion homeostasis,

oxidative stress susceptibility and alterations in membrane biology. Similar to the impact of

cadmium on the dysregulation of *S. pneumoniae* and *S. enterica* metal ion homeostasis, zinc

stress in *A. baumannii* has been shown to result in copper depletion (13).

The *A. baumannii* genome harbours a broad arsenal of both highly-conserved and recently-

acquired efflux systems that have putative functions in metal resistance (13). The P-type

ATPase CopA has been shown to be the primary mediator of cytoplasmic copper efflux in *A.*

*baumannii* and plays a role in its virulence potential (23). The complexity of copper

homeostasis in the periplasm has also been studied (24), but strain-to-strain variation

complicates defining their significance across the species (13, 23). Zinc resistance is mediated

by an efficient efflux pathway which consists of the cation diffusion family (CDF) member

CzcD, a periplasmic zinc chaperone named CzcI and the heavy metal efflux (HME) system

CzcABC (22).

Bacterial cadmium efflux can be mediated by HME, CDF or P-type ATPase family members.

Examples include CadA, a P-type ATPase from *Streptococcus thermophilus* (25), the CDF

member YiiP from *Escherichia coli* (26, 27) and the CznABC HME system from *Helicobacter*

*pylori* (28). These systems may display overlapping substrate profiles within a bacterial

species, which requires experimental verification. Although a recent study has provided

insights into the role of the C-terminal domain in CDF metal specificity (29), overall, the

specificity for transition metals varies greatly between members within the CDF, HME and P-

type ATPase families and is difficult to predict based on protein sequence analyses.

In this study, we define the genome-wide cadmium stress adaptation mechanisms in *A.*

*baumannii*. This work identified both CDF and HME systems that play key roles in cadmium

export in *A. baumannii* which was confirmed by single-gene mutant analyses. We also show

that the bacterium encodes a highly efficient cadmium sensing regulator for the transcriptional

activation of the primary cadmium resistance mechanism. Through the global metallomic

analyses under cadmium stress, this study provides novel insights into cadmium toxicity.

**METHODS**

**Bacterial strains, chemicals, media and growth conditions**

*Acinetobacter baumannii* AB5075_UW mutant derivatives were purchased from the Manoil Laboratory (30) and T26 transposon insertions confirmed by PCR. The *A. baumannii* ATCC 17978 *czcA* and *czcE* mutants were generated in our laboratory for a previous study (22) (See **Tables S1** and **S2** for list of strains and oligonucleotides used in the study, respectively). All chemicals were purchased from Sigma Aldrich (Australia) unless otherwise indicated.

*A. baumannii* strains were routinely grown in Luria Bertani broth (LB), containing 1% tryptone (BD Bacto), 0.5% yeast extract (BD Bacto) and 0.5% sodium chloride. For overnight culturing, a single colony from LB agar (1.5%) was used to inoculate 4 mL of LB medium. Overnight cultures were diluted to an optical density at 600 nm ($OD_{600}$) of 0.01 in 200 µL LB for growth assays or 20 mL LB for all other analyses. For growth assays, cultures in LB media were incubated at 37°C with shaking in a FLUOstar Omega Spectrophotometer (BMG Labtech), with the $OD_{600}$ values presented. The 20 mL cultures used for all other analyses were incubated at 37°C in an Innova 40R shaking incubator (Eppendorf) at 230 rpm until they reached an $OD_{600}$ of 0.7.

**Construction of transposon mutant library**

*A. baumannii* ATCC 17978 was used to generate a dense transposon library. The library was constructed using the protocol as previously described (31). Briefly, transposomes were prepared using an EZ-Tn*5* transposase (Epicentre Biotechnology) and a custom transposon carrying a kanamycin resistance cassette amplified from the pUT_Km plasmid (32) using the primer sets listed in **Table S2**. The transposomes (0.25 µL) were electroporated into 60 µL of freshly prepared electrocompetent *A. baumannii* cells using a Bio-Rad GenePulser II (1.8 kV,

25 µF, and 200 Ω) in a 1 mm electrode gap cuvette (Bio-Rad). Cells were recovered by

resuspension in 1 mL of SOC medium and incubated at 37°C with shaking (200 rpm) for 2 h.

Tn5 insertion recombinants were selected on Mueller Hinton (MH) agar (BD Bacto)

supplemented with 7 µg.mL$^{-1}$ kanamycin. Approximately 12 to 16 transformations were

performed and pooled for each batch, with 10,000 to 50,000 resulting transformants.

Approximately 250,000 mutants were collected and pooled from a total of 10 batches and snap-

frozen and stored in 20% glycerol at -80°C.

**Time kill assay for the selection of cadmium concentration for mutant library treatment**

Approximately $10^9$ CFU from an overnight ATCC 17978 culture was sub-cultured into fresh

10 mL MH broth spiked with different $CdCl_2$ concentrations (0, 3, 60, 80 160, and 320 µM)

and incubated at 37°C with shaking. At 0, 1, 2, 4, 5 and 24 h time points, 100 µL samples were

taken, 10-fold serially diluted in sterile phosphate-buffered saline (PBS) and 10 µL of each

dilution spotted on MH agar.  Colonies were enumerated to determine the surviving cells after

overnight incubation at 37°C.

**Transposon-directed insertion site sequencing (TraDIS) of mutant library and data**

**analysis**

Approximately $10^9$ viable Tn5 mutant cells were inoculated into 10 mL of MH broth (BD

Bacto) and grown at 37°C for 8 h with shaking (200 rpm). A total of 500 µL containing

approximately $10^9$ cells were seeded into 10 mL fresh MH broth with or without 60 µM $CdCl_2$

in duplicate and grown for 16 h at 37°C with shaking (200 rpm). Genomic DNA was then

extracted from approximately $10^{10}$ cells using the DNeasy UltraClean Microbial Kit (Qiagen)

according to the manufacture's protocol. Sequencing and analysis of the transposon mutant

library were performed using the transposon-directed insertion site sequencing (TraDIS)

protocol as described previously (33). The primer sets used for PCR amplification of TraDIS

fragments (Pf5_PCR) and sequencing (Pf5_Seq) are listed in **Table S2**. Samples were

sequenced on a HiSeq2500 Illumina sequencing platform, generating approximately 2 million

50 bp single-end reads per sample as previously described (34). TraDIS sequence reads were

deposited in the European Nucleotide Archive under accession number ERP118051 and

analysed using the BioTraDIS pipeline with default parameters (as described in (33)).

**Cellular metal ion content analysis**

Untreated and metal-stressed bacteria (LB supplemented with 1 μM $CdCl_2$ or 15 μM $CdCl_2$)

were harvested at mid log-phase ($OD_{600}$ 0.7) and washed, by resuspension and centrifugation

at 7,000 × $g$ for 8 mins, first with PBS containing 5 mM ethylenediaminetetraacetic acid

(EDTA), and then with PBS as described previously (22, 23, 35, 36). Bacterial pellets were

desiccated at 95°C overnight, followed by determination of dry cell weight, and resuspension

of pellets in 35% $HNO_3$ and boiling at 95°C for 1 h. Samples were diluted to a final

concentration of 3.5% $HNO_3$ and analysed by inductively coupled plasma-mass spectrometry

(ICP-MS) on an Agilent Solution 8900 QQQ ICP-MS (Adelaide Microscopy, University of

Adelaide).

**qRT-PCR**

For RNA extraction and qRT-PCR analysis, untreated or metal-stressed bacteria (LB

supplemented with 30 μM $CdCl_2$) were harvested when they reached mid-log phase ($OD_{600}$

0.7) and cells lysed in QiaZol (Qiagen) as described previously (37, 38). RNA was extracted

and purified using a PureLink RNA Mini Kit (ThermoFisher Scientific), with on-column

DNase I treatment, according to the manufacturer's instructions. qRT-PCR was performed

using the SuperScript III One-Step RT-PCR kit (ThermoFisher Scientific) on a QuantStudio 7

Flex System (ThermoFisher Scientific). Transcription levels were corrected to those obtained

for *GAPDH* prior to normalization to the transcription levels observed for untreated *A.*

*baumannii* cultures. Oligonucleotide sequences are listed in (**Table S2**).

**Bacterial lipid extraction and gas chromatography/mass spectrometry analysis**

Overnight cultures of *A. baumannii* AB5075_UW were diluted to an $OD_{600}$ of 0.05 in fresh LB

media (20 ml) and grown to mid-log phase ($OD_{600} = 0.7$), with treated cultures supplemented

with 15 μM $CdCl_2$. Cells were harvested by centrifugation at $7,000 \times g$ for 10 minutes and

washed once with PBS, processed pellets were then resuspended in 50 μl of 1.5% NaCl buffer.

For lipid extraction, 1 ml of chloroform:methanol (2:1; v/v) was added to the cell suspension,

mixed vigorously for 2 minutes then incubated at RT for 10 minutes. Following the addition

of 200 μl 1.5% NaCl, the cell suspension was mixed vigorously for 1 minute and centrifuged

at $6,000 \times g$ for phase-separation. The lower phase was recovered and concentrated via nitrogen

evaporation. All samples were stored at -20°C prior to gas chromatography / mass spectrometry

(GC/MS) analysis.

To generate fatty acid methyl esters (FAMEs), concentrated lipid samples were resuspended in

1:1 chloroform and trimethylsulfonium hydroxide (TMSOH) and subsequently analysed using

an Agilent 7890A GC system with a 30 m Agilent DB-FastFAME column (Agilent

Technologies). Mass spectrometry was completed using a coupled Agilent 5975C MSD system

(Agilent Technologies). FAME species were differentiated and determined by comparing to

the FAME mix c4-24 standard (Sigma Aldrich). Data analysis was completed using the Agilent

MassHunter Qualitative Navigator software (Agilent Technologies).

## RESULTS AND DISCUSSION

### *A. baumannii* employs multiple transport systems to resist cadmium stress

To delineate the molecular basis of cadmium toxicity and the mechanisms that aid in cadmium stress resistance we created a Tn*5*-based random transposon mutant library in the laboratory reference strain, *A. baumannii* ATCC 17978. TraDIS sequencing of the base library in rich media revealed a density of at least 113,000 unique mutants. We subjected $10^9$ mutant cells to a subinhibitory concentration (60 μM) of cadmium for 16 h. This concentration was found to present a minor but significant impact on proliferation of this strain (**Fig. S1**), and thus deemed appropriate for TraDIS analysis. Cadmium exposure resulted in 56 genes displaying significantly differential numbers of insertions (**Table S3**): decreased mutant fitness (decreased transposon insertions) in 14 genes and enhanced fitness (increased transposon insertions) in 42 genes (fold change of $\geq 2$; *Q* value, $\leq 0.05$).

The TraDIS analyses identified ACX60_13165, which encodes a putative MerR-like transcriptional regulator, as most affected by cadmium treatment (-4.1 fold change) (**Fig. 1**). The regulatory gene is divergently transcribed to the CDF family transport gene, *czcE* (ACX60_13160), which our analyses defined as the gene with the second greatest loss of represented transposon mutants under cadmium stress (-3.8 fold change). The previously characterized HME-RND zinc efflux system, CzcCBA (ACX60_01405-15) and its metal chaperone *czcI* (ACX60_01400) (22), were also shown to be important in conferring cadmium resistance, where mutant abundance decreased between -2.4 and -3.0 fold change. The CDF member, *czcD*, previously revealed to play a role in zinc efflux, was not found to be significantly impacted by cadmium treatment. We identified that transposon mutants of putative DmeF-like cobalt exporting CDF (ACX60_18165) were enriched under cadmium stress (2.1 fold change). This indicates that cobalt may be accumulated at reduced levels under

cadmium stress and its undesired export by this DmeF-like transporter is subsequently

detrimental to *A. baumannii* fitness.

A solute binding protein (SBP; ACX60_10710) with similarity to methionine and metal ion

binding SBPs such as NlpA (COG1464.1), was impacted by cadmium stress (-3.2 fold change).

The gene encoding this periplasmic protein is located in an operon that also encodes its putative

ATP-binding cassette (ABC) transporter for cargo delivery and subsequent translocation across

the inner membrane, but none of these genes were affected by cadmium stress. Hence, the SBP

may play a role in periplasmic metal buffering instead of facilitating transport.

The relatively large number of genes in which insertional disruption proved to impart a positive

effect included various transcriptional regulators. H-NS (12.6 fold change) is one of *A.*

*baumannii*'s most prominent global transcriptional regulators (39). Although, no role in metal

ion stress has been indicated in previous transcriptomic analyses of an ATCC 17978 *hns*

mutant, in *Salmonella*, the ferric uptake regulator (FUR) regulates the expression of *hns* and

H-NS competes with FUR for some gene targets in *Salmonella* (40, 41). The GacAS regulatory

system, of which mutation is beneficial to survival during cadmium stress (11.7 and 12.1 fold

change, respectively), plays a key role during infection and in lifestyle changes, including

biofilm formation (42). Other regulators include those with putative roles in sensing saturated

fatty acids, including ACX60_05275 (3.3 fold change) and ACX60_17890 (2.0 fold change).

This may indicate that *A. baumannii* adapts to cadmium stress by changing the structure and

subsequently the biophysical properties of its membrane. Fatty acid analysis of *A. baumannii*

under cadmium stress revealed minor but significant changes in 16:1 (1.5% decrease) and 18:0

(0.9% increase) (**Fig. S2**). These changes may indicate a slight increase in the membrane

rigidity following cadmium exposure.

### *A. baumannii* employs a dedicated inner membrane cadmium exporter

To confirm the roles of the putative cadmium transporters in cadmium stress resistance as

observed by TraDIS (**Fig. S3**), we examined mutants in *czcA* (ABUW_0268), *czcD*

(ABUW_0269), *czcE* (ABUW_2851) and *czcF* (ABUW_3664). *A. baumannii* AB5075_UW

and its mutant derivatives were grown with increasing concentrations of cadmium (1, 5, 10, 20

or 30 μM $CdCl_2$) (**Fig. 2A-E**). We found that inactivation of *czcE* rendered the cells hyper-

susceptible to cadmium stress, with significant growth perturbation seen at concentrations as

low as 1 μM cadmium, which is approximately 30-fold lower than the AB5075_UW parental

strain. In addition to a primary role in zinc resistance (22), CzcCBA contributes to cadmium

resistance, with the *czcA*::T26 mutant showing increased sensitivity at 20 μM cadmium or

greater. Consistent with the examination of these efflux systems in strain AB50875_UW, the

ATCC 17978 Δ*czcA* and Δ*czcE* mutants displayed increased susceptibility to cadmium stress

(**Fig. S4**). CzcD and CzcF were found not to play a major role in cadmium resistance. The

concerted efforts from this study and previous work (22) have identified major roles for CzcA,

CzcD and CzcE in zinc and/or cadmium resistance, but no major role has been identified for

CzcF. Hence, we examined the response of CzcF to stress induced by other transition metals

(zinc, copper, nickel, cobalt and iron), which revealed a minor role in zinc and nickel resistance

(**Fig. S5**).

### The impact of cadmium on the *A. baumannii* metallome

Bacterial cells necessitate diligent balancing of metal ions as their hyper-accumulation or

deficit beyond the cellular setpoints has detrimental impacts on metalloprotein function and

has the potential to affect the homeostasis of other metals. First, we examined cadmium

accumulation in *A. baumannii* strain AB5075_UW and the *czcA*::T26 and *czcE*::T26 mutant

strains. These analyses revealed that CzcA inactivation did not impact accumulation following

supplementation with 1 µM cadmium. In contrast, the *czcE*::T26 mutant accumulated cadmium

at nearly 8-fold higher levels compared to the wild-type (**Fig. 3A**). Since growth of the

*czcE*::T26 mutant was dramatically perturbed at concentrations greater than 1 µM cadmium,

only the wild-type and *czcA*::T26 mutant were examined at 15 µM cadmium. The *czcA*::T26

mutant accumulated cadmium at greater levels than those observed for the wild-type (144 vs.

86 µg cadmium.g cells$^{-1}$) (**Fig. 3A**).

Examination of the impact of cadmium on the most abundant *A. baumannii* metal ions revealed

that zinc and copper levels were affected, but not iron (**Fig. 3B-D**). Zinc levels in the wild-type

were significantly lower upon exposure to 1 µM cadmium and below detection when exposed

to 15 µM cadmium. Consistent with these findings, zinc levels in the *czcA*::T26 and *czcA*::T26

mutants decreased to a greater extent upon exposure to 1 µM cadmium compared to the wild-

type (**Fig. 3B**). Interestingly, the wild-type and *czcA*::T26 mutant hyper-accumulated copper

when exposed to 15 µM cadmium (**Fig. 3D**). Despite the similar cellular cadmium levels

observed in the *czcE*::T26 mutant following treatment with 1 µM cadmium and the wild-type

with 15 µM cadmium (**Fig. 2A**), the accumulation of copper was not affected in the *czcE*::T26

mutant (**Fig. 2D**).

**Cadmium stress influences the *A. baumannii* metalloregulator responses**

Transcriptional analyses revealed that *czcE* is up-regulated 480-fold by cadmium stress (**Fig.**

**4A**), while *czcI* was significantly down-regulated by cadmium stress. Analysis of CadR

(ABUW_2852), a MerR-type regulator divergently transcribed from *czcE*, suggested that it

was responsible for activation of *czcE*, which is consistent with the TraDIS analysis (**Fig. 4B**).

We found that *czcF* was significantly up-regulated upon treatment with cadmium, despite it

not playing a major role in cadmium resistance (**Fig. 4I and S6B**). Similar to *czcE*, we found

that *czcF* is regulated by a divergently transcribed MerR-type regulator, ABUW_3665 (**Fig.**

**4C**).

Our metallomic analyses identified that cadmium toxicity impacts zinc and copper homeostasis

(**Fig. 2**). To ascertain the impact of cadmium stress on the expression of metal ion homeostasis

contributors, we analysed transcription of *znuA*, a component of the major zinc acquisition

system ZnuABC (15), *oprC*, a putative copper acquisition system (43), and *copA*, the primary

copper efflux system (22). Under cadmium stress, *A. baumannii* increases expression of *znuA*

(3.3 $\log_2$-fold) (**Fig. 4D**), possibly in an attempt to compensate for zinc starvation. Interestingly,

cadmium-induced copper overaccumulation may be a result of the significant up-regulation of

*oprC* (8.2 $\log_2$-fold) (**Fig. 4D**).

**CONCLUSION**

Cadmium resistance in *A. baumannii* is primarily mediated by a distinct CDF member, CzcE. However, mutation of *czcA* results in *A. baumannii* growth perturbation in the presence of cadmium, which suggests that cytoplasmic cadmium is translocated into the periplasm by CzcE and then exported into the extracellular space by CzcCBA. These findings on the roles of CDFs in conjunction with HME systems in the cytoplasmic and periplasmic translocation of substrates, respectively, are consistent with work performed with *Wautersia metallidurans* (44). Further, previous works have indicated that cadmium exerts greater toxicity in the cytoplasm compared to that of the periplasm (45, 46), which highlights the major significance of CzcE, and the lesser contribution of CzcA in *A. baumannii* cadmium resistance. Interestingly, transcriptional profiling revealed that *czcE* is transcriptionally activated by cadmium, whereas *czcA* transcription is not. Hence, the degree of CzcCBA contribution to cadmium resistance may also be due to the lack of up-regulation, rather than an inability of export from either the cytoplasm or periplasm.

Cadmium stress also results in copper over-accumulation with transcriptional analyses revealing *oprC* to be a possible copper import system. Although *oprC* has been shown to be regulated by OxyR in *Acinetobacter oleivorans* (47), recent analyses of OxyR in *A. baumannii* did not identify such association (48). Hence, the physiological role of OprC warrants further investigation. Zinc homeostasis was also affected by cadmium stress, resulting in a reduction in cellular zinc. However, the molecular basis behind cadmium-induced zinc depletion in *A. baumannii* remains unclear. These results show that cadmium stress results in an inverse dysregulation of copper and zinc homeostasis, which at is at least in part consistent with our previous analysis of these metals (13).

In summary, this study has identified the primary cadmium efflux pathways of *A. baumannii*.

These systems are highly conserved across this species and genus, hence this work is broadly

applicable to *Acinetobacter* biology, exemplified by our parallel analysis of two distinct strains.

We have garnered new knowledge into the environmental stress adaptation strategies, with

systems that have unique and shared metal ion specificities. The analyses of cadmium stress

across a range of distinct metal ion export systems has also provided novel insights into the

alignment of metal sensors and efflux systems, which is of major significance to bacterial

fitness in a broad range of environments.

**Funding**

This work was supported by the National Health and Medical Research Council (Australia) through Project Grant 1159752 to BAE and AKC. AKC was supported by an Australian Research Council (ARC) DECRA fellowship (DE180100929), CAM is an ARC Future Fellow (FT170100006) and KAH is an ARC Future Fellow (FT180100123).

**Authors' contributions**

BAE, KAH, AKC, ITP, JCP and CAM designed the study. SFA, FGA, RM, KAH, NND and BAE performed the experiments. BAE, FGA, SFA, KAH, AKC, RM, ITP, CAM and JCP contributed to the drafting of the manuscript.

**REFERENCES**

1. Malgieri G, Palmieri M, Esposito S, Maione V, Russo L, Baglivo I, de Paola I, Milardi D, Diana D, Zaccaro L, Pedone PV, Fattorusso R, Isernia C. 2014. Zinc to cadmium replacement in the prokaryotic zinc-finger domain. Metallomics 6:96-104.

2. Dietrich N, Tan CH, Cubillas C, Earley BJ, Kornfeld K. 2016. Insights into zinc and cadmium biology in the nematode *Caenorhabditis elegans*. Arch Biochem Biophys 611:120-133.

3. Ammendola S, Cerasi M, Battistoni A. 2014. Deregulation of transition metals homeostasis is a key feature of cadmium toxicity in Salmonella. Biometals 27:703-14.

4. Begg SL, Eijkelkamp BA, Luo Z, Counago RM, Morey JR, Maher MJ, Ong CL, McEwan AG, Kobe B, O'Mara ML, Paton JC, McDevitt CA. 2015. Dysregulation of transition metal ion homeostasis is the molecular basis for cadmium toxicity in *Streptococcus pneumoniae*. Nat Commun 6:6418.

5. Eijkelkamp BA, Morey JR, Ween MP, Ong CL, McEwan AG, Paton JC, McDevitt CA. 2014. Extracellular zinc competitively inhibits manganese uptake and compromises oxidative stress management in *Streptococcus pneumoniae*. PLoS One 9:e89427.

6. WHO. 2017. WHO publishes list of bacteria for which new antibiotics are urgently needed. WHO Media centre.

7. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C, Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M, Cox EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N, Group WHOPPLW. 2018. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis 18:318-327.

8. CDC. 2019. 2019 Antibiotic Resistance Threats Report. CDC.

9. Cerqueira GM, Peleg AY. 2011. Insights into *Acinetobacter baumannii* pathogenicity. IUBMB Life 63:1055-60.

10.    Peleg AY, de Breij A, Adams MD, Cerqueira GM, Mocali S, Galardini M, Nibbering
PH, Earl AM, Ward DV, Paterson DL, Seifert H, Dijkshoorn L. 2012. The success of
acinetobacter species; genetic, metabolic and virulence attributes. PLoS One 7:e46984.

11.    Adams MD, Goglin K, Molyneaux N, Hujer KM, Lavender H, Jamison JJ, MacDonald
IJ, Martin KM, Russo T, Campagnari AA, Hujer AM, Bonomo RA, Gill SR. 2008.
Comparative genome sequence analysis of multidrug-resistant *Acinetobacter*
*baumannii*. J Bacteriol 190:8053-64.

12.    Farrugia DN, Elbourne LD, Hassan KA, Eijkelkamp BA, Tetu SG, Brown MH, Shah
BS, Peleg AY, Mabbutt BC, Paulsen IT. 2013. The complete genome and phenome of
a community-acquired *Acinetobacter baumannii*. PLoS One 8:e58628.

13.    Hassan KA, Pederick VG, Elbourne LD, Paulsen IT, Paton JC, McDevitt CA,
Eijkelkamp BA. 2017. Zinc stress induces copper depletion in Acinetobacter
baumannii. BMC Microbiol 17:59.

14.    Williams CL, Neu HM, Gilbreath JJ, Michel SL, Zurawski DV, Merrell DS. 2016.
Copper resistance of the emerging pathogen *Acinetobacter baumannii*. Appl Environ
Microbiol 82:6174-6188.

15.    Hood MI, Mortensen BL, Moore JL, Zhang Y, Kehl-Fie TE, Sugitani N, Chazin WJ,
Caprioli RM, Skaar EP. 2012. Identification of an *Acinetobacter baumannii* zinc
acquisition system that facilitates resistance to calprotectin-mediated zinc
sequestration. PLoS Pathog 8:e1003068.

16.    Mortensen BL, Rathi S, Chazin WJ, Skaar EP. 2014. *Acinetobacter baumannii* response
to host-mediated zinc limitation requires the transcriptional regulator Zur. J Bacteriol
196:2616-26.

17.    Nairn BL, Lonergan ZR, Wang J, Braymer JJ, Zhang Y, Calcutt MW, Lisher JP, Gilston
BA, Chazin WJ, de Crecy-Lagard V, Giedroc DP, Skaar EP. 2016. The response of
*Acinetobacter baumannii* to zinc starvation. Cell Host Microbe 19:826-36.

18.    Gaddy JA, Arivett BA, McConnell MJ, Lopez-Rojas R, Pachon J, Actis LA. 2012. Role
of acinetobactin-mediated iron acquisition functions in the interaction of *Acinetobacter*

*baumannii* strain ATCC 19606T with human lung epithelial cells, *Galleria mellonella*

caterpillars, and mice. Infect Immun 80:1015-24.

19.    Nwugo CC, Gaddy JA, Zimbler DL, Actis LA. 2011. Deciphering the iron response in

*Acinetobacter baumannii*: A proteomics approach. J Proteomics 74:44-58.

20.    Zimbler DL, Penwell WF, Gaddy JA, Menke SM, Tomaras AP, Connerly PL, Actis

LA. 2009. Iron acquisition functions expressed by the human pathogen *Acinetobacter*

*baumannii*. Biometals 22:23-32.

21.    Eijkelkamp BA, Hassan KA, Paulsen IT, Brown MH. 2011. Investigation of the human

pathogen *Acinetobacter baumannii* under iron limiting conditions. BMC Genomics

12:126.

22.    Alquethamy SF, Adams FG, Naidu V, Khorvash M, Pederick VG, Zang M, Paton JC,

Paulsen IT, Hassan KA, Cain AK, McDevitt CA, Eijkelkamp BA. 2020. The role of

zinc efflux during *Acinetobacter baumannii* infection. ACS Infect Dis 6:150-158.

23.    Alquethamy SF, Khorvash M, Pederick VG, Whittall JJ, Paton JC, Paulsen IT, Hassan

KA, McDevitt CA, Eijkelkamp BA. 2019. The role of the CopA copper efflux system

in *Acinetobacter baumannii* virulence. Int J Mol Sci 20.

24.    Williams CL, Neu HM, Alamneh YA, Reddinger RM, Jacobs AC, Singh S, Abu-Taleb

R, Michel SLJ, Zurawski DV, Merrell DS. 2020. Characterization of *Acinetobacter*

*baumannii* copper resistance reveals a role in virulence. Front Microbiol 11:16.

25.    Schirawski J, Hagens W, Fitzgerald GF, Van Sinderen D. 2002. Molecular

characterization of cadmium resistance in *Streptococcus thermophilus* strain 4134: an

example of lateral gene transfer. Appl Environ Microbiol 68:5508-16.

26.    Lu M, Fu D. 2007. Structure of the zinc transporter YiiP. Science 317:1746-8.

27.    Wei Y, Fu D. 2006. Binding and transport of metal ions at the dimer interface of the

Escherichia coli metal transporter YiiP. J Biol Chem 281:23492-502.

28.    Stahler FN, Odenbreit S, Haas R, Wilrich J, Van Vliet AH, Kusters JG, Kist M,

Bereswill S. 2006. The novel *Helicobacter pylori* CznABC metal efflux pump is

required for cadmium, zinc, and nickel resistance, urease modulation, and gastric
colonization. Infect Immun 74:3845-52.

29. Udagedara SR, La Porta DM, Spehar C, Purohit G, Hein MJA, Fatmous ME, Casas
Garcia GP, Ganio K, McDevitt CA, Maher MJ. 2020. Structural and functional
characterizations of the C-terminal domains of CzcD proteins. J Inorg Biochem
208:111087.

30. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK,
Zurawski DV, Brittnacher MJ, Manoil C. 2015. Resources for genetic and genomic
analysis of emerging pathogen *Acinetobacter baumannii*. J Bacteriol 197:2027-35.

31. Langridge GC, Phan MD, Turner DJ, Perkins TT, Parts L, Haase J, Charles I, Maskell
DJ, Peters SE, Dougan G, Wain J, Parkhill J, Turner AK. 2009. Simultaneous assay of
every Salmonella Typhi gene using one million transposon mutants. Genome Res
19:2308-16.

32. Herrero M, de Lorenzo V, Timmis KN. 1990. Transposon vectors containing non-
antibiotic resistance selection markers for cloning and stable chromosomal insertion of
foreign genes in gram-negative bacteria. J Bacteriol 172:6557-67.

33. Barquist L, Mayho M, Cummins C, Cain AK, Boinett CJ, Page AJ, Langridge GC,
Quail MA, Keane JA, Parkhill J. 2016. The TraDIS toolkit: sequencing and analysis for
dense transposon mutant libraries. Bioinformatics 32:1109-11.

34. Hassan KA, Cain AK, Huang T, Liu Q, Elbourne LD, Boinett CJ, Brzoska AJ, Li L,
Ostrowski M, Nhu NT, Nhu Tdo H, Baker S, Parkhill J, Paulsen IT. 2016.
Fluorescence-Based Flow Sorting in Parallel with Transposon Insertion Site
Sequencing Identifies Multidrug Efflux Systems in Acinetobacter baumannii. MBio 7.

35. Pederick VG, Eijkelkamp BA, Begg SL, Ween MP, McAllister LJ, Paton JC, McDevitt
CA. 2015. ZnuA and zinc homeostasis in *Pseudomonas aeruginosa*. Sci Rep 5:13139.

36. Eijkelkamp BA, Begg SL, Pederick VG, Trapetti C, Gregory MK, Whittall JJ, Paton
JC, McDevitt CA. 2018. Arachidonic acid stress impacts Pneumococcal fatty acid
homeostasis. Front Microbiol 9:813.

37.  Hassan KA, Jackson SM, Penesyan A, Patching SG, Tetu SG, Eijkelkamp BA, Brown
MH, Henderson PJ, Paulsen IT. 2013. Transcriptomic and biochemical analyses
identify a family of chlorhexidine efflux proteins. Proc Natl Acad Sci U S A 110:20254-
9.

38.  Giles SK, Stroeher UH, Eijkelkamp BA, Brown MH. 2015. Identification of genes
essential for pellicle formation in *Acinetobacter baumannii*. BMC Microbiol 15:116.

39.  Eijkelkamp BA, Stroeher UH, Hassan KA, Elbourne LD, Paulsen IT, Brown MH. 2013.
H-NS plays a role in expression of *Acinetobacter baumannii* virulence features. Infect
Immun 81:2574-83.

40.  Prajapat MK, Saini S. 2012. Interplay between Fur and HNS in controlling virulence
gene expression in Salmonella typhimurium. Comput Biol Med 42:1133-40.

41.  Troxell B, Sikes ML, Fink RC, Vazquez-Torres A, Jones-Carson J, Hassan HM. 2011.
Fur negatively regulates hns and is required for the expression of HilA and virulence in
*Salmonella enterica* serovar Typhimurium. J Bacteriol 193:497-505.

42.  Cerqueira GM, Kostoulias X, Khoo C, Aibinu I, Qu Y, Traven A, Peleg AY. 2014. A
global virulence regulator in *Acinetobacter baumannii* and its control of the
phenylacetic acid catabolic pathway. J Infect Dis 210:46-55.

43.  Abdollahi S, Rasooli I, Mousavi Gargari SL. 2018. The role of TonB-dependent copper
receptor in virulence of *Acinetobacter baumannii*. Infect Genet Evol 60:181-190.

44.  Munkelt D, Grass G, Nies DH. 2004. The chromosomally encoded cation diffusion
facilitator proteins DmeF and FieF from *Wautersia metallidurans* CH34 are
transporters of broad metal specificity. J Bacteriol 186:8036-43.

45.  Helbig K, Bleuel C, Krauss GJ, Nies DH. 2008. Glutathione and transition-metal
homeostasis in *Escherichia coli*. J Bacteriol 190:5431-8.

46.  Helbig K, Grosse C, Nies DH. 2008. Cadmium toxicity in glutathione mutants of
*Escherichia coli*. J Bacteriol 190:5439-54.

47.  Kim J, Cho Y, Jang IA, Park W. 2015. Molecular mechanism involved in the response
to hydrogen peroxide stress in *Acinetobacter oleivorans* DR1. Appl Microbiol
Biotechnol 99:10611-26.

48.  Juttukonda LJ, Green ER, Lonergan ZR, Heffern MC, Chang CJ, Skaar EP. 2018.
*Acinetobacter baumannii* OxyR regulates the transcriptional response to hydrogen
peroxide. Infect Immun doi:10.1128/IAI.00413-18.

 **FIGURE LEGENDS**

**Figure 1. The *A. baumannii* cadmium resistome.** The effect of cadmium treatment (60 $\mu M$

for 16 hours) on the abundance of gene mutations (Tn*5* insertions) mapping to the *A. baumannii*

ATCC 17978 chromosome and plasmid pAB3 as determined by TraDIS analysis. The data

represent the mean of biological triplicates. The *Q* value was determined by using the method

described previously (33).

**Figure 2. Cadmium resistance in *A. baumannii*.** The effects of cadmium (1, 5, 10, 20 or 30

$\mu M$ CdCl$_2$) upon growth of the **(A)** wild-type, **(B)** *czcA*::T26*, **(C)** *czcD*::T26, **(D)** *czcE*::T26 or

**(E)** *czcF*::T26 mutant cells was compared to that of untreated (UT) cells. The growth was

examined by measuring the optical density at 600 nm (OD$_{600}$). The data are the mean of at least

biological triplicates ($\pm$ SEM). The visibility of error bars may be occluded by the symbols.

**Figure 3. The impact of cadmium exposure on *A. baumannii* metal ion homeostasis.** The

accumulation of **(A)** cadmium, **(B)** iron, **(C)** zinc and **(D)** copper in wild-type, *czcA*::T26 or

*czcE*::T26 cells with or without cadmium supplementation (1 $\mu M$ or 15 $\mu M$ CdCl$_2$), as

determined by inductively coupled plasma – mass spectrometry. Metal ion levels were

quantified as the weight of the metal ($\mu g$) per the weight of the dry cell pellet (g). For all panels,

the data are the mean of at least biological triplicates ($\pm$ SEM). Statistical analyses were

performed using a two-tailed Student's *t*-test; n.s. = not significant, * = $p < 0.05$, ** = $p < 0.01$

and **** = $p < 0.0001$. The detection limit of zinc is indicated by the dotted line (b.d. = below

detection).

**Figure 4. Regulation of cadmium resistance. (A)** The transcriptional responses of *czcI*, *czcC*,

*czcA*, *czcD*, *czcE* and *czcF* in wild-type AB5075_UW cells treated with 30 $\mu M$ cadmium were

examined by qRT-PCR. The transcription levels of **(B)** *czcE* or **(C)** *czcF* were examined in

regulator mutant and AB5075_UW wild-type cells by qRT-PCR following treatment with 30

µM cadmium. **(D)** The transcription levels of *znuA*, *oprC* and *copA* in wild-type cells were

examined by qRT-PCR following treatment with 30 µM cadmium. For all panels, the data are

the mean (±SEM) of at least biological triplicates. The visibility of error bars may be occluded

by the symbols. Statistical analyses were performed using a two-tailed Student *t*-test; n.s. = not

significant, * = $p < 0.05$, ** = $p < 0.01$ and *** = $p < 0.001$.