The 55K SNP-Based Exploration of QTL for Spikelet Number Per Spike in a Tetraploid Wheat (*Triticum turgidum* L.) Population: Chinese Landrace 'Ailanmai' × Wild Emmer

Ziqiang Mo[1,+], Jing Zhu[1,+], Jiatai Wei[1], Jieguang Zhou[1], Qiang Xu[1], Huaping Tang[1], Yang Mu[1], Mei Deng[1], Qiantao Jiang[1], Yaxi Liu[1], Guoyue Chen[1], Jirui Wang[1], Pengfei Qi[1], Wei Li[2], Yuming Wei[1], Youliang Zheng[1], Xiujin Lan[1,*], Jian Ma[1,*]

[1]State Key Laboratory of Crop Gene Exploration and Utilization in Southwest China, Triticeae Research Institute, Sichuan Agricultural University, Chengdu, 611130, China

[2]College of Agronomy, Sichuan Agricultural University, Chengdu, 611130, China

[+] Contributed equally to this work.

*Authors for correspondence:

Dr. Jian Ma, E-mail: jianma@sicau.edu.cn

Dr. Xiujin Lan, E-mail: lanxiujin@163.com

Abstract

The spikelet number per spike (SNS) is a primary factor determining wheat yield. Common wheat breeding reduces the genetic diversity among elite germplasm resources, leading to a detrimental effect on future wheat production. It is, therefore, necessary to explore new genetic resources for SNS to increase wheat yield. A tetraploid landrace 'Ailanmai' × wild emmer wheat recombinant inbred line (RIL) population was used to construct a genetic map using the wheat 55K single nucleotide polymorphism (SNP) array. The linkage map containing 1150 bin markers with a total genetic distance of 2411.8 cM was obtained. Based on phenotypic data from eight environments and best linear unbiased prediction (BLUP) values, five quantitative trait loci (QTL) for SNS were identified, explaining 6.71-29.40% of the phenotypic variation. Two of them, *QSns.sau-AM-2B.3* and *QSns.sau-AM-3B.2*, were detected as the major and novel QTL. Their effects were further validated in two additional F_2 populations using tightly linked Kompetitive Allele-Specific PCR (KASP) markers. Potential candidate genes within the physical intervals of corresponding QTL were predicted to participate in inflorescence development and spikelet formation. Genetic associations between SNS and other agronomic traits were also detected and analyzed. This study demonstrates the feasibility of wheat 55K SNP array developed for common wheat in genetic mapping of tetraploid population and shows the potential application of wheat related species in wheat improvement programs.

Keywords: Tetraploid wheat, high-quality genetic map, spikelet number per spike, QTL mapping, genetic correlations, potential application

Introduction

Wheat (*Triticum* sp.) is among the most widely planted crops globally that provides about 20% of the calories consumed by humans (http://www.fao.org/faostat). Breeding high-yield wheat cultivars is a sustainable approach to meet the demands of a growing human population. Wheat yield is determined by spike number per unit area (SNPA), kernel number per spike (KNS), and kernel weight (KW). KNS is jointly determined by the spikelet number per spike (SNS) and the kernel number per spikelet (KNL). SNS depends on the number of lateral spikelets produced before the spike meristem transitions to the terminal spikelet. It is a complex quantitative trait greatly affected by genetic and environmental factors (Gao et al., 2019; Kuzay et al., 2019). Therefore, dissecting the genetic mechanism of SNS at the QTL level can provide more insights into the role of SNS in yield formation (Ma et al., 2019a; Yao et al., 2019; Chen et al., 2020; Li et al., 2021).

A few genes related to spikelet number and morphology have been reported. For example, the *WHEAT FRIZZY PANICLE* (*WFZP*) gene encodes a member of the APETALA2/ethylene response transcription factor (AP2/ERF) family, which influences the supernumerary spikelet trait in common wheat (Dobrovolskaya et al., 2015). Regulatory expression of the two pivotal flowering genes, *Photoperiod-1* (*Ppd-1*) and *FLOWERING LOCUS T* (*FT*), increases the number of fertile spikelets (Boden et al., 2015). The inflorescence growth rate and the development of paired spikelets are controlled by the *TEOSINTE BRANCHED1* (*TB1*) expression level (Dixon et al., 2018). Similarly, the ortholog of the rice *APO1, WHEAT ORTHOLOG OF APO1* (*WAPO1*) affects the spikelet number (Kuzay et al., 2019). Overexpression of *TaFL-2D* increases the number of spikelets and florets (Wang et al., 2017).

More loci for SNS should thus be identified and utilized owing to its important role

in yield formation. Nonetheless, common wheat breeding reduces the genetic

diversity among elite germplasm resources, thus adversely affecting wheat yield

(Cavanagh et al., 2013; Sthapit et al., 2020). Notably, the numerous genetic

resources for agronomic traits and disease resistance from wheat related species may

effectively solve future wheat production challenges (Zaïm et al., 2017; El Haddad et

al., 2021). As the progenitor of modern cultivated tetraploid and hexaploid wheat,

wild emmer wheat (*T. turgidum* subsp. *dicoccoides*) is the secondary gene pool of

common wheat. It is resistant to various diseases and possesses important agronomic

traits, thereby playing a potentially significant role in wheat breeding (Xie and Nevo,

2008; Yu et al., 2020). For example, *Yr15* derived from wild emmer wheat confers

resistance to yellow rust (Klymiuk et al., 2018). The reduced-function allele *GNI-A1*

increases the number of fertile florets per spikelet (Sakuma et al., 2019). Cognizant

of this, exploiting the genetic resources of wild emmer wheat is crucial in solving the

existing bottleneck in wheat breeding. Wheat landraces are also primary materials

for wheat improvement. They contain unique gene resources preserved under

long-term natural selection and human intervention with strong adaptability to local

environmental conditions and high production potential (Lopes et al., 2015; Crossa

et al., 2016).

In the current research, a high-quality genetic map was constructed based on a RIL

population developed from the cross between a Chinese landrace 'Ailanmai' and

wild emmer using the wheat 55K SNP array to identify QTL for SNS, and major and

novel QTL were validated in the different genetic backgrounds by KASP markers.

Materials and Methods

Plant Materials

A tetraploid wheat population (AM population) containing 121 F_8 recombinant

inbred lines (RILs, including two parental lines) derived from a Chinese landrace

'Ailanmai' (AL, *T. turgidum* L. cv. Ailanmai, 2n=28, AABB) and a wild emmer

accession (LM001, *T. turgidum* subsp. *dicoccoides*) were used in this study. AL is a

durum wheat landrace native to Sichuan Province, China. It has the advantages of

dwarf, multiple floret and strong environmental adaptability (Liu et al., 1999). AM

RIL population was developed aiming at identifying favorable alleles from AL and

LM001 to speed up their breeding utilization. The spike morphology of the parents

and representative lines are shown in Figure 1. Major and novel SNS QTL identified

in the AM population were validated in two populations including LM001 ×

PI193877 (*T. turgidum* ssp. *dicoccon*) (139 F_2 lines) and AL × AS2268 (*T. carthlicum*

Nevski) (100 F_2 lines). The above plant materials were preserved by Triticeae

Research Institute, Sichuan Agricultural University.

Trait Measurement and Data Analysis

Trait measurements of AM population were performed in eight environments:

Chongzhou (103° 38' E, 30° 32' N) in 2017, 2018, 2019, 2020, and 2021 (E_1, E_2, E_3,

E_4, and E_7), Wenjiang (103° 51' E, 30° 43' N) in 2020 and 2021 (E_5 and E_8), and

Ya'an (103° 0' E, 29° 58' N) in 2020 (E_6). Field experiments adopted a randomized

complete block design, and were managed following the conventional practices of

wheat production. The plot for a single line was space-planted (0.1-m) in a single

1.5-m row with a 0.3-m of row spacing. The agronomic traits were measured using

the methods described by Liu et al. (2020a). Spikelet number per spike (SNS) was

measured by counting the spikelet number of the main-spike per plant; plant height

(PH) was determined by the distance from the soil surface to the tip of the

main-spike (excluding awns) per plant; spike length (SL) was measured by the

length from the base to the tip of the main-spike (excluding awns) per plant; kernel

number per spike (KNS) was counted as the kernel number of the main-spike per

plant; spike density (SD) was calculated by dividing the SNS by SL; thousand kernel

weight (TKW) was calculated as 10 times the average weight of 100 seeds with three

repetitions in each line. Anthesis date (AD) was the number of days between sowing

and half of the plants flowering in each line (Ma et al., 2019a). Spike extension

length (SEL) was measured as the distance between the base of the main-spike and

the petiole of the flag leaf per plant (Li et al., 2020). At least four disease-free plants

with consistent growth status were selected for measurements. The environmental

information of surveyed agronomic traits of AM population is listed in

Supplementary Table 1. Two F_2 validation populations were grown in Chongzhou in

2021 and disease-free plants were selected to measure SNS.

The IBM SPSS Statistics 25 (Armonk, NY, USA) was used to analyze the

phenotypic data, including Pearson's correlation, frequency distribution, standard

error, and the Student's t-test ($P < 0.05$). The broad-sense heritability (H^2) and best

linear unbiased prediction (BLUP) of agronomic traits from multiple environments

were calculated using SAS version 8.0 (Cary, North Carolina).

Construction of the Genetic Map

High-quality genomic DNA was isolated from fresh leaves using the Plant Genomic

DNA Kit (Tiangen Biotech, Beijing, China). The DNA of 119 RILs and two parental

lines was subsequently genotyped by CapitalBio Technology (Beijing, China) using

the wheat 55K SNP array, and RILs were used for linkage map construction.

The genetic map was constructed following the methods described by Liu et al.

(2018) and Lin et al. (2021) . Firstly, the Poly High-Resolution markers from A and

B genomes showing a minor allele frequency (≥ 0.3) among the mapping population

were retained for subsequent analysis. Secondly, the BIN function of QTL

IciMapping 4.1 (Meng et al., 2015) was used to analyze the remaining markers based

on their segregation patterns in the mapping population, with the parameters

'Distortion value' and 'Missing rate' being set as 0.01 and 20%, respectively. A

single marker with the lowest 'Missing rate' in each bin (bin marker) was selected to

construct genetic maps. Finally, the bin markers were grouped and sorted using the

Kosambi mapping function in IciMapping 4.1 with the logarithm of odds (LOD) ≥ 6

after preliminary analysis of SNP markers with LOD values ranging from 2 to 10.

The genetic maps were drawn using MapChart 2.3.2 (Voorrips, 2002). The syntenic

relationships between the genetic and physical maps of the bin markers were further

presented using the Strawberry Perl 5.24.0.1 (Rothwell, 2019).

QTL Analysis

QTL analysis was performed using the inclusive composite interval mapping (ICIM)

in the Biparental Populations (BIP) module of QTL IciMapping 4.1. The LOD

threshold was set as 3.4 based on a test of 1000 permutations in IciMapping which

was used to determine the threshold that corresponded to a genome-wide false

discovery rate of 5% ($P < 0.05$), and loci detected in only single environment were

removed. The QTL-by-Environment interaction of SNS trait was analyzed using the

function of multi-environment trials (MET-ADD) in IciMapping (Step = 1 cM, PIN

= 0.001, and LOD ≥ 3.4). Epistatic interaction of SNS QTL was also identified using

the multi-environmental trials (MET-EPI) pattern of IciMapping. Based on the initial

QTL mapping results, we developed new KASP markers to densify the genetic map

and narrow the mapping interval. The genomic DNA of two parental lines was

collected and genotyped by CapitalBio Technology (Beijing, China) using the wheat

660K SNP array. Polymorphic SNPs detected in initial mapping region of two

parental lines were converted to KASP markers to genotype the mapping population

and integrated into the genetic map.

QTL identified in at least five environments (including BLUP) and explained more

than 10% of the phenotypic variation were regarded as major ones, while those

having common flanking markers were treated as a single one. QTL was then named

following the Genetic Nomenclature provision

(http://wheat.pw.usda.gov/ggpages/wgc/98/Intro.htm). In the naming system, 'Sau'

represented 'Sichuan Agricultural University'. The RILs' name 'AM' was also added

to the QTL's name to distinguish them from others reported previously.

Sequences of the markers were blasted against the durum wheat (Svevo) (Maccaferri et al., 2019), wild emmer (Zavitan; v2.0) (Zhu et al., 2019), and common wheat genotype 'Chinese Spring' (CS; v2.1) (Zhu et al., 2021) genomes to identify their physical positions. The Svevo and Zavitan genomes were used when considering the physical positions of QTL detected from AL and LM001, respectively. Predicted genes mapped between the flanking markers and their function annotations and expression patterns were obtained from the Triticeae Multi-omics Center (http://202.194.139.32/).

QTL Validation

The tightly linked SNP markers of major and novel SNS QTL were subjected to develop KASP markers. KASP primers were designed and applied following the method of Tan et al. (2017). Each validation population was divided into two haplotype groups (with homozygous alleles from any parent) according to their alleles present in parents and progenies. Phenotypic differences between the two groups were analyzed using Student's *t*-test ($P < 0.05$).

Results

SNP Markers and Genetic Map

As suggested by the Affymetrix Company, the high probability probes from the Poly High-Resolution group only were reserved. Thus, 8724 SNPs with minor allele frequency (≥ 0.3) among the mapping population were retained for following analysis. Subsequently, 823 markers from the D genome were rejected. Finally, 7901 SNP markers were classified into 1192 bins, and a single marker with the lowest 'Missing rate' in each bin (bin marker) was screened to construct the genetic map. The linkage analysis results revealed that 1150 bin markers were mapped on 15 linkage groups. One linkage group was constructed for each chromosome except chromosome 3B, which had two (Table 1 and Supplementary Table 2). Based on the bin information, 6537 mapped markers (including 1150 bin markers and 5387

co-segregated markers) were integrated into the genetic map with a total length of

2411.8 cM (Table 1 and Supplementary Table 3). The length of genetic map in each

linkage group ranged between 49.7 cM (3B1) and 225.1 cM (5A). The average

distance of bin markers ranged between 1.09 cM (4B) and 3.64 cM (3B2), with an

average density of one bin marker per 2.10 cM. The number of mapped markers

ranged between 37 on chromosome 3B1 and 944 on chromosome 2A (Table 1 and

Supplementary Table 3). Notably, 56% and 44% of the mapped markers were

distributed on A and B subgenomes, respectively.

Comparison of Genetic and Physical Maps

Considering that the parent accessions AL and LM001 were durum wheat and wild

emmer wheat, respectively, the sequences of the 6537 mapped markers were blasted

against durum wheat and wild emmer genomes to identify their physical positions

(Supplementary Table 3). The genetic chromosomal locations of the 1150 bin

markers were compared with their physical chromosomal locations in the durum

wheat genome. Among them, 1087 (94.52 %) markers showed good consistencies

(Figure 2a), and 63 (5.48%) markers had inconsistent chromosomal locations. In the

wild emmer genome, the genetic chromosomal locations of 71 (6.17%) markers were

inconsistent with physical chromosomal locations, while the remaining markers

(1079, 93.83%) were well matched (Figure 2b). Notably, 1026 (89.22%) bin markers

had consistent chromosomal locations in genetic map, durum wheat genome and

wild emmer genome. Similarly, the genetic maps of the 6537 mapped markers were

compared with their physical maps (Supplementary Table 3). 6242 (95.49%) and

6249 (95.59%) markers showed consistent chromosomal locations in the durum

wheat and wild emmer genomes, respectively, while 6031 (92.26%) markers had

consistent chromosomal locations in genetic map and two physical maps.

Phenotypic Variation

The SNS of AL and LM001 in multiple environments ranged between 25.00 and

27.00, and 23.33 and 26.00, respectively (Table 2). That of AM RILs ranged between

15.33 and 34.00. There were no significant differences in SNS between the two

parental lines. However, continuous variation and bidirectional transgressive

segregation was observed in the RIL population. These results suggested that it was

feasible to analyze the SNS loci in the current population.

The broad-sense heritability (H^2) of SNS ranged between 0.93 and 0.97 in each

environment (Table 2), indicating that the SNS trait was mainly controlled by genetic

factors. Strong interaction was detected between SNS and environments ($P < 0.01$),

with significant correlation coefficients ranging between 0.50 and 0.90 (Figure 3).

The frequency distribution of SNS in multiple environments indicated that it was

polygenically inherited (Figure 3).

Genetic Correlations between SNS and other Agronomic Traits

Pearson's correlations were analyzed between SNS and seven other agronomic traits

based on the BLUP values (Table 3). SNS was significantly and positively correlated

with PH, AD, SL, SEL, KNS, and SD. However, SNS was not significantly

correlated with TKW.

Analysis of QTL Associated with SNS

Five QTL for SNS were identified and distributed on chromosomes 5A (1 QTL), 2B

(2), and 3B (2), with LOD scores ranging between 3.55 and 21.49 (Table 4). They

explained 6.71-29.40% of the phenotypic variation. *QSns.sau-AM-5A*,

QSns.sau-AM-2B.3, and *QSns.sau-AM-3B.2* were detected in at least five

environments and explained more than 10% of the phenotypic variation, which were

treated as major QTL. And the other two QTL were identified as minor QTL.

QSns.sau-AM-5A was detected in five environments and located in a 19.1 cM region

between AX-108790581 and AX-111596791. It accounted for 8.27-12.09% of the

phenotypic variation (Table 4). AM population were divided into two haplotype

groups based on the genotype of QTL's flanking markers, and the effect of

QSns.sau-AY-5A.2 increased SNS significantly by 4.54-10.48% (Figure 4a).

QSns.sau-AM-2B.2 was stably identified in all environments, explaining 9.09-29.40%

of the phenotypic variation (Table 4). It was firstly mapped between AX-111626925

and AX-111588559 (Figure 4b). AX-95072014 and AX-109422178 were

polymorphic SNPs detected in initial mapping interval of two parental lines using

the wheat 660K SNP array. They were converted to KASP markers

(KASP-AX-95072014 and KASP-AX-109422178) to genotype the mapping

population and integrated into the genetic map. Remapping results showed that

QSns.sau-AM-2B.2 was located in a 2.2 cM region between KASP-AX-95072014

and AX-111588559. The additive effect of *QSns.sau-AM-2B.2* was negative,

indicating that the increasing SNS allele was contributed by LM001. The effects of

QSns.sau-AM-2B.2 on increasing SNS in single environment ranged from 7.00% to

12.18% (Figure 4b).*QSns.sau-AM-3B.2* was detected in five environments (Table 4).

It explained up to 15.46% of the phenotypic variation. The positive allele of

QSns.sau-AM-3B.2 was from AL, and RILs containing the positive allele of

QSns.sau-AM-3B.2 had a higher SNS than those containing the negative one. The

effect of *QSns.sau-AM-3B.2* significantly increased SNS by 5.02-10.14% in single

environment (Figure 4c).

The QTL-by-Environment interaction analysis identified 47 QTL (Supplementary

Table 4). Five of them were the same as detected in individual environment analysis,

indicating that they were stably expressed for SNS. In addition, epistatic interaction

were also analyzed, and the results showed that there were no epistasis effects

between five SNS QTL detected in the present study (Supplementary Table 5).

Effects of Major QTL on Increasing SNS

The effects of three QTL (*QSns.sau-AM-5A*, *QSns.sau-AM-2B.2*, and

QSns.sau-AM-3B.2) on increasing the SNS were further revealed in the mapping

population based on the genotypes of the flanking markers (Figure 5). Compared to

the haplotype group that carried the negative alleles of three QTL, haplotype groups

with the favorable alleles from at least one QTL significantly increased SNS. The

haplotype group containing the favorable alleles from three QTL significantly

increased SNS compared with those with two or single one, respectively. No

significant differences were detected among three haplotype groups with any two

positive alleles. *QSns.sau-AM-2B.2* had the largest individual positive effect on SNS.

However, there was no significant difference between the individual effect of three

QTL on increasing SNS.

Validation of Major and Novel SNS QTL

The SNP markers closely linked to *QSns.sau-AM-2B.2* (AX-111588559) and

QSns.sau-AM-3B.2 (AX-110548993) were converted to KASP markers, to verify the

effects of them in two other populations (Supplementary Table 6 and Figure 6).

Since the favorable alleles of *QSns.sau-AM-2B.2* and *QSns.sau-AM-3B.2* were from

LM001 and AL, respectively, haplotype group with homozygous alleles of LM001

showed a significant increase in SNS by 8.33% (Figure 6a) and those from AL

increased SNS significantly by 6.00% (Figure 6b).

Discussion

Exploration of the SNS Loci from Tetraploid Wheat Species Using the Wheat

55K SNP Array

Previous studies have reported on SNS in tetraploid wheat at the QTL level. Peng et

al. (2001) identified four SNS QTL, explaining a total of 49.3% of the phenotypic

variation in a wild emmer × durum wheat F_2 population genotyped using restriction

fragment length polymorphisms (RFLP) markers. Genotyping of the durum ×

cultivated emmer wheat RIL population with the 9 K iSelect SNP chip identified

four SNS QTL on chromosomes 1B, 3B, 7A, and 7B (Faris et al., 2014a). In the

same time, a linkage map of chromosome 2A generated in the Langdon ×

Langdon-wild emmer wheat substitution lines population consisting of 23 simple

sequence repeat (SSR) markers was used to identify an SNS QTL, *QSpn.fcu-2A*,

explaining 19.4% of the phenotypic variation (Faris et al., 2014b). SNS QTL,

QSpn.fcu-1B, and *QSpn.fcu-5A*, have been identified in a durum × cultivated emmer

wheat population genotyped using the 90K iSelect array (Sharma et al., 2019). These

reports enhance our understanding of the genetic mechanism of SNS in tetraploid

wheat.

The wheat 55K SNP array was developed based on the selection and optimization of

the 660K SNP array which was designed based on hexaploid, tetraploid and diploid

wheat, and has the advantages of being efficient and genome-specific (Sun et al.,

2020). It has been widely used for genotyping and genetic loci identification of

agronomic traits and disease resistance in bread wheat (Ren et al., 2018; Ma et al.,

2019b; Liu et al., 2020b; Ma et al., 2020; Tu et al., 2021). To the author's knowledge,

this is the first study to adopt the wheat 55K SNP array to construct the genetic

linkage map of a tetraploid wheat RIL population. Our results confirm the feasibility

of using a 55K SNP array developed for common wheat for genetic mapping of

tetraploid wheat, thus proving its broad utilization.

Comparison of the Major SNS QTL Identified with those of Previous Studies

There was no significant difference for SNS detected between the parental lines in

multiple environments (Table 2). The phenotypic data of SNS in the six

environments showed approximately normal distribution (Figure 4). Three major and

stably expressed QTL for SNS were identified. This phenomenon is attributed to the

transgressive inheritance in progenies that result in the bidirectional segregation of

SNS. Several genes controlling SNS may inhibit each other in a given parental

genotype resulting in the absence of the corresponding phenotype in the parents.

However, a single locus produced by genetic recombination between parental

genotypes can display a conspicuous phenotype in the progenies without interference

from other loci that control the same trait. The major SNS loci in the AM RIL

population can thus be identified. This is not uncommon in the past studies of QTL

identification based on RIL populations (Liu et al., 2020a; Shang et al., 2020).

Physical locations of known SNS QTL were detected to verify whether major QTL

were novel (Supplementary Table 7). In wild emmer genome, *QSns.sau-AM-5A* was

physically located on chromosome arm 5AL at 557.72-571.41 Mb, overlapping with

QSps.ccsu-5A.2 (568.96-662.44 Mb) (Kumar et al., 2007) and *QTL1755_5A*

(569.50-579.91 Mb) (Peng et al., 2003), indicating that they are likely alleles.

QSns.sau-AM-2B.2 was located at 731.25-731.60 Mb on chromosome arm 2BL in

wild emmer genome. Previous studies have detected SNS QTL on chromosome arm

2BS. However, none has been reported on 2BL. For example, *Qspn-2007* was

located on chromosome arm 2BS with the closest marker Xgwm429 (Manickavelu

et al., 2011). *QFsn.sdau-2B-1* was located on chromosome arm 2BS, explaining up

to 28.43% of the phenotypic variation (Deng et al., 2017). Some genes regulating

SNS, such as *Ppd-B1* (Mohler et al., 2004) and *WFZP-2B* (Dobrovolskaya et al.,

2015), have also been detected on chromosome arm 2BS. In the durum wheat

genome, *QSns.sau-AM-3B.2* was located on chromosome arm 3BS at 65.87-72.59

Mb. Similarly, Cui et al. (2012) identified two QTL, *QSpn.WJ.3B.1* and

QSpn.WY.3B.1, all closely linked to the locus Xgwm566. Azadi et al. (2015) detected

three QTL for SNS on chromosome arm 3BL. Comparing the physical positions of

three major QTL with those previously reported indicates that *QSns.sau-AY-2B.2* and

QSns.sau-AY-3B.2 are probably novel loci controlling SNS in tetraploid wheat.

Potential Candidate Genes in the Physical Intervals of *QSns.sau-AM-2B.2*

QSns.sau-AM-2B.2 was physical located in a 0.35 Mb region of wild emmer genome,

and 4 genes and 3 functional orthologs were annotated (Supplementary Table 8a).

QSns.sau-AM-3B.2 was located in a 6.72 Mb interval of durum wheat genome with

62 genes and 38 functional orthologs (Supplementary Table 8b). Since the physical

interval of *QSns.sau-AM-2B.2* contains only 3 orthologs, it is valuable to analyze

function annotations and expression patterns to predict candidate genes.

TRIDC2BG075290 encodes a protein kinase which involves in cell signal

transduction and plant growth. Previous research reported wheat protein kinase gene

TaSnRK2.9□5A was signifcantly correlated with high thousand kernel weight and

high kernel per spike (Ur Rehman et al., 2019), suggesting that protein kinase can

have a positive impact on agronomic traits. In addition, *Stomatal opening factor 1*

(*OST1*, *SnRK2.6*) is a homologous gene of *TRIDC2BG075290* in *Arabidopsis*. *OST1*

can promote the expression of *FLOWERING LOCUS C* (*FLC*) to participate in the

regulation of flowering by ABA (Wang et al., 2013), which may affect inflorescence

development and the formation of spikelets. *TRIDC2BG075310* and

TRIDC2BG075320 all encode SAUR-like auxin-responsive protein. It has been

reported that plant SAUR-like auxin-responsive protein family is involved in the

growth of roots, hypocotyls, leaves and stamens (Chae et al., 2012; Hou et al., 2013).

Expression patterns analyses showed that *TRIDC2BG075290* has higher expression

levels in spike and flag leaf (Supplementary Figure 1). *TRIDC2BG075310* has a high

expression level throughout the spike development stage, while *TRIDC2BG075320*

is only expressed in spike. Therefore, we speculate that these 3 functional orthologs

may affect the wheat inflorescence development, which can be further determined by

fine mapping and gene cloning in future.

Phenotypic Correlations of Agronomic Traits

SNS was significantly and positively correlated with PH in the present study (Table

3). This finding was consistent with that of Ajmal et al. (2009) and Wu et al. (2012).

We speculate that SL is part of PH, and the taller plants have a long spike with more

spikelets (Yao et al., 2011). There was a significant and positive relationship between

SNS and AD (Table 3), consistent with the results of Ma et al. (2019a). It is supposed

that wheat plants that take longer to flower tend to produce longer spikes with more

spikelets (Shaw et al., 2013; Kamran et al., 2014; Boden et al., 2015). These results

indicate that AD plays an important role in spikelet formation.

Similarly, there was a strong significant and positive correlation between SNS and

SL (Table 3). The increase in SL requires a longer spike development stage

combined with optimal temperature and light to promote spikelet germination (Shaw

et al., 2013). On the whole, floret fertility determines the KNS in wheat. More

spikelets produce more fertile florets, thereby increasing the KNS (Gao et al., 2019;

Sakuma et al., 2019). This reason potentially caused the SNS to be significantly and

positively correlated with KNS (Table 3). Moreover, SNS was significantly and

positively correlated with SD because SD is the ratio of SNS to SL (Table 3).

It is postulated that there is a negative correlation between SNS and TKW (Li et al.,

2007; Ma et al., 2019a). The negative correlation is attributed to source limitations

during kernel filling or more kernels with lower KW in the spike's distal positions

(Quintero et al., 2018). Previous studies also suggest that SNS is negatively

correlated with SEL because of competition in product assimilation between SNS

and SEL (Bancal, 2008; Li et al., 2020). However, these conclusions are inconsistent

with the findings of this study. Herein, SNS was positively correlated with TKW and

SEL (Table 3). It probably indicates that the loci controlling SNS play an individual

role without interacting with SEL. Previous studies postulate that SEL has a special

function in photosynthesis and nutrients and water storage (Bridgemohan and

Bridgemohan, 2014; Ávila-Lovera et al., 2017). Therefore, a longer SEL is more

conducive for ventilation and transportation of more nutrients and water to kernels

thereby increasing the KW.

These conclusions are vital in comprehensively understanding the complex

relationships in agronomic traits, and provide new insights into wheat yield

improvement. KNS is one of the primary determinants of grain yield. Therefore,

wheat breeding programs should focus on increasing SNS under moderate SD in the

present study.

Conclusion

Herein, the wheat 55K SNP array was successfully used to construct a high-quality

genetic map that identified five QTL for SNS in a tetraploid wheat RIL population.

The additive effects of the major SNS QTL in increasing SNS were revealed in the

mapping population. *QSns.sau-AM-2B.2* and *QSns.sau-AM-3B.2* were major and

novel QTL, and their effects were successfully validated in other genetic

backgrounds using closely linked KASP markers. The genetic correlations between

SNS and other agronomic traits were also evaluated. The novel genetic loci of SNS

and tightly linked KASP markers will be valuable in further fine mapping and gene

cloning.

Data Availability Statement

All data generated or analyzed during this study are included in this published article

and its supplementary information files, further inquiries can be directed to the

corresponding author.

Author Contributions

ZQM and JZ conducted the study and drafted this manuscript. JTW and JGZ

participated in phenotype measurement and data analysis. QX and HPT helped with

field work and data analysis. YM, MD, QTJ, and YXL participated in data collection

and analysis. GYC, JRW, PFQ, and WL did QTL analysis and manuscript revision.

YMW and YLZ discussed the results and revised the manuscript. XJL guided the

study and revised the manuscript. JM designed the experiments, guided the entire

study, participated in data analysis, wrote and extensively revised this manuscript.

All authors participated in the research and approved the final manuscript.

Funding

This work is supported by the National Natural Science Foundation of China

(31971937 and 31970243), the Key Research and Development Program of Sichuan

Province (2018NZDZX0002), the Applied Basic Research Programs of Science and

Technology Department of Sichuan Province (2020YJ0140 and 2021YJ0503), the

International Science and Technology Cooperation and Exchanges Program of

Science and Technology Department of Sichuan Province (2021YFH0083), and the

Key Projects of Scientific and Technological Activities for Overseas Students of

Sichuan Province. The funders did not participate in the study design, data analysis,

or preparation of the manuscript.

Conflict of Interest

The authors declare that they have no competing interests.

466
467
468 **References**
469

Ajmal, S., Zakir, N., and Mujahid, M. (2009). Estimation of genetic parameters and
character association in wheat. *J. Agric. Biol. Sci.* 1, 15-18. doi:
10.5713/ajas.2011.10456.
Ávila-Lovera, E., Zerpa, A.J., and Santiago, L.S. (2017). Stem photosynthesis and
hydraulics are coordinated in desert plant species. *New Phytol.* 216, 1119-1129.
doi: 10.1111/nph.14737.
Azadi, A., Mardi, M., Hervan, E.M., Mohammadi, S.A., Moradi, F., Tabatabaee,
M.T., et al. (2015). QTL mapping of yield and yield components under normal
and salt-stress conditions in bread wheat (*Triticum aestivum* L.). *Plant Mol. Biol.*
Rep. 33(1), 102-120. doi: 10.1007/s11105-014-0726-0.
Bancal, P. (2008). Positive contribution of stem growth to grain number per spike in
wheat. *Field Crop Res.* 105(1), 27-39. doi: 10.1016/j.fcr.2007.06.008.
Boden, S.A., Cavanagh, C., Cullis, B.R., Ramm, K., Greenwood, J., Jean Finnegan,
E., et al. (2015). *Ppd-1* is a key regulator of inflorescence architecture and paired
spikelet development in wheat. *Nat. Plants* 1(2), 14016. doi:
10.1038/nplants.2014.16.
Bridgemohan, P., and Bridgemohan, R.S.H. (2014). Evaluation of anti-lodging plant
growth regulators on the growth and development of rice (*Oryza sativa*). *J*
Cereals Oilseeds 5, 12-16. doi: 10.5897/JCO14.0128.
Cavanagh, C.R., Chao, S., Wang, S., Huang, B.E., Stephen, S., Kiani, S., et al.
(2013). Genome-wide comparative diversity uncovers multiple targets of
selection for improvement in hexaploid wheat landraces and cultivars. *Proc. Natl.*
Acad. Sci. U. S. A. 110(20), 8057-8062. doi: 10.1073/pnas.1217133110.
Chae, K., Isaacs, C.G., Reeves, P.H., Maloney, G.S., Muday, G.K., Nagpal, P., et al.
(2012). *Arabidopsis* SMALL AUXIN UP RNA63 promotes hypocotyl and stamen
filament elongation. *Plant J.* 71(4), 684-697. doi:

10.1111/j.1365-313X.2012.05024.x.

Chen, Z., Cheng, X., Chai, L., Wang, Z., Du, D., Wang, Z., et al. (2020). Pleiotropic
QTL influencing spikelet number and heading date in common wheat (*Triticum*
aestivum L.). *Theor. Appl. Genet.* 133(6), 1825-1838. doi:
10.1007/s00122-020-03556-6.

Crossa, J., Jarquin, D., Franco, J., Perezrodriguez, P., Burgueno, J., Saintpierre, C., et
al. (2016). Genomic prediction of gene bank wheat landraces. *G3-Genes Genom.*
Genet. 6(7), 1819-1834. doi: 10.1534/g3.116.029637.

Cui, F., Ding, A., Li, J., Zhao, C., Wang, L., Wang, X., et al. (2012). QTL detection
of seven spike-related traits and their genetic correlations in wheat using two
related RIL populations. *Euphytica* 186(1), 177-192. doi:
10.1007/s10681-011-0550-7.

Deng, Z., Cui, Y., Han, Q., Fang, W., Li, J., and Tian, J. (2017). Discovery of
consistent QTLs of wheat spike-related traits under nitrogen treatment at different
development stages. *Front. Plant Sci.* 8, 2120. doi: 10.3389/fpls.2017.02120.

Dixon, L.E., Greenwood, J.R., Bencivenga, S., Zhang, P., Cockram, J., Mellers, G.,
et al. (2018). *TEOSINTE BRANCHED1* regulates inflorescence architecture and
development in bread wheat (*Triticum aestivum*). *Plant Cell* 30(3), 563-581. doi:
10.1105/tpc.17.00961.

Dobrovolskaya, O., Pont, C., Sibout, R., Martinek, P., Badaeva, E., Murat, F., et al.
(2015). *FRIZZY PANICLE* drives supernumerary spikelets in bread wheat. *Plant*
Physiol. 167, 189-199. doi: 10.1104/pp.114.250043.

El Haddad, N., Kabbaj, H., Zaïm, M., El Hassouni, K., Tidiane Sall, A., Azouz, M.,
et al. (2021). Crop wild relatives in durum wheat breeding: Drift or thrift? *Crop*
Sci. 61(1), 37-54. doi: 10.1002/csc2.20223.

Faris, J.D., Zhang, Q., Chao, S., Zhang, Z., and Xu, S.S. (2014a). Analysis of
agronomic and domestication traits in a durum × cultivated emmer wheat
population using a high-density single nucleotide polymorphism-based linkage
map. *Theor. Appl. Genet.* 127, 2333-2348. doi: 10.1007/s00122-014-2380-1.

Faris, J.D., Zhang, Z., Garvin, D.F., and Xu, S.S. (2014b). Molecular and
comparative mapping of genes governing spike compactness from wild emmer
wheat. *Mol. Genet. Genomics* 289(4), 641-651. doi: 10.1007/s00438-014-0836-2.

Gao, X., Wang, N., Wang, X., and Zhang, X. (2019). Architecture of wheat
inflorescence: insights from rice. *Trends Plant Sci.* 24, 802-809. doi:
10.1016/j.tplants.2019.06.002.

Hou, K., Wu, W., and Gan, S. (2013). *SAUR36*, a small auxin up RNA gene, is
involved in the promotion of leaf senescence in *Arabidopsis*. *Plant Physiol* 161(2),
1002-1009. doi: 10.1104/pp.112.212787.

Kamran, A., Iqbal, M., and Spaner, D. (2014). Flowering time in wheat (*Triticum*
aestivum L.): a key factor for global adaptability. *Euphytica* 197, 1-26. doi:
10.1007/s10681-014-1075-7.

Klymiuk, V., Yaniv, E., Huang, L., Raats, D., Fatiukha, A., Chen, S., et al. (2018).
Cloning of the wheat *Yr15* resistance gene sheds light on the plant tandem
kinase-pseudokinase family. *Nat. Commun.* 9(1), 3735. doi:

10.1038/s41467-018-06138-9.

Kumar, N., Kulwal, P.L., Balyan, H.S., and Gupta, P.K. (2007). QTL mapping for
yield and yield contributing traits in two mapping populations of bread wheat.
Mol. Breed. 19(2), 163-177. doi: 10.1007/s11032-006-9056-8.

Kuzay, S., Xu, Y., Zhang, J., Katz, A., Pearce, S., Su, Z., et al. (2019). Identification
of a candidate gene for a QTL for spikelet number per spike on wheat
chromosome arm 7AL by high-resolution genetic mapping. *Theor. Appl. Genet.*
132 (9), 2689-2705. doi: 10.1007/s00122-019-03382-5.

Li, C., Tang, H., Luo, W., Zhang, X., Mu, Y., Deng, M., et al. (2020). A novel,
validated, and plant height-independent QTL for spike extension length is
associated with yield-related traits in wheat. *Theor. Appl. Genet.* 133, 3381-3393.
doi: 10.1007/s00122-020-03675-0.

Li, S., Jia, J., Wei, X., Zhang, X., Li, L., Chen, H., et al. (2007). A intervarietal
genetic map and QTL analysis for yield traits in wheat. *Mol. Breed.* 20(2),
167-178. doi: 10.1007/s11032-007-9080-3.

Li, T., Deng, G., Tang, Y., Su, Y., Wang, J., Cheng, J., et al. (2021). Identification and
validation of a novel locus controlling spikelet number in bread wheat (*Triticum*
aestivum L.). *Front. Plant Sci.* 12, 611106. doi: 10.3389/fpls.2021.611106.

Lin, Y., Jiang, X., Hu, H., Zhou, K., Wang, Q., Yu, S., et al. (2021). QTL mapping for
grain number per spikelet in wheat using a high-density genetic map. *Crop J.* doi:
10.1016/j.cj.2020.12.006.

Liu, D., Yen, C., Yang, J., Zheng, Y., and Lan, X. (1999). The chromosomal locations
of high crossability genes in tetraploid wheat *Triticum turgidum* L. cv. Ailanmai
native to Sichuan, China. *Euphytica* 108(2), 79-82. doi:
10.1023/A:1003691925501.

Liu, H., Ma, J., Tu, Y., Zhu, J., Ding, P., Liu, J., et al. (2020a). Several stably
expressed QTL for spike density of common wheat (*Triticum aestivum*) in
multiple environments. *Plant Breed.* 139, 284-294. doi: 10.1111/pbr.12782.

Liu, J., Luo, W., Qin, N., Ding, P., Zhang, H., Yang, C., et al. (2018). A 55 K SNP
array-based genetic map and its utilization in QTL mapping for productive tiller
number in common wheat. *Theor. Appl. Genet.* 131(11), 2439-2450. doi:
10.1007/s00122-018-3164-9.

Liu, J., Tang, H., Qu, X., Liu, H., Li, C., Tu, Y., et al. (2020b). A novel, major, and
validated QTL for the effective tiller number located on chromosome arm 1BL in
bread wheat. *Plant Mol. Biol.* 104(1), 173-185. doi: 10.1007/s11103-020-01035-6.

Lopes, M.S., Ibrahim, E.B., Baenziger, P.S., Sukhwinder, S., Conxita, R., Kursad, O.,
et al. (2015). Exploiting genetic diversity from landraces in wheat breeding for
adaptation to climate change. *J. Exp. Bot.* 66(12), 3477-3486. doi:
10.1093/jxb/erv122.

Ma, J., Ding, P., Liu, J., Li, T., Zou, Y., Habib, A., et al. (2019a). Identification and
validation of a major and stably expressed QTL for spikelet number per spike in
bread wheat. *Theor. Appl. Genet.* 132(11), 3155-3167. doi:
10.1007/s00122-019-03415-z.

Ma, J., Qin, N., Cai, B., Chen, G., Ding, P., Zhang, H., et al. (2019b). Identification

and validation of a novel major QTL for all-stage stripe rust resistance on 1BL in
the winter wheat line 20828. *Theor. Appl. Genet.* 132(5), 1363-1373. doi:
10.1007/s00122-019-03283-7.

Ma, J., Tu, Y., Zhu, J., Luo, W., Liu, H., Li, C., et al. (2020). Flag leaf size and
posture of bread wheat: genetic dissection, QTL validation and their relationships
with yield-related traits. *Theor. Appl. Genet.* 133(1), 297-315. doi:
10.1007/s00122-019-03458-2.

Maccaferri, M., Harris, N.S., Twardziok, S.O., Pasam, R.K., Gundlach, H., Spannagl,
M., et al. (2019). Durum wheat genome highlights past domestication signatures
and future improvement targets. *Nat. Genet.* 51(5), 885-895. doi:
10.1038/s41588-019-0381-3.

Manickavelu, A., Kawaura, K., Imamura, H., Mori, M., and Ogihara, Y. (2011).
Molecular mapping of quantitative trait loci for domestication traits and β-glucan
content in a wheat recombinant inbred line population. *Euphytica* 177(2), 179-190.
doi: 10.1007/s10681-010-0217-9.

Meng, L., Li, H., Zhang, L., and Wang, J. (2015). QTL IciMapping: Integrated
software for genetic linkage map construction and quantitative trait locus
mapping in biparental populations. *Crop J.* 3(3), 269-283. doi:
10.1016/j.cj.2015.01.001.

Mohler, V., Lukman, R., Ortiz-Islas, S., William, M., Worland, A.J., van Beem, J., et
al. (2004). Genetic and physical mapping of photoperiod insensitive gene *Ppd-B1*
in common wheat. *Euphytica* 138(1), 33-40. doi:
10.1023/B:EUPH.0000047056.58938.76.

Peng, J., Korol, A., Fahima, T., Roder, M., Li, Y., and Nevo, E. (2001). QTLs for
agronomic traits in tetraploid wild emmer wheat, *Triticum dicoccoides*. *J. Sichuan*
Agric. Univ. 19, 317-334. doi: 10.16036/j.issn.1000-2650.2001.04.005.

Peng, J., Ronin, Y.I., Fahima, T., Roder, M.S., Li, Y., Nevo, E., et al. (2003).
Domestication quantitative trait loci in *Triticum dicoccoides*, the progenitor of
wheat. *P. Natl. Acad. Sci. USA* 100(5), 2489-2494. doi: 10.1073/pnas.252763199.

Quintero, A., Molero, G., Reynolds, M.P., and Calderini, D.F. (2018). Trade-off
between grain weight and grain number in wheat depends on GxE interaction: A
case study of an elite CIMMYT panel (CIMCOG). *Eur. J. Agron.* 92, 17-29. doi:
10.1016/j.eja.2017.09.007.

Ren, T., Hu, Y., Tang, Y., Li, C., Yan, B., Ren, Z., et al. (2018). Utilization of a
wheat55K SNP array for mapping of major QTL for temporal expression of the
tiller number. *Front. Plant Sci.* 9, 333. doi: 10.3389/fpls.2018.00333.

Rothwell, W. (2019). Origin of Perl. In: Beginning Perl Programming. Apress,
Berkeley, CA. doi: 10.1007/978-1-4842-5055-6_1.

Sakuma, S., Golan, G., Guo, Z., Ogawa, T., Tagiri, A., Sugimoto, K., et al. (2019).
Unleashing floret fertility in wheat through the mutation of a homeobox gene.
Proc. Natl. Acad. Sci. U. S. A. 116, 5182-5187. doi: 10.1073/pnas.1815465116.

Shang, Q., Zhang, D., Li, R., Wang, K., Cheng, Z., Zhou, Z., et al. (2020). Mapping
quantitative trait loci associated with stem-related traits in maize (*Zea mays* L.).
Plant Mol. Biol. 104, 583-595. doi: 10.1007/s11103-020-01062-3.

Sharma, J.S., Running, K.L.D., Xu, S.S., Zhang, Q., Peters Haugrud, A.R., Sharma,
S., et al. (2019). Genetic analysis of threshability and other spike traits in the
evolution of cultivated emmer to fully domesticated durum wheat. *Mol. Genet.*
Genomics 294(3), 757-771. doi: 10.1007/s00438-019-01544-0.
Shaw, L.M., Turner, A.S., Laurence, H., Simon, G., Laurie, D.A., and Somers, D.E.
(2013). Mutant alleles of *Photoperiod-1* in wheat (*Triticum aestivum* L.) that
confer a late flowering phenotype in long days. *PLoS One* 8(11), e79459. doi:
10.1371/journal.pone.0079459.
Sthapit, S.R., Marlowe, K., Covarrubias, D.C., Ruff, T.M., Eagle, J.D., McGinty,
E.M., et al. (2020). Genetic diversity in historical and modern wheat varieties of
the U.S. Pacific Northwest. *Crop Sci.* 60, 3175-3190. doi: 10.1002/csc2.20299.
Sun, C., Dong, Z., Zhao, L., Ren, Y., Zhang, N., and Chen, F. (2020). The Wheat
660K SNP array demonstrates great potential for marker-assisted selection in
polyploid wheat. *Plant Biotechnol. J.* 18(6), 1354-1360. doi: 10.1111/pbi.13361.
Tan, C.-T., Yu, H., Yang, Y., Xu, X., Chen, M., Rudd, J.C., et al. (2017).
Development and validation of KASP markers for the greenbug resistance gene
Gb7 and the Hessian fly resistance gene *H32* in wheat. *Theor. Appl. Genet.* 130(9),
1867-1884. doi: 10.1007/s00122-017-2930-4.
Tu, Y., Liu, H., Liu, J., Tang, H., Mu, Y., Deng, M., et al. (2021). QTL mapping and
validation of bread wheat flag leaf morphology across multiple environments in
different genetic backgrounds. *Theor. Appl. Genet.* 134(1), 261-278. doi:
10.1007/s00122-020-03695-w.
Ur Rehman, S., Wang, J., Chang, X., Zhang, X., Mao, X., and Jing, R. (2019). A
wheat protein kinase gene *TaSnRK2.9*-5A associated with yield contributing traits.
Theor. Appl. Genet. 132(4), 907-919. doi: 10.1007/s00122-018-3247-7.
Voorrips, R.E. (2002). MapChart: software for the graphical presentation of linkage
maps and QTLs. *J. Hered.* 93(1), 77-78. doi: 10.1093/jhered/93.1.77.
Wang, Y., Lin, L., Ye, T., Lu, Y., Chen, X., and Wu, Y. (2013). The inhibitory effect
of ABA on floral transition is mediated by ABI5 in Arabidopsis. *J. Exp. Bot.* 64(2),
675-684. doi: 10.1093/jxb/ers361.
Wang, Y., Yu, H., Tian, C., Sajjad, M., Gao, C., Tong, Y., et al. (2017). Transcriptome
association identifies regulators of wheat spike architecture. *Plant Physiol.* 175(2),
746-757. doi: 10.1104/pp.17.00694.
Wu, X., Chang, X., and Jing, R. (2012). Genetic insight into yield-associated traits of
wheat grown in multiple rain-fed environments. *PloS One* 7, e31249. doi:
10.1371/journal.pone.0031249.
Xie, W., and Nevo, E. (2008). Wild emmer: genetic resources, gene mapping and
potential for wheat improvement. *Euphytica* 164(3), 603-614. doi:
10.1007/s10681-008-9703-8.
Yao, H., Xie, Q., Xue, S., Luo, J., Lu, J., Kong, Z., et al. (2019). *HL2* on
chromosome 7D of wheat (*Triticum aestivum* L.) regulates both head length and
spikelet number. *Theor. Appl. Genet.* 132(6), 1789-1797. doi:
10.1007/s00122-019-03315-2.
Yao, J., Ren, L., Zhang, P., Yang, X., and Zhou, M. (2011). Genetic and correlation

analysis of plant height and its components in wheat. *J. Triticeae Crop* 31,

604-610 (in Chinese with English abstract).

Yu, J., Zhao, Y., Ding, M., Yu, Z., Jiang, Y., Ma, W., et al. (2020). Wild emmer

chromosome arm substitution lines: Useful resources for wheat genetic study and

breeding. *Crop Sci.* 60, 1761-1769. doi: 10.1002/csc2.20022.

Zaïm, M., El Hassouni, K., Gamba, F., Filali-Maltouf, A., Belkadi, B., Sourour, A., et

al. (2017). Wide crosses of durum wheat (*Triticum durum* Desf.) reveal good

disease resistance, yield stability, and industrial quality across Mediterranean sites.

Field Crop Res. 214, 219-227. doi: 10.1016/j.fcr.2017.09.007.

Zhu, T., Wang, L., Rimbert, H., Rodriguez, J.C., Deal, K.R., De Oliveira, R., et al.

(2021). Optical maps refine the bread wheat *Triticum aestivum* cv. Chinese Spring

genome assembly. *Plant J.* doi: 10.1111/tpj.15289.

Zhu, T., Wang, L., Rodriguez, J.C., Deal, K.R., Avni, R., Distelfeld, A., et al. (2019).

Improved Genome Sequence of Wild Emmer Wheat Zavitan with the Aid of

Optical Maps. *G3: Genes. Genom. Genet.* 9(3), 619-624. doi:

10.1534/g3.118.200902.

Table 1 Information of SNP markers in the constructed genetic map

Chromosome	Group	Number of bin markers	Number of mapped markers	Length (cM)	Density (cM/bin)
1A	1	101	586	153.3	1.52
1B	1	69	488	149.3	2.16
2A	1	66	944	161.8	2.45
2B	1	92	424	183.7	2.00
3A	1	99	517	161.0	1.63
3B	1	17	37	49.7	2.92
	2	27	174	98.4	3.64
4A	1	80	282	199.9	2.50
4B	1	77	422	83.7	1.09
5A	1	92	298	225.1	2.45
5B	1	121	674	207.5	1.72
6A	1	60	636	167.0	2.78
6B	1	58	160	162.5	2.80
7A	1	92	395	214.2	2.33
7B	1	99	500	194.8	1.97
A genome	7	590	3658	1282.2	2.17
B genome	8	560	2879	1129.6	2.02
Total	15	1150	6537	2411.8	2.10

Table 2 Phenotypic data and heritability (H^2) of spikelet number per spike (SNS) for the AM population in multiple environments

Environment	Parents		AL × LM001 (AM)			
	AL	**LM001**	**Min-Max**	**Mean**	**SD**	H^2
E_1	25.00	24.33	20.00-31.67	25.96	2.56	0.95
E_2	26.00	23.33	15.33-33.00	24.41	3.22	0.97
E_3	26.33	26.00	17.00-31.25	24.81	2.80	0.96
E_4	26.00	25.40	18.00-32.00	26.21	2.87	0.96
E_5	27.00	24.60	18.50-31.00	24.99	2.42	0.95
E_6	25.80	25.40	20.00-34.00	26.76	2.69	0.96
E_7	23.40	24.25	15.75-31.00	23.68	2.62	0.95
E_8	25.44	24.67	18.50-30.67	24.93	2.55	0.95
BLUP	25.60	24.78	18.41-30.10	25.22	2.12	0.93

BLUP phenotype values based on the best linear unbiased prediction, *SD* standard deviation, H^2 the broad-sense heritability

Table 3 Correlations between spikelet number per spike (SNS) and other agronomic traits in the AM population

Trait	Spikelet number per spike (SNS)
Plant height (PH)	0.30**
Anthesis date (AD)	0.34**
Spike length (SL)	0.70**
Spike extension length (SEL)	0.23*
Kernel number per spike (KNS)	0.36**
Spike density (SD)	0.32**
Thousand kernel weight (TKW)	0.14

*Significance level at $P < 0.05$, **Significance level at $P < 0.01$

Table 4 Quantitative trait loci (QTL) for spikelet number per spike (SNS) identified in multiple environments in the AM population

QTL	Env.	Interval (cM)	Left Marker	Right Marker	LOD	PVE (%)	Add	Physical position (Mb)
QSns.sau-AM-5A	E[1]	116.1-131.9	AX-108790581	AX-111499991	5.62	12.07	-0.95	557.72-571.41[a]
	E[3]	131.9-135.2	AX-111499991	AX-111596791	4.01	8.27	-0.83	
	E[4]	116.1-131.9	AX-108790581	AX-111499991	3.55	10.34	-0.95	
	E[7]	116.1-131.9	AX-108790581	AX-111499991	8.17	9.1812.0	-0.85	
	BLUP	116.1-131.9	AX-108790581	AX-111499991	6.61	9	-0.75	
QSns.sau-AM-2B.1	E[4]	113.7-114.6	AX-109421413	AX-109506426	3.90	15.66	-0.75	710.11-717.87[a]
	E[8]	113.7-114.6	AX-109421413	AX-109506426	3.82	7.54	-0.63	
QSns.sau-AM-2B.2	E[1]	131.3-132.2	KASP-AX-109422178	AX-111588559	9.02	17.99	-1.16	731.25-731.60[a]
	E[2]	131.3-132.2	KASP-AX-109422178	AX-111588559	5.10	9.09	-1.06	
	E[3]	131.3-132.2	KASP-AX-109422178	AX-111588559	11.14	26.57	-1.48	
	E[4]	131.3-132.2	KASP-AX-109422178	AX-111588559	7.35	15.79	-1.17	
	E[5]	130.0-131.3	KASP-AX-95072014	KASP-AX-109422178	4.97	20.39	-0.85	
	E[7]	130.0-131.3	KASP-AX-95072014	KASP-AX-109422178	21.49	25.60	-1.41	
	E[8]	131.3-132.2	KASP-AX-109422178	AX-111588559	8.87	19.32	-1.00	
	BLUP	131.3-132.2	KASP-AX-109422178	AX-111588559	15.97	29.40	-1.16	25.95-47.17[b]
QSns.sau-AM-3B.1	E[4]	22.7-42.5	AX-110399975	AX-109050064	6.61	17.54	1.26	
	E[6]	22.7-42.5	AX-110399975	AX-109050064	3.97	11.85	1.02	
	E[8]	22.7-42.5	AX-110399975	AX-109050064	7.63	16.78	0.95	65.87-72.59[b]
QSns.sau-AM-3B.2	E[1]	48.3-49.7	AX-109393346	AX-110548993	4.82	9.04	0.85	
	E[2]	47.0-48.3	AX-109876826	AX-109393346	3.86	6.71	0.94	
	E[3]	48.3-49.7	AX-109393346	AX-110548993	5.84	12.67	1.05	
	E[7]	47.0-48.3	AX-109876826	AX-109393346	14.77	15.46	1.12	
	BLUP	48.3-49.7	AX-109393346	AX-110548993	8.64	14.27	0.83	

Env. Environment, *BLUP* best linear unbiased prediction; *LOD* logarithm of odds; *PVE* phenotype variance explained; *Add* additive effect, positive values: positive alleles from AL, negative values: positive alleles from LM001. *Physical position*[a] the flanking marker sequences of QTL which positive alleles detected from LM001 were blasted against wild emmer genome to get physical positions; *Physical position*[b] those detected from AL were blasted against durum wheat genome to get physical positions

Figure Legends

Figure 1 The spike morphology of the parents and representative lines from the AM population. (scale bar = 2 cm).

Figure 2 The syntenic relationships between the genetic and physical maps of bin markers. Gen-1A to Gen-7B represented the 15 chromosomal genetic maps used in the current study; Phy-1A to Phy-7B represented the 14 chromosomal physical maps of (**a**) durum wheat and (**b**) wild emmer genomes.

Figure 3 The phenotype, frequency distribution, and correlation coefficients of spikelet number per spike (SNS) in multiple environments. **Significance level at $P < 0.01$.

Figure 4 AM RILs were divided into two haplotype groups based on the genotype of flanking markers, and the SNS differences caused by the corresponding QTL were represented. Genetic maps of (**a**) *QSns.sau-AM-5A*, (**b**) *QSns.sau-AM-2B.2*, (**c**) *QSns.sau-AM-3B.2* and their effects. *Significance level at $P < 0.05$, **Significance level at $P < 0.01$.

Figure 5 The additive effects of *QSns.sau-AM-5A*, *QSns.sau-AM-2B.2*, and *QSns.sau-AM-3B.2* on increasing SNS. '+' and '-' represented haplotype group with and without the positive allele of the corresponding QTL based on the genotype of flanking markers, respectively. Letters a, b, c, and d represent significant difference among haplotype group.

Figure 6 Validation of (**a**) *QSns.sau-AM-2B.2* (KASP-AX-111588559) and (**b**) *QSns.sau-AM-3B.2* (KASP-AX-110548993) in LM001 × PI193877 and AL × AS2268 populations by Student's *t*-test, respectively. **Significance level at $P < 0.01$.

Supplementary Materials

Supplementary Figure 1 Expression patterns at the different development stage of 3 functional orthologs in the physical interval of *QSns.sau-AM-2B.2*. (**a**) *TRIDC2BG075290*; (**b**) *TRIDC2BG075310*; (**c**) *TRIDC2BG075320*.

Supplementary Table 1 Environmental information of surveyed agronomic traits in AM population.

Supplementary Table 2 The 1150 bin markers were used for QTL mapping.

Supplementary Table 3 Comparison of the genetic and physical positions of 6537 mapped markers (including bin markers and co-segregated markers).

Supplementary Table 4 Quantitative trait loci (QTL) detected in the QTL-by-Environment interaction module.

Supplementary Table 5 Epistasis interaction analysis of QTL for spikelet number per spike (SNS).

Supplementary Table 6 Details of KASP markers.

Supplementary Table 7 Comparison of the major quantitative trait loci (QTL) with loci previously reported for spikelet number per spike (SNS).

Supplementary Table 8 Predicted genes in the physical interval of *QSns.sau-AM-2B.2* and *QSns.sau-AM-3B.2*.

AL LM001 #71 #99 #118 #26 #132

(a)

(b)

(c)

QSns.sau-AM-5A	+	+	−	−	+	−	+	−
QSns.sau-AM-2B.2	+	+	+	+	−	−	−	−
QSns.sau-AM-3B.2	+	−	+	−	+	+	−	−
Lines	7	14	7	8	11	4	10	13