Title: Inhibitory interneurons show early dysfunction in a SOD1 mouse model of ALS

Abbreviated Title: Dysfunction of inhibitory interneurons in ALS

C. F. CAVARSAN[1,2], P. R. STEELE[1,2,3], L. M. McCANE[3], K. J. LaPRE[1,2], A. C. PURITZ[4], N. KATENKA[5],
AND K. A. QUINLAN[1,2,4]

1 George and Anne Ryan Institute for Neuroscience, University of Rhode Island, Kingston, RI, 02881 USA
2 Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island,
Kingston, RI, 02881 USA
3 Interdisciplinary Neuroscience Program, University of Rhode Island, Kingston, RI, 02881 USA
4 Department of Physiology, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611 USA
5 Department of Computer Science and Statistics, University of Rhode Island, Kingston, RI, 02881 USA

11                                                                    Corresponding Author:
12                                                                    Katharina A. Quinlan
13                                                                    kaquinlan@uri.edu
14                                        George and Anne Ryan Institute for Neuroscience
15                                                                    University of Rhode Island
16                                                                    130 Flagg Rd
17                                                                    Kingston RI 02881

Number of pages: 30
Number of figures: 5
Number of tables: 4
Number of words in abstract: 150
Number of words in introduction: 619
Number of words in discussion: 1545
24

**Abstract:** Few studies in amyotrophic lateral sclerosis (ALS) focus on the premotor interneurons

synapsing onto motoneurons (MNs). We hypothesized inhibitory interneurons contribute to

dysfunction, particularly if altered before MN neuropathology. We directly assessed excitability and

morphology of ventral lumbar glycinergic interneurons from SOD1$^{G93A}$GlyT2eGFP (SOD1) and wildtype

GlyT2eGFP (WT) mice. SOD1 interneurons were smaller but density was unchanged. Patch clamp

revealed dampened excitability in SOD1 interneurons, including depolarized PICs and voltage threshold.

Renshaw cells (RCs; confirmed with immunohistochemistry) showed similar dampened excitability.

Morphology and electrophysiology were used to create a "random forest" statistical model to predict

RCs when histological verification was not possible. Predicted SOD1 RCs were less excitable (consistent

with experimental results); predicted SOD1 non-RCs were *more* excitable. In summary, inhibitory

interneurons show very early perturbations poised to impact MNs, modify motor output, and provide

early biomarkers of ALS. Therapeutics like riluzole that universally reduce CNS excitability could

exacerbate this dysfunction.

**Key words**: inhibition, glycine, Renshaw cells, ALS, spinal cord, patch clamp, random forest

**Introduction**

Amyotrophic lateral sclerosis (ALS) is a rapidly evolving adult-onset neurological disease characterized by a progressive loss of corticospinal neurons and motoneurons (MNs). There has been considerable debate in the field over the role of hyperexcitability in neurodegenerative processes in ALS. While drug treatment for ALS based on nonspecific reduction of neuronal excitability (with riluzole, for example) (Bellingham 2011),  does result in a modest increase in lifespan (Bensimon et al., 1994), similar treatment paradigms have been disappointing clinically. A more nuanced understanding of the excitability of vulnerable neurons could help in creating a more targeted and effective approach for treatment of ALS. For example, if inhibitory pathways are failing in ALS and neuronal excitability is universally reduced with riluzole, this could further exacerbate inhibitory dysfunction by reducing activity not only in vulnerable MNs but also in inhibitory interneurons presynaptic to MNs. Thus it is important to consider all aspects of neuronal excitability, including intrinsic excitability of vulnerable neurons, synaptic drive, and neuromodulation (Gunes et al., 2020). Intrinsic properties of MNs have been well studied but the same is not true for interneurons that are synaptically connected to them. In fact, no studies thus far have directly assessed activity of spinal premotor interneurons in an ALS model. Despite evidence that ALS patients have disrupted inhibition at spinal levels (Raynor and Shefner, 1994; Shefner and Logigian, 1998; Sangari et al., 2016; Howells et al., 2020), much is still unknown concerning the involvement of inhibitory circuitry in ALS. Morphological alterations in inhibitory circuits have been demonstrated in animal models of ALS. These include degeneration of spinal interneurons, fewer neurons expressing markers of inhibitory neurotransmitters including GlyT2 prior to loss of MNs (Martin et al., 2007; Hossaini et al., 2011), and loss of glycinergic boutons onto MNs prior to symptom onset (Chang and Martin, 2009a). Typically, fast motor neurons have greater numbers of inhibitory synaptic contacts, but these are largely lost in SOD1 mice beginning when motor unit atrophy is first observed (Pun et al., 2006; Hegedus et al., 2007; Allodi et al., 2020). Recurrent inhibitory circuits mediated by Renshaw cells (RCs) are impaired before symptom onset by both loss of MN collaterals which provide synaptic drive, and complex changes to RC-MN synaptic structures (Casas et al., 2013; Wootz et al., 2013). A few studies have suggested activity is decreased in inhibitory interneurons by indirect measurements. Quantification of synaptic inputs to MNs has shown that frequency of inhibitory postsynaptic potentials in spinal MNs is decreased embryonically in both SOD1 mouse and zebrafish models (McGown et al., 2013; Branchereau et al., 2019) and glycinergic inputs to MNs decay faster in SOD1 MNs (Medelin et al., 2016). On a larger scale, blocking the activity of V1 inhibitory interneurons during *in vivo* treadmill running was recently found to mimic the early locomotor deficits in SOD1 mice

(Allodi et al., 2020), suggesting that inhibitory interneurons could be inactive or under-active. However,

none of these studies directly examined electrophysiological activity in inhibitory interneurons.

We hypothesized that inhibitory spinal interneurons could contribute to the pathogenesis of ALS

through a depression of MN inhibition. Decreased activity of inhibitory interneurons could result in

synaptically-driven hyperexcitability of MNs, and other long-term changes in network function. In this

study, we examined electrical and morphological properties of glycinergic interneurons in the spinal

cord of SOD$^{G93A}$ GlyT2-eGFP (SOD1) mice compared to GlyT2-eGFP (WT) using whole cell patch clamp

and three-dimensional reconstructions, immunohistochemistry for the RC marker calbindin (CBD), and

statistical modeling. We show here that significant dysfunction is present in SOD1 glycinergic

interneurons, including RCs and other subclasses of inhibitory interneurons. Impairment in glycinergic

interneurons should be explored as both a mechanism of vulnerability of MNs and a potential biomarker

of early dysfunction of spinal circuits.

**Results**

**Morphology of glycinergic interneurons**

Ventrally located glycinergic interneurons in the lumbar enlargement of WT and SOD1 mice postnatal

day (P)6-10 were reconstructed from image stacks. There was no change in soma size and density of

ventral glycinergic interneurons between WT and SOD1 mice when interneurons throughout the ventral

horn were assessed. However, when we narrowed our examination to only those glycinergic

interneurons located adjacent to MNs within lamina IX (most likely premotor interneurons), SOD1

interneuron somata were smaller in several measures. Typical ventrolateral glycinergic neurons in the

region of the MN pools in lamina IX are shown in the areas delimited with dotted lines in **Figure 1A**.

Complete reconstructions of soma morphology were performed on 3902 GlyT2 interneurons (1918 WT

and 1984 SOD1 interneurons) from fixed lumbar spinal cords (image stacks obtained using confocal

microscopy) and 682 GlyT2 interneurons from only lamina IX (362 WT and 320 SOD1 interneurons).

Reconstructions revealed that glycinergic interneurons in lamina IX were significantly smaller in every

measurement of the soma (soma perimeter, largest cross-sectional area, soma surface area, and soma

volume) (Fig 1B). Cell counts performed in the ventral lumbar spinal cord of 8 WT mice and 9 SOD1 mice

showed no significant differences (Fig 1C). Dendritic morphology was not possible in these images due

to the high number of GFP+ neurons and processes but was completed on a smaller subset of neurons

that were imaged live after patch clamp electrophysiology with a 2-photon microscope.

**Figure 1: GlyT2 interneurons in the ventral horn and lamina IX from WT and SOD1 mice were counted and morphologically analyzed**. Representative images of GFP interneurons in **A**, WT on the top and SOD1 on the bottom. Lamina IX delimited with dotted lines in images of ventral horns. (**B**) Morphological analysis showed no differences between glycinergic neurons in the ventral horn (top panels), but within lamina IX (bottom panels), SOD1 GlyT2 interneurons were smaller than WT in soma perimeter, soma largest cross-sectional area, soma surface and soma volume. (**C**) Density of GlyT2 interneurons were not significantly different between WT and SOD in the ventral horn or in lamina IX. Significance denoted with * indicates $p \leq 0.05$. Scale bar in A: 100 µm.

In these experiments, 59 glycinergic neurons (34 WT and 25 SOD1) were recorded using visually guided

patch clamp with Texas Red dye in the electrode as shown in **Figure 2A.** The location of patched neurons

was distributed throughout the ventral horn, not limited to lamina IX (Fig 2B). Thus, similar to the larger

dataset above, soma sizes of patched SOD1 interneurons were not significantly smaller than WT (Fig 2C).

However, dendrites were larger in patched SOD1 GlyT2 interneurons, including greater dendritic surface

area and volume (see **Table 1** for complete results).

**Electrophysiology of glycinergic interneurons**

Whole cell patch clamp revealed very different intrinsic properties of ventral glycinergic interneurons.

Recording was performed on 59 ventral interneurons from throughout the ventral horn (not limited to

lamina IX) in transverse lumbar spinal cord slices from P6-10 mice. SOD1 inhibitory interneurons (n = 25)

were found to have diminished intrinsic excitability compared to WT interneurons (n = 34).

Measurements of intrinsic excitability included depolarized onset and peak voltage of persistent inward

currents (PICs) and depolarized threshold for action potential firing, as shown in **Figure 3**. In voltage

clamp, slowly depolarizing voltage ramps were used to measure PICs (typical leak subtracted traces

shown in Fig3A). Voltage sensitivity is determined by measuring voltage at PIC onset and peak. Both PIC

onset and peak were significantly depolarized in SOD1 interneurons, indicating less intrinsic excitability.

Amplitude of the PIC was unchanged. Depolarizing current ramps were used to measure the input-

output relationship of the neurons in current clamp. Inhibitory interneurons from SOD1 mice were

found to have significantly higher threshold voltage than control interneurons (Fig 3E-F). This shift is

likely driven by the changes in PIC. The current at firing onset, or I-ON, was not significantly changed ($p$ =

0.15), similar to the current at firing offset, or I-OFF ($p$ = 0.07). After-spike afterhyperpolarization (AHP)

was also significantly different in SOD1 interneurons. The duration (at half AHP amplitude) was shorter

in SOD1 neurons, as shown in Fig 3H-I. Amplitude of the AHP was unchanged. While changes in the PIC

and threshold indicate less excitability in SOD1 interneurons, the shortened AHP suggests an increased

ability to fire at higher rates in SOD1 interneurons, a property that is associated with increased

excitability. However, no changes in firing rates were detected in SOD1 interneurons. All other

properties were found to be similar in SOD1 and WT glycinergic interneurons, including maximum firing

rates, sag/rebound currents ($I_H$), action potential parameters and membrane properties (see **Table 2** for

all electrical properties). See discussion for further interpretation of these findings.

132
133

**Figure 2 GlyT2 interneurons from WT and SOD1 mice were patched throughout the ventral horn**. Images of typical interneurons in **A**, and locations of all patched interneurons in **B** (WT = blue, SOD1 = red), with dotted line indicating area of lamina IX. **(C)** In patched interneurons, soma surface area was unchanged, while dendrite surface area was larger in SOD1 interneurons than in WT. Significance denoted with * indicates $p \leq 0.05$.

134
135

136 **Table 1:** Reconstruction data of GlyT2-GFP+ interneurons in WT and SOD1 animals.

| | | WT (Mean ± SE) | SOD (Mean ± SE) | p |
|---|---|---|---|---|
| **GlyT2-GFP+ Interneurons** | | | | |
| **Ventral horn** | | N = 1918 | N = 1984 | |
| **Soma** | Perimeter (μm) | 70.8 (± 4.4) | 64.3 (± 2.5) | 0.200 |
| | Cross Sectional Area (μm²) | 229.2 (± 17.3) | 215.1 (± 14.4) | 0.535 |
| | Surface Area (μm²) | 413.4 (± 21.7) | 348.9 (± 22.8) | 0.115 [K] |
| | Volume (μm³) | 714.0 (± 64.3) | 552.1 (± 54.6) | 0.115 [K] |
| **Lamina IX** | | N = 362 | N = 320 | |
| **Soma** | Perimeter (μm) | 70.0 (± 4.4) | 58.1 (± 2.5) * | 0.034 [K] |
| | Cross Sectional Area (μm²) | 240.2 (± 17.0) | 186.9 (± 15.9) * | 0.031 |
| | Surface Area (μm²) | 421.7 (± 13.9) | 358.2 (± 21.9) * | 0.025 |
| | Volume (μm³) | 731.7 (± 44.1) | 568.4 (± 49.5) * | 0.022 |
| **Patched GlyT2-GFP+ Interneurons** | | | | |
| | | N = 34 | N = 25 | |
| **Soma** | Perimeter (μm) | 68.4 (± 3.4) | 61.6 (± 3.0) | 0.082 |
| | Cross Sectional Area (μm²) | 174.4 (± 13.4) | 183.1 (± 16.9) | 0.813 [K] |
| | Surface Area (μm²) | 637.5 (± 40.3) | 644.3 (± 47.6) | 0.903 [K] |
| | Volume (μm³) | 1472.4 (± 165.8) | 1611.1 (± 188.9) | 0.523 [K] |
| **Dendrites** | Length (μm) | 101.4 (± 17.3) | 106.3 (± 10.0) | 0.116 [K] |
| | Surface Area (μm²) | 512.7 (± 70.6) | 633.6 (± 52.1) * | 0.024 [K] |
| | Volume (μm³) | 264.7 (± 39.6) | 375.3 (± 37.9) * | 0.010 [K] |

137 Significance denoted with * indicates $p \leq 0.05$.

138 [K] Indicates Kruskall Wallis analysis for non-normal distributions; all others were ANOVA analysis.

**Figure 3: Electrophysiology of SOD1 glycinergic interneurons.** The most prominent difference in SOD1 interneurons was the shift in voltage dependence of PICs. (**A**) Representative, leak-subtracted current-voltage relationship of PICs from WT (P8) and SOD1 (P6) interneurons; arrows indicate voltage at PIC onset and voltage at PIC peak. (**B-D**) Mean onset, peak and amplitude of PICs in WT and SOD1 interneurons. (**E-F**) Threshold voltage (as measured by current ramps as shown) was depolarized in SOD1 interneurons. Starting potential for both interneurons was -65mV. The current at firing onset (I-ON) was not significantly different as shown in **G.** (**H**) The AHP in SOD1 interneurons was shorter, as shown by the smaller mean duration at half amplitude in a representative P8 WT and P7 SOD1 interneuron in **I.** Neither the AHP amplitude nor the AHP decay time constant, tau, were significantly altered in SOD1 glycinergic interneurons as shown in **J-K.** Vertical scale bars in E: top = 50pA, and bottom = 20mV. Horizontal scale bar in E = 0.5s and all scale bars apply to both left and right panels. Vertical scale bar in H = 10mV (APs were truncated from image), and horizontal scale bar = 50ms. Significance denoted with * indicates *p* ≤ 0.05.

143 **Table 2:** Electrophysiological properties of GlyT2-GFP+ interneurons in WT and SOD1 animals.

| Parameters | Groups | | |
|---|---|---|---|
| | WT (Mean ± SE) | SOD (Mean ± SE) | *p* |
| | *N* = 34 | *N* = 25 | |
| Resting membrane potential (mV) | -46.7 ± 1.1 | -45.7 ± 1.4 | 0.524[K] |
| Capacitance (pF) | 66 ± 7 | 55 ± 5 | 0.529[K] |
| Input resistance (MΩ) | 295 ± 32 | 299 ± 41 | 0.794[K] |
| **PIC properties** | | | |
| PIC onset (mV) | -50.5 ± 1.0 | -46.3 ± 1.0* | 0.004 |
| PIC peak (mV) | -35.1 ± 0.8 | -31.2 ± 0.9* | 0.003 |
| PIC amplitude (pA) | 100 ± 13 | 97 ± 12 | 0.890[K] |
| Normalized PIC (pA/pF) | 1.79 ± 0.24 | 1.92 ± 0.22 | 0.226[K] |
| **Firing properties** | | | |
| Inst max firing freq range max (Hz) | 112 ± 8 | 113 ± 8 | 0.961 |
| Inst max firing freq range min (Hz) | 55 ± 3 | 63 ± 7 | 0.442[K] |
| Steady state firing freq range max (Hz) | 36 ± 2 | 40 ± 3 | 0.355[K] |
| Steady state firing freq range min (Hz) | 6 ± 0.4 | 6 ± 0.7 | 0.453[K] |
| I-ON (pA) | -33 ± 12 | -14 ± 11 | 0.148[K] |
| I-OFF (pA) | 5 ± 9 | 36 ± 14 | 0.074[K] |
| Δ I (pA) | 38 ± 8 | 50 ± 10 | 0.331[K] |
| **AP properties** | | | |
| AP overshoot past 0 (mV) | 20.0 ± 1.7 | 21.4 ± 1.6 | 0.553 |
| AP duration at half peak (ms) | 1.53 ± 0.10 | 1.48 ± 0.08 | 0.944 |
| AP rate of rise (V/s) | 87 ± 5 | 83 ± 3 | 0.428[K] |
| AP rate of fall (V/s) | 45 ± 3 | 43 ± 2 | 0.924[K] |
| Threshold on ramp (mV) | -40.6 ± 0.8 | -37.9 ± 1.1* | 0.052[K] |
| Threshold on step (mV) | -47.3 ± 0.8 | -45.1 ± 1.3 | 0.199[K] |
| **AHP properties** | | | |
| AHP amplitude (mV) | 11.2 ± 0.6 | 11.0 ± 0.8 | 0.704[K] |
| AHP duration at half amplitude (s) | 0.12 ± 0.01 | 0.10 ± 0.01* | 0.045[K] |
| AHP tau (s$^{-1}$) | 0.12 ± 0.02 | 0.10 ± 0.02 | 0.980[K] |
| **I$_H$ properties** | | | |
| Hyperpolarizing Sag (%) | 32 ± 3 | 32 ± 4 | 0.537[K] |
| Depolarizing Rebound (%) | 17 ± 3 | 19 ± 5 | 0.536[K] |

144 Significance denoted with * indicates *p* ≤ 0.05.

145 [K] Indicates Kruskall Wallis analysis for non-normal distributions; all others were ANOVA analysis.

**Renshaw cells: electrophysiology and morphology**

Glycinergic interneurons in the ventral spinal cord are not a homogenous population and we sought to determine more specifically what subtype of ventral inhibitory interneurons were morphologically and electrophysiologically altered in SOD1 mice. Since we recorded glycinergic interneurons in the region of lamina IX, we expected our population to include RCs. To distinguish them, we performed immunohistochemistry for the RC marker CBD on GlyT2 neurons labeled with dye during recording (see **Figure 4A**). We also examined density of RCs in lamina IX of the spinal cords, which was unchanged between WT (N = 8 mice)  and SOD1 (N = 7 mice; Figure 4B). Similar to the results above for all GlyT2 interneurons, electrophysiological parameters of SOD1 RCs were very different compared to WT (7 WT and 6 SOD interneurons). The voltage dependence of the PIC (both onset and peak) was significantly depolarized in SOD1 RCs, indicating they are less excitable. Threshold voltage was also significantly higher in SOD1 RCs when measured from step protocols, though it was not significant when measured from current ramp protocols (Figure 4C). All RC electrophysiological properties are included in **Table 3**.

**Figure 4: Renshaw cell (RC) density and electrophysiology.** (**A**) Image of an interneuron which was patched and filled with Texas Red (left panel), and immunoreactive for CBD (middle panel); overlay provided in right panel. The RC is indicated by a white arrow in middle panel. (**B**) No differences were found in the density of CBD+ cells in lamina IX of WT and SOD1 mice. (**C**) Several electrophysiological parameters were measured from the cells. PIC onset, PIC peak and voltage threshold showed higher values in RCs from SOD1 animals, compared to WT RCs. There were no significant differences between groups in PIC amplitude, I-on and AHP duration. Significance denoted with * indicates $p \leq 0.05$. Scale bar in A: 100 μm.

163 **Table 3:** Electrophysiological properties of Renshaw cells, in both groups, WT and SOD1.

| Parameters | Groups | | |
|---|---|---|---|
| | WT (Mean ± SE) | SOD (Mean ± SE) | *p* |
| | N = 7 | N = 6 | |
| **PIC properties** | | | |
| PIC onset (mV) | -53.7 ± 2.5 | -46.3 ± 1.9* | 0.040 |
| PIC peak (mV) | -39.4 ± 2.2 | -31.8 ± 2.0* | 0.027 |
| PIC amplitude (pA) | 109.4 ± 41.6 | 94.6 ± 18.7 | 0.775 [K] |
| Normalized PIC (pA/pF) | 1.76 ± 0.41 | 2.13 ± 0.47 | 0.475 [K] |
| **Firing properties** | | | |
| Inst max firing freq range max (Hz) | 120.8 ± 24.9 | 122.3 ± 17.4 | 0.962 |
| Inst max firing freq range min (Hz) | 55.8 ± 9.2 | 62.0 ± 7.1 | 0.608 |
| Steady state firing freq range max (Hz) | 43.0 ± 8.1 | 43.7 ± 7.6 | 0.954 |
| Steady state firing freq range min (Hz) | 4.8 ± 0.5 | 7.8 ± 1.7 | 0.097 |
| I-ON (pA) | -34.8 ± 15.9 | -35.1 ± 29.7 | 0.985 |
| I-OFF (pA) | 7.9 ± 20.8 | 12.6 ± 37.7 | 0.475 [K] |
| Δ I (pA) | 42.7 ± 14.9 | 48.0 ± 16.1 | 0.567 [K] |
| **AP properties** | | | |
| AP overshoot past 0 (mV) | 21.3 ± 4.0 | 23.9 ± 4.1 | 0.661 |
| AP duration at half peak (ms) | 1.2 ± 0.1 | 1.4 ± 0.1 | 0.452 |
| AP rate of rise (V/s) | 99.1 ± 5.7 | 94.1 ± 4.0 | 0.501 |
| AP rate of fall (V/s) | 54.2 ± 5.7 | 47.5 ± 3.6 | 0.359 |
| Threshold on ramp (mV) | -42.6 ± 1.4 | -39.8 ± 1.9 | 0.262 |
| Threshold on step (mV) | -48.7 ± 1.1 | -45.0 ± 1.0* | 0.032 |
| **AHP properties** | | | |
| AHP amplitude (mV) | 12.6 ± 1.1 | 11.2 ± 1.7 | 0.481 |
| AHP duration at half amplitude (s) | 0.118 ± 0.016 | 0.103 ± 0.017 | 0.516 |
| AHP tau ($s^{-1}$) | 0.103 ± 0.022 | 0.082 ± 0.014 | 1.000 [K] |
| **I$_H$ properties** | | | |
| Hyperpolarizing Sag (%) | 35.3 ± 6.7 | 44.8 ± 11.5 | 0.471 |
| Depolarizing Rebound (%) | 19.1 ± 4.1 | 30.1 ± 17.4 | 0.568 [K] |

Significance denoted with * indicates $p \leq 0.05$.

[K] Indicates Kruskall Wallis analysis for non-normal distributions; all others were ANOVA analysis.

**Predictive statistical modeling**

Since it was not possible to identify all recorded GlyT2 neurons as RCs or non-RCs using

immunohistochemistry, we performed predictive statistical modeling. We created a series of statistical

models using random forest and principal components analysis (see methods for more details on

creation of the models). All variables, including morphology and electrical properties, were grouped into

8 categories of related variables. For example, all variables pertaining to dendritic morphology were

grouped into a single category. Importance of each variable was determined by the decrease in accuracy

of the model when variables were removed. The variables that were used by the random forest model

are listed in **Table 4** in descending order of importance. Variables with large positive values were crucial

for the model's accuracy in predicting RC vs non-RCs, while variables that have negative values proved

detrimental to the predictive value of the model. For comparison, the grouping that provided the best

prediction of CBD-tested neurons using PCA were action potential features (LOOCV accuracy 70%, AIC

=28, p=0.11 and 0.08, PC1 and PC2, respectively). Variables in this grouping that had effect sizes ≥ 0.5

were I-off, AP threshold, AP rise and AP fall. The most accurate random forest model correctly predicted

the identity of the CBD-tested interneurons with 79% accuracy; thus we used this model to predict RCs

and non-RCs in all recorded GlyT2+ interneurons in which RC identification using immunohistochemistry

was not possible. The model predicted that 80% of our 59 patched interneurons were RCs. This

percentage is greater than that measured from our immunohistochemistry results: 65% of our patched

interneurons (15/23) were positive for CBD.  Predicted RCs were then analyzed as a group for

differences between WT and SOD interneurons, while predicted non-RCs were analyzed separately. We

found that in predicted RCs, there was a depression of excitability very consistent with our experimental

data. The predicted RC grouping had significant depolarization of the PIC and significantly depolarized

threshold. PIC parameters of the predicted groups are shown in **Figure 5**. Interestingly, in predicted non-

RCs excitability increased. Both the PIC amplitude and the PIC normalized to capacitance were larger in

the group of predicted non-RCs from SOD1 mice. Further studies will be needed to investigate more fully

the properties of different subtypes of non-RC glycinergic interneurons in this and other ALS models at

this and other time points in the lead up to MN dysfunction and degeneration.

**Table 4:** Variables used to create the random forest predictive model. Variables are arranged in descending order of importance for prediction of RCs and non-RCs, based on the mean decrease in accuracy test.

| Variable | Mean Decrease in Accuracy |
|---|---|
| Dendrite Mean Length ($\mu$m) | 8.27508593 |
| Cell Body Surface Area ($\mu$m$^2$) | 3.98158541 |
| Capacitance (pF) | 3.82644995 |
| AHP tau (s$^{-1}$) | 3.42103087 |
| Cell Body Length ($\mu$m) | 2.97501036 |
| Perimeter ($\mu$m) | 2.15259564 |
| Dendrite Surface Area ($\mu$m$^2$) | 1.58220831 |
| Cell Body Volume ($\mu$m$^3$) | 1.30757721 |
| Cell Body Surface Area ($\mu$m$^2$) | 1.17777719 |
| V threshold (mV) | 0.99130144 |
| Cell Body Area ($\mu$m$^2$) | 0.89067177 |
| AP size (mV) | 0.59738713 |
| Enclosed Volume ($\mu$m$^3$) | 0.57054623 |
| Steady state firing freq max (Hz) | 0.28434868 |
| I-ON (pA) | 0.13715380 |
| $I_H$ Sag (mV) | -0.06241806 |
| Dendrite Volume ($\mu$m$^3$) | -0.41041638 |
| Dendrite Length ($\mu$m) | -0.91989565 |
| Instantaneous max firing rate (Hz) | -0.96322954 |
| Inst max firing freq range max (Hz) | -1.40411698 |
| AP overshoot (mV) | -1.53244473 |
| $I_H$ Rebound (mV) | -1.95402459 |

## PIC parameters for predicted Renshaw cells

## PIC parameters for predicted non-Renshaw cells

**Figure 5** Genotype is predicted to have opposite effects on predicted RCs and non-RCs. **A – D** Predicted RCs have depolarized PIC onset, similar to overall experimental results. This is accompanied by depolarized threshold (data not shown). **E – H** Predicted non-RC glycinergic interneurons have larger PIC amplitudes, including larger normalized PIC amplitudes. * Significance denoted with * indicates $p \leq 0.05$.

**Discussion**

This study shows for the first time that RC are depressed in excitability and glycinergic interneurons may

be contributing to ALS pathology. Alterations in both morphology and excitability of glycinergic

interneurons occurs at a very early stage. RCs are depressed in activity and lamina IX interneurons are

smaller in SOD1 mice. Based on our statistical modeling, we predict the remaining non-RC inhibitory

interneurons have increased excitability, including larger PICs. We speculate that these shifts in

interneuron excitability could alter the excitability of postsynaptic MNs, modify motor output, and

provide an early biomarker of ALS.

*Intrinsic and synaptic hyperexcitability*

A core issue this study brings to light is potential long-term disruption to the balance of excitatory and

inhibitory synaptic transmission in ALS. Spinal cord and cortex both show imbalance between excitatory

and inhibitory synaptic transmission in SOD$^{G93A}$ mice (Avossa et al., 2006; Saba et al., 2016). Clinically,

ALS patients show evidence of altered synaptic activity as well, including a reduction in inhibition. A

reduction of glycinergic receptor binding in the anterior gray matter of the spinal cord (Hayashi et al.,

1981; Whitehouse et al., 1983) has been demonstrated along with abnormal glycine and gamma amino

butyric acid (GABA) levels in blood serum (Malessa et al., 1991; Niebroj-Dobosz and Janik, 1999).

Electrophysiology also suggests disruption in spinal inhibitory circuits in ALS patients (Raynor and

Shefner, 1994; Shefner and Logigian, 1998; Sangari et al., 2016). An interesting question is whether the

neurons are altered in excitability only within the motor circuit (i.e. corticospinal neurons and the

synaptically-connected spinal inhibitory and cholinergic interneurons, and MNs), or if disturbances are

more widespread throughout the nervous system. In the spinal cord, interneurons that serve as conduits

between vulnerable corticospinal and spinal MNs include cholinergic interneurons, Ia inhibitory

interneurons and RCs. There is abundant evidence of disruption in these cholinergic, GABAergic and

glycinergic synapses from animal models (Martin et al., 2007; Chang and Martin, 2009b; Pullen and

Athanasiou, 2009; Hossaini et al., 2011; Herron and Miles, 2012; Casas et al., 2013; Wootz et al., 2013;

Saxena et al., 2013; Milan et al., 2015; Dukkipati et al., 2016; Medelin et al., 2016). Similarly, in the

cortex, neurons that are not themselves vulnerable to neurodegeneration in ALS also exhibit altered

excitability and morphology in a parallel time course to the vulnerable corticospinal neurons (Clark et al.,

2017, 2018; Kim et al., 2017). The changes in excitability of other neuronal populations must contribute

to the imbalance of both excitatory and inhibitory neurotransmission that has been clearly

demonstrated in ALS patients and animal models.

A case for hyperexcitability and glutamate excitotoxicity involvement in ALS pathogenesis has been

more controversial. Altered intrinsic excitability has been consistently reported in both corticospinal

projection neurons and spinal MNs, but these perturbations appear to fluctuate based on age/disease

progression. Corticospinal projection neurons have increased firing and other signs of hyperexcitability

during the first week of postnatal development (Pieri et al., 2009; Saba et al., 2016, 2019; Kim et al.,

2017), but excitability is normal in young and adult pre-symptomatic ages (Kim et al., 2017; Saba et al.,

2019), and may become hypoexcitable at the age of symptom onset (Saba et al., 2019).  Similarly, spinal

MNs show intrinsic hyperexcitability at very early (embryonic) stages (Kuo et al., 2004, 2005; Martin et

al., 2013). At a postnatal age similar to the inhibitory interneurons in this study, spinal MNs do *not* have

an altered threshold, rheobase or frequency-current relationship, though other abnormalities are

present in PICs, action potential duration, AHP and dendritic $Ca^{2+}$ entry (Pambo-Pambo et al., 2009;

Quinlan et al., 2011, 2015; Leroy et al., 2014). Thus, at the time point studied here, inhibitory

interneurons are more disrupted in excitability than the vulnerable spinal MNs. At symptom onset, MNs

may finally succumb to hypo excitability, though not all studies agree on this point (Delestrée et al.,

2014; Martinez-Silva et al., 2018; Jensen et al., 2020). Notably, MNs derived from ALS patients' induced

pluripotent stem cells also show initial hyperexcitability followed by hypoexcitability (Wainger et al.,

2014; Devlin et al., 2015), and increasing excitability of MNs enhanced neuroprotection rather than

neurodegeneration (Saxena et al., 2013). A unifying theme throughout the findings is that vulnerable

MNs show excessive homeostatic gain in response to perturbation (Kuo et al., 2020), and not only

excitability but other cellular properties wildly fluctuate over the lifespan of the animal (Irvin et al.,

2015). In light of that, the present findings that inhibitory interneurons are less excitable should be

further explored to fully characterize their contribution to neurodegenerative processes.

*Changes in neuron morphology*

In ALS models, altered morphology in spinal MNs occurs so early that it could be viewed as a deficit in

normal development. Embryonically, SOD1 MNs have shorter projections but this is reversed during

postnatal development.  At 1-2 weeks of age, MNs begin to show more dendritic branching and larger

soma sizes (Amendola and Durand, 2008; Filipchuk and Durand, 2012; Martin et al., 2013). Spinal MNs

remain larger through adulthood (Dukkipati et al., 2018). In contrast to the spinal MNs at this age, we

found that inhibitory interneurons in lamina IX were smaller and patched interneurons had expanded

dendritic surface area and volume. This is reminiscent of changes in corticospinal neurons:

presymptomatic corticospinal neurons also have increased arborization but with smaller soma

diameters (Ozdinler et al., 2011; Saba et al., 2016). In both sporadic and familial ALS patients,

postmortem tissue shows degeneration of apical dendrites in corticospinal neurons and smaller soma

size (Genç et al., 2017). Changes in neuronal size over the life span could greatly increase the impact of

other stressors (e.g., proteostasis, metabolic deficits, intracellular transport) on vulnerable neuron

populations.

*After spike afterhyperpolarization*

The AHP can contribute to a neuron's firing rate such that a neuron with a large, long-lasting AHP will

typically fire at a slower rate (typical of slow/type I MNs) than a neuron with small, fast-decaying AHP

(typical of fast/type II MNs). In ALS patients, AHP is shortened in MNs of patients with little force deficits

and later elongated in patients with large force deficits (Piotrkiewicz and Hausmanowa-Petrusewicz,

2011), which could reflect early changes in physiology of vulnerable fast MNs and later, the remaining,

less-vulnerable population of slow MNs. Similarly, SOD1 mouse MNs show shorter AHPs early, and

longer AHPs around symptom onset (Quinlan et al., 2011; Jensen et al., 2020). Here we show that AHP is

shorter in duration in SOD1 interneurons very early, however we did not find changes in any measure of

the minimum or maximum firing rates in GlyT2 interneurons or RCs, so the physiological importance of

this finding is not clear.

*Inhibitory interneurons: Subtype specific effects*

The interneurons in this study were mostly composed of RCs, based on both our positivity rate with

immunohistochemistry for CBD (65% positive) and our statistical modeling (80% RC / 20% non-RC).

Interneurons located in lamina IX likely reflect a mix of RCs and Ia inhibitory interneurons. Based on

highly significant results in our RC group and the prediction of higher excitability in non-RC inhibitory

interneurons, further studies are warranted to examine how inhibitory interneurons may be differently

modulated in a subtype-specific manner in animal models of ALS.

*Functional implications for impaired inhibitory circuits*

Outward signs of improperly functioning inhibitory interneurons may be apparent from subtle changes

in motor patterning and could potentially be used as a biomarker of early ALS. Locomotor disturbances

have been demonstrated *in vivo* in presymptomatic SOD1 mice during walking/running (Vinsant et al.,

2013a, 2013b; Akay, 2014; Quinlan et al., 2017; Allodi et al., 2020). In the embryonic and neonatal spinal

cord *in vitro*, increased duration of depolarizing events in MNs and a slower period of locomotor-related

bursting has been described (Medelin et al., 2016; Branchereau et al., 2019), which indicate that even

during development the functioning of locomotor circuits is already impaired. In presymptomatic adult

mice capable of running on a treadmill, subtle locomotor differences are observed including advanced

intermuscular phasing and a slower speed (Quinlan et al., 2017; Allodi et al., 2020) which indicate that

before any large scale neurodegeneration occurs, there are changes in patterns of activity in MNs and

interneurons that could be exploited for use as a biomarker of ALS. At initial symptom onset, the first

failure of the tibialis anterior manifests as abnormality in the swing phase of the ankle (Akay, 2014).

Interestingly, silencing different interneuron populations (including inhibitory interneurons and

cholinergic interneurons) reveals the contribution of each population: silencing cholinergic interneurons

increases locomotor deficits in SOD1 mice (Landoni et al., 2019), while silencing inhibitory interneurons

has no effect suggesting these interneurons are already impaired (Allodi et al., 2020).

*Clinical-translational implications*

Evidence of MN hyperexcitability in ALS patients is present alongside signs of dysfunction in inhibition

(Rothstein et al., 1992; Plaitakis and Constantakakis, 1993; Raynor and Shefner, 1994; Shefner and

Logigian, 1998; Andreadou et al., 2008; Bae et al., 2013). Perhaps reducing excitability (with riluzole, for

example) only results in a modest increase in lifespan (Bensimon et al., 1994) because it could be helpful

to MNs while exacerbating inhibitory dysfunction by further depressing excitability of inhibitory

interneurons like RCs. Thus, a more targeted approach to reduce MN excitability could be beneficial

early in disease pathology. Stimulation of inhibitory nerves or certain protocols of transcranial

stimulation could be explored, along with augmentation of inhibitory neurotransmission.

**Materials and Methods**

**Ethics Statement**

Experiments were performed in accordance with the United States National Institutes of Health Guide for

Care and Use of Laboratory Animals. Approval of the Northwestern University's Animal Care and Use

Committee was obtained for all experiments performed in this study. All efforts were made to minimize

animal suffering and to reduce the number of animals used.

**Animals and Tissue harvest**

Transgenic B6SJL mice overexpressing the human *SOD1$^{G93A}$* gene (strain 002726, Jackson Labs, Bar Harbor,

ME, USA) and their wild type littermates were used (nontransgenic for the human *SOD1$^{G93A}$* gene).

Transgenic animals were identified using standard PCR techniques by Transnetyx (Cordova, TN, USA) and

were bred with GlyT2-eGFP mice (Zeilhofer et al., 2005) generating SOD1$^{G93A}$ GlyT2-eGFP mice, here called

SOD1. Inhibitory glycinergic interneurons express the Na$^+$ and Cl$^-$ coupled glycine transporter 2, or GlyT2.

GlyT2-eGFP expression is driven by the GlyT2 promotor. Specifically, GlyT2-eGFP females were bred with

SOD1$^{G93A}$ GlyT2-eGFP males, and progeny were used for experiments without knowing if they were

SOD1$^{G93A}$ GlyT2-eGFP or GlyT2-eGFP. Genotyping was performed post-experimentally. For the following

studies juvenile mouse pups were used between postnatal day (P) 6 - 10. Mice were deeply anesthetized

with isoflurane (Henry Schein Animal Health, Dublin, OH, USA), decapitated and eviscerated. The lumbar

spinal cord from L1 - L6 was removed and embedded in 2.5% w/v agar (No. A-7002, Sigma-Aldrich, St

Louis, MO, USA). The agar block was then superglued with Loctite 401 (Henkel Corporation, Rocky Hill,

CN, USA) to a stainless steel slicing chuck and 350 μm transverse slices were made using the Leica 1000

vibratome (Leica Microsystems, Buffalo Grove, IL, USA) as described previously (Quinlan et al., 2011).

During both spinal cord isolation and slicing, the spinal cord was immersed in 1–4°C high osmolarity

dissecting solution containing (mM): sucrose 234.0, KCl 2.5, CaCl$_2$ · 2H$_2$O 0.1, MgSO$_4$ · 7H$_2$O 4.0, HEPES

15.0, glucose 11.0, and Na$_2$PO$_4$ 1.0. The pH was adjusted to 7.35 when bubbled with 95% O$_2$/5% CO$_2$ using

1 M KOH (Fluka Analytical, Sigma-Aldrich). After cutting, the slices were incubated for >1 h at 30°C in

incubating solution containing (mM): NaCl 126.0, KCl 2.5, CaCl$_2$ · 2H$_2$O 2.0, MgCl$_2$ · 6H$_2$O 2.0, NaHCO$_3$ 26.0,

glucose 10.0, pH 7.4 when bubbled with 95% O$_2$/5% CO$_2$ (all reagents for solutions were purchased from

Sigma-Aldrich).

**Electrophysiology**

Whole cell patch clamp was performed on MNs from the lumbar segments using 2–4 MΩ glass electrodes

pulled from glass capillary tubes (Item #TW150F-4, World Precision Instruments, Sarasota, FL, USA) with

a Flaming-Brown P-97 (Sutter Instrument Company, Novato, CA, USA). Electrodes were positioned using

a Sutter Instrument MP-285 motorized micromanipulator (Sutter Instrument Company). Whole-cell patch

clamp measurements were performed at room temperature using the Multiclamp700B amplifier

(Molecular Devices, Burlingame, CA, USA) and Winfluor software (University of Strathclyde, Glasgow,

Scotland). Briefly, slices were perfused with a modified Ringer's solution containing (in mM): 111 NaCl,

3.09 KCl, 25.0 NaHCO$_3$, 1.10 KH$_2$PO$_4$, 1.26 MgSO$_4$, 2.52 CaCl$_2$, and 11.1 glucose. The solution was

oxygenated with 95% O$_2$/5% CO$_2$ and the perfusion rate was 2.5 – 3.0 ml/min. Patch electrodes contained

(in mM) 138 K-gluconate, 10 HEPES, 5 ATP-Mg, 0.3 GTP-Li and Texas Red dextran (150 μM, 3000 MW,

from Invitrogen, Life Technologies, Grand Island, NY, USA). In voltage-clamp mode, fast and slow

capacitance transients, as well as whole-cell capacitance, were compensated using the automatic

capacitance compensation on the Multiclamp. Holding potential was set at −90 mV, and neurons were

subjected to slow, depolarizing voltage ramps of 22.5 mV s$^{-1}$, bringing the cell to 0 mV in 4 s, and then

back to the holding potential in the following 4 s. In current clamp, neurons were subjected to depolarizing

current ramps for testing I-on (the current level at firing onset), I-off (the current level at cessation of

firing), and the slope of the frequency–current relationship. Negative current was often necessary to

prevent action potential (AP) firing.

**Neuron Selection**

Glycinergic interneurons were visually selected for recording based on 1) expression of GFP, and 2)

location near the MN pools. Neurons that did not repetitively fire or did not maintain a resting membrane

potential below -35 mV were excluded from electrophysiological analysis. Neurons that were not clearly

stained with Texas Red after patch clamp and/or could not be imaged clearly were not included in

anatomical analysis.

**Electrophysiological analysis**

Persistent inward current (PIC) parameters were measured after subtraction of leak current. Onset was

defined as the voltage at which the current began to deviate from the horizontal, leak-subtracted trace.

PIC peak was the voltage at which the PIC reached peak amplitude. The linear portion of leak current

(usually between -90 and -75 mV) was used to calculate whole cell input resistance. The first action

potential in the train evoked by a depolarizing current ramp in current clamp mode was used to measure

all parameters relating to action potentials. Threshold voltage was defined as the voltage at which the

slope exceeded 10 V/s. Threshold for action potential firing was tested in two ways. The first was to use

the voltage at firing onset from the current ramps up to 130 pA/s (in current clamp). The second was the

voltage at which a single action potential could be evoked with a current step. Action potential overshoot

is the voltage past 0 mV the first spike reaches. Duration of the action potential is measured at half of

action potential height (height is defined as overshoot – threshold voltage). Rates of rise and fall are

defined as the peak and the trough of the first derivative of the action potential profile.

**Immunohistochemistry**

Immediately after electrophysiological recording, slices were fixed in (4%) paraformaldehyde for 24 h. As

described by Kupferschmidt and colleagues (Kupferschmidt et al., 2015), sections were then incubated

in sodium borohydride ($NaBH_4$; 5mg/mL in $diH_2O$) to quench aldehyde-induced catecholamine

fluorescence and reduce free aldehyde groups, followed by Sudan Black B solution (0.2% in 70%

ethanol). Slices were washed twice for 30 min in room temperature PBS to extinguish the

immunohistochemistry reaction. Slices were blocked using 2% normal horse serum in PBST (0.2% Triton-

X100 in PBS) for 4 h, and incubated in PBST with rabbit anti-Texas Red antibody (polyclonal IgG, 1:2000,

ThermoFisher Scientific®, A-6399); mouse anti-Calbindin-D-28K (monoclonal; 1:3000, Sigma, C9848) for

72 h at 4°C. Following four washes in PBST over a total of 16–24 h, slices were incubated in the

appropriate fluorescent dye-conjugated secondary antibodies (goat anti-rabbit IgG, Texas Red-

conjugated, and goat anti-mouse IgG, Cyanine5-conjugated, both ThermoFisher Scientific®, 1:500, T-

6391, and 1:500, A10524, respectively) for 48 h. Then they were washed again in 0.1M PBS and DAPI

treated for 5 min, washed and covered using Fluoromount® and glass coverslips.

**Morphological analysis**

Neuron morphology was assessed from two types of images: 1) 2 photon image stacks of live neurons

taken immediately after obtaining electrophysiological parameters, and 2) confocal image stacks taken

after fixation and tissue processing. Images collected using these two methods were analyzed

separately. Measurements from 2-photon images included measurements from both soma and

dendrites (soma volume, maximum soma cross-sectional area, soma perimeter in the largest plane,

soma surface area, number of primary dendrites, and average dendrite length, dendrite surface area

and dendrite volume). Measurements from confocal images only included soma measurements.

*Two photon imaging:* An Olympus BX-51WIF microscope fitted with an Olympus 40x/0.8NA water-

dipping objective lens was used. Two-photon excitation fluorescence microscopy was performed with a

galvanometer-based Coherent Chameleon Ultra II laser (Coherent, Santa Clara, CA, USA) tuned to 900

nm. A red Bio-Rad 2100MPD photomultiplier tube (Bio-Rad, Hercules, CA, USA) (570 – 650 nm) was used

to collect emission in the red channel. Z-stacks were obtained for each interneuron at 1024 x 1024 pixels

(308 x 308 µm) resolution and roughly 100 µm in depth (step size 1 µm). From these Z-stacks, Texas

Red® -filled neurons were three-dimensionally reconstructed using Neurolucida software (MBF

Bioscience, Williston, VT, USA).

*Confocal imaging:* Sections were observed with a confocal laser microscope (LSM510, Zeiss,

Oberkochen, Darmstadt, Germany) with argon (488 nm) and helium-neon (543 nm) lasers and a 20X

objective. Each confocal image obtained in triple staining experiments was split into three channels. Co-

localization analysis of CBD, Texas Red and GlyT2 positive cells included visual inspection of labeling,

size, and shape of the cell through the stack. From these Z-stacks, GlyT2 positive interneurons were

three-dimensionally reconstructed using Neurolucida software (MBF Bioscience, Williston, VT, USA).

CBD positive cells were counted using ImageJ (National Institutes of Health, Bethesda, MD).

**Statistical analysis**

The assumptions of homogeneity of variances and the normality of the distribution of values for each

measured characteristic were evaluated with Levene's and Shapiro-Wilk tests, respectively.  Group

comparisons (WT vs SOD1) were performed using ANOVA on characteristics that satisfied both

conditions and using the Kruskal-Wallis test for the rest of the characteristics. Results are presented as

means ± SEM. Significance level was set at $p \leq 0.05$.

*Statistical modeling: Random Forest and Principal Component Analysis*

As it was not possible to confirm or rule out RC identity in the majority of our neurons using

immunohistochemistry, we created statistical models based on those we had tested to predict RC

identity among remaining untested neurons. To do this, we employed a random forest method of

statistical learning that generates a large number of decision trees, each constructed using a different,

randomly selected (with replacement) subset of the training data set. Thus, all variables in the dataset

were first split into 8 subgroups of related variables. Groups included membrane properties, action

potential measurements, PIC properties, firing frequency measurements, AHP parameters, $I_H$

characteristics, soma measurements and dendritic properties. For example, all variables pertaining to

PIC characteristics (PIC onset, peak voltage, PIC amplitude and normalized amplitude) were grouped.

Next, a random forest model (Breiman, 2001) was created for each subgroup of variables varying the

number of measurements to include in each tree. This method determined the number of variables that

yielded the most accurate model. The reason this was performed (instead of including all measured

parameters) is that models with larger numbers of variables tend to overfit the training dataset and fail

to classify accurately unseen/test cases. Therefore, it was essential to distill down our parameters to

only include the most important variables for classifying neurons as RC or non-RC. The decision trees are

then used to classify an observation by selecting the most common label among multiple trees. Random

forest allows varying two parameters: the number of variables to consider when building each tree and

the number of generated trees. One can choose a set of these two parameters that yields the most

optimal model. The random forest model with the lowest error was then used to evaluate the most

important variables in each subgroup for distinguishing between RCs and non-RCs. Here we used Mean

Decrease in Accuracy Index to evaluate the importance of the various measured characteristics of the

neurons in predicting RCs vs non-RCs. After the important variables in each subgroup were identified, a

dataset was created with only these variables and the above process was repeated using these

important variables.  Model accuracy was evaluated by using a leave-one-out cross validation (LOOCV).

The accuracy for the best random forest model created from the previously determined important

variables was 79%. The best random forest model was then used to predict the RC/non-RC classification

for untested cells in the dataset.  Once the RC/non-RC classification was determined, normality and

homogeneity tests were run on each group followed by ANOVA and Kruskal Wallis in order to determine

group differences between WT RCs and SOD1 RCs as well as between WT non-RCs and SOD1 non-RCs.

Principal components analysis (PCA) (Hotelling, 1933) of each previously created subgroup of variables

was also evaluated for predictive value. The scores from the first two principal components in each

subgroup were used in a logistic regression model to predict the known CBD-tested cells. Using PCA, the

most important variables were identified as the same as those identified by the random forest method.

The model accuracy was also tested using LOOCV.  The best PCA model accuracy was 70%, therefore,

the random forest model (with 79% accuracy based on LOOCV) was selected to predict classification of

the remaining untested neurons rather than this method.

**Acknowledgments**
This project was funded by a springboard fellowship from Target ALS to KAQ and a University of Rhode
Island Graduate School Tuition Scholarship to LMM. The authors thank Drs Emily Reedich and Claudia
Fallini for comments on the initial draft of the manuscript.
471

**References**

Akay T (2014) Long-term measurement of muscle denervation and locomotor behavior in individual wild-type and ALS model mice. J Neurophysiol 111:694–703.

Allodi I, Montañana-Rosell R, Selvan R, Löw P, Kiehn O (2020) Locomotor deficits in ALS mice are paralleled by loss of V1-interneuron-connections onto fast motor neurons. bioRxiv:2020.06.23.166389 Available at: http://biorxiv.org/content/early/2020/06/23/2020.06.23.166389.abstract.

Amendola J, Durand J (2008) Morphological differences between wild-type and transgenic superoxide dismutase 1 lumbar motoneurons in postnatal mice. J Comp Neurol 511:329–341.

Andreadou E, Kapaki E, Kokotis P, Paraskevas GP, Katsaros N, Libitaki G, Petropoulou O, Zis V, Sfagos C, Vassilopoulos D (2008) Plasma glutamate and glycine levels in patients with amyotrophic lateral sclerosis. In Vivo (Brooklyn) 22:137–142.

Avossa D, Grandolfo M, Mazzarol F, Zatta M, Ballerini L (2006) Early signs of motoneuron vulnerability in a disease model system: Characterization of transverse slice cultures of spinal cord isolated from embryonic ALS mice. Neuroscience 138:1179–1194.

Bae JS, Simon NG, Menon P, Vucic S, Kiernan C, Doyle AC, Case-book T, Holmes S (2013) The Puzzling Case of Hyperexcitability in Amyotrophic Lateral Sclerosis What Really is Hyperexcitability in ALS ? How Can We Objectively Assess. J Clin Neurol 9:65–74.

Bensimon G, Lacomblez L, Meininger V (1994) A Controlled Trial of Riluzole in Amyotrophic Lateral Sclerosis. N Engl J Med 330:585–591.

Branchereau P, Martin E, Supiot L, Hodeib F, Laupénie A, Dalvi U, Zhu H, Cazenave W, Cattaert D (2019) Relaxation of synaptic inhibitory events as a compensatory mechanism in fetal SOD spinal motor networks. Elife 8:1–28.

Breiman L (2001) Random forests. Mach Learn 45:5–32.

Casas C, Herrando-Grabulosa M, Manzano R, Mancuso R, Osta R, Navarro X (2013) Early presymptomatic cholinergic dysfunction in a murine model of amyotrophic lateral sclerosis. Brain Behav 3:145–158.

Chang Q, Martin LJ (2009a) Glycinergic Innervation of Motoneurons Is Deficient in Amyotrophic Lateral Sclerosis Mice A Quantitative Confocal Analysis. Am J Pathol 174:574–585.

Chang Q, Martin LJ (2009b) Glycinergic innervation of motoneurons is deficient in amyotrophic lateral sclerosis mice: A quantitative confocal analysis. Am J Pathol 174:574–585.

Clark RM, Blizzard CA, Young KM, King AE, Dickson TC (2017) Calretinin and Neuropeptide y interneurons are differentially altered in the motor cortex of the SOD1 G93A mouse model of ALS. Sci Rep 7:1–13

Available at: http://dx.doi.org/10.1038/srep44461.

Clark RM, Brizuela M, Blizzard CA, Dickson TC (2018) Reduced excitability and increased neurite

complexity of cortical interneurons in a familial mouse model of amyotrophic lateral sclerosis. Front Cell

Neurosci 12:1–7.

Delestrée N, Manuel M, Iglesias C, Elbasiouny SM, Heckman CJ, Zytnicki D (2014) Adult spinal

motoneurones are not hyperexcitable in a mouse model of inherited amyotrophic lateral sclerosis. J

Physiol 592:1687–1703.

Devlin AC, Burr K, Borooah S, Foster JD, Cleary EM, Geti I, Vallier L, Shaw CE, Chandran S, Miles GB

(2015) Human iPSC-derived motoneurons harbouring TARDBP or C9ORF72 ALS mutations are

dysfunctional despite maintaining viability. Nat Commun 6:1–12 Available at:

http://dx.doi.org/10.1038/ncomms6999.

Dukkipati SS, Chihi A, Wang Y, Elbasiouny SM (2016) Experimental Design and Data Analysis Issues

Contribute to Inconsistent Results of C-Bouton Changes in Amyotrophic Lateral Sclerosis. eNeuro 4:1–

13.

Dukkipati SS, Garrett TL, Elbasiouny SM (2018) The vulnerability of spinal motoneurons and soma size

plasticity in a mouse model of amyotrophic lateral sclerosis. J Physiol 596:1723–1745.

Filipchuk AA, Durand J (2012) Postnatal dendritic development in lumbar motoneurons in mutant

superoxide dismutase 1 mouse model of amyotrophic lateral sclerosis. Neuroscience 209:144–154

Available at: http://dx.doi.org/10.1016/j.neuroscience.2012.01.046.

Genç B, Jara JH, Lagrimas AKB, Pytel P, Roos RP, Mesulam MM, Geula C, Bigio EH, Özdinler PH (2017)

Apical dendrite degeneration, a novel cellular pathology for Betz cells in ALS. Sci Rep 7:1–10 Available at:

http://dx.doi.org/10.1038/srep41765.

Gunes ZI, Kan VWY, Ye XQ, Liebscher S (2020) Exciting Complexity: The Role of Motor Circuit Elements in

ALS Pathophysiology. Front Neurosci 14:1–30.

Hayashi H, Suga M, Satake M, Tsubaki T (1981) Reduced glycine receptor in the spinal cord in

amyotrophic lateral sclerosis. Ann Neurol 9:292–294.

Hegedus J, Putman CT, Gordon T (2007) Time course of preferential motor unit loss in the SOD1G93A

mouse model of amyotrophic lateral sclerosis. Neurobiol Dis 28:154–164.

Herron LR, Miles GB (2012) GENDER-SPECIFIC PERTURBATIONS IN MODULATORY INPUTS TO

MOTONEURONS IN A MOUSE MODEL OF AMYOTROPHIC LATERAL SCLEROSIS. Neuroscience 226:313–

323 Available at: http://dx.doi.org/10.1016/j.neuroscience.2012.09.031.

Hossaini M, Cano SC, Van Dis V, Haasdijk ED, Hoogenraad CC, Holstege JC, Jaarsma D (2011) Spinal

inhibitory interneuron pathology follows motor neuron degeneration independent of glial mutant

superoxide dismutase 1 expression in SOD1-ALS mice. J Neuropathol Exp Neurol 70:662–677.

Hotelling H (1933) Analysis of a complex of statistical variables into Principal Components. J Educ

Psychol 24:417–441.

Howells J, Sangari S, Matamala JM, Kiernan MC, Marchand-Pauvert V, Burke D (2020) Interrogating

interneurone function using threshold tracking of the H reflex in healthy subjects and patients with

motor neurone disease. Clin Neurophysiol.

Irvin CW, Kim RB, Mitchell CS (2015) Seeking homeostasis: Temporal trends in respiration, oxidation, and

calcium in SOD1 G93A amyotrophic lateral sclerosis mice. Front Cell Neurosci 9:1–11.

Jensen DB, Kadlecova M, Allodi I, Meehan CF (2020) Spinal motoneurones are intrinsically more

responsive in the adult G93A SOD1 mouse model of Amyotrophic Lateral Sclerosis. J Physiol.

Kim J, Hughes EG, Shetty AS, Arlotta P, Goff LA, Bergles DE, Brown SP (2017) Changes in the Excitability

of Neocortical Neurons in a Mouse Model of Amyotrophic Lateral Sclerosis Are Not Specific to

Corticospinal Neurons and Are Modulated by Advancing Disease. J Neurosci 37:9037–9053 Available at:

http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0811-17.2017.

Kuo JJ, Schonewille M, Siddique T, Schults AN, Fu R, Bar PR, Anelli R, Heckman C, Kroese AB (2004)

Hyperexcitability of Cultured Spinal Motoneurons From Presymptomatic ALS Mice. J Neurophysiol

91:571–575 Available at: http://jn.physiology.org/cgi/doi/10.1152/jn.00665.2003.

Kuo JJ, Siddique T, Fu R, Heckman CJ (2005) Increased persistent Na+ current and its effect on

excitability in motoneurones cultured from mutant SOD1 mice. J Physiol 563:843–854.

Kuo SW, Binder MD, Heckman CJ (2020) Excessive Homeostatic Gain in Spinal Motoneurons in a Mouse

Model of Amyotrophic Lateral Sclerosis. Sci Rep 10.

Kupferschmidt DA, Cody PA, Lovinger DM, Davis MI (2015) Brain BLAQ: Post-hoc thick-section

histochemistry for localizing optogenetic constructs in neurons and their distal terminals. Front

Neuroanat 9:1–9.

Landoni LM, Myles JR, Wells TL, Mayer WP, Akay T (2019) Cholinergic modulation of motor neurons

through the C-boutons are necessary for the locomotor compensation for severe motor neuron loss

during amyotrophic lateral sclerosis disease progression. Behav Brain Res 369.

Leroy F, Lamotte d'Incamps B, Imhoff-Manuel RD, Zytnicki D (2014) Early intrinsic hyperexcitability does

not contribute to motoneuron degeneration in amyotrophic lateral sclerosis. Elife 3:1–25.

Malessa S, Leigh PN, Bertel O, Sluga E, Hornykiewicz O (1991) Amyotrophic lateral sclerosis: glutamate

dehydrogenase and transmitter amino acids in the spinal cord. J Neurol Neurosurg Psychiatry 54:984–

988.

Martin E, Cazenave W, Cattaert D, Branchereau P (2013) Embryonic alteration of motoneuronal

morphology induces hyperexcitability in the mouse model of amyotrophic lateral sclerosis. Neurobiol Dis

54:116–126.

Martin LJ, Liu Z, Chen K, Price AC, Pan Y, Swaby JA, Golden WC (2007) Motor Neuron Degeneration in

Amyotrophic Lateral Sclerosis Mutant Superoxide Dismutase-1 Transgenic Mice: Mechanisms of

Mitochondriopathy and Cell Death. J Comp Neurol 500:20–46.

Martinez-Silva M de L, Imhoff-Manuel RD, Sharma A, Heckman C, Shneider NA, Roselli F, Zytnicki D,

Manuel M (2018) Hypoexcitability precedes denervation in the large fast-contracting motor units in two

unrelated mouse models of ALS. Elife 7:1–26.

McGown A, McDearmid JR, Panagiotaki N, Tong H, Al Mashhadi S, Redhead N, Lyon AN, Beattie CE, Shaw

PJ, Ramesh TM (2013) Early interneuron dysfunction in ALS: Insights from a mutant sod1 zebrafish

model. Ann Neurol 73:246–258.

Medelin M, Rancic V, Cellot G, Laishram J, Veeraraghavan P, Rossi C, Muzio L, Sivilotti L, Ballerini L (2016)

Altered development in GABA co-release shapes glycinergic synaptic currents in cultured spinal slices of

the SOD1G93Amouse model of amyotrophic lateral sclerosis. J Physiol 594:3827–3840.

Milan L, Courtand G, Cardoit L, Masmejean F, Barrière G, Cazalets JR, Garret M, Bertrand SS (2015) Age-

related changes in pre- and postsynaptic partners of the cholinergic C-boutons in wild-type and

SOD1G93Alumbar motoneurons. PLoS One 10:1–21.

Niebroj-Dobosz I, Janik P (1999) Amino acids acting as transmitters in amyotrophic lateral sclerosis (ALS).

Acta Neurol Scand 100:6–11.

Ozdinler PH, Benn S, Yamamoto TH, Guzel M, Brown RH, Macklis JD (2011) Corticospinal Motor Neurons

and Related Subcerebral Projection Neurons Undergo Early and Specific Neurodegeneration in

hSOD1G93A Transgenic ALS Mice. J Neurosci 31:4166–4177 Available at:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4184-10.2011.

Pambo-Pambo A, Durand J, Gueritaud J-P (2009) Early excitability changes in lumbar motoneurons of

transgenic SOD1G85R and SOD1G(93A-Low) mice. J Neurophysiol 102:3627–3642 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/19828728.

Pieri M, Carunchio I, Curcio L, Mercuri NB, Zona C (2009) Increased persistent sodium current

determines cortical hyperexcitability in a genetic model of amyotrophic lateral sclerosis. Exp Neurol

215:368–379 Available at: http://dx.doi.org/10.1016/j.expneurol.2008.11.002.

Piotrkiewicz M, Hausmanowa-Petrusewicz I (2011) Motoneuron afterhyperpolarisation duration in

amyotrophic lateral sclerosis. J Physiol 589:2745–2754.

Plaitakis A, Constantakakis E (1993) Altered metabolism of excitatory amino acids, N-acetyl-aspartate

and N-acetyl-aspartylglutamate in amyotrophic lateral sclerosis. Brain Res Bull 30:381–386.

Pullen AH, Athanasiou D (2009) Increase in presynaptic territory of C-terminals on lumbar motoneurons

of G93A SOD1 mice during disease progression. Eur J Neurosci 29:551–561.

Pun S, Santos AF, Saxena S, Xu L, Caroni P (2006) Selective vulnerability and pruning of phasic

motoneuron axons in motoneuron disease alleviated by CNTF. Nat Neurosci 9:408–419.

Quinlan KA, Kajtaz E, Ciolino JD, Imhoff-Manuel RD, Tresch MC, Heckman CJ, Tysseling VM (2017)

Chronic electromyograms in treadmill running SOD1 mice reveal early changes in muscle activation. J

Physiol 595:5387–5400.

Quinlan KA, Lamano JB, Samuels J, Heckman CJ (2015) Comparison of dendritic calcium transients in

juvenile wild type and SOD1G93A mouse lumbar motoneurons. Front Cell Neurosci 9:1–13 Available at:

http://journal.frontiersin.org/article/10.3389/fncel.2015.00139/abstract.

Quinlan KA, Schuster JE, Fu R, Siddique T, Heckman CJ (2011) Altered postnatal maturation of electrical

properties in spinal motoneurons in a mouse model of amyotrophic lateral sclerosis. J Physiol 589:2245–

2260 Available at: http://doi.wiley.com/10.1113/jphysiol.2010.200659.

Raynor EM, Shefner JM (1994) Recurrent inhibition is decreased in patients with amyotrophic lateral

sclerosis. Neurology 44:2148–2153.

Rothstein JD, Martin LJ, Kuncl RW (1992) Decreased Glutamate Transport by the Brain and Spinal Cord in

Amyotrophic Lateral Sclerosis. N Engl J Med 326:1464–1468.

Saba L, Viscomi MT, Caioli S, Pignataro A, Bisicchia E, Pieri M, Molinari M, Ammassari-Teule M, Zona C

(2016) Altered Functionality, Morphology, and Vesicular Glutamate Transporter Expression of Cortical

Motor Neurons from a Presymptomatic Mouse Model of Amyotrophic Lateral Sclerosis. Cereb Cortex

26:1512–1528.

Saba L, Viscomi MT, Martini A, Caioli S, Mercuri NB, Guatteo E, Zona C (2019) Modified age-dependent

expression of NaV1.6 in an ALS model correlates with motor cortex excitability alterations. Neurobiol Dis

130:104532 Available at: https://doi.org/10.1016/j.nbd.2019.104532.

Sangari S, Iglesias C, El Mendili MM, Benali H, Pradat PF, Marchand-Pauvert V (2016) Impairment of

sensory-motor integration at spinal level in amyotrophic lateral sclerosis. Clin Neurophysiol 127:1968–

1977.

Saxena S, Roselli F, Singh K, Leptien K, Julien JP, Gros-Louis F, Caroni P (2013) Neuroprotection through

Excitability and mTOR Required in ALS Motoneurons to Delay Disease and Extend Survival. Neuron

80:80–96 Available at: http://dx.doi.org/10.1016/j.neuron.2013.07.027.

Shefner JM, Logigian EL (1998) The mixed nerve silent period in normal subjects and patients with

amyotrophic lateral sclerosis. Electromyogr Clin Neurophysiol 38:505–510.

Vinsant S, Mansfield C, Jimenez-Moreno R, Moore VDG, Yoshikawa M, Hampton TG, Prevette D, Caress

J, Oppenheim RW, Milligan C (2013a) Characterization of early pathogenesis in the SOD1G93A mouse

model of ALS: Part II, results and discussion. Brain Behav 3:431–457.

Vinsant S, Mansfield C, Jimenez-Moreno R, Moore VDG, Yoshikawa M, Hampton TG, Prevette D, Caress

J, Oppenheim RW, Milligan C (2013b) Characterization of early pathogenesis in the SOD1G93A mouse

model of ALS: Part I, background and methods. Brain Behav 3:335–350.

Wainger BJ, Kiskinis E, Mellin C, Wiskow O, Han SSW, Sandoe J, Perez NP, Williams LA, Lee S, Boulting G,

Berry JD, Brown RH, Cudkowicz ME, Bean BP, Eggan K, Woolf CJ (2014) Intrinsic membrane

hyperexcitability of amyotrophic lateral sclerosis patient-derived motor neurons. Cell Rep 7:1–11

Available at: http://dx.doi.org/10.1016/j.celrep.2014.03.019.

Whitehouse PJ, Wamsley JK, Zarbin MA, Price DL, Tourtellotte WW, Kuhar MJ (1983) Amyotrophic

lateral sclerosis: Alterations in neurotransmitter receptors. Ann Neurol 14:8–16.

Wootz H, Fitzsimons-Kantamneni E, Larhammar M, Rotterman TM, Enjin A, Patra K, André E, Van

Zundert B, Kullander K, Alvarez FJ (2013) Alterations in the motor neuron-renshaw cell circuit in the

Sod1G93Amouse model. J Comp Neurol 521:1449–1469.

Zeilhofer HU, Studler B, Arabadzisz D, Schweizer C, Ahmadi S, Layh B, Bösl MR, Fritschy JM (2005)

Glycinergic neurons expressing enhanced green fluorescent protein in bacterial artificial chromosome

transgenic mice. J Comp Neurol 482:123–141.