Title: Inhibitory interneurons show early dysfunction in a SOD1 mouse model of amyotrophic

lateral sclerosis

Running Title: Dysfunction of inhibitory interneurons in ALS

C. F. CAVARSAN[1,2], P. R. STEELE[1,2,3], L. T. GENRY[1,2,3], E.J. REEDICH[1,2], L. M. McCANE[3], K. J.

LaPRE[1,2], A. C. PURITZ[4], M. MANUEL[1,2], N. KATENKA[5], AND K. A. QUINLAN[1,2,4]

1 George and Anne Ryan Institute for Neuroscience, University of Rhode Island, Kingston, RI, 02881 USA

2 Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island,

Kingston, RI, 02881 USA

3 Interdisciplinary Neuroscience Program, University of Rhode Island, Kingston, RI, 02881 USA

4 Department of Physiology, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611 USA

5 Department of Computer Science and Statistics, University of Rhode Island, Kingston, RI, 02881 USA

Corresponding Author:

Katharina A. Quinlan (https://orcid.org/0000-0002-6177-4849)

kaquinlan@uri.edu

George and Anne Ryan Institute for Neuroscience

University of Rhode Island

130 Flagg Rd

Kingston RI 02881

Category according to the table of contents: Neuroscience

Key words: inhibition, glycine, Renshaw cells, amyotrophic lateral sclerosis, spinal cord, patch

clamp.

Key Points Summary:

- Spinal inhibitory interneurons could contribute to amyotrophic lateral sclerosis (ALS) pathology,

but their excitability has never been directly measured.

- We studied the excitability and morphology of glycinergic interneurons in early postnatal

transgenic mice (SOD1^{G93A}GlyT2eGFP).

- Interneurons were less excitable and had marginally smaller somas but larger primary neurites

in SOD1 mice.

- GlyT2 interneurons were analyzed according to their localization within the ventral spinal cord.

Interestingly, the greatest differences were observed in the most ventrally-located interneurons.

- We conclude that inhibitory interneurons show presymptomatic changes that may contribute to

excitatory / inhibitory imbalance in ALS.

Abstract: Few studies in amyotrophic lateral sclerosis (ALS) measure effects of the disease on inhibitory

interneurons synapsing onto motoneurons (MNs). However, inhibitory interneurons could contribute to

dysfunction, particularly if altered before MN neuropathology, and establish a long-term imbalance of

inhibition / excitation. We directly assessed excitability and morphology of glycinergic (GlyT2 expressing)

ventral lumbar interneurons from SOD1G93AGlyT2eGFP (SOD1) and wildtype GlyT2eGFP (WT) mice on

postnatal days 6-10. Patch clamp revealed dampened excitability in SOD1 interneurons, including

depolarized persistent inward currents (PICs), increased voltage and current threshold for firing action

potentials, along with a marginal decrease in afterhyperpolarization (AHP) duration. Primary neurites of

ventral SOD1 inhibitory interneurons were larger in volume and surface area than WT. GlyT2

interneurons were then divided into 3 subgroups based on location: (1) interneurons within 100 μm of

the ventral white matter, where Renshaw cells (RCs) are located, (2) interneurons interspersed with

MNs in lamina IX, and (3) interneurons in the intermediate ventral area including laminae VII and VIII.

Ventral interneurons in the RC area were the most profoundly affected, exhibiting more depolarized

PICs and larger primary neurites. Interneurons in lamina IX had depolarized PIC onset. In lamina VII-VIII,

interneurons were least affected. In summary, inhibitory interneurons show very early region-specific

perturbations poised to impact excitatory / inhibitory balance of MNs, modify motor output, and

provide early biomarkers of ALS. Therapeutics like riluzole which universally reduce CNS excitability

could exacerbate the inhibitory dysfunction described here.

Abstract Figure:

Abstract Figure: SOD1 glycinergic interneurons in the ventral horn show altered morphology and excitability, including depolarization of PICs, depolarized threshold, shorter AHPs, smaller somata and larger primary neurites. Ventrally located interneurons are the most prominently affected.

Introduction

Amyotrophic lateral sclerosis (ALS) is a rapidly evolving adult-onset neurological disease characterized by progressive loss of corticospinal neurons and spinal motoneurons (MNs). There has been considerable debate in the field over the role of hyperexcitability in neurodegenerative processes in ALS. While drug treatment for ALS based on nonspecific reduction of neuronal excitability (with riluzole, for example) (Bellingham, 2011), results in a modest increase in lifespan (Bensimon et al., 1994), similar treatment paradigms have been clinically disappointing. A more nuanced understanding of the excitability of vulnerable neurons could help to create a more targeted and effective approach for treatment of ALS. For example, if inhibitory pathways are failing in ALS and neuronal excitability is universally reduced with riluzole, this could further exacerbate inhibitory dysfunction by reducing activity not only in vulnerable MNs but also in inhibitory interneurons presynaptic to MNs. Thus, it is important to consider all aspects of neuronal excitability, including intrinsic excitability of vulnerable neurons, synaptic drive, and neuromodulation (Gunes et al., 2020). Intrinsic properties of MNs have been well studied but the same is not true for interneurons that are synaptically connected to them. In fact, no studies thus far have directly assessed activity of spinal premotor interneurons in an ALS model.

Despite evidence that ALS patients have disrupted inhibition at spinal levels (Raynor and Shefner, 1994; Shefner and Logigian, 1998; Sangari et al., 2016; Howells et al., 2020; Özyurt et al., 2020), much is still unknown concerning the involvement of inhibitory circuitry in ALS. Morphological alterations in inhibitory circuits have been demonstrated in animal models of ALS. These include degeneration of spinal interneurons, fewer neurons expressing markers of inhibitory neurotransmitters (including GlyT2) prior to MN loss (Martin et al., 2007; Hossaini et al., 2011), and loss of glycinergic boutons onto MNs prior to symptom onset (Chang and Martin, 2009a). Typically, fast MNs have greater numbers of inhibitory synaptic contacts, but these are largely lost in SOD1 mice beginning when motor unit atrophy is first observed (Pun et al., 2006; Hegedus et al., 2007; Allodi et al., 2021). Recurrent inhibitory circuits mediated by Renshaw cells (RCs) are impaired before symptom onset by both loss of MN collaterals which provide synaptic drive, and complex changes to RC-MN synaptic structures (Casas et al., 2013; Wootz et al., 2013). A few studies have suggested activity is decreased in inhibitory interneurons by indirect measurements. Quantification of synaptic inputs to MNs has shown that frequency of inhibitory postsynaptic potentials in spinal MNs is decreased embryonically in both SOD1 mouse and zebrafish models (McGown et al., 2013; Branchereau et al., 2019) and glycinergic inputs to MNs decay faster in

SOD1 MNs (Medelin et al., 2016). On a larger scale, blocking the activity of V1 inhibitory interneurons

during *in vivo* treadmill running was recently found to mimic the early locomotor deficits in SOD1 mice

(Allodi et al., 2021), suggesting that inhibitory interneurons could be inactive or under-active in ALS.

However, none of these studies directly examined electrophysiological activity in inhibitory

interneurons.

We hypothesized that inhibitory spinal interneurons could contribute to the pathogenesis of ALS

through a depression of MN inhibition. Decreased activity of inhibitory interneurons could result in

synaptically-driven hyperexcitability of MNs and other long-term changes in network function. In this

study, we examined electrical and morphological properties of glycinergic interneurons in the spinal

cord of SOD^{G93A} GlyT2-eGFP (SOD1) mice compared to GlyT2-eGFP (WT) mice using whole cell patch

clamp and three-dimensional reconstructions. We show here that significant dysfunction is present in

SOD1 glycinergic interneurons, which likely comprise several subclasses of inhibitory interneurons. In

general, SOD1 glycinergic interneurons are less excitable and have altered primary neurite morphology.

Impairment in glycinergic interneurons should be explored as both a mechanism of vulnerability of MNs

and a potential biomarker of early dysfunction of spinal circuits.

Materials and Methods

Ethical Approval

Experiments were performed in accordance with the United States National Institutes of Health Guide for

Care and Use of Laboratory Animals and Northwestern University's Animal Care and Use Committee.

Approval was obtained for all experiments performed in this study (IS00001228), and all experiments

conformed to the principles and regulations described in (Grundy, 2015). All efforts were made to

minimize animal suffering and to reduce the number of animals used.

Animals and Tissue harvest

Transgenic B6SJL mice overexpressing the human *SOD1^{G93A}* gene (strain 002726, The Jackson Laboratory,

Bar Harbor, ME, USA) and their wild type littermates (nontransgenic for the human *SOD1^{G93A}* gene) were

used. Transgenic animals were identified using standard PCR techniques by Transnetyx (Cordova, TN, USA)

and were bred with GlyT2-eGFP mice obtained from Ole Kiehn's lab at Karolinska Institute in Stockholm,

Sweden, with permission of Hans Zeilhofer (Zeilhofer et al., 2005), generating SOD1^{G93A} GlyT2-eGFP mice,

here called SOD1. Inhibitory glycinergic interneurons express the Na^+ and Cl^- coupled glycine transporter

2, or GlyT2, and GlyT2-eGFP expression is driven by the GlyT2 promotor. Specifically, GlyT2-eGFP females

were bred with SOD1[G93A] GlyT2-eGFP males, and progeny were used for experiments before genotyping

was performed. GlyT2-eGFP mice are referred to as wild type (WT), not carrying the SOD1[G93A] mutation.

All mice used in this study were housed at Northwestern's Center for Comparative medicine and had *ad*

libitum access to food and water. In total, 21 WT and 12 SOD1 juvenile mouse pups were used for the

following studies. Mice were deeply anesthetized with isoflurane (Henry Schein Animal Health, Dublin,

OH, USA) by inhalation until insensate to toe pinch, decapitated and eviscerated. The lumbar spinal cord

from L1 - L6 was removed and embedded in 2.5% w/v agar (No. A-7002, Sigma-Aldrich, St Louis, MO, USA).

The agar block was then superglued with Loctite 401 (Henkel Corporation, Rocky Hill, CN, USA) to a

stainless-steel slicing chuck and 350 μm transverse slices were made using the Leica 1000 vibratome (Leica

Microsystems, Buffalo Grove, IL, USA) as described previously (Quinlan et al., 2011). During both spinal

cord isolation and slicing, the spinal cord was immersed in 1–4 °C high osmolarity dissecting solution

containing (mM): sucrose 234.0, KCl 2.5, $CaCl_2 \cdot 2H_2O$ 0.1, $MgSO_4 \cdot 7H_2O$ 4.0, HEPES 15.0, glucose 11.0, and

Na_2PO_4 1.0. The pH was adjusted to 7.35 when bubbled with 95% O_2/5% CO_2 using 1 M KOH (Fluka

Analytical, Sigma-Aldrich). After cutting, the slices were incubated for >1 h at 30 °C in incubating solution

containing (mM): NaCl 126.0, KCl 2.5, $CaCl_2 \cdot 2H_2O$ 2.0, $MgCl_2 \cdot 6H_2O$ 2.0, $NaHCO_3$ 26.0, glucose 10.0, pH

7.4 when bubbled with 95% O_2/5% CO_2 (all reagents for solutions were purchased from Sigma-Aldrich).

Electrophysiology

Whole cell patch clamp was performed on interneurons from the lumbar segments using 2–4 MΩ glass

electrodes pulled from glass capillary tubes (Item #TW150F-4, World Precision Instruments, Sarasota, FL,

USA) with a Flaming-Brown P-97 (Sutter Instrument Company, Novato, CA, USA). Electrodes were

positioned using a Sutter Instrument MP-285 motorized micromanipulator (Sutter Instrument Company).

Whole-cell patch clamp measurements were performed at room temperature using the Multiclamp700B

amplifier (Molecular Devices, Burlingame, CA, USA) and Winfluor software (University of Strathclyde,

Glasgow, Scotland). Briefly, slices were perfused with a modified Ringer's solution containing (in mM): 111

NaCl, 3.09 KCl, 25.0 $NaHCO_3$, 1.10 KH_2PO_4, 1.26 $MgSO_4$, 2.52 $CaCl_2$, and 11.1 glucose. The solution was

oxygenated with 95% O_2/5% CO_2, and the perfusion rate was 2.5 – 3.0 ml/min. Patch electrodes contained

(in mM) 138 K-gluconate, 10 HEPES, 5 ATP-Mg, 0.3 GTP-Li and Texas Red dextran (150 μM, 3000 MW,

from Invitrogen, Life Technologies, Grand Island, NY, USA). In voltage-clamp mode, fast and slow

capacitance transients, as well as whole-cell capacitance, were compensated using the automatic

capacitance compensation on the Multiclamp. Whole cell capacitance was recorded from the Multiclamp.

Neuron Selection: Glycinergic interneurons were visually selected for recording based on 1) expression of

GFP, and 2) location in the ventral horn. Neurons that did not repetitively fire or did not maintain a resting

membrane potential below -35 mV were excluded from electrophysiological analysis. Based on these

criteria, 34/37 WT interneurons and 25/29 SOD1 interneurons were included in this analysis from mice

between postnatal day (P) 6 - 10.

Electrophysiological analysis: Holding potential was set at −90 mV, and neurons were subjected to slow,

depolarizing voltage ramps of 22.5 mV s^{-1}, bringing the cell to 0 mV in 4 s, and then back to the holding

potential in the following 4 s. In current clamp, neurons were subjected to depolarizing current ramps for

testing I-on (the current level at firing onset), I-off (the current level at cessation of firing), and the slope

of the frequency–current relationship. The difference between I-on and I-off is calculated for ΔI. Negative

current was often necessary to prevent action potential (AP) firing. Resting membrane potential was

recorded as the potential when 0 current was injected in voltage clamp. Persistent inward current (PIC)

parameters were measured after subtraction of leak current. PIC onset was defined as the voltage at

which the current began to deviate from the horizontal, leak-subtracted trace. PIC peak was the voltage

at which the PIC reached peak amplitude. The linear portion of leak current (usually between -90 and -75

mV) was used to calculate whole cell input resistance. The first action potential in the train evoked by a

depolarizing current ramp in current clamp mode was used to measure all parameters relating to action

potentials. Threshold voltage was defined as the voltage at which the slope exceeded 10 V/s. Threshold

for action potential firing was tested in two ways. The first was to use the voltage at firing onset from the

current ramps up to 130 pA/s (in current clamp). The second was the voltage at which a single action

potential could be evoked with a current step. Action potential overshoot is the voltage past 0 mV the

first spike (from a ramp) reaches. Duration of the action potential is measured at half of action potential

height (height is defined as overshoot – threshold voltage). Rates of rise and fall are defined as the peak

and the trough of the first derivative of the action potential profile. Instantaneous firing frequency range

max was the maximum firing rate that could be evoked from a neuron. It was measured from the first or

second firing frequencies on a large (>150 pA) depolarizing step. Steady state firing frequency range

minimum was the lowest firing frequency evoked from a small (~50 pA) depolarizing step. Hyperpolarizing

steps were used to measure I_H. The sag and rebound were measured from largest evoked response. Sag

potential was calculated as a % of the total amplitude of the hyperpolarization. If a spike was evoked on

rebound, the voltage threshold was used as amplitude of rebound. AHP duration was measured from a

single spike fired during a period in which the membrane potential was stable. Amplitude was measured

as the downward deflection from the membrane potential. Duration of the AHP was measured from the

falling phase of the spike at baseline potential to the time of recovery to baseline membrane potential.

Half amplitude duration was measured as the duration at half of the measured AHP amplitude from the

falling phase of the spike at baseline potential. Tau is measured as the rate of decay from the last third of

the AHP.

Regional classification: Regions were determined from spinal cord landmarks. In the case of patched

neurons, photos were taken of the position of the patch electrode within the spinal cord slices after

recording was complete. These photos were used to recreate a map of the position of the interneurons.

For GFP+ neurons that were not patched but were reconstructed, z projections were used to group cells

based on landmarks. The dorsal boundary of the ventral horn was identified by the central canal, the

dorsal edge of lateral motoneuron pools and/or 300µm distance from the ventral white matter.

Interneurons within the ventral-most region of the spinal cord (within 100 µm of the ventral white

matter) were grouped into the RC region described previously (Siembab et al., 2010). Neurons that were

located within and on the lateral edges of the motor pools (in between motoneurons and the white

matter) were classified as lamina IX interneurons. Since the location of the motor pools varied

throughout the lumbar enlargement, this classification was determined by examination of the z stacks

and photos. Neurons that were not found within lamina IX and were not within the ventral RC region

were grouped in the intermediate region of Lamina VII/VIII. All patched neurons were filled with Texas

Red dextran to allow their 3D reconstruction. Reconstructions were limited to the first dendritic node

(more details in following section). Unpatched GFP+ glycinergic interneurons that touched the boundary

lines between zones were excluded from analysis to prevent double counting. When the position of

GFP+ interneurons was not clear from the z stack, they were not included in the analysis.

Morphological analysis

Neuron morphology was assessed from two types of images: 1) 2-photon image stacks of live neurons

taken immediately after obtaining electrophysiological parameters, and 2) confocal image stacks taken

after fixation and tissue processing. Images collected using these two methods were analyzed separately

due to tissue shrinkage during the fixation process. Measurements from patched neurons included both

soma and neurites (soma volume, soma surface area; and average neurite length [per cell], total neurite

length [per cell], average and total neurite surface area and average and total neurite volume). Neurite

reconstructions were performed up to the first node for several reasons, including variable dye filling of

fine/distal processes and potential extension of neurites outside of the areas imaged. Neurites are

modeled as the somata but by treating each branch as a frustum. Measurements of GFP+ glycinergic

interneurons from fixed tissue samples included soma volume and soma surface area. Z-stacks acquired

by confocal microscopy were used to generate 3D soma reconstructions in Neurolucida 360 (MBF

Bioscience, Williston, VT, USA). Values shown throughout this study were calculated using Neurolucida's

branched structure analysis, specifically from the cell bodies summary page, including "enclosed

volume" (calculated as the shell made from the cell body contours), and "surface area" (the sum of each

section's perimeter of the reconstruction multiplied by the thickness).

Two-photon imaging: An Olympus BX-51WIF microscope fitted with an Olympus 40x/0.8NA water-

dipping objective lens was used. Two-photon excitation fluorescence microscopy was performed with a

galvanometer-based Coherent Chameleon Ultra II laser (Coherent, Santa Clara, CA, USA) tuned to 900

nm. A red Bio-Rad 2100MPD photomultiplier tube (Bio-Rad, Hercules, CA, USA) (570 – 650 nm) was used

to collect emission in the red channel. Z-stacks were obtained for each interneuron at 1024 x 1024 pixels

(308 x 308 μm) resolution and roughly 100 μm in depth (step size 1 μm). From these Z-stacks, Texas

Red® -filled neurons were three-dimensionally reconstructed, as described above, using Neurolucida

software.

Confocal imaging: After patch clamp electrophysiology was complete, spinal cord sections were fixed,

washed, and mounted with a coverslip for confocal microscopy. Z-stack photomicrographs (step size 1

μm) of glycinergic (GlyT2-eGFP+) interneurons in the lumbar ventral horn were imaged using a Nikon

Eclipse Ti2 inverted confocal microscope at 20x magnification. From these Z-stacks, GlyT2 positive

interneurons were three-dimensionally reconstructed using Neurolucida software as described above.

The total number of analyzed interneurons was over 2000, including over 1000 from 6 WT mice and

over 1000 from 6 SOD1 mice.

Statistical analysis

Each patched neuron is treated as an independent observation. The assumptions of homogeneity of variances and the normality of the distribution of values for each measured characteristic were evaluated with Levine's and Shapiro-Wilk tests, respectively. Group comparisons (WT vs SOD1) for patch clamp data were performed using one-way ANOVA on parameters that satisfied both conditions (homogeneity and normality of distribution) and using the Kruskal-Wallis test for those that did not. All results are presented as means +/- standard deviation (SD). Rather than basing our interpretation of results solely on *p* value, we have presented here the effect size of each measurement based on Hedges' *g* value (Hedges, 1981). Effect size is presented using Cumming plots (Cumming, 2012) generated using DABEST, as in (Huh et al., 2021) for all patched neurons. Effect sizes are presented along with 95% confidence intervals (CI) based on a bootstrap distribution of the mean, resampled 5000 times and bias corrected. To compare anatomical data from unpatched neurons we used a random intercept linear mixed-effects model to account for the random effects of the mouse and the fixed effect of genotype; this was performed due to the correlation between data points created by the large number of repeated measurements from individual mice. The linear mixed model was used to avoid errors of pseudoreplication (Lazic, 2010; Yu et al., 2022) and dependency of *p* value on sample size (Lin et al., 2013; Gómez-de-Mariscal et al., 2021).

Results

Morphology of glycinergic interneurons

Ventrally located glycinergic interneurons in the lumbar enlargement of WT and SOD1 mice at postnatal day (P) 6-10 were reconstructed from z-stack images obtained using confocal microscopy. Typical ventrolateral glycinergic neurons are shown in **Figure 1**. Complete reconstructions of soma morphology were performed on over 2000 GlyT2 interneurons (1050 interneurons from 6 WT mice and 1170 interneurons from 6 SOD1 mice). Overall, soma size was not different: results of a linear mixed model did not indicate significance in either the soma surface area or enclosed volume, as shown in **Table 1**, and as can be appreciated by the lack of difference in distribution between the two populations as shown in Fig 1B. Analyzing dendritic morphology was not possible in these images due to the high number of GFP+ neurons and processes but was performed on a smaller subset of neurons that were imaged live after patch clamp electrophysiology with a 2-photon microscope.

Figure 1: Comparing size of GlyT2 interneurons in the ventral horn. Representative photomicrographs of spinal cord showing GFP interneurons (green) in **A**. (**B**) Morphological analysis showed there was little difference in size between SOD1 glycinergic interneurons in ventral horn (N = 1170 neurons from 6 mice) and WT (N = 1050 neurons from 6 mice). Based on a linear mixed model differences are not significant. Scale bar in **A**: 100 μm applies to both images.

		Ventral horn GlyT2-GFP+ Interneurons (somata only)		
		WT (Mean ± SD)	SOD (Mean ± SD)	p
		N = 1050	N = 1170	
Soma	Surface Area (µm²)	450 (± 208)	390 (± 178)	0.387[LM]
	Volume (µm³)	509 (± 306)	414 (± 247)	0.335[LM]
		Patched GlyT2-GFP+ Interneurons			
		WT (Mean ± SD)	SOD (Mean ± SD)	p	g (95% CI low, high)
		N = 33	N = 24		
Soma	Surface Area (µm²)	638 (± 232)	653 (± 239)	0.728[K]	0.065 (-0.467, 0.621)
	Volume (µm³)	1472 (± 952)	1643 (± 951)	0.414[K]	0.177 (-0.350, 0.731)
Primary Neurites	Total Neurite Length (µm)	101 (± 100)	104 (± 49)	0.905	0.029 (-0.441, 0.649)
	Average Neurite length (µm)	29 (± 20)	31 (± 13)	0.663	0.109 (-0.387, 0.745)
	Total Surface Area (µm²)	513 (± 405)	634 (± 266)	0.032[K]	0.338 (-0.230, 0.978)
	Average Surface Area (µm²)	152 (± 81)	195 (± 81)	0.025[K]	0.523 (-0.035, 1.092)
	Total Volume (µm³)	265 (± 227)	382 (± 190)	0.009[K]	0.546 (-0.023, 1.202)
	Average Volume (µm³)	79 (± 54)	119 (± 61)	0.008[K]	0.679 (0.097, 1.245)

Table 1: Reconstruction data of GlyT2-GFP+ interneurons in WT and SOD1 animals.

N is the number of cells included in analysis; SD is standard deviation. Values of *p* are calculated from ANOVA except when marked with [K] indicating Kruskal-Wallis Test for nonparametric distributions or [LM] indicating linear mixed effects model. Values of *g* are determined using Hedge's statistic to calculate effect size from raw data and include the 95% confidence interval of the effect size

To study electrophysiological properties of GlyT2 interneurons, 59 glycinergic neurons (34 WT and 25 SOD1) were recorded using visually guided patch clamp with Texas Red dye in the electrode as shown in **Figure 2.** The location of patched neurons was distributed throughout the ventral horn as shown in Fig 2B. In this sample, we were able to analyze morphology of neurites up to the first node (branching point) and found that neurites were larger in patched SOD1 GlyT2 interneurons, including a modest increase in both neurite surface area and volume (see Table 1 for complete results). The number of primary neurites was not different in SOD1 vs WT interneurons. Please note that dimensions of these neurons cannot be directly compared to the previous section since these neurons were imaged in living tissue while the previous analysis was performed in fixed tissue, which shrinks.

Figure 2: GlyT2 interneurons from WT and SOD1 mice were patched throughout the ventral horn. Images of typical interneurons, filled with Texas Red as they were recorded in **A**, and locations of all patched interneurons in **B** (WT = blue circles, SOD1 = red circles). Consistent with previous results, soma volume (**C**) and soma surface area (**D**) were unchanged in patched neurons, while total surface area and total volume of primary neurites (**E** and **F**) were modestly larger in SOD1 interneurons than in WT. Effect sizes are shown to the right of each swarm plot, with distribution of the 95% CI of the effect sizes in grey to the right of the mean difference (SOD1 − WT).

Electrophysiology of glycinergic interneurons

Whole cell patch clamp revealed very different intrinsic properties of ventral glycinergic interneurons.

Recording was performed on 59 ventral interneurons from throughout the ventral horn in transverse

lumbar spinal cord slices from P6-10 mice. SOD1 inhibitory interneurons (n = 25 from 10 mice) were

found to have diminished intrinsic excitability compared to WT interneurons (n = 34 from 15 mice).

Measurements of intrinsic excitability included depolarized onset and peak voltage of PICs, resulting in a

small change in threshold for action potential firing, and the current at firing onset (I-ON) and offset (I-

OFF) as shown in **Figure 3**. In voltage clamp, slowly depolarizing voltage ramps were used to measure

PICs (typical leak subtracted traces shown in Fig 3A). Voltage sensitivity is determined by measuring

voltage at PIC onset and peak. Both PIC onset and peak were significantly depolarized in SOD1

interneurons, while amplitude of the PIC was unchanged. Depolarizing current ramps were used to

measure the input-output relationship of the neurons in current clamp. Inhibitory interneurons from

SOD1 mice were found to have significantly higher threshold voltage than WT interneurons (Fig 3E-F).

This shift is likely driven by changes in PICs. The current at firing onset, or I-ON, and firing offset (I-OFF)

was similarly larger in SOD1 interneurons. The AHP appeared to be shorter duration in SOD1

interneurons, however corresponding changes in the decay rate (tau) were not present. The total AHP

duration and the duration at half amplitude were slightly shorter in SOD1 neurons, as shown in Fig 3J

and **Table 2**, though amplitude of the AHP was unchanged. While changes in the PIC and threshold

indicate less excitability in SOD1 interneurons, the shortened AHP suggests an increased ability to fire at

higher rates in SOD1 interneurons, a property that is associated with increased excitability. However, no

changes in firing rates were detected in SOD1 interneurons. All other properties were found to be

similar in SOD1 and WT glycinergic interneurons, including maximum firing rates, sag/rebound currents

(I_H), action potential parameters and membrane properties (see Table 2 for all electrical properties). See

discussion for further interpretation of these findings.

Figure 3: Electrophysiology of SOD1 glycinergic interneurons. The most prominent difference in SOD1 interneurons was the shift in voltage dependence of PICs. (**A**) Representative, leak-subtracted current-voltage relationship of PICs from WT (P8) and SOD1 (P6) interneurons. (**B-D**) Mean onset, peak and amplitude of PICs in WT (blue symbols on left) and SOD1 interneurons (red symbols on right). (**E-F**) Threshold for action potentials evoked with current ramps was higher in SOD1 interneurons (compare dotted lines indicating voltage threshold). Starting potential for both interneurons was -65 mV. The current at firing onset and offset (I-ON and I-OFF, respectively) was also increased as shown in **G-H.** (**I**) The duration of the AHP in SOD1 interneurons appeared shorter, as shown in a representative P8 WT and P7 SOD1 interneuron and cumulative data for total duration of AHP in **J.** However, neither the AHP amplitude nor the AHP decay time constant, tau, were significantly altered in SOD1 glycinergic interneurons, as shown in **K-L.** Vertical scale bars in E: top = 50 pA, and bottom = 20 mV. Horizontal scale bar in **E** = 0.5s and all scale bars apply to both left and right panels. Vertical scale bar in H = 10 mV (APs were truncated from image), and horizontal scale bar = 50 ms.

Parameters	Groups			Statistics
	WT (Mean ± SD)	SOD (Mean ± SD)	*p*	*g* (95% CI low, high)
	N = 34	*N* = 25		
Resting membrane potential (mV)	-46.7 ± 6.4	-45.7 ± 6.9	0.524[K]	0.155 (-0.385, 0.705)
Capacitance (pF)	66 ± 42	55 ± 27	0.529[K]	-0.289 (-0.725, 0.210)
Input resistance (MΩ)	205 ± 104	220 ± 131	0.794[K]	0.127 (-0.398, 0.657)
PIC properties				
PIC onset (mV)	-50.5 ± 6.0	-46.2 ± 5.0	0.004	0.765 (0.254, 1.217)
PIC peak (mV)	-35.2 ± 4.9	-31.2 ± 4.6	0.003	0.820 (0.299, 1.285)
PIC amplitude (pA)	100 ± 73	97 ± 62	0.890[K]	-0.037 (-0.520, 0.494)
Normalized PIC (pA/pF)	1.79 ± 1.4	1.92 ± 1.1	0.226[K]	0.100 (-0.354, 0.666)
Firing properties				
Inst firing freq range max (Hz)	112 ± 44	113 ± 42	0.961	0.013 (-0.493, 0.557)
Steady state firing freq range min (Hz)	6 ± 2.4	6 ± 3.2	0.453[K]	0.277 (-0.282, 0.831)
I-ON (pA)	-51 ± 111	-14 ± 53	0.093[K]	0.392 (-0.113, 0.672)
I-OFF (pA)	-5 ± 66	36 ± 69	0.038[K]	0.286 (-0.324, 0.852)
Δ I (pA)	47 ± 59	50 ± 52	0.530[K]	-0.140 (-0.555, 0.453)
AP properties				
AP overshoot past 0 (mV)	19.9 ± 9.8	21.4 ± 7.9	0.529	0.156 (-0.346, 0.670)
AP duration at half peak (ms)	1.54 ± 0.54	1.48 ± 0.40	0.944[K]	-0.118 (-0.559, 0.425)
AP rate of rise (V/s)	87 ± 29	83 ± 13	0.446[K]	0.172 (-0.616, 0.328)
AP rate of fall (V/s)	45 ± 17	43 ± 11	0.995[K]	0.113 (-0.398, 0.593)
Threshold on ramp (mV)	-40.6 ± 4.4	-37.9 ± 5.7	0.076[K]	0.542 (-0.011, 1.050)
Threshold on step (mV)	-47.3 ± 4.4	-45.1 ± 6.2	0.199[K]	0.413 (-0.191, 0.879)
AHP properties				
AHP amplitude (mV)	11.2 ± 3.4	11.0 ± 3.9	0.704[K]	-0.047 (-0.565, 0.480)
AHP duration (s)	0.37 ± 0.21	0.29 ± 0.19	0.207[K]	-0.353 (-0.867, 0.171)
AHP duration at half amplitude (s)	0.12 ± 0.06	0.10 ± 0.06	0.045[K]	-0.370 (-0.873, 0.221)
AHP tau (s^{-1})	0.117 ± 0.115	0.105 ± 0.083	0.980[K]	-0.121 (-0.552, 0.426)
I$_H$ properties				
Hyperpolarizing Sag (%)	32 ± 18	32 ± 21	0.537[K]	-0.011 (-0.563, 0.556)
Depolarizing Rebound (%)	17 ± 15	19 ± 24	0.536[K]	0.086 (-0.454, 0.698)

Table 2: Electrophysiological properties of GlyT2-GFP+ interneurons in WT and SOD1 animals.
N is the number of cells included in analysis.
[K] Indicates Kruskal-Wallis Test for nonparametric distributions; all other *p* values were calculated using ANOVA
analysis.
Values of *g* are determined using Hedge's statistic to calculate effect size and include the 95% confidence interval
of the effect size

Regional analysis of glycinergic interneurons

Since ventral glycinergic interneurons are not a homogenous population, we tested for regional
differences. Interneurons were divided into three groups based on location: 1) the "ventral RC region":
within 100 μm of the ventral white matter where Renshaw cells are located, 2) lamina IX, and 3) the
"intermediate region": ventral horn region including lamina VII and VIII, excluding the previous two
regions. These regions are represented in **Figure 4**. The first region corresponds to the location of RCs
(but could contain other types of glycinergic interneurons). The second region contains interneurons
that are interspersed with the MNs within the motor pools, likely to be composed of premotor
interneurons (more on this in the discussion). The third region contains interneurons that are a mix of
premotor interneurons and other classes of ventral interneurons. There was overlap of the first two
regions, so interneurons within the ventral most 100 μm that also were located within motor pools were
included in both the ventral RC region and lamina IX, as shown in open circles in Figure 4.

Figure 4: Regional divisions of interneurons. (**A**) Three regions were used for analysis of regional differences in glycinergic interneurons. (**B**) Map of the patched interneurons coded for regions. Ventral region with RC cells (green circles) overlapped with lamina IX (hollow circles). The remaining interneurons that were not within the RC region or lamina IX were grouped into an intermediate region of lamina VII and VIII (red triangles). Scale bar in **A** = 100 μm.

GlyT2 interneurons in Ventral RC region

Within the ventral region where RCs are located, we found that glycinergic SOD1 interneurons there showed the most significant changes in both electrophysiology and morphology. In patch clamped interneurons from this region (N = 8 WT from 7 mice and 13 SOD1 from 9 mice), there was a significant depolarization in PIC onset and peak in glycinergic SOD1 interneurons compared to WT interneurons in the same location. Additionally, primary neurites were found to be longer and larger in volume and surface area in SOD1 interneurons in the ventral-most 100 μm. Unpatched GlyT2+ SOD1 interneurons in this region (188 WT interneurons from 6 WT mice and 225 SOD1 interneurons from 6 SOD1 mice), analyzed from fixed spinal cords were smaller than WT glycinergic interneurons. These changes (summarized in **Figure 5** and **Table 3**) show that glycinergic neurons in the region where RCs are located are very affected by the SOD1 mutation. Primary neurites, soma volume, PIC onset and PIC peak were all as affected or more affected in the RC region than the larger group of ventral horn interneurons (for example compare primary neurite measurements in Fig 2E-F and Fig 5G-H).

GlyT2 interneurons in Lamina IX

The SOD1 interneurons located in lamina IX, intermixed with MNs were modestly altered in excitability and morphology. As shown in **Figure 6** and **Table 4**, PIC onset was depolarized in SOD1 interneurons in this region, compared to interneurons in this region from WT mice (n = 14 WT interneurons from 8 mice and 11 SOD1 interneurons from 9 mice). Longer total length of primary neurites was also found in patched interneurons in this region. Within the larger number of unpatched interneurons (310 WT interneurons from 6 WT mice and 258 SOD1 interneurons from 6 SOD1 mice), there was no significant change in soma size.

GlyT2 interneurons in the Intermediate region

The third regional group of glycinergic interneurons was defined as the intermediate region. These interneurons were located in laminae VII and VIII, excluding the ventral 100 μm and lamina IX. Within this area, 16 WT and 8 SOD1 interneurons were patched from 11 WT and 5 SOD1 mice respectively, and 655 unpatched interneurons were analyzed from 6 WT mice and 812 unpatched SOD1 interneurons from 6 SOD1 mice. As shown in **Figure 7** and **Table 5**, the PIC peak was the only parameter that was altered in SOD1 neurons in this region. Overall, the GlyT2 interneurons located here were the least affected by the SOD1 mutation.

Figure 5: SOD1 GlyT2 interneurons in the ventral RC region were altered in electrophysiology and morphology. Both onset (**A**) and peak (**B**) of the PIC were depolarized. Threshold voltage and current at firing onset and offset were more variable as shown by the effect sizes (**C-E**). Soma volume of patched neurons was not significantly altered but neurites were larger in total volume and total length (**F-H**) in SOD1 interneurons. Larger numbers of unpatched neurons were analyzed for soma volume (**I**) and surface area (**J**) in the RC region. A linear mixed model (including inter-mouse variability) revealed that soma volume was significantly smaller in SOD1 mice in this region.

	WT (Mean ± SD)	SOD (Mean ± SD)	p	g (95% CI low, high)
	N = 8	N = 13		
PIC Onset (mV)	-51.3 (± 5.7)	-45.6 (± 5.5)	0.036	0.971 (-0.058, 1.768)
PIC Peak (mV)	-36.8 (± 7.5)	-30.8 (± 4.7)	0.034	0.989 (0.026, 1.919)
PIC Amplitude (pA)	77 (± 44)	115 (± 71)	0.218[K]	0.579 (-0.299, 1.210)
I-ON (pA)	-66 (± 110)	-18 (± 51)	0.096[K]	0.582 (-0.276, 1.398)
I-OFF (pA)	-32 (± 58)	7 (± 54)	0.060[K]	0.671 (-0.293, 1.467)
Δ I (pA)	36 (± 55)	25 (± 25)	0.717[K]	-0.280 (-1.354, 0.679)
Threshold on ramp (mV)	-42.4 (± 4.0)	-38.0 (± 6.1)	0.070[K]	0.775 (-0.035, 1.393)
Threshold on step (mV)	-47.0 (± 3.1)	-44.1 (± 5.6)	0.261[K]	0.580 (-0.396, 1.150)
AHP amplitude (mV)	11.6 (± 3.9)	11.4 (± 3.5)	0.921	-0.044 (-0.958, 0.904)
AHP duration (s)	0.39 (± 0.19)	0.36 (± 0.18)	0.711	-0.162 (-1.163, 0.676)
AHP tau (s^{-1})	0.117 (± 0.14)	0.109 (± 0.07)	0.346[K]	-0.078 (-1.329, 0.935)
Morphology of Patched interneurons in RC region				
	N = 8	N = 13		
Soma				
Soma Surface Area (μm^2)	589 (± 167)	700 (± 285)	0.334	0.428 (-0.402, 1.180)
Soma Volume (μm^3)	1326 (± 553)	1841 (± 1150)	0.253	0.508 (-0.294, 1.138)
Primary Neurites				
Total Neurite Length (μm)	76 (± 24)	118 (± 50)	0.044	0.931 (0.079, 1.623)
Average Neurite Length (μm)	26 (± 8)	35 (± 14)	0.161[K]	0.710 (-0.082, 1.456)
Total Neurite Surface Area (μm^2)	408 (± 162)	681 (± 237)	0.010[K]	1.234 (0.182, 2.320)
Average Neurite Surface Area (μm^2)	137 (± 41)	206 (± 77)	0.025[K]	1.002 (0.199, 1.708)
Total Neurite Volume (μm^3)	196 (± 110)	391 (± 179)	0.013[K]	1.191 (0.279, 2.049)
Average Neurite Volume (μm^3)	65 (± 27)	120 (± 59)	0.016[K]	1.060 (0.320, 1.753)
Morphology of Unpatched interneurons in RC region				
	N = 188	N = 225		
Soma Surface Area (μm^2)	464 (± 180)	362 (± 168)	0.106[LM]	
Soma Volume (μm^3)	515 (± 247)	358 (± 215)	0.040[LM]	

381 **Table 3**: Electrophysiological properties of WT and SOD1 GlyT2-GFP+ interneurons in ventral RC region. [K] Indicates
382 Kruskal-Wallis Test for nonparametric distributions; [LM] Indicates linear mixed effects model; all others were ANOVA
383 analysis. Values of *g* are determined using Hedge's statistic to calculate effect size and include the 95% confidence
384 interval of the effect size

Figure 6: Modest changes in electrophysiology were present in glycinergic SOD1 interneurons in lamina IX. PIC onset (**A**) was depolarized in SOD1 interneurons without reliable changes in PIC peak (**B**), voltage threshold, I-ON or I-OFF (**C-E**). Soma (**F**) and primary neurites (**G-H**) did not show consistent changes. Unpatched glycinergic neurons were not significantly altered in soma enclosed volume (**I**) and surface area (**J**) in SOD1 interneurons compared to WT in Lamina IX.

	WT (Mean ± SD)	SOD (Mean ± SD)	*p*	*g* (95% CI low, high)
	N = 14	N = 11		
PIC Onset (mV)	-50.5 (± 6.7)	-44.0 (± 5.0)	0.013	1.046 (0.340, 1.652)
PIC Peak (mV)	-33.1 (± 3.9)	-31.1 (± 4.5)	0.248	0.462 (-0.349, 1.165)
PIC Amplitude (pA)	116 (± 81)	98 (± 76)	0.412[K]	-0.218 (-0.944, 0.655)
I-ON (pA)	-69 (± 143)	-11 (± 64)	0.155[K]	0.485 (-0.439, 0.928)
I-OFF (pA)	-23 (± 72)	22 (± 73)	0.080[K]	0.602 (-0.401, 1.207)
Δ I (pA)	48 (± 76)	33 (± 30)	0.702[K]	-0.226 (-0.564, 1.021)
Threshold on ramp (mV)	-40.7 (± 4.9)	-37.8 (± 6.8)	0.244	0.473 (-0.370, 1.209)
Threshold on step (mV)	-47.4 (± 4.2)	-43.9 (± 7.4)	0.112[K]	0.595 (-0.437, 1.377)
AHP amplitude (mV)	11.4 (± 3.9)	10.2 (± 3.5)	0.421	-0.319 (-1.09, 0.514)
AHP duration (s)	0.41 (± 0.24)	0.25 (± 0.16)	0.062	-0.764 (-1.509, 0.053)
AHP tau (s^{-1})	0.129 (± 0.118)	0.076 (± 0.067)	0.274[K]	-0.511 (-1.088, 0.390)
Morphology of Patched interneurons in lamina IX				
	N = 14	N = 10		
Soma				
Soma Surface Area (µm²)	703 (± 242)	734 (± 284)	0.771	0.118 (-0.740, 0.959)
Soma Volume (µm³)	1732 (± 1080)	1989 (± 1149)	0.412[K]	0.224 (-0.620, 1.076)
Primary Neurites				
Total Neurite Length (µm)	101 (± 130)	129 (± 60)	0.046[K]	0.258 (-0.610, 1.855)
Average Neurite Length (µm)	29 (± 24)	37 (± 16)	0.069[K]	0.335 (-0.544, 1.442)
Total Neurite Surface Area (µm²)	545 (± 524)	721 (± 293)	0.089[K]	0.384 (-0.522, 1.722)
Average Neurite Surface Area (µm²)	164 (± 99)	208 (± 80)	0.128[K]	0.466 (-0435, 1.381)
Total Neurite Volume (µm³)	309 (± 283)	388 (± 161)	0.114[K]	0.316 (-0.498, 1.474)
Average Neurite Volume (µm³)	95 (± 65)	113 (± 46)	0.178[K]	0.307 (-0.496, 1.233)
Morphology of Unpatched interneurons in lamina IX				
	N = 310	N = 258		
Soma Surface Area (µm²)	459 (± 164)	372 (± 169)	0.069[LM]	
Soma Volume (µm³)	525 (± 257)	392 (± 248)	0.067[LM]	

Table 4: Electrophysiological properties of WT and SOD1 GlyT2-GFP+ interneurons in lamina IX.

[K] Indicates Kruskal-Wallis Test for nonparametric distributions; [LM] Indicates linear mixed effects model; all others were ANOVA analysis. Values of *g* are determined using Hedge's statistic to calculate effect size and include the 95% confidence interval of the effect size

Figure 7: SOD1 interneurons in the intermediate regions of laminae VII and VIII were largely unaffected. PIC onset (**A**) was not significantly different, though PIC peak (**B**) was marginally more depolarized in SOD1 neurons. Voltage and current threshold (**C-E**) were not significantly affected. Morphology of soma (**F**) and neurites (**G-H**) in patched SOD1 interneurons were unchanged, and larger numbers of unpatched neurons in this region (**I-J**) were not significantly smaller.

Parameters	Groups			
	WT (Mean ± SD)	SOD (Mean ± SD)	*p*	*g* (95% CI low, high)
	N = 16	N = 8		
PIC Onset (mV)	-49 (± 5)	-47 (± 4)	0.435	0.332 (-0.451, 1.240)
PIC Peak (mV)	-36 (± 4)	-32 (± 5)	0.046	0.884 (-0.210, 1.796)
PIC Amplitude (pA)	92 (± 67)	77 (± 50)	0.881[K]	-0.235 (-0.914, 0.667)
I-ON (pA)	-18 (± 45)	-16 (± 54)	0.776[K]	0.035 (-1.219, 0.762)
I-OFF (pA)	21 (± 47)	74 (± 76)	0.101[K]	0.877 (-0.038, 1.904)
Δ I (pA)	39 (± 35)	90 (± 69)	0.076[K]	1.006 (-0.077, 2.145)
Threshold on ramp (mV)	-38.9 (± 3.1)	-37.0 (± 6.3)	0.525[K]	0.439 (-0.701, 1.495)
Threshold on step (mV)	-46.7 (± 4.7)	-46.3 (± 6.4)	0.490[K]	0.080 (-0.850, 1.319)
AHP amplitude (mV)	11.3 (± 3.2)	10.5 (± 5.3)	0.658	-0.198 (-1.375, 0.853)
AHP duration (s)	0.34 (± 0.21)	0.25 (± 0.23)	0.341[K]	-0.401 (-1.189, 0.843)
AHP tau (s^{-1})	0.119 (± 0.127)	0.127 (± 0.125)	1.000[K]	0.062 (-0.657, 1.381)

Morphology of Patched interneurons in laminae VII and VIII.				
	N = 15	N = 8		
Soma				
Soma Surface Area (μm^2)	574 (± 244)	559 (± 160)	0.699[K]	-0.066 (-0.733, 0.847)
Soma Volume (μm^3)	1232 (± 919)	1228 (± 570)	0.776[K]	0.005 (-0.645, 0.956)
Primary Neurites				
Total Neurite Length (μm)	110 (± 80)	77 (± 33)	0.220[K]	-0.466 (-0.888, 0.435)
Average Neurite Length (μm)	30 (± 18)	24 (± 8)	0.606[K]	-0.352 (-0.813, 0.700)
Total Neurite Surface Area (μm^2)	506 (± 321)	514 (± 265)	0.699[K]	0.026 (-0.665, 0.994)
Average Neurite Surface Area (μm^2)	139 (± 72)	168 (± 85)	0.366[K]	0.365 (-0.419, 1.393)
Total Neurite Volume (μm^3)	234 (± 189)	330 (± 210)	0.272[K]	0.474 (-0.394, 1.45)
Average Neurite Volume (μm^3)	64 (± 46)	109 (± 70)	0.071[K]	0.782 (-0.150, 1.730)

Morphology of Unpatched interneurons in laminae VII and VIII.				
	N = 655	N = 812		
Soma Surface Area (μm^2)	448 (± 225)	394 (± 178)	0.495[LM]	
Soma Volume (μm^3)	510 (± 329)	425 (± 253)	0.455[LM]	

Table 5: Electrophysiological properties of WT and SOD1 GlyT2-GFP+ interneurons in Intermediate Region of
lamina VII and VIII.

[K] Indicates Kruskal-Wallis Test for nonparametric distributions; [LM] Indicates linear mixed effects model; all others
were ANOVA analysis. Values of *g* are determined using Hedge's statistic to calculate effect size and include the
95% confidence interval of the effect size

Taken together, these results show that glycinergic interneurons are affected by the SOD1 mutation,

particularly those in the ventral-most region of the spinal cord. These neurons are altered very early in

life in both electrophysiology and morphology.

Discussion

414
This study shows for the first time that glycinergic interneurons are depressed in excitability in the SOD1

mouse model and may be contributing to ALS pathology. Alterations in both morphology and excitability

of glycinergic interneurons occur at a very early, presymptomatic stage, and are particularly prominent

in those interneurons located within 100 µm of the ventral white matter. We speculate that these shifts

in interneuron excitability and morphology could have functional consequences that alter inhibition of

MNs, modify motor output, and could even provide an early biomarker of ALS.

Intrinsic and synaptic hyperexcitability

A core issue this study brings to light is potential long-term disruption to the balance of excitatory and

inhibitory synaptic transmission in ALS. Spinal cord and cortex both show an imbalance between

excitatory and inhibitory synaptic transmission in SOD^{G93A} mice (Avossa et al., 2006; Saba et al., 2016).

Clinically, ALS patients show evidence of altered synaptic activity as well, including a reduction in

inhibition. A reduction of glycinergic receptor binding in the anterior gray matter of the spinal cord

(Hayashi et al., 1981; Whitehouse et al., 1983) has been demonstrated along with abnormal glycine and

gamma-aminobutyric acid (GABA) levels in blood serum (Malessa et al., 1991; Niebroj-Dobosz and Janik,

1999). Electrophysiology also suggests disruption in spinal inhibitory circuits in ALS patients (Raynor and

Shefner, 1994; Shefner and Logigian, 1998; Sangari et al., 2016; Özyurt et al., 2020). An interesting

question is whether the neurons are altered in excitability only within the motor circuit (i.e.,

corticospinal neurons and the synaptically-connected spinal inhibitory and cholinergic interneurons, and

MNs), or if disturbances are more widespread throughout the nervous system. In the spinal cord,

interneurons that serve as conduits between vulnerable corticospinal and spinal MNs include cholinergic

interneurons, Ia inhibitory interneurons and RCs. There is abundant evidence of disruption in these

cholinergic, GABAergic and glycinergic synapses from animal models (Martin et al., 2007; Chang and

Martin, 2009b; Pullen and Athanasiou, 2009; Hossaini et al., 2011; Herron and Miles, 2012; Casas et al.,

2013; Saxena et al., 2013; Wootz et al., 2013; Milan et al., 2015; Dukkipati et al., 2016; Medelin et al.,

2016). Similarly, in the cortex, neurons that are not themselves vulnerable to neurodegeneration in ALS

also exhibit altered excitability and morphology in a parallel time course to the vulnerable corticospinal

neurons (Clark et al., 2017, 2018; Kim et al., 2017). The changes in excitability of other neuronal

populations must contribute to the imbalance of both excitatory and inhibitory neurotransmission that

has been clearly demonstrated in ALS patients and animal models.

A case for hyperexcitability and glutamate excitotoxicity in ALS pathogenesis has been more

controversial. Altered intrinsic excitability has been consistently reported in both corticospinal

projection neurons and spinal MNs, but these perturbations appear to fluctuate based on age/disease

progression. Corticospinal projection neurons have increased firing and other signs of hyperexcitability

during the first week of postnatal development (Pieri et al., 2009; Saba et al., 2016, 2019; Kim et al.,

2017), but excitability is normal in young and adult pre-symptomatic ages (Kim et al., 2017; Saba et al.,

2019), and may become hypoexcitable at the age of symptom onset (Saba et al., 2019). Similarly, spinal

MNs show intrinsic hyperexcitability at very early (embryonic) stages (Kuo et al., 2004, 2005; Martin et

al., 2013). At postnatal ages (similar to the inhibitory interneurons in this study), spinal MNs do *not* have

an altered threshold, rheobase or frequency-current relationship, though other abnormalities are

present in PICs, action potential duration, AHP and dendritic Ca^{2+} entry (Pambo-Pambo et al., 2009;

Quinlan et al., 2011, 2015; Leroy et al., 2014). Thus, at the age studied here, inhibitory interneurons are

affected differently than the vulnerable spinal MNs. At symptom onset, MNs may finally succumb to

hypoexcitability, though not all studies agree on this point (Delestrée et al., 2014; Martinez-Silva et al.,

2018; Jensen et al., 2020). Notably, MNs derived from ALS patients' induced pluripotent stem cells also

show initial hyperexcitability followed by hypoexcitability (Wainger et al., 2014; Devlin et al., 2015), and

increasing excitability of MNs enhanced neuroprotection rather than neurodegeneration (Saxena et al.,

2013). A unifying theme throughout the findings is that vulnerable MNs show excessive homeostatic

gain in response to perturbation (Kuo et al., 2020), and not only excitability but other cellular properties

fluctuate over the lifespan of the animal (Irvin et al., 2015). In light of that, whether inhibitory

interneurons are less excitable at other ages should be explored in future studies to fully characterize

their contribution to neurodegenerative processes.

Changes in neuron morphology

In ALS models, altered morphology in spinal MNs occurs so early that it could be viewed as a deficit in

normal development. Embryonically, SOD1 MNs have shorter projections but this is reversed during

postnatal development. At 1-2 weeks of age, SOD1 MNs begin to show more dendritic branching and

larger soma sizes than WT MNs (Amendola and Durand, 2008; Filipchuk and Durand, 2012; Martin et al.,

2013). Interestingly, this could be caused by lack of inhibitory input to MNs (Fogarty et al., 2016). SOD1

spinal MNs remain larger than WT MNs through adulthood (Dukkipati et al., 2018). In contrast to the

spinal MNs at this age, we found that SOD1 inhibitory interneurons in the ventral-most region had

smaller soma size and yet had expanded surface area and volume of primary neurites compared to WT

glycinergic interneurons. This is reminiscent of changes in corticospinal neurons: presymptomatic

corticospinal neurons also have increased arborization but with smaller soma diameters (Ozdinler et al.,

2011; Saba et al., 2016). In both sporadic and familial ALS patients, postmortem tissue shows

degeneration of apical neurites in corticospinal neurons and smaller soma size (Genç et al., 2017). In

SOD1 spinal interneurons in the present study, increased neurite surface area and volume of primary

neurites were most prominent in the ventral interneurons in the RC region. This region also showed

increased total neurite length and based on reconstructed, unpatched neurons, smaller soma volume.

Altered neuronal size / arborization could alter the impact of other stressors (e.g., proteostasis,

metabolic deficits, intracellular transport).

After spike afterhyperpolarization

The AHP can contribute to a neuron's firing rate such that a neuron with a large, long-lasting AHP will

typically fire at a slower rate (typical of slow/type I MNs) than a neuron with small, fast-decaying AHP

(typical of fast/type II MNs). In ALS patients, the AHP is shortened in MNs of patients with little force

deficits and later elongated in patients with large force deficits (Piotrkiewicz and Hausmanowa-

Petrusewicz, 2011), which could reflect early changes in physiology of vulnerable fast MNs and later, the

remaining, less-vulnerable population of slow MNs. Similarly, SOD1 mouse MNs show shorter AHPs

early, and longer AHPs around symptom onset (Quinlan et al., 2011; Jensen et al., 2020). Here we show

that AHP is modestly shorter in duration in glycinergic SOD1 interneurons very early in life. In the

regional analysis, AHP parameters were not found to be significantly altered in any of the 3 regions, but

it is possible this was due to a loss in statistical power in the smaller groups. We did not find

concomitant changes in minimum or maximum firing rates in GlyT2 interneurons, so the physiological

importance of the altered AHP duration in interneurons is unclear.

Inhibitory interneurons: Subtype specific effects

The interneurons in this study were heterogenous, and based on their location and electrophysiology,

they likely were composed of V1 and V2b interneurons including commissural interneurons (V2b

interneurons), RCs, and Ia inhibitory interneurons (both V1 interneurons), and perhaps inhibitory

propriospinal interneurons (Willis and Willis, 1964; Restrepo et al., 2009; Bikoff et al., 2016; Flynn et al.,

2017). Because of this heterogeneity, interneurons were separated into regions for further analysis, to

examine more specifically which inhibitory interneurons were perturbed. Based on our regional analysis,

interneurons in the ventral-most 100 μm from the white matter were altered the most in both

excitability and morphology, with the most strongly depolarized PICs and larger primary neurites, more

so than glycinergic interneurons within lamina IX and intermediate lamina VI and VII. Interneurons

located in the region within 100 μm from the ventral white matter likely contain more RCs based on the

location (Siembab et al., 2010) and electrophysiological characteristics (Perry et al., 2015; Bikoff et al.,

2016). Glycinergic interneurons located in lamina IX are reminiscent of the locomotor-related GABAergic

interneurons characterized by Nishimaru and colleagues, which were located within the ventrolateral

spinal cord, surrounded MNs and were active during locomotor activity (Nishimaru et al., 2011). Indeed,

these may be the same neurons: colocalization of GABA and glycine is entirely possible (Jonas et al.,

1998; Allain et al., 2006). If the lamina IX interneurons are indeed locomotor-related, even the modest

dampening of excitability observed here in SOD1 mice could contribute to changes in locomotor

patterning as has been described in other studies (Quinlan et al., 2017; Allodi et al., 2021). Further

studies are warranted to examine how inhibitory interneurons may be differently modulated in a

subtype-specific manner in animal models of ALS.

Functional implications for impaired inhibitory circuits

Outward signs of improperly functioning inhibitory interneurons may be apparent from subtle changes

in motor patterning and could potentially be used as a biomarker of early ALS. Locomotor disturbances

have been demonstrated *in vivo* in presymptomatic SOD1 mice during walking/running (Vinsant et al.,

2013; Akay, 2014; Quinlan et al., 2017; Allodi et al., 2021). In the embryonic and neonatal spinal cord *in*

vitro, increased duration of depolarizing events in MNs and a slower period of locomotor-related

bursting has been described (Medelin et al., 2016; Branchereau et al., 2019), which indicate that even

during development the functioning of locomotor circuits is already impaired. In presymptomatic adult

mice capable of running on a treadmill, subtle locomotor differences are observed including advanced

intermuscular phasing and a slower speed (Quinlan et al., 2017; Allodi et al., 2021) which indicate that

before any large-scale neurodegeneration occurs, there are changes in patterns of activity in MNs and

interneurons that could be exploited for use as a biomarker of ALS. At initial symptom onset, the first

failure of the tibialis anterior manifests as abnormality in the swing phase of the ankle (Akay, 2014).

Interestingly, silencing different interneuron populations (including inhibitory interneurons and

cholinergic interneurons) reveals the contribution of each population: silencing cholinergic interneurons

increases locomotor deficits in SOD1 mice (Landoni et al., 2019), while silencing inhibitory interneurons

has no effect suggesting these interneurons are already impaired (Allodi et al., 2021).

Clinical-translational implications

Evidence of MN hyperexcitability in ALS patients is present alongside signs of dysfunction in inhibition

(Rothstein et al., 1992; Plaitakis and Constantakakis, 1993; Raynor and Shefner, 1994; Shefner and

Logigian, 1998; Andreadou et al., 2008; Bae et al., 2013). Perhaps reducing excitability (with riluzole, for

example) only results in a modest increase in lifespan (Bensimon et al., 1994) because it could be helpful

to MNs while exacerbating inhibitory dysfunction by further depressing excitability of inhibitory

interneurons like RCs. Thus, a more targeted approach to reduce MN excitability could be beneficial

early in disease pathology. Stimulation of inhibitory nerves or certain protocols of transcranial

stimulation could be explored, along with augmentation of inhibitory neurotransmission.

Data Availability Statement

All data presented in this article were included in the Figures and Tables. Data were excluded only based

on the criteria stated in the methods.

Competing interests

Authors have no competing interests/conflicts of interest.

Author contributions

CFC: acquisition of images and 3D reconstruction data, analysis and statistics, writing the manuscript

PRS: analysis of electrophysiology and statistical data, and revising the manuscript

EJR: 3D reconstructions and analysis, revisions to the manuscript

LTG: statistical analysis of electrophysiology and morphology, revisions to the manuscript

LMM: analysis of statistical data, revising the manuscript

KJL: analysis of 3D reconstruction data, revising the manuscript

ACP: analysis of electrophysiology data, revising the manuscript

MM: statistical analysis and effect size analysis, revisions to the manuscript

NK: conception and design of the work, revising the manuscript

KAQ: conception and design of the work, acquisition of electrophysiology and imaging data, analysis of

electrophysiological data, statistics and writing the manuscript

All authors approved the final version of the manuscript; agreed to be accountable for all aspects of the

work in ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved; and all persons designated as authors qualify for authorship,

and all those who qualify for authorship are listed.

Funding

This project was funded by a springboard fellowship from Target ALS and R01NS104436 to KAQ and a

University of Rhode Island Graduate School Tuition Scholarship to LMM.

Acknowledgments

The authors made use of equipment supported by the Institutional Development Award (IDeA) Network

for Biomedical Research Excellence from the National Institute of General Medical Sciences of the

National Institutes of Health under grant number P20GM103430. The authors thank Drs. Ole Kiehn and

Hans Zeilhofer for providing access and permission (respectively) to GlyT2eGFP mice for this study. The

authors thank Claudia Fallini for comments on the initial drafts of the manuscript.

References

Akay T (2014) Long-term measurement of muscle denervation and locomotor behavior in individual wild-type and ALS model mice. J Neurophysiol 111:694–703.

Allain A-E, Bairi A, Meyrand P, Branchereau P (2006) Expression of the Glycinergic System During the Course of Embryonic Development in the Mouse Spinal Cord and its Co-Localization with GABA Immunoreactivity. Journal of Comparative Neurology 496:832–846.

Allodi I, Montañana-Rosell R, Selvan R, Löw P, Kiehn O (2021) Locomotor deficits in a mouse model of ALS are paralleled by loss of V1-interneuron connections onto fast motor neurons. Nat Commun 12:1–18.

Amendola J, Durand J (2008) Morphological differences between wild-type and transgenic superoxide dismutase 1 lumbar motoneurons in postnatal mice. Journal of Comparative Neurology 511:329–341.

Andreadou E, Kapaki E, Kokotis P, Paraskevas GP, Katsaros N, Libitaki G, Petropoulou O, Zis V, Sfagos C, Vassilopoulos D (2008) Plasma glutamate and glycine levels in patients with amyotrophic lateral sclerosis. In Vivo (Brooklyn) 22:137–142.

Avossa D, Grandolfo M, Mazzarol F, Zatta M, Ballerini L (2006) Early signs of motoneuron vulnerability in a disease model system: Characterization of transverse slice cultures of spinal cord isolated from embryonic ALS mice. Neuroscience 138:1179–1194.

Bae JS, Simon NG, Menon P, Vucic S, Kiernan C, Doyle AC, Case-book T, Holmes S (2013) The Puzzling Case of Hyperexcitability in Amyotrophic Lateral Sclerosis What Really is Hyperexcitability in ALS ? How Can We Objectively Assess. J of clinical neurology 9:65–74.

Bellingham MC (2011) A Review of the Neural Mechanisms of Action and Clinical Efficiency of Riluzole in Treating Amyotrophic Lateral Sclerosis: What have we Learned in the Last Decade? CNS Neurosci Ther 17:4–31.

Bensimon G, Lacomblez L, Meininger V (1994) A Controlled Trial of Riluzole in Amyotrophic Lateral Sclerosis. New England Journal of Medicine 330:585–591.

Bikoff JB, Gabitto MI, Rivard AF, Drobac E, Machado TA, Miri A, Brenner-Morton S, Famojure E, Diaz C, Alvarez FJ, Mentis GZ, Jessell TM (2016) Spinal Inhibitory Interneuron Diversity Delineates Variant Motor Microcircuits. Cell 165:207–219.

Branchereau P, Martin E, Supiot L, Hodeib F, Laupénie A, Dalvi U, Zhu H, Cazenave W, Cattaert D (2019) Relaxation of synaptic inhibitory events as a compensatory mechanism in fetal SOD spinal motor networks. Elife 8:1–28.

Casas C, Herrando-Grabulosa M, Manzano R, Mancuso R, Osta R, Navarro X (2013) Early presymptomatic cholinergic dysfunction in a murine model of amyotrophic lateral sclerosis. Brain Behav 3:145–158.

Chang Q, Martin LJ (2009a) Glycinergic Innervation of Motoneurons Is Deficient in Amyotrophic
Lateral Sclerosis Mice A Quantitative Confocal Analysis. American Journal of Pathology 174:574–
585.

Chang Q, Martin LJ (2009b) Glycinergic innervation of motoneurons is deficient in amyotrophic
lateral sclerosis mice: A quantitative confocal analysis. American Journal of Pathology 174:574–
585.

Clark RM, Blizzard CA, Young KM, King AE, Dickson TC (2017) Calretinin and Neuropeptide y
interneurons are differentially altered in the motor cortex of the SOD1 G93A mouse model of ALS.
Sci Rep 7:1–13.

Clark RM, Brizuela M, Blizzard CA, Dickson TC (2018) Reduced excitability and increased neurite
complexity of cortical interneurons in a familial mouse model of amyotrophic lateral sclerosis.
Front Cell Neurosci 12:1–7.

Cumming G (2012) Understanding the new statistics: effect sizes, confidence intervals, and meta-
analysis. New York: Routledge.

Delestrée N, Manuel M, Iglesias C, Elbasiouny SM, Heckman CJ, Zytnicki D (2014) Adult spinal
motoneurones are not hyperexcitable in a mouse model of inherited amyotrophic lateral sclerosis.
Journal of Physiology 592:1687–1703.

Devlin AC, Burr K, Borooah S, Foster JD, Cleary EM, Geti I, Vallier L, Shaw CE, Chandran S, Miles GB
(2015) Human iPSC-derived motoneurons harbouring TARDBP or C9ORF72 ALS mutations are
dysfunctional despite maintaining viability. Nat Commun 6:1–12.

Dukkipati SS, Chihi A, Wang Y, Elbasiouny SM (2016) Experimental Design and Data Analysis Issues
Contribute to Inconsistent Results of C-Bouton Changes in Amyotrophic Lateral Sclerosis. eNeuro
4:1–13.

Dukkipati SS, Garrett TL, Elbasiouny SM (2018) The vulnerability of spinal motoneurons and soma
size plasticity in a mouse model of amyotrophic lateral sclerosis. Journal of Physiology 596:1723–
1745.

Filipchuk AA, Durand J (2012) Postnatal dendritic development in lumbar motoneurons in mutant
superoxide dismutase 1 mouse model of amyotrophic lateral sclerosis. Neuroscience 209:144–154.

Flynn JR, Conn VL, Boyle KA, Hughes DI, Watanabe M, Velasquez T, Goulding MD, Callister RJ,
Graham BA (2017) Anatomical and molecular properties of long descending propriospinal neurons
in mice. Front Neuroanat 11:1–13.

Fogarty MJ, Kanjhan R, Bellingham MC, Noakes PG (2016) Glycinergic neurotransmission: A potent
regulator of embryonic motor neuron dendritic morphology and synaptic plasticity. Journal of
Neuroscience 36:80–87.

Genç B, Jara JH, Lagrimas AKB, Pytel P, Roos RP, Mesulam MM, Geula C, Bigio EH, Özdinler PH
(2017) Apical dendrite degeneration, a novel cellular pathology for Betz cells in ALS. Sci Rep 7:1–10.

Gómez-de-Mariscal E, Guerrero V, Sneider A, Jayatilaka H, Phillip JM, Wirtz D, Muñoz-Barrutia A
(2021) Use of the p-values as a size-dependent function to address practical differences when
analyzing large datasets. Sci Rep 11.

Grundy D (2015) Principles and standards for reporting animal experiments in The Journal of
Physiology and Experimental Physiology. J Physiol 593:2547–2549 Available at:
http://www.physoc.org/animal-experiments.

Gunes ZI, Kan VWY, Ye XQ, Liebscher S (2020) Exciting Complexity: The Role of Motor Circuit
Elements in ALS Pathophysiology. Front Neurosci 14:1–30.

Hayashi H, Suga M, Satake M, Tsubaki T (1981) Reduced glycine receptor in the spinal cord in
amyotrophic lateral sclerosis. Ann Neurol 9:292–294.

Hedges L v (1981) DISTRIBUTION THEORY FOR GLASS'S ESTIMATOR OF EFFECT SIZE AND RELATED
ESTIMATORS. Journal oi Educational Statistics 6:107–128.

Hegedus J, Putman CT, Gordon T (2007) Time course of preferential motor unit loss in the
SOD1G93A mouse model of amyotrophic lateral sclerosis. Neurobiol Dis 28:154–164.

Herron LR, Miles GB (2012) GENDER-SPECIFIC PERTURBATIONS IN MODULATORY INPUTS TO
MOTONEURONS IN A MOUSE MODEL OF AMYOTROPHIC LATERAL SCLEROSIS. Neuroscience
226:313–323.

Hossaini M, Cano SC, Van Dis V, Haasdijk ED, Hoogenraad CC, Holstege JC, Jaarsma D (2011) Spinal
inhibitory interneuron pathology follows motor neuron degeneration independent of glial mutant
superoxide dismutase 1 expression in SOD1-ALS mice. J Neuropathol Exp Neurol 70:662–677.

Howells J, Sangari S, Matamala JM, Kiernan MC, Marchand-Pauvert V, Burke D (2020) Interrogating
interneurone function using threshold tracking of the H reflex in healthy subjects and patients with
motor neurone disease. Clinical Neurophysiology 131:1986–1996.

Huh S, Heckman CJ, Manuel M (2021) Time course of alterations in adult spinal motoneuron
properties in the SOD1(G93A) mouse model of ALS. eNeuro.

Irvin CW, Kim RB, Mitchell CS (2015) Seeking homeostasis: Temporal trends in respiration,
oxidation, and calcium in SOD1 G93A amyotrophic lateral sclerosis mice. Front Cell Neurosci 9:1–
11.

Jensen DB, Kadlecova M, Allodi I, Meehan CF (2020) Spinal motoneurones are intrinsically more
responsive in the adult G93A SOD1 mouse model of Amyotrophic Lateral Sclerosis. J Physiol.

Jonas P, Bischofberger J, Sandkühler J (1998) Corelease of two fast neurotransmitters at a central
synapse. Science (1979) 281:419–424.

Kim J, Hughes EG, Shetty AS, Arlotta P, Goff LA, Bergles DE, Brown SP (2017) Changes in the
Excitability of Neocortical Neurons in a Mouse Model of Amyotrophic Lateral Sclerosis Are Not
Specific to Corticospinal Neurons and Are Modulated by Advancing Disease. The Journal of
Neuroscience 37:9037–9053.

Kuo JJ, Schonewille M, Siddique T, Schults AN, Fu R, Bar PR, Anelli R, Heckman C, Kroese AB (2004)
Hyperexcitability of Cultured Spinal Motoneurons From Presymptomatic ALS Mice. J Neurophysiol
91:571–575.

Kuo JJ, Siddique T, Fu R, Heckman CJ (2005) Increased persistent Na+ current and its effect on
excitability in motoneurones cultured from mutant SOD1 mice. Journal of Physiology 563:843–854.

Kuo SW, Binder MD, Heckman CJ (2020) Excessive Homeostatic Gain in Spinal Motoneurons in a
Mouse Model of Amyotrophic Lateral Sclerosis. Sci Rep 10.

Landoni LM, Myles JR, Wells TL, Mayer WP, Akay T (2019) Cholinergic modulation of motor
neurons through the C-boutons are necessary for the locomotor compensation for severe motor
neuron loss during amyotrophic lateral sclerosis disease progression. Behavioural Brain Research
369.

Lazic SE (2010) The problem of pseudoreplication in neuroscientific studies: is it affecting your
analysis? BMC Neurosci 11:1–17 Available at: http://www.biomedcentral.com/1471-2202/11/5.

Leroy F, Lamotte d'Incamps B, Imhoff-Manuel RD, Zytnicki D (2014) Early intrinsic hyperexcitability
does not contribute to motoneuron degeneration in amyotrophic lateral sclerosis. Elife 3:1–25.

Lin M, Lucas HC, Shmueli G (2013) Too big to fail: Large samples and the p-value problem.
Information Systems Research 24:906–917.

Malessa S, Leigh PN, Bertel O, Sluga E, Hornykiewicz O (1991) Amyotrophic lateral sclerosis:
glutamate dehydrogenase and transmitter amino acids in the spinal cord. J Neurol Neurosurg
Psychiatry 54:984–988.

Martin E, Cazenave W, Cattaert D, Branchereau P (2013) Embryonic alteration of motoneuronal
morphology induces hyperexcitability in the mouse model of amyotrophic lateral sclerosis.
Neurobiol Dis 54:116–126.

Martin LJ, Liu Z, Chen K, Price AC, Pan Y, Swaby JA, Golden WC (2007) Motor Neuron Degeneration
in Amyotrophic Lateral Sclerosis Mutant Superoxide Dismutase-1 Transgenic Mice: Mechanisms of
Mitochondriopathy and Cell Death. Journal of Comparative Neurology 500:20–46.

Martinez-Silva M de L, Imhoff-Manuel RD, Sharma A, Heckman C, Shneider NA, Roselli F, Zytnicki D,
Manuel M (2018) Hypoexcitability precedes denervation in the large fast-contracting motor units in
two unrelated mouse models of ALS. Elife 7:1–26.

McGown A, McDearmid JR, Panagiotaki N, Tong H, Al Mashhadi S, Redhead N, Lyon AN, Beattie CE,
Shaw PJ, Ramesh TM (2013) Early interneuron dysfunction in ALS: Insights from a mutant sod1
zebrafish model. Ann Neurol 73:246–258.

Medelin M, Rancic V, Cellot G, Laishram J, Veeraraghavan P, Rossi C, Muzio L, Sivilotti L, Ballerini L
(2016) Altered development in GABA co-release shapes glycinergic synaptic currents in cultured
spinal slices of the SOD1G93Amouse model of amyotrophic lateral sclerosis. Journal of Physiology
594:3827–3840.

Milan L, Courtand G, Cardoit L, Masmejean F, Barrière G, Cazalets JR, Garret M, Bertrand SS (2015)
Age-related changes in pre- and postsynaptic partners of the cholinergic C-boutons in wild-type
and SOD1G93Alumbar motoneurons. PLoS One 10:1–21.

Niebroj-Dobosz I, Janik P (1999) Amino acids acting as transmitters in amyotrophic lateral sclerosis
(ALS). Acta Neurol Scand 100:6–11.

Nishimaru H, Sakagami H, Kakizaki M, Yanagawa Y (2011) Locomotor-related activity of GABAergic
interneurons localized in the ventrolateral region in the isolated spinal cord of neonatal mice. J
Neurophysiol 106:1782–1792.

Ozdinler PH, Benn S, Yamamoto TH, Guzel M, Brown RH, Macklis JD (2011) Corticospinal Motor
Neurons and Related Subcerebral Projection Neurons Undergo Early and Specific
Neurodegeneration in hSOD1G93A Transgenic ALS Mice. Journal of Neuroscience 31:4166–4177.

Özyurt MG, Topkara B, İşak B, Türker KS (2020) Amyotrophic lateral sclerosis weakens spinal
recurrent inhibition and post-activation depression. Clinical Neurophysiology 131:2875–2886.

Pambo-Pambo A, Durand J, Gueritaud J-P (2009) Early excitability changes in lumbar motoneurons
of transgenic SOD1G85R and SOD1G(93A-Low) mice. J Neurophysiol 102:3627–3642.

Perry S, Gezelius H, Larhammar M, Hilscher MM, Lamotte d'Incamps B, Leao KE, Kullander K (2015)
Firing properties of Renshaw cells defined by Chrna2 are modulated by hyperpolarizing and small
conductance ion currents I_h and I_{SK}. European Journal of Neuroscience
41:887–898.

Pieri M, Carunchio I, Curcio L, Mercuri NB, Zona C (2009) Increased persistent sodium current
determines cortical hyperexcitability in a genetic model of amyotrophic lateral sclerosis. Exp
Neurol 215:368–379.

Piotrkiewicz M, Hausmanowa-Petrusewicz I (2011) Motoneuron afterhyperpolarisation duration in
amyotrophic lateral sclerosis. Journal of Physiology 589:2745–2754.

Plaitakis A, Constantakakis E (1993) Altered metabolism of excitatory amino acids, N-acetyl-
aspartate and N-acetyl-aspartylglutamate in amyotrophic lateral sclerosis. Brain Res Bull 30:381–
386.

Pullen AH, Athanasiou D (2009) Increase in presynaptic territory of C-terminals on lumbar
motoneurons of G93A SOD1 mice during disease progression. European Journal of Neuroscience
29:551–561.

Pun S, Santos AF, Saxena S, Xu L, Caroni P (2006) Selective vulnerability and pruning of phasic
motoneuron axons in motoneuron disease alleviated by CNTF. Nat Neurosci 9:408–419.

Quinlan KA, Kajtaz E, Ciolino JD, Imhoff-Manuel RD, Tresch MC, Heckman CJ, Tysseling VM (2017)
Chronic electromyograms in treadmill running SOD1 mice reveal early changes in muscle
activation. Journal of Physiology 595:5387–5400.

Quinlan KA, Lamano JB, Samuels J, Heckman CJ (2015) Comparison of dendritic calcium transients
in juvenile wild type and SOD1G93A mouse lumbar motoneurons. Front Cell Neurosci 9:1–13.

Quinlan KA, Schuster JE, Fu R, Siddique T, Heckman CJ (2011) Altered postnatal maturation of
electrical properties in spinal motoneurons in a mouse model of amyotrophic lateral sclerosis.
Journal of Physiology 589:2245–2260.

Raynor EM, Shefner JM (1994) Recurrent inhibition is decreased in patients with amyotrophic
lateral sclerosis. Neurology 44:2148–2153.

Restrepo CE, Lundfald L, Szabó G, Erdélyi F, Zeilhofer HU, Glover JC, Kiehn O (2009) Transmitter-
phenotypes of commissural interneurons in the lumbar spinal cord of newborn mice. Journal of
Comparative Neurology 517:177–192.

Rothstein JD, Martin LJ, Kuncl RW (1992) Decreased Glutamate Transport by the Brain and Spinal
Cord in Amyotrophic Lateral Sclerosis. New England Journal of Medicine 326:1464–1468.

Saba L, Viscomi MT, Caioli S, Pignataro A, Bisicchia E, Pieri M, Molinari M, Ammassari-Teule M,
Zona C (2016) Altered Functionality, Morphology, and Vesicular Glutamate Transporter Expression
of Cortical Motor Neurons from a Presymptomatic Mouse Model of Amyotrophic Lateral Sclerosis.
Cerebral Cortex 26:1512–1528.

Saba L, Viscomi MT, Martini A, Caioli S, Mercuri NB, Guatteo E, Zona C (2019) Modified age-
dependent expression of NaV1.6 in an ALS model correlates with motor cortex excitability
alterations. Neurobiol Dis 130:104532.

Sangari S, Iglesias C, El Mendili MM, Benali H, Pradat PF, Marchand-Pauvert V (2016) Impairment of
sensory-motor integration at spinal level in amyotrophic lateral sclerosis. Clinical Neurophysiology
127:1968–1977.

Saxena S, Roselli F, Singh K, Leptien K, Julien JP, Gros-Louis F, Caroni P (2013) Neuroprotection
through Excitability and mTOR Required in ALS Motoneurons to Delay Disease and Extend Survival.
Neuron 80:80–96.

Shefner JM, Logigian EL (1998) The mixed nerve silent period in normal subjects and patients with
amyotrophic lateral sclerosis. Electromyogr Clin Neurophysiol 38:505–510.

Siembab VC, Smith CA, Zagoraiou L, Berrocal MC, Mentis GZ, Alvarez FJ (2010) Target selection of
proprioceptive and motor axon synapses on neonatal V1-derived Ia inhibitory interneurons and
Renshaw cells. Journal of Comparative Neurology 518:4675–4701.

Vinsant S, Mansfield C, Jimenez-Moreno R, Moore VDG, Yoshikawa M, Hampton TG, Prevette D,
Caress J, Oppenheim RW, Milligan C (2013) Characterization of early pathogenesis in the
SOD1G93A mouse model of ALS: Part II, results and discussion. Brain Behav 3:431–457.

Wainger BJ, Kiskinis E, Mellin C, Wiskow O, Han SSW, Sandoe J, Perez NP, Williams LA, Lee S,
Boulting G, Berry JD, Brown RH, Cudkowicz ME, Bean BP, Eggan K, Woolf CJ (2014) Intrinsic

membrane hyperexcitability of amyotrophic lateral sclerosis patient-derived motor neurons. Cell
Rep 7:1–11.

Whitehouse PJ, Wamsley JK, Zarbin MA, Price DL, Tourtellotte WW, Kuhar MJ (1983) Amyotrophic
lateral sclerosis: Alterations in neurotransmitter receptors. Ann Neurol 14:8–16.

Willis W, Willis JC (1964) Location of Renshaw Cells. Nature 204:1214–1215.

Wootz H, Fitzsimons-Kantamneni E, Larhammar M, Rotterman TM, Enjin A, Patra K, André E, Van
Zundert B, Kullander K, Alvarez FJ (2013) Alterations in the motor neuron-renshaw cell circuit in the
Sod1G93Amouse model. Journal of Comparative Neurology 521:1449–1469.

Yu Z, Guindani M, Grieco SF, Chen L, Holmes TC, Xu X (2022) Beyond t test and ANOVA:
applications of mixed-effects models for more rigorous statistical analysis in neuroscience
research. Neuron 110:21–35.

Zeilhofer HU, Studler B, Arabadzisz D, Schweizer C, Ahmadi S, Layh B, Bösl MR, Fritschy JM (2005)
Glycinergic neurons expressing enhanced green fluorescent protein in bacterial artificial
chromosome transgenic mice. Journal of Comparative Neurology 482:123–141.