Polyploidy-associated paramutation in Arabidopsis is determined by small RNAs, temperature, and allele structure

Heinrich Bente[1], Andrea M. Foerster[1], Nicole Lettner[1], Ortrun Mittelsten Scheid[1]

[1]Gregor Mendel Institute of Molecular Plant Biology (GMI), Austrian Academy of Sciences, Vienna Biocenter (VBC), 1030 Vienna, Austria

ABSTRACT

Paramutation is a form of non-Mendelian inheritance in which the expression of a paramutable allele changes when it encounters a paramutagenic allele. This change in expression of the paramutable alleles is stably inherited even after segregation of both alleles. While the discovery of paramutation and studies of its underlying mechanism were made with alleles that change plant pigmentation, paramutation-like phenomena are known to modulate the expression of other traits and in other eukaryotes, and many cases have probably gone undetected. It is likely that epigenetic mechanisms are responsible for the phenomenon, as paramutation forms epialleles, genes with identical sequences but different expression states. This could account for the intergenerational inheritance of the paramutated allele, providing profound evidence that triggered epigenetic changes can be maintained over generations. Here, we use a case of paramutation that affects a transgenic selection reporter gene in tetraploid *Arabidopsis thaliana*. Our data suggest that different types of small RNA are associated with paramutable and paramutagenic epialleles. In addition, deletion of a repeat within the epiallele changes its paramutability. Further, the temperature during the growth of the epiallelic hybrids determines the degree and timing of the allelic interaction. The data further make it plausible why paramutation in this system becomes evident only in the segregating F2 population of tetraploid plants containing both epialleles. In summary, the results support a model for polyploidy-associated paramutation, with similarities as well as distinctions from other cases of paramutation.

Keywords: Paramutation, epialleles, small RNAs, polyploidy, RNA-directed DNA methylation, Arabidopsis thaliana

AUTHOR SUMMARY

A curiosity in the history of genetics is the fact that experiments with peas, Mendel's most studied plants, resulted in one of the first observations of non-Mendelian genetics published already in 1915. Crossings with plants showing a "rogue" leaf phenotype made normal leaf shape disappear in all progeny, without recovery. This phenomenon was later demonstrated for more traits in other plants and termed paramutation. Paramutation is due to the epigenetic switch of aa active gene to a silenced version which is then maintained in the inactive state in later generations. This demonstrates that acquired epigenetic changes can become permanent. Despite its early observation and numerous studies in mainly maize and tomato, it is barely understood how paramutation is established and which parameters influence the process. We investigated a case of paramutation in *Arabidopsis thaliana*, crossing plants with genetically identical but epigenetically different alleles that result in resistance or sensitivity to hygromycin in the growth medium. Unexpectedly, paramutation did not become manifest in the hybrids but only in their progeny and only in plants with a doubled chromosome set. These features make this paramutation distinct from other cases and our studies revealed the involvement of several parameters: an important role for sRNAs to initiate silencing, the sequence of the allele itself, the environmental conditions during growth of the hybrids, the developmental stage, and the copy number ratio between the epialleles.

INTRODUCTION

Paramutation is the term for a specific interaction between alleles of a gene, in which one paramutable allele becomes heritably modified by encounter with a paramutagenic allele. It violates the Mendelian rule of independent segregation, as the paramutated allele is maintained in the new state and even can acquire paramutagenicity itself. This led to its description as para-(resembling)-mutation or somatic gene conversion by the pioneers working on maize and tomato in the 1950's [1-4]. Even earlier genetic experiments with peas [5] are now interpreted as evidence of paramutation [6]. Numerous cases of paramutation have been reported in several species including metazoans and plants [reviewed in 7, 8, 9], indicating that paramutation is not just a bizarre exception but may be a general phenomenon among eukaryotes [10].

Paramutation has an epigenetic basis and does not change the DNA sequence of participating alleles. Instead, specific, newly acquired epigenetic states can be installed and maintained over generations independent of the trigger of the change. It is likely the best example for true transgenerational inheritance of defined epigenetic changes connected with a specific trigger, namely the encounter with the paramutagenic allele. Although many principles of epigenetic regulation are now known, the molecular basis of paramutation is still

not well understood. Genetic screens, mainly in maize, have identified several genes whose

function is necessary for establishing and/or maintenance of paramutation, e.g. *"Required to*

Maintain Repression" (*RMR*) [11] or *"Mediator Of Paramutation"* (*MOP*) [12]. The nature of

their gene products [reviewed in 13] suggested that paramutation requires components of the

RNA-directed DNA methylation (RdDM) pathway [reviewed in 14, 15, 16], which is responsible

for silencing repetitive sequences via small RNAs (sRNAs) that guide the RNA-induced

silencing complex (RISC) to corresponding genomic loci, leading to DNA methylation and the

formation of heterochromatin. However, the majority of RdDM targets are not paramutable,

suggesting that additional factors, such as *cis*-acting determinants [17] or other *trans*-acting

elements are involved. To identify these, additional paramutation systems will be informative,

especially those that differ in penetrance, frequency, and stability of the epigenetic change.

Paramutation has also been demonstrated in *Arabidopsis thaliana* [18]. It involves

epialleles of a transgene that, in an active state, expresses hygromycin phosphotransferase

(HPT) conferring resistance (R epiallele) to hygromycin. The inactive state confers sensitivity

(S epiallele) to the antibiotic. *In vitro* culture of protoplasts from diploid plants with an R epiallele

and subsequent regeneration resulted in several tetraploid plants, some of which were

hygromycin-resistant while others had spontaneously acquired a silent epiallele. These

silenced plants acquired DNA methylation and histone marks characteristic of silenced

chromatin [19, 20], no longer transcribed the *HPT* gene, and rendered plants hygromycin-

sensitive. Diploid R and S derivatives were generated by backcrossing to diploids, and progeny

that were homozygous for either R or S alleles inherited their respective expression state and

phenotype over numerous generations, in both, diploid and tetraploid lines. Molecular analysis

confirmed that the sequence of the R and S allele were identical, and the *HPT* gene is inserted

in an intergenic region far from transposons. Therefore, it was concluded that the R and S

alleles differ in their epigenetic state and are true epialleles [21]. A mutant screen scoring for

reactivation of the silent epiallele led to identification of the *trans*-acting mutants *ddm1* and

hog1, confirming the importance of DNA and histone methylation [19]. It also demonstrated

that *cis*-acting mutations with structural variants of the epiallele (RΔ) led to restored

transcription and hygromycin resistance [21]. Together these data demonstrated that the S

allele results from epigenetic silencing of the *HPT* gene.

Crossing diploid R and S plants resulted in 100% hygromycin-resistant F1 plants, and

selfed F1 plants produced F2 progeny with close to 75% (3:1) resistant plants, as expected for

a dominant trait and confirming independent segregation if the epialleles. However, while F1

hybrids from crosses between tetraploid R and S plants were 100% resistant, their F2 progeny

contained significantly fewer resistant plants than the expected 97% (35:1), although there was

variation between independent tetraploid F2 populations. This argued against a genetic

determinant of the lost resistance and suggested that epigenetic components were interfering

with the independence of segregation, similar to paramutation [18]. However, there were
interesting differences from the classical paramutation cases. While the loss of gene
expression was visible in maize and tomato in the F1 hybrid [22] of these diploid species, the
phenotype indicating paramutation in the R/S interaction was restricted to the F2 generation
of tetraploid Arabidopsis hybrids only. These data led to the hypothesis that this system's late
onset of the interaction and its requirement for polyploidy could provide more insight into the
mechanisms that determine the occurrence and degree of paramutation.

Here, we investigated the role of the epialleles' structure, the connection with
transcriptional activity, and the potential for secondary paramutation. We examined the effect
of temperature during growth of the F1 hybrids, and profiled small RNAs associated with the
epialleles in seedlings and flower buds of diploid and tetraploid plants. We show that the
degree of paramutation depends on environmental conditions, developmental stage, and an
interplay between the structure of the epialleles and the nature of the small RNAs associated
with them. Based on these data, we present a model that explains the role of copy number in
the epiallelic interaction in polyploid plants.

RESULTS

Nature and nomenclature of the epialleles involved in paramutation

The phenotype revealing paramutation in Arabidopsis is based on expression or silencing of the hygromycin phosphotransferase (*HPT*) gene conferring resistance to the antibiotic hygromycin. A construct with *HPT* under control of the CaMV 35S promoter was used for PEG-mediated protoplast transformation [23] and inserted in an intergenic region of chromosome three in the Arabidopsis genome. Plasmid copies recombined before integration, resulting in an insert consisting of a vector fragment upstream of the promoter and the *HPT* open reading frame, a partially deleted CaMV 35S terminator followed by a second copy of the vector and the CaMV 35S promoter, and approximately 500 bp non-coding non-plant carrier DNA used during the protoplast transformation. This structure has been previously described [18, 21]; a schematic illustration of all variants used in this study (considering the transcription direction opposite to the reference genome) is shown in Fig 1A. Transcribed and silenced epialleles have identical DNA sequence and are hereafter referred to as R (resistant) or S (sensitive). The "empty" insertion site in the wild type is termed W. Diploid and tetraploid genotypes will be called RR/SS/WW and RRRR/SSSS/WWWW, respectively. An overview of the origin and crosses of each epiallele and the respective mutants is illustrated in Fig 1B.

Paramutation depends on temperature during growth of tetraploid F1 hybrids

Plants homozygous for either R or S inherit hygromycin resistance or sensitivity as stable traits. Hybrids obtained by crossing diploid plants and selfing them resulted in F2 progeny with the expected 3:1 ratio, regardless whether R was combined with S or with the wild type lacking an epiallele (RS or RW). In contrast, paramutation occurs in RRSS (tetraploid) hybrids and is revealed by a reduced number of resistant F2 plants, significantly lower than in the control crosses of R with W (RRWW), independent of whether the S allele came from the maternal or paternal side, but with variation between F2 populations from different F1 parents [18]. This variation raised the question whether external factors during the growth of the F1 hybrids could influence the penetrance of paramutation. We hypothesized that temperature is an important parameter, because its influence on a paramutation-resembling inheritance in Drosophila had been reported [24]. We grew diploid and tetraploid F1 plants obtained from crosses between R and S (or W) at 10°C, 19°C, or 24°C during their flowering and seed set period (Fig 2A). We germinated F2 seeds on hygromycin selection medium and scored the ratio of resistant seedlings (Fig 2B and 2C). F2 from tetraploid RRSS hybrids grown at 19°C (the standard growth conditions) confirmed the results from earlier experiments, with a reduced ratio of resistant plantlets compared to the RRWW controls. This difference was even more significant in the progeny from parents grown at 24°C, indicating an enhanced interaction between R and S. Unexpectedly, progeny from RRSS and RRWW plants grown at 10°C showed no difference

in the resistance assay, indicating that the presence of the S epiallele did not lead to paramutation of R at the lower temperature (Fig 2B). All F2 plants from diploid hybrids contained approximately 75% resistant seedlings, excluding any influence of the S epiallele under all tested temperature conditions (Fig 2C). Therefore, paramutation occurs at 19°C but not in plants grown at 10°C. These results confirm that paramutation in this system is associated with tetraploidy and indicate that its penetrance is indeed affected by external factors, exemplified here by temperature.

Epialleles involved in Arabidopsis paramutation have different sRNA signatures

Multiple lines of evidence indicate that *trans*-silencing phenomena, such as paramutation, involve the production of small RNAs (sRNA) that silence the paramutated allele. The presence of sRNAs at loci participating in paramutation has been shown for genes in maize [25, 26], tomato [10], and Arabidopsis [27], including a very limited analysis of the S allele in this study [21]. To address the possible mechanistic role of sRNA, we generated sRNA libraries from total RNA of 14 d-old seedlings of tetraploid and diploid homozygous R and S lines and mapped reads with a length from 18-26 nt to the Arabidopsis TAIR10 reference genome extended *in silico* with the inserted *HPT* epiallele. Whereas the size distribution of sRNAs mapping to the whole genome were similar in all libraries, there was a striking difference regarding the size classes between lines carrying the different epialleles: we found mainly 21 nt sRNAs at R and mostly 24 nt sRNAs at S, regardless of ploidy (tetraploids: Fig 3A, diploids: S1A Fig). Both size classes of sRNAs mapped predominantly to the duplicated region (promoter and vector sequence) of the epiallele (Fig 3B, S1B Fig), indicating that these sRNAs may originate from either copy. Libraries prepared from wild type plants without the reporter do not contain reads mapping to the reporter (S2B Fig), confirming the origin of the 21/24 nt reads in Fig 3B from the epialleles. The abundance of the 24 nt sRNAs was similar between diploid and tetraploid S lines, and they were distributed along the whole duplicated promoter/vector region. In contrast, tetraploid R lines had higher levels of 21 nt sRNAs than diploids. Many of them mapped to a specific region of the promoter. The *HPT*-ORF and the non-coding region downstream of the second promoter copy were almost devoid of sRNAs in libraries from all samples (Fig 3B, S1B Fig). These data indicate that the active and silent epialleles are characterized by matching sRNAs of different length, with specific profiles and quantities.

The frequency of the four bases at the 5'end of sRNAs correlates with their length and serves as an indicator for their potential association with specific Argonaute (AGO) proteins [28]. In all lines, sRNAs mapping to the wild-type genome confirmed the known preference for a 5'U in 21 nt sRNAs and a 5'A in 24 nt sRNAs (S2A Fig). The 5' base distribution of epiallelic sRNAs in both tetraploids (Fig 3C) and diploids (Fig S1C) matches this preference: 5'

194 nucleotides of the 21 nt sRNAs mapping to the R epiallele were biased for 5'U, indicating a
195 preference for AGO1, while half of the 24 nt sRNAs from the S epiallele had an A at the 5' end,
196 suggesting association with either AGO4 [28] or AGO3 [29].

sRNA signature and mutant analysis support involvement of RdDM

The relatively high percentage of 5'A in the 24 nt sRNAs from the S epiallele (Fig 3C and
S1C Fig) suggested the involvement of RNA-directed DNA methylation (RdDM) [30, 31], as
suggested by several mutational studies in maize paramutation [32]. The limitation of the R/S
interaction to tetraploid plants made any forward-directed screen impractical, as recovery of
recessive mutations could only be expected in M3 generations and with extremely low
probability. However, the efficiency of CRISPR-Cas9- based mutagenesis in Arabidopsis [33]
allowed us to test the hypothesis that RdDM is involved in this case of paramutation.
Paramutation between the B-I and B' allele in maize requires functional RDR2 [34] and several
subunits of the plant-specific DNA-dependent RNA polymerases IV and V [35-37]. Therefore,
we targeted two corresponding Arabidopsis genes, *RDR2* (At4g11130) and the largest subunit
of PolIV (*NRPD1*, At1g63020) in tetraploid R and S plants for CRISPR-Cas9-based
mutagenesis. We selected lines homozygous for the desired mutations and the absence of the
Cas9 gene by genotyping. Loss of function was demonstrated by the genome-wide absence
of endogenous 24 nt small interfering RNAs (S3A Fig). Furthermore, the mutation in *NRPD1*
led to depletion of 24 nt sRNAs in SSSS (S3B Fig). Unlike the *NRPD1* mutant *rmr7* in maize,
which reactivated the silent B' allele [34], the newly generated Arabidopsis *cas9nrpd1* mutant
did not lead to reactivation of the S epiallele. We tested the potential for paramutation by
generating F1 hybrids homozygous for the mutation and scoring F2 seedlings for hygromycin
resistance, together with progeny from corresponding wild type combinations grown in parallel.
The ratio of resistant to sensitive progeny from mutant RRSS hybrids was much closer to that
of the RRWW crosses (Fig 4A) than in the wild-type background (Fig 4B). This indicates a
reduced establishment of paramutation when RdDM is impaired, but that RdDM does not play
a role in the maintenance of the silent state of the S epiallele.

The active epiallele is sensitive to silencing by *trans*-acting small RNAs

The results of the mutant analysis suggested that sRNAs are important for establishing
paramutation. We therefore tested the hypothesis that sRNAs were sufficient to silence the
otherwise stable R epiallele. We designed three hairpin (hp) constructs that would induce
strong sRNA formation targeting different regions within the duplicated vector/promoter region
(hp1/2/3; Fig 1A) and transformed RR and RRRR plants. Seeds from individual T1 plants were
sorted for the presence or absence of the hairpin construct via seed-specific GFP expression,
which was encoded in the vector, and grown with or without hygromycin. Under selection, all
hairpin variants caused reduced root growth in seedlings, compared to plants without hairpins

(Fig 5A and 5B), indicating silencing of the R allele in diploid and tetraploid lines. The degree
of silencing was highest with hp3 which is homologous to the promoter. However, hp1 and hp2
which are homologous to the vector sequence further away from the promoter within the
epiallele also resulted in significant growth inhibition in seedlings under selection (Fig 5A). To
quantify the effect of the hairpins on transcriptional levels at a later stage of development, we
determined *HPT* expression by qRT-PCR in flower buds from diploid and tetraploid T1 plants
with and without hairpins sorted as above. In diploid plants, all three hairpin variants reduced
HPT expression to a similar level (Fig 5C left panel), whereas the effect increases with
proximity of the hp-matching region to the transcription start site of the *HPT* gene in tetraploid
plants (Fig 5C right panel). The expression levels of the hairpin constructs are expected to be
averaged as we sampled many T1 plants representing independent insertion sites. Therefore,
the hairpin constructs appear to silence the two R copies in diploids more efficiently than the
four R copies in tetraploids. These data indicate that stoichiometry between sRNAs and the
target genes could explain the role of ploidy in the response to silencing cues.

New silenced epialleles can exert secondary paramutation

The epialleles do not contain an inverted repeat that could form hairpins to serve as a
source of sRNAs. However, tandem-oriented repeats can affect paramutation, depending on
their copy number [17, 38]. Evidence for a role of the direct repeat of the vector/promoter region
in our paramutation system came from the previously described structural variants (RΔ,
derived from the S epiallele by mutagenesis), in which rearrangements downstream of the
HPT-ORF led to reactivation of the upstream promoter [21]. We made use of two different,
independent deletions to ask whether and how they would affect the paramutability. In RΔ2,
the whole transgenic sequence downstream of the defective terminator (including the
vector/promoter repeat) was deleted, whereas RΔ4 lacked only the terminal non-coding carrier
sequence, maintaining the repeat (Fig 1A). Tetraploid, homozygous derivatives of both lines
(RΔ2RΔ2RΔ2RΔ2 and RΔ4RΔ4RΔ4RΔ4, selected after colchicine treatment) were crossed
with SSSS or WWWW lines (Fig. 1B) and the F2 progenies screened for hygromycin
resistance. Only F2 offspring from reciprocal crosses between RΔ4RΔ4RΔ4RΔ4 x SSSS
(RΔ4RΔ4SS), but not from RΔ2RΔ2RΔ2RΔ2 x SSSS (RΔ2RΔ2SS) crosses, showed reduced
hygromycin resistance like crosses with full length RRRR (Fig 6A). This indicates that RΔ2 is
a neutral allele, whereas RΔ4 is still paramutable.

The most persuasive evidence for paramutation is acquired paramutagenicity of a
paramutated allele, i.e. the ability to exert secondary paramutation. In the initial paramutation
cross RRRR x SSSS, paramutated R alleles do not have any sequence polymorphism that
distinguishes them from S alleles in the segregating F2. To enable identification of the

paramutated allele after its conversion, we used the deletion in RΔ4 for PCR-based genotyping
in the progeny of RΔ4RΔ4SS plants. We selected individual F2 plants that were either
homozygous (+/+) or heterozygous (+/-) for the deletion and quantified *HPT* expression with
qRT-PCR. As expected from previous studies and in accordance with the hygromycin
resistance assay, RRRR and RΔ4RΔ4RΔ4RΔ4 showed strong expression, while there were
no detectable transcripts in SSSS and WWWW. Moreover, only progeny from RΔ4RΔ4WW
hybrid parents expressed *HPT* in comparable amounts (Fig 6B). In contrast, progeny from
RΔ4RΔ4SS parents had very low *HPT* expression, including plants homozygous for the
previously active RΔ4 allele. This demonstrates that they contained only paramutated copies
(Fig 6B, black star), indicating strong suppression by the S allele in the previous generation.
This silencing lasted after segregation. Therefore, all RΔ4 alleles had undergone paramutation
to SΔ4SΔ4SΔ4SΔ4 (Fig 1B). We used these plants to assay secondary paramutation.
Tetraploid hybrids from crosses between plants with the fully paramutated alleles
(SΔ4SΔ4SΔ4SΔ4) and plants with the initial active R alleles (RRRR) (Fig 1B) resulted in F2
populations with a similarly reduced ratio of hygromycin resistance as obtained after the
primary paramutation (Fig 6C). This indicates that once an *HPT* allele is paramutated, it can
reduce *HPT* expression from a naïve R allele in the process of secondary paramutation.

**Removal of the downstream duplication does not reactivate a silent upstream promoter
but boosts expression from active epialleles**

RΔ2 and RΔ4 alleles had been identified in a forward genetic screen selecting for
reactivation of the S epiallele [21]. Both deletions resulted in hygromycin resistance, but RΔ2
lost paramutagenicity while RΔ4 maintained this feature. The different, randomly induced
structural changes (Fig 1A) suggested that the removal of the duplicated downstream region
in RΔ2, but not in RΔ4, might explain the difference. To precisely recapitulate this event, we
applied CRISPR gene editing to eliminate the downstream repeat as well as the non-coding
sequence from R and S alleles. Cas9-encoding transgenes with suitable sgRNAs were
transformed into diploid and tetraploid plants. T1 plants were selfed or outcrossed and the
progeny genotyped for homozygosity of the wanted deletion and the absence of the Cas9
transgene. To distinguish the new deletions from the previously described alleles, the Cas-9
induced alleles are denoted as Rγ and Sγ (Fig 1A). We screened diploid (RγRγ, SγSγ) and
tetraploid (RγRγRγRγ, SγSγSγSγ) plants for hygromycin resistance. R-derived plants with the
deletion grew longer roots on hygromycin (Fig 7A and 7B). Northern blots with RNA from these
lines confirmed a shorter transcript of ~1.6 kb (as excepted for the deletion), with ~3-4x more
signal strength than in R plants, in both diploids and tetraploids (Fig 7C). These data indicate
that, despite transcription and stable hygromycin resistance from full length R epialleles, the
associated 21 nt RNAs contribute to partial suppression of the upstream promoter

In contrast to the deletions after random mutagenesis [21], S-derived plants with the
targeted deletions (SγSγ, SγSγSγSγ) remained completely sensitive and without detectable
HPT transcripts (Fig 7D). Therefore, the removal of the duplicated regions was not sufficient
to restore the *HPT* expression. The sRNA profiles of the deletion lines were quite similar
between diploid and tetraploid lines, and between S and Sγ (S4 Fig B and D). Unexpectedly,
the profiles of both *HPT*-expressing deletion alleles (RΔ2, Rγ) lost the 21 nt sRNAs
characteristic for the full-length R allele but gained 24 nt sRNAs along the remaining vector-
promoter region (S4 C and F). This demonstrates that removal of the downstream repeat
eliminates the 21 nt class and further implies that the occurrence of 24 nt sRNA does not *per*
se lead to silencing. The main difference in the profiles of active (RΔ2, Rγ) versus silent (Sγ)
deletion alleles is the presence of sRNAs along the *HPT* coding region in the latter (S4 Fig C,
D, and F). Those at the vector-promoter regions are obviously not interfering with expression
and might be remnants from the previously present duplication, resembling the peaks of 24 nt
sRNAs at the tandemly orientated LTR regions of transposons in several plants [39-42].

To determine if Sγ and Rγ could participate in paramutation, we performed reciprocal crosses
between tetraploids with full length or deleted R and S alleles and screened F2 segregation
under hygromycin selection. Resistance in progeny from RRSγSγ hybrids was higher than for
RRSS parents (Fig 7E and 7F), indicating reduced paramutagenicity of S without the repeat.
The full-length S, paramutagenic in RRSS, was no longer paramutagenic in RγRγSS and
RγRγSγSγ (Fig 7A and 7B). Taken together, the removal of the duplicate and the absence of
21 nt sRNAs boosts expression from the active epiallele and renders it less or not at all
sensitive to paramutation, while the removal of the duplication from the silent epiallele weakens
its paramutagenicity. This implies that the structure of the epialleles on both partners, in
combination with the specific small RNA profiles, determines whether and how the epialleles
interact.

DISCUSSION

In this study, we investigated a case of paramutation in *Arabidopsis thaliana* that shares the main features, including secondary paramutation, with other examples, while being different with regard to its timing and link to polyploidy. We documented important roles for sRNAs to initiate silencing, the sequence of the allele itself, the environmental conditions during growth of the hybrids, the developmental stage, and the copy number ratio between the epialleles. We have described a model that makes these features plausible. The potentially strong *HPT* expression from the upstream Cauliflower Mosaic Virus 35S promoter is weakened in the active epiallele by 21 nt sRNAs connected with the downstream repeat, but still remains strong enough to produce sufficient HPT protein to detoxify hygromycin. Regular F2 resistance segregation after crosses with the "empty" wild type imply that one active allele, even with the repeat, is enough to render a plant resistant, regardless of whether the dosage is 1 out of 2 (diploid) or 1 out of 4 (tetraploid). If the cross introduced the inactive epiallele, *HPT* expression is still maintained high enough to allow resistance as long as the associated 24 nt sRNA are present in limited amounts. The ectopic overexpression of sRNA varieties from the hairpin constructs demonstrated that they are triggers of rapid, efficient, and lasting trans-silencing. However, in R and S lines without hairpin constructs, 21 nt and 24 nt sRNA, respectively, are not detectable on northern blots, have very low read counts in the small RNA libraries and are therefore present in only tiny amounts. In diploid hybrids, both epialleles are present in a 1:1 ratio, which likely does not tilt the balance towards silencing. In tetraploids, 22.2% of the F2 progeny will contain one active and three silent copies; here, the abundance of 24 nt sRNAs cross a threshold for initiating the trans-silencing effect. The "missing" portion of resistant plants in paramutation cross progeny is usually between 15-30%, compatible with the assumption that especially plants with more silent than active epialleles (never occurring in diploids) are undergoing the switch. A similar higher incidence for paramutation with a triplex genotype for the paramutagenic *sulfurea* allele is apparent in tetraploid tomato hybrids [43], and paramutation can occur independently at single alleles in subsequent generations [44].

The Arabidopsis/*HPT* and the tomato/*sulf* system are the only cases for which a role of tetraploidy in paramutation has been described. In neither case is the silent state of the paramutagenic epiallele connected with polyploidy, as both alleles are stable in diploid background, but paramutation seems to be supported by the 4x karyotype. Genome duplications after formation of polyploids has often been associated with epigenetic changes [45-49] and might trigger the origin of epialleles. However, most studies were performed with allopolyploids, in which combining different genomes modifies chromatin, DNA methylation, or small RNA profiles. In these cases, changes can be due to either increased chromosome numbers, increased genetic diversity, increased gene copy numbers, incompatibility of regulatory components, or a mixture of these parameters. The situation in autopolyploids is

simpler, as it excludes genetic determinants. Newly created autopolyploids can contain
epigenetic modifications, e.g. switchgrass [50] and Arabidopsis [51], but effects on gene
expression are much smaller than in allopolyploids [52] and occur rather stochastically [53].
We did not find prominent differences between diploid and tetraploid lines, besides the
divergent *HPT* expression and small RNA profiles at the epialleles. Therefore, the limitation of
paramutation to the tetraploid hybrids is likely not due to the chromosome number as such, or
to indirect consequences like larger nuclei [54], cell cycle regulation [55], or differences in
meiosis [56]. It is more plausible to assume that rather the copy number of the epialleles, or
better their ratio, determine the epigenetic change. Copy numbers, not varied by ploidy but by
allelic variation of repetitive sequences, are also a major determinant of paramutation in maize
[38, 57, 58], again mostly in tandem configuration but sometimes far upstream of the coding
sequence.

Another paradigm for a dosage effect is transcriptional silencing of a transposon, which
occurs only after the copy number of the genomic source has increased beyond a threshold
by new insertions [59]. A striking similarity with *HPT* paramutation is the role of 24 nt sRNAs
that characterize the paramutagenic epiallele and that appear in connection with the switch to
transcriptional transposon silencing. 24 nt sRNAs were also reported in very low amounts for
the B-I/B' paramutation in maize, depending on the RDR2 ortholog MOP1, but were also found
in neutral alleles and therefore not sufficient for paramutation [60]. As in our study, their ectopic
expression in larger amounts can mimic paramutation [61], again supporting a role for dosage
dependency. There is no report of 21 nt sRNAs at paramutable maize alleles. This is in contrast
to the *HPT* system, where their presence might be connected to the downstream location of
the repeat and its presence in the transcript. However, the repeats are in tandem orientation,
so that hairpin formation as a precursor for double-stranded RNA is unlikely. The requirement
forMOP1 in maize paramutation [12] and the reduced paramutation in *rdr2* mutants in our study
strongly support a dependence on RNA-dependent RNA polymerase for the creation of the
24 nt small RNAs. As MOP2 and MOP3 are also components of RdDM based silencing [35,
36], it is clear that this epigenetic regulatory pathway is a core component of paramutation.
Even outside of paramutation siRNA-induced epigenetic changes can persist over several
generations [62]. However, the interference with RdDM in the Arabidopsis system did not result
in hygromycin resistance, in contrast to restored BI and PI expression and regain of
pigmentation e.g. in mop1 [12]. However, this could be due to the exceptionally strong silencing
at the S epiallele, which needs simultaneous removal of DNA methylation and repressive
histone modifications [19].

The role of the 21 nt sRNAs in the Arabidopsis system remains elusive. They are only
present together with the full-length active R allele regardless of ploidy and clearly dependent
on the presence of the downstream repeat, as they disappear after its removal in Δ and γ

deletion lines. They are usually associated with post-transcriptional silencing (PTGS) [63],
characterized by the 5' preference for U and associated with AGO1 [28] and guide mRNA
cleavage or translational repression [64]. The much stronger *HPT* expression in the deletion
lines lacking 21 nt sRNAs indicates that, indeed, they could be responsible for partial
suppression, but against expectation, the plants remain hygromycin-resistant. In fact, while the
role of 24 nt sRNAs in paramutation is becoming more and more obvious, the stability of *HPT*
expression over more than 20 years and numerous rounds of seed amplification, despite the
presence of the repeat and 21 nt sRNAs, is becoming the bigger enigma.

These plants have been propagated in different locations, over decades, and under a
variety of environmental conditions. Among thousands of scored seedlings of the R and S
lines, we never observed a spontaneous switch of the epialleles from silent to active or from
active to silent state. The latter might have gone undetected but certainly remains below 0.1%.
However, the temperature dependence of paramutation indicates that, after combination of the
epialleles in the same plant, external factors can become highly relevant, with striking similarity
to earlier observations with the R paramutation system in maize [65]. However, involvement of
small RNAs was not described in this maize system, and the expression of the R gene itself is
regulated by light and temperature, which is not the case for the *HPT* gene. In our system, the
prevention of paramutation at lower temperatures resembles the inhibition of RNAi-based viral
defense in the cold [4], and the enhanced paramutation at higher temperatures supports an
important role for the quantity of 24 nt sRNAs whose production is enhanced at higher
temperature in both plants [66] and Paramecium, resulting in heterochromatin formation [67].

In summary, the data presented here for the *HPT* epialleles suggest that the occurrence
and degree of paramutation are determined by multiple factors. The dependence on functional
RdDM, the involvement of repeats, and the influence of environmental conditions overlap with
features described for other paramutation phenomena. This study revealed an additional role
for the copy number of the paramutable allele in the polyploid configuration and determination
of paramutation through a delicate balance between transcription on one side and the nature
and amount of small RNAs on the other. The example of repeated switches between epigenetic
states at the same locus, as reflected in the pedigree of the lines, provides evidence that
spontaneous changes, targeted triggers, as well as genetic mutation events contribute to
heritable stable yet reversible states and thereby to epigenetic diversity. With growing evidence
for the existence of epialleles [68], it is tempting to speculate that paramutation between them
often goes unnoticed and that paramutation may play an underestimated role in population
genetics and evolution.

MATERIAL AND METHODS

Plant material

Arabidopsis thaliana ecotype Zürich (Zh) without (W) or with the transgenic epialleles (R and S) in diploid and tetraploid background were previously described [18]. Derivative lines (Δ) with genetic alterations leading to activation of the S epiallele were identified after failed T-DNA insertion [21]. RdDM mutants in tetraploid background and targeted deletion of the duplicated region were generated by CRISPR mutagenesis (see below). To generate tetraploid derivatives, 14 d-old diploid seedlings grown on plates were submerged in 0.1% colchicine (w/v) for 2 h, washed extensively in tap water and transferred to soil. Seeds from individual plants were harvested separately and sieved through a 350 µm mesh. Larger seeds retained in the sieve were more likely to be polyploid and were grown on soil to maturity. Ploidy of the progeny was evaluated by flow cytometry, by chopping approx. 100 mg of 14 d-old seedlings with a razor blade in 2 ml Galbraith buffer [69]. Nuclei were filtered through a 30 µm nylon mesh (CellTrics, Sysmex Partec, Görlitz, Germany) and pelleted by 10 min centrifugation at 4000 g and 4°C. The pellet was washed in 1 ml Galbraith buffer and centrifuged again. The pellet was resuspended in 500 µl Galbraith buffer with 5 µl of a solution containing 0.1 mg/ml 4',6-diamidino-2-phenylindole (DAPI). After 10 min incubation on ice in the dark, samples were profiled for DAPI signal per nucleus on a BD LSRFortessa' (BD Biosciences, Franklin Lakes, New Jersey, US) and compared to that of diploid and tetraploid control samples. An overview of all plant lines in this study is provided in Fig. 1.

Growth conditions

Plants were regularly grown under long day (LD) conditions (16 h light 19°C, 8 h dark 16°C) and 60% relative humidity. For the temperature experiments with F1 hybrids, plants were grown on soil at standard conditions until development of the first inflorescences (approx. 21 days after germination) and then split into three cohorts: one remained in the standard conditions (19°C), the others were moved to chambers with either 10°C or 24°C under the same LD until seed set. Seeds were harvested, dried, and stratified at 4°C for 48 h before use.

Crossings

All crosses were made after manual emasculation of immature flower buds and pollination by transfer of mature pollen onto the stigma of the prepared recipient. Contaminations were avoided by using sterilized tools and optical control under a binocular. All crosses were made in reciprocal orientation to exclude potential parent-of-origin effects.

Hygromycin resistance assay

Seeds were aliquoted in 2 ml tubes and surface-sterilized for 10 min in a closed box with chlorine gas, generated by mixing 50 ml 14% sodium hypochlorite and 10 ml 37% hydrochloric acid. Seeds were sown on plates containing germination medium (GM,

with or without 20 µg/ml hygromycin B (Calbiochem) and grown for 14 d under long day (LD) conditions (16 h light, 8 h dark 21°C). Plates were incubated horizontally or vertically for root growth assays. Resistance was evaluated by counting green plants with extended leaves and well-developed roots. Resistance ratios were calculated with reference to the number of all germinated seeds. Root length of plants grown on vertical plates was determined with Fiji ((Version 1.51e) [70] and the corresponding SmartRoot plugin (Version 4.21) [71] relative to a ruler and size of the plate.

DNA and RNA extraction

For genotyping, DNA was extracted from one young leaf in a 2 ml tube containing approx. 10 glass beads and 500 µl extraction buffer (200 mM Tris pH 7.5, 250 mM NaCl, 25 mM EDTA) and ground in an MM400 homogenizer (Retsch, Düsseldorf, Germany) for 1 min at 30 Hz. Samples were centrifuged 1 min at 16000 g and 450 µl supernatant was transferred into a 1.5 ml tube containing 450 µl isopropanol and 50 µl sodium acetate pH 5.2. After >1 h at -20°C, precipitates were collected by 20 min centrifugation at 16000 g, washed once in 75% EtOH and air-dried. Pellets were dissolved in 70 µl TE-buffer and stored at 4°C until usage.

For PCR analysis and bisulfite sequencing, DNA was prepared from 0.1 – 1 g plant material ground in liquid nitrogen and suspended in 0.3 – 3 ml CTAB elution buffer (A4150; PanReac AppliChem, Darmstadt, Germany, supplied with 1% polyvinyl pyrrolidone (PVP) and 1% β-mercaptoethanol (β-ME)). After adding RNase A (EN0531, Thermo Fisher Scientific, Waltham, Massachusetts, US), samples were shaken 2 h at 65°C followed by phenol-chloroform extraction as described [72]. DNA was precipitated as above and dissolved in an appropriate amount of TE buffer. Concentrations were determined on a Nanodrop ND1000 (Thermo Fisher Scientific, Waltham, Massachusetts, US). DNA samples were stored at -20°C.

RNA was extracted from either 100 mg 14 d-old seedlings or 30 mg 35 d-old flower buds with TRI-Reagent (T9424, Sigma Aldrich, St. Louis, Missouri, US) according to the manufacturer's protocol. Total RNA was eluted in 50 µl RNase-free ddH2O and the concentration determined as for DNA. RNA integrity was analyzed by running aliquots on a 1.5% agarose-TAE gel, before treating 2.5 µg with DNaseI (EN0525, Thermo Fisher Scientific, Waltham, Massachusetts, US) according to the manual. DNAseI was heat-inactivated at 65°C for 10 min in the presence of 10 mM EDTA. DNA-free total RNA (RNA+) was precipitated by adding 1/10 volume sodium acetate pH 5.2 and 3 volumes 100% cold EtOH for > 2 h at -20°C. RNA was pelleted and washed once in 700 µl cold 75% EtOH. Air-dried pellets were dissolved in 40 µl RNase-free ddH2O and stored at -70°C until use.

Quantitative RT-PCR

cDNA was synthesized from total RNA with RevertAid H Minus Reverse Transcriptase (EP0451; Thermo Fisher Scientific, Waltham, Massachusetts, US), with Random Hexamer primers (S0142; Thermo Fisher Scientific, Waltham, Massachusetts, US) according to the manufacturer's recommendation, but extending cDNA synthesis at 42°C for 90 min in the presence of 20 U RiboLock RNase Inhibitor (EO0381; Thermo Scientific). Prior to quantitative real-time PCR (qRT-PCR), absence of genomic DNA contaminations was controlled by 40 cycles PCR without reverse transcriptase (RT) at 60°C annealing and 1 min elongation, or with RT and primers spanning the intron of the UBC28 gene (At1g64230), in 30 cycles with 20 sec elongation. qRT-PCR was performed with FastStart Essential DNA Green Master (Roche; Rotkreuz, Switzerland) with approx. 5 ng of cDNA on a LightCycler96 system (Roche) according to the manufacturer's instructions. Amplification was performed in a two-step protocol with pre-incubation at 95°C for 10 min and 45 cycles alternating between 95°C 10 sec/60°C 30 sec. A final melting cycle at 97°C preceded melting curve analysis with the ΔΔCt method [73] relative to the housekeeping genes *SAND* (At2g28390) or *EIF4A1* (At3g13920). The following primers were used: fw 5' GGGTAAATAGCTGCGCCGATGGTT and rv 5' CACGGCGGGAGATGCAATAGGTC (expression of *HPT*); fw 5' AACTCTATGCAGCATTTGATCCACT and rv 5' TGATTGCATATCTTTATCGCCATC (qRT-PCR normalization to *SAND*/At2g28390); fw 5' ATCCAAGTTGGTGTGTTCTCC and rv 5' GAGTGTCTCGAGCTTCCACTC (qRT-PCR normalization to *EIF4a*/At3g13920). fw 5' TCCAGAAGGATCCTCCAACTTCCTGCAGT and rv 5' ATGGTTACGAGAAAGACACCGCCTGAATA (spanning a small intron in *UBC28*/At1g64230 to exclude contamination of cDNA with genomic DNA).

Northern blotting

mRNA gels were blotted onto Hybond NX membranes (Amersham, UK) by capillary blotting with 20 x SSC overnight. Membranes were washed once in 2 x SSC, air-dried and crosslinked in a UV Stratalinker 2400 (Stratagene, La Jolla, California, US) in Autocrosslink mode (DNA and mRNA blots). To visualize size markers and to confirm equal loading, crosslinked membranes (DNA and mRNA blots) were stained for 5 min at RT with methylene blue staining solution (0.04% methylene blue [w/v] in 0.5 M sodium acetate pH 5.2) and washed in ddH2O. Methylene blue stain was removed by incubation at RT for 20 min in 2 x SSC with 1% SDS on a shaker and then air-dried.

Probes for nucleic acid blots were either made from oligonucleotides, generated by PCR amplification of template DNA of the corresponding region, or obtained from plasmids via restriction digest. Twenty-five mM oligonucleotide were end-labelled with T4 Polynucleotide Kinase (PNK; EK0031; Thermo Scientific) according to the manual with 25 µCi [γ-32P]ATP (SRP-501; Hartmann Analytic; Braunschweig; Germany) for 45 min at 37°C. Radioactive

labelled oligonucleotide probes were purified with Illustra MicroSpin G-25 micro columns (GE

Healthcare, Chalfont St Giles, UK). Amplicons or restrictions fragments were gel-purified

(D4002; Zymo, Irvine, California, US) and diluted to 25 ng/µl. Twenty-five ng were labelled with

50 µCi [α-32P]dCTP (SRP-305; Hartmann Analytic; Braunschweig; Germany) with the

Amersham Rediprime II Random Prime Labeling System (RPN1633; GE Healthcare, Chalfont

St Giles, UK) and purified on Illustra G-50 ProbeQuant micro columns (GE Healthcare,

Chalfont St Giles, UK). Prior to hybridization, the longer, double-stranded probes were

denatured for 5 min at 99°C.

Blot were hybridized as described [74] in an hybridization oven overnight at 42°C for oligo-

probes or 60°C for longer probes. Membranes were washed twice for 20 min each in 2 x SSC

+ 2% SDS at 50°C for oligo-probes or 65°C for longer probes, followed by 15 min in 2 x SSC

+0.2% SDS at the respective temperatures. Membranes were wrapped in Saran foil and

exposed to phosphoscreens (GE Healthcare, Chalfont St Giles, UK) for up to 3 days. Screens

were scanned on a phosphoimager (Typhoon FLA 9500, GE Healthcare, Chalfont St Giles,

UK).

Preparation and sequencing of sRNAs

Sequencing libraries were prepared from either 500 ng total RNA or 3 µl TraPR-enriched [75]

sRNA with the NEBNext Multiplex Small RNA Library Prep Set for Illumina (E7300; NEB,

Ipswich, Massachusetts) according to the manufacturer's protocol. Libraries were amplified in

14 cycles and cleaned up with AMPure XP beads (Beckman Coulter, Brea, California, US)

prior to size selection on 6% polyacrylamide gels. cDNA libraries from approx. 140 bp to 150 bp

(miRNAs/siRNAs including adapters) were gel-eluted as described in the manufacturer's

protocol, EtOH-precipitated at -20°C overnight and eluted in 12 µl TE buffer. Size range of the

libraries was confirmed on a Fragment Analyzer (Agilent formerly Advanced Analytical, Santa

Clara, California, US) with the HS NGS Fragment Kit (1-6000 bp, DNF-474, Agilent, formerly

Advanced Analytical, Santa Clara, California, US). sRNAs were sequenced on a HiSeq

2500' system in single end 50 bp mode (Illumina, San Diego, Santa Clara, California, US) by

the VBCF NGS facility.

Plasmid construction and plant transformation

Cloning for subsequent Sanger sequencing and subcloning steps was made with the

CloneJET PCR cloning kit (K1231, Thermo Fisher Scientific, Waltham, Massachusetts, US).

For plant transformation with the recloned epiallele, we used a pSUN vector containing an

oleosin promoter upstream of GFP [76]. For Cas9-expressing vectors, the ubiquitin4-2

promoter (PcUbi4-2) of *Petroselinum crispum* in the binary vector pDE-Cas9 [33] was replaced

with the egg cell-specific promoter EC1.2p [77] and the previously described seed-specific

GFP marker was inserted to generate the pDEECO vector backbone. Two specific sgRNAs

each were expressed from an Arabidopsis U6-26 promoter and processed via the tRNA-based multiplex system [78]. For ectopic silencing constructs, the PPT cassette of pDE-Cas9 containing a minimal 35S promoter sequence was removed, and the SpCas9 coding sequence was replaced by the PDK/catalase intron sequence with flanking multiple cloning sites from the pHELLSGATE12 vector [79], resulting in the pDEPPi backbone. Sequences matching different regions of the epiallele were inserted in sense and antisense direction to result in hairpin-forming transcripts. All vectors were transformed into E. coli strains DH5α. Plasmids were prepared with Zyppy Plasmid Miniprep Kit (D4020, Zymo, Irvine, California, US) and eluted in 50 µl ddH2O. Plasmids were controlled by Sanger sequencing before being transformed into electrocompetent *Agrobacterium tumefaciens* strain GV3101 pMP90. Arabidopsis plants were transformed with the floral dip method [80] and transformants selected among seeds of the T0 plants based on the GFP signal in seeds as described [76].

CRISPR mutagenesis

sgRNAs were designed with Chopchop [81-83] and ranked based on their efficiency score and suitable target sites. Prior to cloning the vectors for deletion mutagenesis, we tested several sgRNA candidates for efficient cleavage at the target site in an *in vitro* assay, incubating reconstituted sgRNA-Cas9 ribonucleoprotein with amplicons containing the target site [84]. For genotyping potential mutant candidates, the *in vitro* assay was modified by replacing the wild type amplicon with that of pooled genomic DNA from candidate plants [84]. PCR products were analyzed by electrophoresis, and the analysis repeated for individual plants in positive pools. Amplicons around the mutated site were further analyzed by Sanger sequencing to determine the type of mutation.

sRNA mapping and profiling

From the reads of the sRNAs libraries, adapters were trimmed and reads were size-selected to the length of 18 nt–26 nt by using cutadapt (Version 1.9.1; https://doi.org/10.14806/ej.17.1.200), with options -a AGATCGGA-g CGACGATC -m 18 -M 26 -discard-untrimmed. Read quality and size distribution was controlled using FastQC (Version 0.11.5; http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were aligned to the TAIR10 genome [85] including an extra contig containing the sequence of the transgenic epiallele plus upstream and downstream flanking regions. Alignment was done using Bowtie2 [86] with the options -k 500 -no-unal.

Annotation was done by comparison of genomic positions of small RNA sequences using the Araport11 and repeat-masker databases. For small RNAs with multiple genomic locations, an annotation was attributed given a priority order.

For small RNA profiles, reads aligned to the transgenic epiallele and flanking regions were retrieved and used to calculate the normalized read counts (reads per millions mapped reads) for each nucleotide and to produce a graphical representation using R.

Procedures and scripts used for bioinformatic analysis and graphical representation are available upon request.

sRNA sequencing data are deposited in the Gene Expression Omnibus database (https://www.ncbi.nlm.nih.gov/geo/) under the project number (waiting for verification).

ACKNOWLEDGEMENTS

We are grateful for helpful discussions within our group and colleagues on campus, especially Arturo Marí-Ordóñez, Peter Schlögelhofer, and Michael Nodine. We thank the lab of Olivier Voinnet for suggestions and discussions, and in particular Thomas Grentzinger for supplying the TraPR columns used for sRNA purification ahead of publication and Alexis Sarazin who provided an in-house sRNA-sequencing data processing pipeline. We are also grateful to Helene Fasching and Emi Miyakoda for help with resistance assays and genotyping. We thank the campus facilities, especially Next Generation Sequencing and Plant Facilities from the VBCF and MolBiol Service and BioOptics from the IMP/IMBA/GMI facilities for their excellent support. We thank Arturo Marí-Ordóñez, Liam Dolan, Vincent Castric, and J. Matthew Watson for valuable comments on the manuscript. We further appreciate the financial support by the Austrian Science Fund (FWF DK1238) to the Doctoral Program "Chromosome Dynamics".

REFERENCES

1. Brink RA. Paramutation at the R locus in maize. Cold Spring Harb Symp Quant Biol. 1958;23:379-91. PubMed PMID: 13635569.

2. Coe EH. A regular and continuing conversion-type phenomenon at the B-locus in maize. Proceedings of the National Academy of Sciences of the United States of America. 1959;45(6):828-32. doi: 10.1073/pnas.45.6.828. PubMed PMID: WOS:A1959WJ53600010.

3. Hagemann R. Somatische Konversion bei *Lycopersicon esculentum* Mill. Zeitschrift für Vererbungslehre. 1958;89:587-613.

4. Szittya G, Silhavy D, Molnar A, Havelda Z, Lovas A, Lakatos L, et al. Low temperature inhibits RNA silencing-mediated defence by the control of siRNA generation. Embo J. 2003;22(3):633-40. PubMed PMID: 12554663.

5. Bateson W, Pellew C. On the genetics of "rogues" among culinary peas (Pisum sativum). Journal of Genetics. 1915;5(I):13-36. doi: 10.1007/bf02982150. PubMed PMID: WOS:000200607000002.

6. Santo TE, Pereira RJ, Leitao JM. The pea (Pisum sativum L.) rogue paramutation is accompanied by alterations in the methylation pattern of specific genomic sequences. Epigenomes. 2017;1(1). doi: 10.3390/epigenomes1010006.

7. Pilu R. Paramutation phenomena in plants. Semin Cell Dev Biol. 2015;44:2-10. PubMed PMID: WOS:000365186700002.

8. Ronsseray S. Paramutation phenomena in non-vertebrate animals. Semin Cell Dev Biol.
2015;44:39-46. doi: 10.1016/j.semcdb.2015.08.009. PubMed PMID: WOS:000365186700006.

9. Gabriel JM, Hollick JB. Paramutation in maize and related behaviors in metazoans. Semin Cell
Dev   Biol.   2015;44:11-21.   doi: 10.1016/j.semcdb.2015.08.008.   PubMed   PMID:
WOS:000365186700003.

10. Gouil Q, Baulcombe DC. Paramutation-like features of multiple natural epialleles in tomato. BMC
Genomics. 2018;19(1):203. Epub 2018/03/21. doi: 10.1186/s12864-018-4590-4. PubMed PMID:
29554868; PubMed Central PMCID: PMCPMC5859443.

11. Hollick JB, Chandler VL. Genetic factors required to maintain repression of a paramutagenic
maize pl1 allele. Genetics. 2001;157(1):369-78. PubMed PMID: 11139517.

12. Dorweiler JE, Carey CC, Kubo KM, Hollick JB, Kermicle JL, Chandler VL. mediator of
paramutation1 is required for establishment and maintenance of paramutation at multiple maize
loci. Plant Cell. 2000;12(11):2101-18. PubMed PMID: 11090212.

13. Giacopelli BJ, Hollick JB. Trans-homolog interactions facilitating paramutation in maize. Plant
Physiol. 2015;168(4):1226-36. Epub 2015/07/08. doi: 10.1104/pp.15.00591. PubMed PMID:
26149572; PubMed Central PMCID: PMCPMC4528761.

14. Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in
plants and animals. Nat Rev Genet. 2010;11(3):204-20. PubMed PMID: 20142834.

15. Cuerda-Gil D, Slotkin RK. Non-canonical RNA-directed DNA methylation. Nat Plants.
2016;2(11):16163. doi: 10.1038/nplants.2016.163. PubMed PMID: 27808230.

16. Erdmann RM, Picard CL. RNA-directed DNA Methylation. PLoS Genet. 2020;16(10):e1009034.
Epub 2020/10/09. doi: 10.1371/journal.pgen.1009034. PubMed PMID: 33031395; PubMed
Central PMCID: PMCPMC7544125.

17. Hovel I, Pearson NA, Stam M. Cis-acting determinants of paramutation. Semin Cell Dev Biol.
2015;44:22-32. doi: 10.1016/j.semcdb.2015.08.012. PubMed PMID: WOS:000365186700004.

18. Mittelsten Scheid O, Afsar K, Paszkowski J. Formation of stable epialleles and their paramutation-
like interaction in tetraploid Arabidopsis thaliana. Nat Genet. 2003;34(4):450-4. PubMed PMID:
12847525.

19. Baubec T, Dinh HQ, Pecinka A, Rakic B, Rozhon W, Wohlrab B, et al. Cooperation of multiple
chromatin modifications can generate unanticipated stability of epigenetic States in Arabidopsis.
Plant Cell. 2010;22(1):34-47. Epub 2010/01/26. doi: 10.1105/tpc.109.072819. PubMed PMID:
20097869; PubMed Central PMCID: PMC2828703.

20. Hetzl J, Foerster AM, Raidl G, Mittelsten Scheid O. CyMATE: a new tool for methylation analysis
of plant genornic DNA after bisulphite sequencing. Plant Journal. 2007;51(3):526-36. PubMed
PMID: WOS:000248790500015.

21. Foerster AM, Dinh HQ, Sedman L, Wohlrab B, Mittelsten Scheid O. Genetic rearrangements can
modify chromatin features at epialleles. Plos Genetics. 2011;7(10):e1002331. PubMed PMID:
MEDLINE:22028669.

22. Patterson GI, Chandler VL. Paramutation in maize and related allelic interactions. Curr Top
Microbiol Immunol. 1995;197:121-41. PubMed PMID: 7493488.

23. Karesch H, Bilang R, Mittelsten Scheid O, Potrykus I. Direct gene transfer to protoplasts of
Arabidopsis thaliana. Plant Cell Reports. 1991;9:571-4.

24. Ciabrelli F, Comoglio F, Fellous S, Bonev B, Ninova M, Szabo Q, et al. Stable Polycomb-
dependent transgenerational inheritance of chromatin states in Drosophila. Nat Genet.
2017;49(6):876-86. doi: 10.1038/ng.3848. PubMed PMID: 28436983; PubMed Central PMCID:
PMCPMC5484582.

25. Chandler VL, Eggleston WB, Dorweiler JE. Paramutation in maize. Plant Mol Biol. 2000;43(2-
3):121-45. PubMed PMID: 10999400.

26. Hollick JB, Patterson GI, Asmundsson IM, Chandler VL. Paramutation alters regulatory control of
the maize pl locus. Genetics. 2000;154(4):1827-38. PubMed PMID: 10747073.

27. Jiang W, Li Z, Yao X, Zheng B, Shen WH, Dong A. jaw-1D: a gain-of-function mutation responsive
to paramutation-like induction of epigenetic silencing. J Exp Bot. 2019;70(2):459-68. doi:
10.1093/jxb/ery365. PubMed PMID: 30346598; PubMed Central PMCID: PMCPMC6322565.

28. Mi SJ, Cai T, Hu YG, Chen Y, Hodges E, Ni FR, et al. Sorting of small RNAs into Arabidopsis
argonaute complexes is directed by the 5 ' terminal nucleotide. Cell. 2008;133(1):116-27. doi:
10.1016/j.cell.2008.02.034. PubMed PMID: WOS:000254641300013.

29. Zhang Z, Liu X, Guo X, Wang XJ, Zhang X. Arabidopsis AGO3 predominantly recruits 24-nt small
RNAs to regulate epigenetic silencing. Nat Plants. 2016;2(5):16049. doi:
10.1038/nplants.2016.49. PubMed PMID: 27243648.

30. Wassenegger M, Heimes S, Riedel L, Sanger HL. RNA-directed de novo methylation of genomic
sequences in plants. Cell. 1994;76(3):567-76. doi: 10.1016/0092-8674(94)90119-8. PubMed
PMID: WOS:A1994MX20700016.

31. Matzke MA, Mosher RA. RNA-directed DNA methylation: an epigenetic pathway of increasing
complexity. Nat Rev Genet. 2014;15(6):394-408. doi: 10.1038/nrg3683. PubMed PMID:
24805120.

32. Hollick JB. Paramutation and related phenomena in diverse species. Nat Rev Genet.
2017;18(1):5-23. Epub 2016/11/01. doi: 10.1038/nrg.2016.115. PubMed PMID: 27748375.

33. Fauser F, Schiml S, Puchta H. Both CRISPR/Cas-based nucleases and nickases can be used
efficiently for genome engineering in Arabidopsis thaliana. Plant Journal. 2014;79(2):348-59. doi:
10.1111/tpj.12554. PubMed PMID: WOS:000339439200014.

34. Stonaker JL, Lim JP, Erhard KF, Jr., Hollick JB. Diversity of Pol IV function is defined by mutations
at the maize rmr7 locus. PLoS Genet. 2009;5(11):e1000706. doi: 10.1371/journal.pgen.1000706.
PubMed PMID: 19936246; PubMed Central PMCID: PMCPMC2775721.

35. Sidorenko L, Dorweiler JE, Cigan AM, Arteaga-Vazquez M, Vyas M, Kermicle J, et al. A dominant
mutation in mediator of paramutation2, one of three second-largest subunits of a plant-specific
RNA polymerase, disrupts multiple siRNA silencing processes. PLoS Genet.
2009;5(11):e1000725. Epub 2009/11/26. doi: 10.1371/journal.pgen.1000725. PubMed PMID:
19936058; PubMed Central PMCID: PMCPMC2774164.

36. Sloan AE, Sidorenko L, McGinnis KM. Diverse gene-silencing mechanisms with distinct
requirements for RNA polymerase subunits in Zea mays. Genetics. 2014;198(3):1031-42. Epub
2014/08/29. doi: 10.1534/genetics.114.168518. PubMed PMID: 25164883; PubMed Central
PMCID: PMCPMC4224150.

37. Erhard KF, Jr., Stonaker JL, Parkinson SE, Lim JP, Hale CJ, Hollick JB. RNA polymerase IV
functions in paramutation in Zea mays. Science. 2009;323(5918):1201-5. PubMed PMID:
19251626.

38. Panavas T, Weir J, Walker EL. The structure and paramutagenicity of the R-marbled haplotype
of Zea mays. Genetics. 1999;153(2):979-91. Epub 1999/10/08. PubMed PMID: 10511572;
PubMed Central PMCID: PMCPMC1460783.

39. Cantu D, Vanzetti LS, Sumner A, Dubcovsky M, Matvienko M, Distelfeld A, et al. Small RNAs,
DNA methylation and transposable elements in wheat. BMC Genomics. 2010;11:408. Epub
2010/06/30. doi: 10.1186/1471-2164-11-408. PubMed PMID: 20584339; PubMed Central
PMCID: PMCPMC2996936.

40. Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, et al. The Arabidopsis
nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing
heterochromatin. Cell. 2013;153(1):193-205. Epub 2013/04/02. doi: 10.1016/j.cell.2013.02.033.

41. McCue AD, Panda K, Nuthikattu S, Choudury SG, Thomas EN, Slotkin RK. ARGONAUTE 6
bridges transposable element mRNA-derived siRNAs to the establishment of DNA methylation.
Embo j. 2015;34(1):20-35. Epub 2014/11/13. doi: 10.15252/embj.201489499. PubMed PMID:
25388951; PubMed Central PMCID: PMCPMC4291478.

42. Li C, Xu H, Fu FF, Russell SD, Sundaresan V, Gent JI. Genome-wide redistribution of 24-nt
siRNAs in rice gametes. Genome Res. 2020;30(2):173-84. Epub 2020/01/04. doi:
10.1101/gr.253674.119. PubMed PMID: 31896557; PubMed Central PMCID: PMCPMC7050532.

43. Berg W, Hagemann R. Paramutation am *sulfurea* Locus von *Lycopersicon esculentum* Mill. VIII.
Der Dosiseffekt in Tetraploiden. Biologisches Zentralblatt. 1979;98:307-14.

44. Berg W, Hagemann R. Paramutation aom *sulfurea* Locus von *Lycopersicon esculentum* Mill. IX.
Schrittweise Paramutation in Tetraploiden. Biologisches Zentralblatt. 1980;99:25-36.

45. Comai L, Tyagi AP, Winter K, Holmes-Davis R, Reynolds SH, Stevens Y, et al. Phenotypic
instability and rapid gene silencing in newly formed arabidopsis allotetraploids. Plant Cell.
2000;12(9):1551-68. PubMed PMID: 11006331.

46. Madlung A, Masuelli RW, Watson B, Reynolds SH, Davison J, Comai L. Remodeling of DNA
methylation and phenotypic and transcriptional changes in synthetic Arabidopsis allotetraploids.
Plant Physiol. 2002;129(2):733-46. PubMed PMID: 12068115.

47. Wendel JF, Lisch D, Hu GQ, Mason AS. The long and short of doubling down: polyploidy,
epigenetics, and the temporal dynamics of genome fractionation. Current Opinions in Genetics
and Development. 2018;49:1-7.

48. Ding M, Chen ZJ. Epigenetic perspectives on the evolution and domestication of polyploid plant
and crops. Current Opinions in Genetics and Development. 2018;42:37-48.

49. Tayale A, Parisod C. Natural pathways to polyploidy in plants and consequences for genome
reorganization. Cytogenetic and Genome Research. 2013;140:79-96.

50. Yan H, Bombarely A, Xu B, Wu B, Frazier TP, Zhang X, et al. Autopolyploidization in switchgrass
alters phenotype and flowering time via epigenetic and transcription regulation. Journal of
Experimental Botany. 2019;70:5673-86.

51. Yu Z, Haberer G, Matthes M, Rattei T, Mayer KF, Gierl A, et al. Impact of natural genetic variation
on the transcriptome of autotetraploid Arabidopsis thaliana. Proc Natl Acad Sci U S A.
2010;107(41):17809-14. Epub 2010/09/30. doi: 10.1073/pnas.1000852107. PubMed PMID:
20876110; PubMed Central PMCID: PMCPMC2955106.

52. Wang JL, Tian L, Madlung A, Lee HS, Chen M, Lee JJ, et al. Stochastic and epigenetic changes
of gene expression in arabidopsis polyploids. Genetics. 2004;167(4):1961-73. PubMed PMID:
WOS:000223720300039.

53. Aversano R, Caruso I, Aronne G, De Micco V, Scognamiglio N, Carputo D. Stochastic changes
affect Solanum wild species following autopolyploidization. J Exp Bot. 2013;64(2):625-35. Epub
2013/01/12. doi: 10.1093/jxb/ers357. PubMed PMID: 23307917; PubMed Central PMCID:
PMCPMC3542052.

54. Sas-Nowosielska H, Bernas T. Spatial relationship between chromosomal domains in diploid and
autotetraploid Arabidopsis thaliana nuclei. Nucleus-Phila. 2016;7(2):216-31. Epub 2016/06/17.
doi: 10.1080/19491034.2016.1182277. PubMed PMID: 27310308; PubMed Central PMCID:
PMCPMC4916897.

55. Li X, Yu E, Fan C, Zhang C, Fu T, Zhou Y. Developmental, cytological and transcriptional analysis
of autotetraploid Arabidopsis. Planta. 2012;236(2):579-96. Epub 2012/04/06. doi:
10.1007/s00425-012-1629-7. PubMed PMID: 22476290.

56. Lloyd A, Bomblies K. Meiosis in autopolyploid and allopolyploid Arabidopsis. Curr Opin Plant Biol.
2016;30:116-22. Epub 2016/03/08. doi: 10.1016/j.pbi.2016.02.004. PubMed PMID: 26950252.

57. Belele CL, Sidorenko L, Stam M, Bader R, Arteaga-Vazquez MA, Chandler VL. Specific tandem
repeats are sufficient for paramutation-induced trans-generational silencing. PLoS Genetics.
2013;9(10). doi: 10.1371/journal.pgen.1003773.

58. Stam M, Belele C, Dorweiler JE, Chandler VL. Differential chromatin structure within a tandem
array 100 kb upstream of the maize b1 locus is associated with paramutation. Genes Dev.
2002;16(15):1906-18. PubMed PMID: 12154122.

59. Marí-Ordóñez A, Marchais A, Etcheverry M, Martin A, Colot V, Voinnet O. Reconstructing de novo
silencing of an active plant retrotransposon. Nature Genetics. 2013;published online.

60. Barbour JE, Liao IT, Stonaker JL, Lim JP, Lee CC, Parkinson SE, et al. required to maintain
repression2 is a novel protein that facilitates locus-specific paramutation in maize. Plant Cell.
2012;24(5):1761-75. Epub 2012/05/09. doi: 10.1105/tpc.112.097618. PubMed PMID: 22562610;
PubMed Central PMCID: PMCPMC3442568.

61. Arteaga-Vazquez M, Sidorenko L, Rabanal FA, Shrivistava R, Nobuta K, Green PJ, et al. RNA-
mediated trans-communication can establish paramutation at the b1 locus in maize. Proc Natl
Acad Sci U S A. 2010;107(29):12986-91. PubMed PMID: 20616013.

62. Regulski M, Lu Z, Kendall J, Donoghue MT, Reinders J, Llaca V, et al. The maize methylome
influences mRNA splice sites and reveals widespread paramutation-like switches guided by small
RNA. Genome Res. 2013;23(10):1651-62. Epub 2013/06/07. doi: 10.1101/gr.153510.112.
PubMed PMID: 23739895; PubMed Central PMCID: PMCPMC3787262.

63. Vaucheret H, Beclin C, Fagard M. Post-transcriptional gene silencing in plants. Journal of Cell
Science. 2001;114(17):3083-91. PubMed PMID: WOS:000171266100006.

64. Bologna NG, Voinnet O. The diversity, biogenesis, and activities of endogenous silencing small
RNAs in Arabidopsis. Annu Rev Plant Biol. 2014;65:473-503. Epub 2014/03/04. doi:
10.1146/annurev-arplant-050213-035728. PubMed PMID: 24579988.

65. Mikula BC. Environmental programming of heritable epigenetic changes in paramutant r-gene
expression using temperature and light at a specific stage of early development in maize
seedlings. Genetics. 1995;140:1379-87.

66. Baev V, Milev I, Naydenov M, Vachev T, Apostolova E, Mehterov N, et al. Insight into small RNA
abundance and expression in high- and low-temperature stress response using deep sequencing
in Arabidopsis. Plant Physiol Biochem. 2014;84:105-14. Epub 2014/09/28. doi:
10.1016/j.plaphy.2014.09.007. PubMed PMID: 25261853.

67. Pirritano M, Götz U, Karunanithi S, Nordström K, Schulz MH, Simon M. Environmental
temperature controls accumulation of transacting siRNAs involved in heterochromatin formation.
Genes. 2018;9(2). Epub 2018/02/22. doi: 10.3390/genes9020117. PubMed PMID: 29466322;
PubMed Central PMCID: PMCPMC5852613.

68. Minow MAA, Colasanti J. Does variable epigenetic inheritance fuel plant evolution? Genome.
2020;63(5):253-62. Epub 2020/02/14. doi: 10.1139/gen-2019-0190. PubMed PMID: 32053387.

69. Galbraith DW, Harkins KR, Maddox JM, Ayres NM, Sharma DP, Firoozabady E. Rapid flow
cytometric analysis of the cell cycle in intact plant tissues. Science. 1983;220(4601):1049-51.

70. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub 2012/06/30.
doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID:
PMCPMC3855844.

71. Lobet G, Pagès L, Draye X. A novel image-analysis toolbox enabling quantitative analysis of root
system architecture. Plant Physiol. 2011;157(1):29-39. Epub 2011/07/21. doi:
10.1104/pp.111.179895. PubMed PMID: 21771915; PubMed Central PMCID:
PMCPMC3165877.

72. Clarke JD. Cetyltrimethyl ammonium bromide (CTAB) DNA miniprep for plant DNA isolation. Cold
Spring Harb Protoc. 2009;2009(3). doi: 10.1101/pdb.prot5177.

73. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative
PCR and the 2(T)(-Delta Delta C) method. Methods. 2001;25(4):402-8. doi:
10.1006/meth.2001.1262. PubMed PMID: WOS:000173949500003.

74. Church GM, Gilbert W. Genomic sequencing. Proceedings of the National Academy of Sciences
of the United States of America-Biological Sciences. 1984;81(7):1991-5. PubMed PMID:
WOS:A1984SQ01800013.

75. Grentzinger T, Oberlin S, Schott G, Handler D, Svozil J, Barragan-Borrero V, et al. A universal
method for the rapid isolation of all known classes of functional silencing small RNAs. Nucleic
Acids Res. 2020;48(14):e79. Epub 2020/06/05. doi: 10.1093/nar/gkaa472. PubMed PMID:
32496553.

76. Shimada TL, Shimada T, Hara-Nishimura I. A rapid and non-destructive screenable marker,
FAST, for identifying transformed seeds of Arabidopsis thaliana. Plant J. 2010;61(3):519-28.
Epub 2009/11/07. doi: 10.1111/j.1365-313X.2009.04060.x. PubMed PMID: 19891705.

77. Wang ZP, Xing HL, Dong L, Zhang HY, Han CY, Wang XC, et al. Egg cell-specific promoter-
controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple target genes in
Arabidopsis in a single generation. Genome Biology. 2015;16. doi: 10.1186/s13059-015-0715-0.
PubMed PMID: WOS:000358117600002.

78. Xie KB, Minkenberg B, Yang YN. Boosting CRISPR/Cas9 multiplex editing capability with the
endogenous tRNA-processing system. Proceedings of the National Academy of Sciences of the
United States of America. 2015;112(11):3570-5. doi: 10.1073/pnas.1420294112. PubMed PMID:
WOS:000351060000101.

79. Helliwell C, Waterhouse P. Constructs and methods for high-throughput gene silencing in plants.
Methods. 2003;30(4):289-95. Epub 2003/06/28. doi: 10.1016/s1046-2023(03)00036-7. PubMed
PMID: 12828942.

80. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated transformation
of Arabidopsis thaliana. Plant Journal. 1998;16(6):735-43. doi: 10.1046/j.1365-
313x.1998.00343.x. PubMed PMID: WOS:000078001900010.

81. Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool for the
next generation of CRISPR genome engineering. Nucleic Acids Research. 2016;44(W1):W272-
W6. doi: 10.1093/nar/gkw398. PubMed PMID: WOS:000379786800045.

82. Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E. CHOPCHOP v3:
expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Res.
2019;47(W1):W171-w4. Epub 2019/05/21. doi: 10.1093/nar/gkz365. PubMed PMID: 31106371;
PubMed Central PMCID: PMCPMC6602426.

83. Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a CRISPR/Cas9 and
TALEN web tool for genome editing. Nucleic Acids Research. 2014;42(W1):W401-W7. doi:
10.1093/nar/gku410. PubMed PMID: WOS:000339715000067.

84. Bente H, Mittelsten Scheid O, Donà M. Versatile in vitro assay to recognize Cas9-induved
mutations. Plant Direct. 2020. doi: 10.1002/pld3.269.

85. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al. The Arabidopsis information
resource: Making and mining the "gold standard" annotated reference plant genome. Genesis.
2015;53(8):474-85. Epub 2015/07/24. doi: 10.1002/dvg.22877. PubMed PMID: 26201819;
PubMed Central PMCID: PMCPMC4545719.

86. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.
2012;9(4):357-U54. doi: 10.1038/nmeth.1923. PubMed PMID: WOS:000302218500017.

    FIGURE LEGENDS

Figure 1. Overview over structure of epialleles and origin of plant material.

(A) Location of the epialleles on chromosome three, their structural variants, and the location

of primers and probes. HPT: ORF of hygromycin phosphotransferase; ΔT: incomplete CaMV

35S terminator; P1, P2: CaMV 35S promoter; V1, V2: vector plasmid sequence; NC: non-

coding sequence; S: silent epiallele rendering plants sensitive (red); R: active epiallele

rendering plants resistant (blue) to hygromycin; Δ, γ: deletion derivatives. (B) Pedigree of lines.

S, R, Δ, γ as in (A), W: wild type allele without insert (black); 2 letters: diploids; 4 letters:

tetraploids; x: crossing; ⊗ selfing; colchicine: generation of tetraploids, * converted by

paramutation. Links to other figures refer to the use of the material in different context.

Figure 2. The degree of paramutation in tetraploids depends on growth temperature in

F1. (A) Experimental setup: Diploid or tetraploid plants with R epialleles were crossed with

those containing S or W, and F1 hybrids from reciprocal crossings were grown at 19°C for

three weeks before transfer to either 10°C, 19°C, or 24°C until seed maturity. F2 seeds were

germinated at 19°C on GM plates containing 20 mg/L hygromycin B and resistance ratios

determined after 14 days for tetraploids (B) and diploids (C). Data for reciprocal crosses were

combined, as no parent-of-origin difference was observed. N = number of tested seedlings in

each group. Bars represent the mean from two biological with three technical replicates each

(B) and one biological with three technical replicates (C), with 100 plated seeds (B) each, or

50 plated seeds (C). Error bars indicate standard deviation. Dashed lines represent the

expected segregation for tetraploid (B, 35:1 ≙ 97.2%) and diploid F2 populations (C, 3:1 ≙

75%). F1 growth temperature is indicated by colour: 10°C (blue), 19°C (green), 24°C (red);

light colours: F2 of R/S hybrids, dark colours: F2 of R/W hybrids. Statistical analysis was

performed by summed Chi-square goodness-of-fit test with the indicated values.

Figure 3. Epialleles are associated with distinct classes of sRNAs. (A) Proportion of sRNA

length from all reads mapped to the epialleles in 14 day-old seedlings with tetraploid active

(RRRR) or silenced (SSSS) epialleles. See S1 Fig for the corresponding data from diploids.

(B) Coverage plots of 18 – 26 nt sRNA along the RRRR or SSSS epialleles. (C) Proportion of

5' prime nucleotides of epiallele-specific sRNAs in RRRR (left) and SSSS (right) plotted by

size. See S1 Fig for the corresponding data for reads mapped to the genome.

Figure 4. Lack of paramutation in the background of RdDM mutants. (A) F2 seeds

obtained by selfing tetraploid F1 hybrids were germinated on GM plates containing 20 mg/L

hygromycin B and resistance ratios determined after 14 days. RRWW control (white),

paramutation test hybrid RRSS in wild type (grey), *nrpd1* mutant (blue) or *rdr2* mutant (red).

Data for reciprocal crosses were combined. N = number of tested seedlings in each group.

The dashed line represents the expected F2 segregation for tetraploids (97.2%), statistical

analysis is based on a summed Chi-square goodness-of-fit test with indicated values. (B) Representative images for resistance assays quantified in (A). Orange asterisks indicate sensitive seedlings, scale bar = 3 cm.

Figure 5. Ectopic RNAi expression reduces *HPT* expression to different extent in diploids and tetraploids. (A) Three constructs with constitutive expression of hairpin-forming palindromes homologous to either the vector region of the R epiallele (hp1, green; hp2, grey) or the promoter region (hp3, blue) were inserted in ectopic positions, and their transcripts are processed into sRNAs. (A) Box plots of root length (measured with SmartRoot software) as indicator of resistance in the presence of 20 mg/L hygromycin in / d-old T1 seedlings grown from seeds of T0 plants with or without the hairpin construct (sorted by seed-specific GFP fluorescence provided by the construct and grown on the same plate). White diamonds indicate the mean of each group. Statistical analysis was made by Student's T-test, p-values of relevant comparisons are indicated. N = number of tested seedlings in each group. (B) Representative images for root growth in lines without (upper panel) and with (lower panel) the hairpin construct, scale bar = 1 cm. (C) Relative expression of *HPT* in flower bud tissue of hairpin-containing T1 plants in tetraploids (left panel) and diploids (right panel), compared to sibling plants without hairpins. Measurements were done in three biological and three technical replicates. Statistical analysis was performed by doing Student's T-test (* = p ≤ 0.05; ** = p ≤ 0.01; *** = p ≤ 0.001).

Figure 6. Paramutation depends on epiallele structure, and paramutated epialleles can exert secondary paramutation. (A) Tetraploid F2 seeds obtained by selfing F1 hybrids of crosses between S/W/R and RΔ2 (light grey) or RΔ4 (dark grey) were germinated on GM plates containing 20 mg/L hygromycin B and resistance ratios determined after 14 days. Data for reciprocal crosses were combined. N = number of tested seedlings in each group. The dashed line represents the expected F2 segregation for tetraploids (97.2%), statistical analysis is based on a summed Chi-square goodness-of-fit test with indicated values. (B) RΔ4 deletion alleles were combined with either W or S partners (all tetraploid). Individual plants in F2 progeny were genotyped for RΔ4, sorted into groups heterozygous (+/-) or homozygous (+/+) for the deletion and subjected to *HPT* expression analysis relative to homozygous RRRR. Bars indicate the mean of two biological and three technical replicates. Error bars indicate standard deviation. Statistical analysis was performed with Student's T-test, p values for relevant comparisons are indicated. (C) Plants from primary paramutation crosses, homozygous for RΔ4 alleles (excluding the presence of the old S allele) and only residual *HPT* expression (black star in B, now called SΔ4SΔ4SΔ4SΔ4, Fig. 1) were crossed to plants with the active, full length R alleles and analysed for hygromycin as described before in F2, in parallel to controls. The dashed line represents expected F2 segregation for tetraploid populations at

97.2%; statistical analysis is based on a summed Chi-square goodness-of-fit test with indicated

values. N = number of tested seedlings in each group.

Figure 7. Deletion of the downstream repeated region leads to higher expression at

active epialleles and reduces paramutation. (A) Representative images of root growth

assays in the presence of 20 mg/L hygromycin with 7 d-old plants containing full length or

truncated active epialleles. Scale bar = 3 cm. (B) Box plots of root length as in (A) (measured

with SmartRoot software). White diamonds indicate the mean of each group. Statistical

analysis was performed with Student's T-test with indicated p-value. N = number of tested

seedlings in each group. (C) Northern blots of *HPT* transcripts in independent lines of tetraploid

(1-6) and diploid (7-12) plants with full length (1,3,5,7,9,11) or truncated (2,4,6,8,10,12) active

epialleles. (D) Northern blots of *HPT* transcripts in tetraploid (1-7) and diploid (8-14) plants. 1:

WWWW; 2: RRRR; 3: RΔ2RΔ2RΔ2RΔ2; 4: RΔ4RΔ4RΔ4RΔ4; 5: RγRγRγRγ; 6: SSSS; 7:

SγSγSγSγ; 8: WW; 9: RR; 10: RΔ2RΔ2; 11: RΔ4RΔ4; 12: RγRγ; 13: SS; 14: SγSγ. Location

of the probe for blots in C and D is illustrated in Fig. 1. (E) Representative image for hygromycin

resistance assays with 14 d-old F2 plants after crossing parents with full length or truncated

epialleles. Scale bar = 3 cm. (F) Resistance ratios calculated from assays like in (E). Data from

reciprocal crosses were combined, statistical analysis is based on a summed Chi-square

goodness-of-fit test with indicated values. N = number of tested seedlings in each group.

Figure 8. Features of polyploidy-dependent paramutation. (A) The expressed epiallele

(blue) confers hygromycin resistance, although the downstream repeat and the 21 nt sRNA

limit transcription from the upstream promoter to some extent. The silent epiallele (red) is

associated with 24 nt sRNAs and exerts strong silencing *in trans*, involving temperature-

dependent RdDM. (B) Distribution of epigenotypes in F2 populations. Paramutation occurs in

a dosage-dependent way and may be restricted to hybrids with more silent than active copies.

This combination does not occur in diploids and explains the limitation of paramutation to

tetraploids.

SUPPORTING FIGURE LEGENDS

S1 Figure. Epialleles associated with distinct classes of sRNAs in diploids. (A)

Proportion of sRNA length from all mapped reads in 14 day-old seedlings with diploid active

(RR) or silenced (SS) epialleles. (B) Coverage plots of 18 - 26 nt sRNA along the RR or SS

epialleles. (C) Proportion of 5' prime nucleotides of epiallele-specific sRNAs in RR (left) and

SS (right) plotted by size.

S2 Figure. Equal sRNA distribution genome-wide and no sRNAs mapping to the

epiallele sequence in wild type. (A) Proportions of 5' nucleotides in different size classes in

sRNA libraries from tetraploids (left) and diploids (right) in lines with active epialleles (top),

silent epialleles (middle), and wildtype lines (bottom). (B) RNA libraries from tetraploid

(WWWW, left) or diploid (WW, right) plants were attempted to map to the full-length sequence

of the epiallele.

S3 Figure. Effects of the Cas9-generated PolIV mutant on sRNA profiles. (A) Strong

reduction of 24 nt sRNAs in a homozygous loss-of-function mutation in the *NRPD1* gene

(encoding the largest subunit of PolIV) generated by CRISPR in the background of the

tetraploid silent epiallele; including (B) those mapping to the epiallele.

S4 Figure. sRNA profiles along the epialleles in F1 hybrids and in deletion mutants. (A)

Profile from flower bud material in a pair of reciprocal paramutation test hybrids. (B – F) Profiles

of tetraploid (left) and diploid (right) lines with the indicated alleles; (B) silent full length allele;

(C) active allele missing downstream repeat after random mutagenesis and hygromycin

screen; (D) CRISPR-generated deletion of the downstream repeat in the background of the

silent epiallele; (E) active full length allele; (F) CRISPR-generated deletion of the downstream

repeat in the background of the active epiallele. All values are reads per million mapped reads

(RPM), and the y-axes are all set to the same maximum of ± 50 RPM.

1012

Bente et al. Figure 1

Bente et al. Figure 2

Bente et al. Figure 3

A

χ^2 27.43   743.13   41.33   79.45

p $1.6*10^{-7}$ $< 2.2*10^{-16}$ $1.3*10^{-10}$ $< 2.2*10^{-16}$

Hyg resistance [%]

N = 540   N = 580   N = 589   N = 290

F1:   RRWW   RRSS   RRSS *nrpd1*   RRSS *rdr2*

B

F1:   RRSS (control)   RRSS (*nrpd1*)   RRSS (*rdr2*)

Bente et al. Figure 4

Bente et al. Figure 5

Bente et al. Figure 6

Bente et al. Figure 7

A

Repeat-dependent suppression

Silencing of R epiallele

- NRPD1/RDR2 - dependent
- Temperature - dependent

∼ 21 nt sRNAs
∼ 24 nt sRNAs

B

Diploid			Tetraploid		
Epigenotype	Frequency	Paramutation probability	Epigenotype	Frequency	Paramutation probability
‖	25%		‖‖	2.8%	
‖	50%	---	‖‖	22.2%	---
			‖‖	50%	Low
			‖‖	22.2%	High
‖	25%		‖‖	2.8%	

Bente et al. Figure 8

Bente et al. Supplemental Figure 1

Bente et al. Supplemental Figure 2

Bente et al. Supplemental Figure 4

■ 20-21 nt   ■ 22-23 nt   ■ 24-25 nt   ■ Other length