Diet has no effect on susceptibility to an RNA virus across host species

K. E. Roberts* and B. Longdon

Centre for Ecology & Conservation, College of Life and Environmental Sciences,

University of Exeter, Penryn Campus, Penryn, Cornwall, TR10 9FE, UK

*Corresponding author: k.roberts@exeter.ac.uk

Abstract

The likelihood of a successful host shift of a parasite to a novel host

species can be influenced by environmental factors that can act on both the host

and parasite. Factors such as resource availability or temperature can drive

changes in species distributions and abundances making encounters between

infected and susceptible individuals more likely. They can also act as stressors

that can change the physiology of the interacting parties, including altering the

host immune response or varying parasite virulence. Changes in nutritional

resource availability have been shown to alter pathogen susceptibility and the

outcome of infection in a range of systems. Here we infected 27 species of

Drosophilidae with an RNA virus on three different food types of differing

protein to carbohydrate ratios. We then measured how viral load and mortality

across the species was affected by changes in diet. We found that changes in the

ratio of protein to carbohydrate in the diet did not alter the outcomes of

infection, with strong positive inter-species correlations in both viral load and

mortality across diets, and a strong positive correlation between mortality and

viral load.

Introduction

A key driver of pathogen host shifts – where a pathogen jumps from one host

species to another – is environmental change (Carlson et al., 2020; Hoberg &

Brooks, 2015). Ecological factors can influence the likelihood of host shifts by

altering species distributions and abundances making encounters more likely, or

by acting as stressors that alter physiological factors including immunity or

virulence. The main ecological factor studied has been temperature, which can

have asymmetrical impacts on hosts and parasites and potentially altering the

likelihood of host shifts (Brooks & Hoberg, 2007; Hoberg & Brooks, 2015; Kirk et

al., 2018; Roberts, Hadfield, Sharma, & Longdon, 2018). The role of other

ecological traits such as resource availability, humidity, population density and

geographical range, or within host ecological traits such as metabolic rate, have

been less well studied in explaining the outcomes of host shifts, despite an

increasing understanding of the role such factors play in effecting the outcomes

of more simple host parasite interactions (Blanford & Thomas, 1999; Cumnock et

al., 2018; Harvell et al., 2002; Hayman, Pulliam, Marshall, Cryan, & Webb, 2016;

Ponton et al., 2013).

Nutrition can shape the outcome of host-parasite interactions through its ability

to moderate both parasite virulence and host resistance (Pike, Lythgoe, & King,

2019; Ponton, Wilson, Cotter, Raubenheimer, & Simpson, 2011; Ponton et al.,

2013). The nutritional resources of a host can impact its ability to resist infection

as immune responses are thought to be costly to both maintain and activate

(Sheena C. Cotter, Simpson, Raubenheimer, & Wilson, 2011; Knutie, Wilkinson,

Wu, Ortega, & Rohr, 2017; Kraaijeveld & Godfray, 1997; Kutzer & Armitage,

2016a; Lochmiller & Deerenberg, 2000; McKean, Yourth, Lazzaro, & Clark, 2008).

Nutrition is known to have long term consequences, with developmental

nutritional status been known to have latent or even trans-generational effects

on immune responses (Fellous & Lazzaro, 2010; Grueber, Gray, Morris, Simpson,

& Senior, 2018; Savola, Vale, & Walling, 2020). Hosts can also show behavioural

modifications in feeding upon infection; parasite-induced anorexia is thought to

be an adaptive host response (Ayres & Schneider, 2009; Rao et al., 2017) and in

some cases hosts actively increase the consumption of certain nutrients in their

diet (Povey, Cotter, Simpson, Lee, & Wilson, 2009). Nutrition may also constrain

the amount of investment that a host can allocate to an energetically demanding

immune response (Sheena C. Cotter et al., 2011; Knutie et al., 2017; Kraaijeveld &

Godfray, 1997; Lochmiller & Deerenberg, 2000), and subsequent ability to cope

with costs associated with a parasite burden if infection does become established

(Sheldon & Verhulst, 1996). A suboptimal nutritional status may lead a host to be

unable to suppress or tolerate a parasite challenge they may otherwise have

been able to resist; or suffer reduced fitness imposed by trade-offs with

resources from life history traits (Sheena C. Cotter et al., 2011; Knutie et al.,

2017; Kraaijeveld & Godfray, 1997; Lochmiller & Deerenberg, 2000).

From a parasite perspective infecting a host of suboptimal nutritional status may

mean they encounter a weaker immune response and therefore infection and

establishment is easier. However, once established the parasite may encounter

its own resource limitations due to competition with an already depleted host,

causing suboptimal growth and potential onward transmission. Therefore,

predicting the outcome of the interaction between nutrition, host immunity and

subsequent resistance is complex as the effects on the two parties may be

divergent.

Multiple life history traits are moderated by resource availability, with condition

dependency across reproductive traits, aging and lifespan (Camus, Fowler, Piper,

& Reuter, 2017; Y. Henry, Overgaard, & Colinet, 2020; Youn Henry, 2018; Lee,

Simpson, & Wilson, 2008; Maklakov et al., 2009). Laboratory experiments on

dietary restriction, where individuals experienced a reduction in nutrition

without inducing malnutrition (differentiated from Calorie Restriction) have

been found to extend life span in a range of organisms (Anderson et al., 2003;

Klass, 1983; Nakagawa, Lagisz, Hector, & Spencer, 2012; Weindruch & Walford,

1982). The effects of dietary restriction appear to be explained by resource-

mediated trade-offs between longevity and reproductive effort (but see review

by Moatt *et al.*, 2020). Geometric frameworks – the use of artificial diets with

known compositions of specific nutrients that develop an understanding of

dimensional nutrient space – have been used to examine the consequences of

different ratios of macronutrients across a range of organisms (Fanson, Weldon,

Pérez-Staples, Simpson, & Taylor, 2009; Jensen, McClure, Priest, & Hunt, 2015; Le

Couteur et al., 2016; Lee et al., 2008; Raubenheimer & Simpson, 1999; S. J.

Simpson & Raubenheimer, 1995; Stephen J. Simpson & Raubenheimer, 2011;

Skorupa, Dervisefendic, Zwiener, & Pletcher, 2008). Across multiple species low

protein to carbohydrate ratios reduce reproductive rates but appear to maximise

lifespan (Nakagawa et al., 2012). However, individuals with diets higher in

protein and lower in carbohydrates have greater reproductive rates but shorter

life spans. Additionally when given a choice of diet, individuals have been shown

to optimise their reproduction over lifespan (Bunning et al., 2016).

To investigate the effect that host diet has on the likelihood of pathogen host

shifts we infected 27 species of Drosophilidae with Drosophila C Virus (DCV;

Dicistroviridae) fed on three diets with varying ratios of protein to

carbohydrates but comparable calorie content. We then measured the change in

viral load and host mortality across these different diets. DCV can cause

nutritional stress in the midgut of flies causing similar pathologies to those seen

in starvation (Chtarbanova et al., 2014). DCV shows specific tissue tropism of

infection in the visceral muscle cells around the midgut and food storage organ

(crop) causing reduced defecation, food blockage and dehydration/starvation

(Arnold, Johnson, & White, 2013; Chtarbanova et al., 2014). If hosts are in a

nutritional environment that allows for investment in immune function, they

may be more able to resist establishment off, or potentially tolerate, a novel

infection. This could then lead to different outcomes following a host shift

depending on resource environment. Either the host could manage to avoid

infection entirely or could become infected and minimise parasite damage,

tolerating infection whilst withstanding a level of parasite growth. Alternatively

hosts may be fully susceptible to infection, and enriched resources may act to

enhance pathogen virulence by enabling within host pathogen growth. We

therefore examined whether changes in nutritional environment alter the

susceptibility to DCV following a host shift, measuring both viral load and host

mortality, and establishing whether patterns are consistent across species and

diet.

Methods

Diet preparation

Three different cornmeal diets were used in order to establish experimental

diets (Table 1). The standard cornmeal diet used in our lab comprised a 1:10

protein to carbohydrate ratio and became our "Medium" - protein: carbohydrate

ratio diet treatment. We also developed two further diets that were

approximately isocaloric (Table 1), a low protein: carbohydrate food (1: 20

protein to carbohydrate) and a high protein: carbohydrate food (1: 5 protein

carbohydrate). These were based around previous findings that suggested that

in *D. melanogaster* lifespan was maximized on a protein: carbohydrate ratio of

around 1:16, and fitness - measured as lifetime egg production at an

intermediate ratio of 1:4 (Lee et al., 2008). All diets were manipulated by altering

the dextrose and yeast amounts whilst maintaining as close as possible the same

calorie content at 142 Calories g/100 m. Yeast was manipulated as it provides

the majority of the protein as well as other non-caloric nutritional requirements

(Piper, 2017). Values were confirmed using the Drosophila Dietary Composition

Calculator (Lesperance & Broderick, 2020).

Viral Infections

Twenty-seven different species of Drosophilidae were maintained in multi

generation populations, in Drosophila stock bottles (Dutscher Scientific) on 50

ml of their respective food medium at 22℃ and 70% relative humidity with a 12-

hour light-dark cycle (See *Supplementary for species and food*). Each day, two

vials of 0-1 day old male flies (15-22 flies per vial) were randomly assigned to

one of three potential food types; low, medium or high, protein: carbohydrate

ratio (Table 1). Flies were kept on their respective food treatments for 5 days,

and tipped onto fresh vials of food every day (Blum, Fischer, Miles, &

Handelsman, 2013; Broderick & Lemaitre, 2012). After 5 days of acclimatisation

on their food treatment flies were experimentally infected with DCV. These

collections and inoculations were carried out over three replicate blocks, with

each block being completed over consecutive days. The order that the species

were infected was randomized each day, as was food treatment for each species.

We used Drosophila C virus (DCV) strain B6A (Longdon et al., 2018), which is

derived from an isolate collected from *D. melanogaster* in Charolles, France

(Jousset, Plus, Croizier, & Thomas, 1972). The virus was prepared as described

previously (Longdon, Cao, Martinez, & Jiggins, 2013). Briefly, DCV was grown in

Schneider's Drosophila line 2 cells and the Tissue Culture Infective Dose 50

(TCID50) per ml was calculated using the Reed-Muench end-point method. Flies

were anesthetized on CO_2 and inoculated using a 0.0125 mm diameter stainless

steel needle that was bent to a right angle ~0.25mm from the end (Fine Science

Tools, CA, USA). The bent tip of the needle was dipped into the DCV solution

(TCID50 = 6.32×10^9) and pricked into the pleural suture on the thorax of the flies

in order to standardize infection (Longdon et al., 2015). We selected this route of

infection as oral inoculation has been shown to lead to stochastic infection

outcomes in *D. melanogaster*, and injection has been show to follow a similar

course of infection (Ferreira et al., 2014). One vial of inoculated flies was

immediately snap frozen in liquid nitrogen to provide time point zero samples to

be used as a reference and therefore control for relative viral dose. The second

vial of flies were infected and then placed back into a new vial of their respective

food treatment. After 2 days (+/- 1 hour) flies were snap frozen in liquid

nitrogen. This time point was chosen based on previous studies that show a clear

increase in viral growth but little mortality at 2 days post infection (Longdon et

al., 2015; Roberts et al., 2018). Each experimental block contained a day 0 and

day 2 replicate for each species, at each diet (27 species × 3 diet treatments × 3

experimental blocks). In total, we quantified viral load in 7580 flies in 474

biological replicates (biological replicate = change in viral load from day 0 to day

2 post-infection), with a mean of 16 flies per replicate (range across species = 8-

28).

Survival

In order to measure the effect of diet on virulence we also carried out a survival

assay where mortality was recorded following infection. The same infection

protocol was carried out as above; one vial of flies was infected with DCV whilst

another was injected using a clean virus free needle dipped in *Drosophila*

Ringer's solution as a control treatment (Sullivan, Ashburner, & Hawley, 2000).

Flies were maintained in vials as described above and tipped onto their

respective fresh food every 2 days. The number of dead flies was counted every

day for 20 days. The survival assay was carried out across three blocks with

infections carried out over consecutive days, to obtain a control and virus

challenged vial per species each day. Treatment (virus challenged or control

sham infected) and the order in which fly species were inoculated were

randomized between blocks. Diet was randomized across days, so for a given

food type a control and viral challenged vial was completed each day, and this

was repeated over consequent days until there was a control and challenged for

each species on each food type (27 species × 2 treatments (control / challenged)

× 3 diet treatments × 3 experimental blocks). In total, we measured mortality in

9222 flies with a mean of 20 flies per replicate (range across species: 6–30 flies).

Measuring the change in viral load

The change in RNA viral load from day 0 to day 2-post infection was measured

using quantitative Reverse Transcription PCR (qRT-PCR). Frozen flies were

homogenised in Trizol reagent (Invitrogen) using a bead homogeniser for 30

seconds (Bead Ruptor 24; Omni international) and stored at -80°C for later

extraction. Total RNA was extracted from the Trizol homogenised flies in a

chloroform isopropanol extraction, reverse-transcribed with Promega GoScript

reverse transcriptase and random hexamer primers. Viral RNA load was

expressed relative to the endogenous control housekeeping gene *RpL32*. Primers

were designed to match the homologous sequence in each species and crossed

an intron-exon boundary so will only amplify mRNA. The primers in *D.*

melanogaster were *RpL32* qRT-PCR F (5'-TGCTAAGCTGTCGCACAAATGG -3') and

RpL32 qRT-PCR R (5'- TGCGCTTGTTCGATCCGTAAC -3'). DCV primers were 599F

(5'-GACACTGCCTTT GATTAG-3') and 733R (5'CCCTCTGGGAACTAAATG-3') as

previously described (Longdon et al., 2015). Two qRT-PCR reactions (technical

replicates) were carried out per sample with both the viral and endogenous

control primers, with replicates distributed across plates in a randomised block

design. qRT-PCR was performed on an Applied Biosystems StepOnePlus system

using Sensifast Hi-Rox Sybr kit (Bioline) with the following PCR cycle: 95°C for 2

min followed by 40 cycles of: 95°C for 5 sec followed by 60°C for 30 sec. Each

qRT-PCR plate contained four standard samples. A linear model was used to

correct the cycle threshold (Ct) values for differences between qRT-PCR plates.

Samples where the technical replicates had Ct values more than 2 cycles apart

after plate correction were repeated. To estimate the change in viral load, we

first calculated ΔCt as the difference between the cycle thresholds of the DCV

qRT-PCR and the RpL32 endogenous control. For each species the viral load of

day 2 flies relative to day 0 flies was calculated as $2^{-\Delta\Delta Ct}$; where ΔΔCt = ΔCt day0 –

ΔCt day2. The ΔCt day 0 and ΔCt day 2 is a pair of ΔCt values from a day 0

biological replicate and a day 2 replicate.

Diet	Ratio Protein: Carb	Cornmeal (g)	Dextrose (g)	Yeast (g)	Agar (g)	Nipagin (ml)	dH20 (L) Calories	Calories per 100ml
High	1:5	176	131.2	84	22	29	1	142.60
Normal	1:10	176	176	38	22	29	1	142.30
Low	1:20	176	203	10	22	29	1	142.02

Table 1: Ingredients for the experimental diet treatments. Amounts given

are enough to produce ~100 vials of food, with calculated calories per 100ml.

Effect of Body Size

To account for any potential differences in body size between species, we

measured wing length as a proxy for body size (Huey et al., 2006). During the

collections for the viral load assay males of each species were collected and

immediately stored in ethanol. Subsequently, wings were removed and

photographed under a dissecting microscope. Using ImageJ software (version

1.48) the length of the IV longitudinal vein from the tip of the proximal segment

to where the distal segment joins vein V was recorded, and the mean taken for

each species, overall there was a mean of 28 wings measured per species (range

20–35).

Host phylogeny

The host phylogeny was inferred as described in Longdon *et al.* (Longdon et al.,

2015) using seven genes (mitochondrial; *COI, COII*, ribosomal; *28S* and nuclear;

A*dh, SOD, Amyrel, RpL32*). Publicly available sequences were downloaded from

Genbank or were Sanger sequenced. In total we had *RpL32* sequences for all 27

species, *28S* from 24 species, *Adh* from 24 species, *Amyrel* from 15 species, *COI*

from 27 species, *COII* from 27 species and *SOD* from 12 species. For each gene

the sequences were aligned in Geneious (version 9.1.8) (Kearse et al., 2012)

using the global alignment setting, with free end gaps and a cost matrix of 70%

similarity (*See supplementary for table of sequences used*). The phylogeny was

inferred using BEAST (v1.10.4) (Drummond, Suchard, Xi, & Rambaut, 2012),

with genes partitioned into three groups; mitochondria, ribosomal and nuclear,

with their own molecular clock models. A random starting tree was used, with a

relaxed uncorrelated lognormal molecular clock. Each of the partitions used a

HKY substitution model with a gamma distribution of rate variation with 4

categories and estimated base frequencies. Additionally, the mitochondrial and

nuclear data sets were partitioned into codon positions 1+2 and 3, with unlinked

substitution rates and base frequencies across codon positions. The tree-shape

prior was set to a birth-death process. The BEAST analysis was run twice to

ensure convergence for 1000 million MCMC generations sampled every 10000

steps. On completion the MCMC process was examined using the program

Tracer (version 1.7.1) (Rambaut, Suchard, Xie, & Drummond, 2014) to ensure

convergence and adequate sampling, and the constructed tree was then

visualised using FigTree (v1.4.4) (Rambaut, 2006).

Statistical analysis

We used phylogenetic mixed models to look at the effects of host relatedness on

mortality and viral load across the three diet treatments. We fitted all models

using a Bayesian approach in the R package MCMCglmm version 2.29 (Hadfield,

2010) in RStudio (R version 3.5.1) (R Development Core Team, 2005). We used a

multivariate model with mortality of the controls, mortality of the virus infected

flies and viral load at each of the diets as the response variables. Mortality was

calculated as the mean portion of flies alive each day for each vial. The model

took the following form:

$y_{hit} = \beta_{1:t} + wingsize\beta_{2:t} + u_{p:ht} + e_{hit}$

Where y is the change in viral load of the i^{th} biological replicate of host species h,

for trait t. β are the fixed effects, with β_1 being the intercepts for each trait and β_2

being the effect of wing size. u_p are the random phylogenetic species effects and e

the model residuals. The models were also run with a non-phylogenetic random

species effect ($u_{s:ht}$) that allow us to estimate the proportion of variation the

host phylogeny explains (Longdon, Hadfield, Webster, Obbard, & Jiggins, 2011).

The main model was run without this term as it struggled to separate the

phylogenetic and non-phylogenetic terms. Our main model therefore assumes a

Brownian motion model of evolution (Felsenstein, 1973). The random effects

and the residuals are assumed to be multivariate normal with a zero mean and a

covariance structure $\mathbf{V_p} \otimes \mathbf{A}$ for the phylogenetic affects and $\mathbf{V_e} \otimes \mathbf{I}$ for the

residuals ($\otimes$ here is the Kronecker product). $\mathbf{A}$ is the phylogenetic relatedness

matrix, $\mathbf{I}$ is an identity matrix and the $\mathbf{V}$ are 9×9 (co)variance matrices describing

the (co)variances between viral load, mortality in virus infected, and mortality in

controls each at the 3 different diet levels. The phylogenetic covariance matrix,

$\mathbf{V_p}$ - describes the inter-specific variances in each trait and the inter-specific

covariances between them. The residual covariance matrix, $\mathbf{V_e}$ describes the

within-species variance that includes both any actual within-species effects and

also any measurement or experimental error. The off-diagonal elements in $\mathbf{V_e}$

(the covariances) are unable to be estimated because no vial has multiple

measurements - so were set to zero. The MCMC chain was run for 1300 million

iterations with a burn-in of 30 million iterations and a thinning interval of 1

million. Results were tested for sensitivity to the use of different priors by being

run with different prior structures (see supplementary R code), which gave

qualitatively similar results. We also ran models with the data subset into viral

load and mortality that gave similar results. All confidence intervals (CI's)

reported are 95% highest posterior density intervals.

Results

In order to investigate the effect that host diet may have on the likelihood of

virus host shifts we quantified DCV viral load in 27 species of Drosophilidae that

had been housed on three different diets (Figure 1). Viral loads differed between

species, with a thousand million times more virus in the most susceptible

compared to the least susceptible species, consistent with previous studies

(Longdon et al., 2015; Roberts et al., 2018). Viral loads were highly repeatable,

with the inter-specific phylogenetic component (v_p), explaining a high proportion

of the variation in viral load across diets (Repeatability = $v_p /(v_p + v_r)$; Low =

0.92, (95% CI: 0.86, 0.96); Medium = 0.90, (95% CI: 0.84,0.96); High = 0.83, (95%

CI: 0.75, 0.92).

Figure 1. Change in RNA viral load (log$_2$) for 27 Drosophilidae species infected with DCV across the three different diets of differing protein: carbohydrate ratios. Individual points are plotted with a small x-axis jitter for clarity of points and represent the change in viral load between day 0 and day 2-post infection.

The proportion of the between species variance that can be explained by the phylogeny can also be calculated, using $v_p/(v_p + v_s)$ (Freckleton, Harvey, & Pagel, 2002), and can be equated to the phylogenetic heritability or Pagel's lambda (Housworth, Martins, & Lynch, 2004; Pagel, 1999). We found that the host phylogeny explained a modest amount of the inter-specific variation in viral loads across diets, however these estimates had broad confidence intervals (Low = 0.20, (95% CI: 3.5 x10^{-6}, 0.63); Medium = 0.34, (95% CI: 2.0 x 10^{-6}, 0.80); High =

0.51, (95% CI: 3.2 x 10⁻⁶, 0.88), due to the model struggling to separate out the

phylogenetic and non-phylogenetic components.

In order to examine if the susceptibility of species responded in the same or

different ways to the changes in diet we examined viral load across the different

protein: carbohydrate ratios. We found strong positive inter-specific correlations

between viral loads across the diet treatments, suggesting the species are

responding in similar ways to the changes in dietary ratios (Table 2). There was

a decline in the between species variance in the high diet compared to low and

medium – but this was not found to be significantly different– (v_p: Low =77.13

(95% CI: 35.09, 125.50); Medium =82.33 (95% CI: 37.55, 135.26); High =45.59

(95% CI: 19.61, 76.39) and mean viral loads were similar across the diets (Low =

11.4 (95% CI: 5.3, 17.7); Medium = 10.6 (95% CI: 3.6, 16.6); High = 10.6, (95%

CI: 3.6, 16.7). Residual variance did not differ significantly between treatments

(Low = 6.45 (95% CI: 4.87, 8.04); Medium = 8.23 (95% CI: 6.45, 10.35); High =

8.32, (95% CI: 6.54, 10.6).

		Interspecific Correlation	95% CIs
Viral load	Low - Medium	0.93	0.85, 0.98
	Medium - High	0.83	0.66, 0.96
	Low - High	0.80	0.63, 0.96
Survival in virus challenged	Low - Medium	0.93	0.81, 0.99
	Medium - High	0.88	0.71, 0.99
	Low - High	0.90	0.73, 0.99

Table 2. Interspecific correlations between viral load and mortality

measures across the different diet treatments.

As similar pathogen loads can cause different levels of harm to their hosts

(tolerance rather than resistance) (Boots, 2008; Råberg, Graham, & Read, 2009;

Roy & Kirchner, 2000) we examined if virus induced mortality differed across

diets over a 20 day period after viral challenge (Figure 2). We found differences

in the virulence (mortality) caused by DCV between host species, with some

species seeing no apparent increase in mortality over the experimental period

compared to sham infected controls, (e.g. *S. pattersoni* and *D. saltans*), whilst

other species show higher susceptibility with up to 50% of flies dead by day 10

post infection (e.g. *D. simulans* and *D. melanogaster*). As with the viral load data

we calculated the repeatability of survival in these virus challenged flies which

was high in all cases (Repeatability = v_p /(v_p + v_r); Low = 0.90, (95% CI: 0.78,

0.98); Medium = 0.87, (95% CI: 0.71,0.97); High = 0.98, (95% CI: 0.85, 1.00). We

also calculated the proportion of between species variance that can be explained

by the phylogeny for the virus infected flies (Low = 0.16, (95% CI: 2.58×10^{-6},

0.62); Medium = 0.18, (95% CI: 5.75×10^{-7}, 0.78); High = 0.32, (95% CI: 7.75×10^{-8}

$^{-8}$, 0.87), which – like the viral load data – had broad confidence intervals due to

the model struggling to separate the phylogenetic and non-phylogenetic

components.

We found strong positive inter-specific correlations between the survival of

virus challenged flies across the diets, suggesting the species are responding in

similar ways to the dietary changes (Table 2). Among species variance in

mortality of virus infected flies was consistent across diets (Low = 0. 18 (95% CI:

0.07, 0.31); Medium = 0.16 (95% CI: 0.04, 0.30); High = 0.12 (95% CI: 0.04, 0.23)

as was the mean mortality (Low = 0.64 (95% CI: 0.47, 0.82); Medium = 0.58

(95% CI: 0.38, 0.75); High = 0.65, (95% CI: 0.47, 0.82). The residual variance was

also consistent across the diets (Low = 0.02 (95% CI: 0.01, 0.03); Medium = 0.02

(95% CI: 0.01, 0.04); High = 0.02, (95% CI: 0.01, 0.03).

Figure 2. Mortality in 27 species of Drosophilidae housed on three

different diets of varying protein: carbohydrate ratios. High- red circles,

Medium - blue crosses and Low- green triangles and either control sham infected

(dashed line) or virally challenged with DCV (solid lines).

We found that there were strong positive correlations between mortality and

RNA viral load (interspecific correlations between viral load and survival of virus

infected flies: Low = 0.89 (95% CI: 0.78, 0.98); Medium = 0.85 (95% CI: 0.67,

0.97); High = 0.67, (95% CI: 0.35, 0.90). To confirm that these differences are due

to changes in mortality caused by the virus rather than intrinsic differences in

the survivorship of the different species, we also inoculated flies with a control

solution. There was far less mortality in the controls than the virus infected flies

(Figure. 2 dashed lines). There was inter-specific variation in control mortality

(Low = 0.18, (95% CI: 0.01, 0.59); Medium = 0.43, (95% CI: 0.01,0.76); High =

0.55, (95% CI: -0.72, 1.00) but this was not significantly correlated with survival

of the virus infected flies (survival of control versus virus infected on: Low = -

0.11 (95% CI: -0.92, 0.75); Medium = 0.34 (95% CI: -0.48 0.97); High = 0.12,

(95% CI: -0.82, 0.89). We found no effect of wing length as a proxy for body size,

(mean: -0.05, 95% CI: -0.13, 0.05).

Discussion

We found dietary treatments of differing protein to carbohydrate ratios did not

alter the outcome of infection in 27 species of Drosophilidae infected with DCV.

We found strong positive inter-specific correlations across diets in both viral

load and mortality, suggesting that the species are responding in similar ways to

nutritional changes. Despite there being among species variation in

susceptibility, changes in diet did not affect viral loads. Nor did it appear to alter

the likelihood of tolerating or surviving an infection, as we found strong positive

correlations between mortality and viral load on each of the diets.

Both mounting and maintaining an immune response requires energy and

nutrients. During an acute immune challenge the provisioning of nutrients may

become more demanding for a host, with pathogen induced malabsorption

through damage to, or obstruction of, digestive tissues (Lochmiller &

Deerenberg, 2000). DCV is known to cause severe pathology of the tissues of the

digestive tract with subsequent accumulation of food in the crop (food storage

organ) and obstruction in the intestine (Chtarbanova et al., 2014). These physical

symptoms alter an infected hosts energy stores with infected flies showing

significantly reduced glycogen and triglyceride levels three to four days post

infection (Chtarbanova et al., 2014). DCV infected flies also increase in body

mass, with a reduced food intake and reduced metabolism, suggesting that they

may be experiencing increased water retention (Arnold et al., 2013; Chtarbanova

et al., 2014; Thomas-Orillard, 1984). We therefore hypothesised that changing

the ratio of protein to carbohydrate in the diet may alter outcome of infection,

and as species may all have their own "optimal diet", that species may respond in

different ways to such changes. However our current study does not appear to

support this.

Geometric frameworks for nutrition were developed in response to the fact that

what is "optimal" will depend on a balance of particular nutrients in the

organism and trait being investigated (Archer, Royle, South, Selman, & Hunt,

2009; Sheena C. Cotter et al., 2019; S. J. Simpson & Raubenheimer, 1995). For

example mice infected with *Salmonella* were found to survive better on diets

containing a higher ratio of protein to carbohydrate (Peck, Babcock, & Alexander,

1992) as were bacterially infected army worm caterpillars - with survival

increasing with dietary protein: carbohydrate, suggesting high protein

requirements are associated with bacterial resistance (Povey et al., 2009). A

recent study using 10 different protein: carbohydrate diets challenged flies with *P.*

entomophila bacteria (Savola, Montgomery, et al., 2020). Survival on low protein

diets was found to be very low in infected flies suggesting protein was key for

tolerating a bacterial challenge. This study also monitored lifespan and reproduction

in flies and found that regardless of injury and infection, dietary restriction extended

lifespan and reduced reproductive output (Savola, Montgomery, et al., 2020).

However a potential mechanism of the interaction of diet and infection has been

suggested in research using a different model system with *in vivo* and *in vitro*

elements (Wilson et al., 2020). Caterpillars of *S. littoralis* challenged with the

bacteria *X. nematophila in vivo* and on high dietary protein had slower bacterial

growth and higher survival. When this was combined with *in vitro* experiments the

results suggested this was being driven by the osmolality of the hosts' blood

(hemolymph) being altered by an increase in solutes in the high protein diets that

acted to slow bacterial growth (Wilson et al., 2020). Although these results are

using a bacterial system whilst we use a viral host shift, it highlights a significant

area for further mechanistic studies of the interaction of diet and immunity.

Resources are critical in determining immune investment and the level of

response to a pathogen challenge in invertebrates (Traniello, Rosengaus, &

Savoie, 2002) and development time, longevity, feeding behaviour and

reproduction are all affected by dietary modifications (Moatt et al., 2020). In

particular protein is fundamental for producing amino acid rich immunological

components needed for resistance, particularly to viral pathogens (S. C. Cotter &

Kilner, 2010; Lee et al., 2008; Ponton, Lalubin, et al., 2011) whilst carbohydrates

are essential for energetically costly immune responses (Arnold et al., 2013;

Kutzer & Armitage, 2016b). A recent study of bacterial infection in *D.*

melanogaster found that a low protein: carbohydrate diet can improve survival

upon infection, and the expression levels of genes associated with antimicrobial

peptides were significantly affected by diet (Mondotte & Saleh, 2017; Xu &

Cherry, 2014). This study also found that infected flies choose to actively

decrease their protein intake whilst maintaining their carbohydrate

consumption, highlighting the importance of the correct balance of key nutrients.

Flies can respond to the absence of just one essential amino acid from their diet;

increasing their appetite for foods rich in such amino acids (Kay et al., 2014;

Keaton Wilson, Ruiz, & Davidowitz, 2019; Lochmiller & Deerenberg, 2000;

McKean et al., 2008), suggesting they that they are constantly modifying their

intake in response to internal cues. This mirrors the result of an experiment in

rats, upon immune stimulation the rats actively ate more carbohydrates than fat,

suggesting they wanted a direct source of energy and not just increased calories

(Ponton et al., 2020). Whereas caterpillars infected with a nucleopolyhedrovirus,

actively choose to feed on higher protein foods and as a result had stronger

immune responses, including encapsulation, lysozyme-like antimicrobial activity

and phenol-oxidase activity and decreased levels of infection (Leitão-Gonçalves

et al., 2017).

Other nutrients may also interact with the ability of a host to respond to

infection. For example, lipid and fats have been associated with *D. melanogaster*

response to DCV viral infection; peroxisomes were found to be required for host

defense to infection, through their primary function in lipid metabolism (Aubert,

Goodall, & Dantzer, 1995). Indeed recent understanding of the lipidome in

mosquito cells co-infected with *Wolbachia* and Dengue or Zika virus pointed to

the importance of fatty acids in the mechanism underlying *Wolbachia* -mediated

pathogen blocking (Manokaran et al., 2020). The lipid levels across our

experimental diets were however held constant and this is therefore a potential

area for further study.

There has been an increased use of a chemically defined (holidic) diet in order

to manipulate individual nutrients present in fly diets (Lee, Cory, Wilson,

Raubenheimer, & Simpson, 2006). Exome matched diets have been shown to

alleviate trade-offs in fecundity and longevity (Merkling, Riahi, Overheul,

Schenck, & van Rij, 2019). A possible extension of this would be to look at the

effect of matching diets to transcriptional changes during infection, and seeing if

this alleviates (or exacerbates) pathology.

Changes in diet have been shown to alter pathogen susceptibility in a number of

systems. We hypothesised that changes in diet could alter the potential outcomes

of virus host shifts. We found that changes in the ratio of protein to carbohydrate

did not alter susceptibility to DCV across host species in this instance.

Author Contributions: KER conceived, designed and carried out the

experiment, both KER and BL analysed, wrote and edited the manuscript and

agreed to the final submitted version.

Conflict of Interest statement: Authors declare no conflict of interest

Acknowledgments

KER and BL are supported by a Sir Henry Dale Fellowship jointly funded by the

Wellcome Trust and the Royal Society (grant no. 109356/Z15/Z).

References

Anderson, R. M., Latorre-Esteves, M., Neves, A. R., Lavu, S., Medvedik, O., Taylor, C., … Sinclair, D. A. (2003). Yeast Life-Span Extension by Calorie Restriction Is Independent of NAD Fluctuation. *Science*, *302*(5653), 2124–2126. doi:10.1126/science.1088697

Archer, C. R., Royle, N., South, S., Selman, C., & Hunt, J. (2009). Nutritional geometry provides food for thought. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, *64*(9), 956–959. doi:10.1093/gerona/glp069

Arnold, P. A., Johnson, K. N., & White, C. R. (2013). Physiological and metabolic consequences of viral infection in Drosophila melanogaster. *Journal of Experimental Biology*, *216*(17), 3350–3357. doi:10.1242/jeb.088138

Aubert, A., Goodall, G., & Dantzer, R. (1995). Compared effects of cold ambient temperature and cytokines on macronutrient intake in rats. *Physiology and Behavior*, *57*(5), 869–873. doi:10.1016/0031-9384(94)00391-H

Ayres, J. S., & Schneider, D. S. (2009). The Role of Anorexia in Resistance and Tolerance to Infections in Drosophila. *PLoS Biology*, *7*(7), e1000150. doi:10.1371/journal.pbio.1000150

Blanford, S., & Thomas, M. B. (1999). Host thermal biology: the key to understanding host-pathogen interactions and microbial pest control? *Agricultural and Forest Entomology*, *1*(3), 195–202. doi:10.1046/j.1461-9563.1999.00027.x

Blum, J. E., Fischer, C. N., Miles, J., & Handelsman, J. (2013). Frequent replenishment sustains the beneficial microbiome of Drosophila melanogaster. *MBio*, *4*(6), 1–8. doi:10.1128/mBio.00860-13

Boots, M. (2008). Fight or learn to live with the consequences? *Trends in Ecology & Evolution*, *23*(5), 248–250. doi:10.1016/j.tree.2008.01.006

Broderick, N. A., & Lemaitre, B. (2012). Gut-associated microbes of Drosophila melanogaster. *Gut Microbes*, *3*(4), 307–321. doi:10.4161/gmic.19896

Brooks, D. R., & Hoberg, E. P. (2007). How will global climate change affect parasite-host assemblages? *Trends in Parasitology*, *23*(12), 571–574. doi:10.1016/j.pt.2007.08.016

Bunning, H., Bassett, L., Clowser, C., Rapkin, J., Jensen, K., House, C. M., … Hunt, J. (2016). Dietary choice for a balanced nutrient intake increases the mean and reduces the variance in the reproductive performance of male and female cockroaches. *Ecology and Evolution*, *6*(14), 4711–4730. doi:10.1002/ece3.2243

Camus, M. F. F., Fowler, K., Piper, M. W. D., & Reuter, M. (2017). Sex and genotype effects on nutrient-dependent fitness landscapes in Drosophila melanogaster. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1869), 20172237. doi:10.1098/rspb.2017.2237

Carlson, C. J., Albery, G. F., Merow, C., Trisos, C. H., Zipfel, C. M., Eskew, E. A., … Bansal, S. (2020). Climate change will drive novel cross-species viral transmission. *BioRxiv*, 2020.01.24.918755. doi:10.1101/2020.01.24.918755

Chtarbanova, S., Lamiable, O., Lee, K.-Z., Galiana, D., Troxler, L., Meignin, C., … Imler, J.-L. (2014). Drosophila C virus systemic infection leads to intestinal obstruction. *Journal of Virology*, *88*(24), 14057–69. doi:10.1128/JVI.02320-14

Cotter, S. C., & Kilner, R. M. (2010). Personal immunity versus social immunity.

Behavioral Ecology, *21*(4), 663–668. doi:10.1093/beheco/arq070
Cotter, Sheena C., Reavey, C. E., Tummala, Y., Randall, J. L., Holdbrook, R., Ponton,
F., … Wilson, K. (2019). Diet modulates the relationship between immune
gene expression and functional immune responses. *Insect Biochemistry and*
Molecular Biology, *109*, 128–141. doi:10.1016/j.ibmb.2019.04.009
Cotter, Sheena C., Simpson, S. J., Raubenheimer, D., & Wilson, K. (2011).
Macronutrient balance mediates trade-offs between immune function and
life history traits. *Functional Ecology*, *25*(1), 186–198. doi:10.1111/j.1365-
2435.2010.01766.x
Cumnock, K., Gupta, A. S., Lissner, M., Chevee, V., Davis, N. M., & Schneider, D. S.
(2018). Host Energy Source Is Important for Disease Tolerance to Malaria.
Current Biology, *28*(10), 1635-1642.e3. doi:10.1016/j.cub.2018.04.009
Drummond, A., Suchard, M., Xi, E. D., & Rambaut, A. (2012). Bayesian
phylogenetics with BEAUti and the BEAST 1.7. Molecular Biology And
Evolution 29.
Fanson, B. G., Weldon, C. W., Pérez-Staples, D., Simpson, S. J., & Taylor, P. W.
(2009). Nutrients, not caloric restriction, extend lifespan in Queensland fruit
flies (Bactrocera tryoni). *Aging Cell*, *8*(5), 514–523. doi:10.1111/j.1474-
9726.2009.00497.x
Fellous, S., & Lazzaro, B. P. (2010). Larval food quality affects adult (but not
larval) immune gene expression independent of effects on general
condition. *Molecular Ecology*, *19*(7), 1462–1468. doi:10.1111/j.1365-
294X.2010.04567.x
Felsenstein, J. (1973). Maximum-likelihood estimation of evolutionary trees from
continuous characters. *American Journal of Human Genetics*, *25*(5), 471–492.
doi:10.1038/ncomms15810
Ferreira, Á. G., Naylor, H., Esteves, S. S., Pais, I. S., Martins, N. E., & Teixeira, L.
(2014). The Toll-Dorsal Pathway Is Required for Resistance to Viral Oral
Infection in Drosophila. *PLoS Pathogens*, *10*(12).
doi:10.1371/journal.ppat.1004507
Freckleton, R. P., Harvey, P. H., & Pagel, M. (2002). Phylogenetic analysis and
comparative data: a test and review of evidence. *The American Naturalist*,
160(6), 712–26. doi:10.1086/343873
Grueber, C. E., Gray, L. J., Morris, K. M., Simpson, S. J., & Senior, A. M. (2018).
Intergenerational effects of nutrition on immunity: a systematic review and
meta-analysis. *Biological Reviews of the Cambridge Philosophical Society*,
93(2), 1108–1124. doi:10.1111/brv.12387
Hadfield, J. D. (2010). MCMC methods for multi-respoinse generalized linear
mixed models: The MCMCglmm R package. *Journal of Statistical Software*,
33(2), 1–22. doi:10.1002/ana.22635
Harvell, C. D., Mitchell, C. E., Ward, J. R., Altizer, S., Dobson, A. P., Ostfeld, R. S., &
Samuel, M. D. (2002). Climate warming and disease risks for terrestrial and
marine biota. *Science (New York, N.Y.)*, *296*(5576), 2158–62.
doi:10.1126/science.1063699
Hayman, D. T. S., Pulliam, J. R. C., Marshall, J. C., Cryan, P. M., & Webb, C. T. (2016).
Environment, host, and fungal traits predict continental-scale white-nose
syndrome in bats. *Science Advances*, *2*(1), e1500831.
doi:10.1126/sciadv.1500831
Henry, Y., Overgaard, J., & Colinet, H. (2020). Dietary nutrient balance shapes

619   phenotypic traits of Drosophila melanogaster in interaction with gut
620   microbiota. *Comparative Biochemistry and Physiology Part A: Molecular &*
621   *Integrative Physiology*, *241*, 110626. doi:10.1016/j.cbpa.2019.110626
622   Henry, Youn. (2018). Microbiota disruption leads to reduced cold tolerance in
623   Drosophila flies. doi:10.1007/s00114-018-1584-7
624   Hoberg, E. P., & Brooks, D. R. (2015). Evolution in action: climate change,
625   biodiversity dynamics and emerging infectious disease. *Philosophical*
626   *Transactions of the Royal Society of London. Series B, Biological Sciences*,
627   *370*(1665), 20130553–20130553. doi:10.1098/rstb.2013.0553
628   Housworth, E. A., Martins, E. P., & Lynch, M. (2004). The phylogenetic mixed
629   model. *The American Naturalist*, *163*(1), 84–96. doi:10.1086/380570
630   Huey, R. B., Moreteau, B., Moreteau, J.-C., Gibert, P., Gilchrist, G. W., Ives, A. R., …
631   David, J. R. (2006). Sexual size dimorphism in a Drosophila clade, the D.
632   obscura group. *Zoology*, *109*(4), 318–330. doi:10.1016/j.zool.2006.04.003
633   Jensen, K., McClure, C., Priest, N. K., & Hunt, J. (2015). Sex-specific effects of
634   protein and carbohydrate intake on reproduction but not lifespan in
635   *Drosophila melanogaster*. *Aging Cell*, *14*(4), 605–615.
636   doi:10.1111/acel.12333
637   Jousset, F. X., Plus, N., Croizier, G., & Thomas, M. (1972). [Existence in Drosophila
638   of 2 groups of picornavirus with different biological and serological
639   properties]. *Comptes Rendus Hebdomadaires Des Seances de l'Academie Des*
640   *Sciences. Serie D: Sciences Naturelles*, *275*(25), 3043–6. Retrieved from
641   http://www.ncbi.nlm.nih.gov/pubmed/4631976
642   Kay, A. D., Bruning, A. J., van Alst, A., Abrahamson, T. T., Hughes, W. O. H., &
643   Kaspari, M. (2014). A carbohydrate-rich diet increases social immunity in
644   ants. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1778).
645   doi:10.1098/rspb.2013.2374
646   Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., …
647   Drummond, A. (2012). Geneious Basic: an integrated and extendable
648   desktop software platform for the organization and analysis of sequence
649   data. *Bioinformatics*, *28*(12), 1647–1649.
650   Keaton Wilson, J., Ruiz, L., & Davidowitz, G. (2019). Dietary Protein and
651   Carbohydrates Affect Immune Function and Performance in a Specialist
652   Herbivore Insect (Manduca sexta). *Physiological and Biochemical Zoology*,
653   *92*(1), 58–70. doi:10.1086/701196
654   Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P. K., Krkošek, M., & Luijckx, P.
655   (2018). Empirical evidence that metabolic theory describes the temperature
656   dependency of within-host parasite dynamics. *PLOS Biology*, *16*(2),
657   e2004608. doi:10.1371/journal.pbio.2004608
658   Klass, M. R. (1983). A method for the isolation of longevity mutants in the
659   nematode Caenorhabditis elegans and initial results. *Mechanisms of Ageing*
660   *and Development*, *22*(3–4), 279–286. doi:10.1016/0047-6374(83)90082-9
661   Knutie, S. A., Wilkinson, C. L., Wu, Q. C., Ortega, C. N., & Rohr, J. R. (2017). Host
662   resistance and tolerance of parasitic gut worms depend on resource
663   availability. *Oecologia*, *183*(4), 1031–1040. doi:10.1007/s00442-017-3822-
664   7
665   Kraaijeveld, A. R., & Godfray, H. C. (1997). Trade-off between parasitoid
666   resistance and larval competitive ability in Drosophila melanogaster.
667   *Nature*, *389*(6648), 278–80. doi:10.1038/38483

Kutzer, M. A. M., & Armitage, S. A. O. (2016a). The effect of diet and time after
bacterial infection on fecundity, resistance, and tolerance in
Drosophila melanogaster. *Ecology and Evolution*, *6*(13), 4229–4242.
doi:10.1002/ece3.2185
Kutzer, M. A. M., & Armitage, S. A. O. (2016b). The effect of diet and time after
bacterial infection on fecundity, resistance, and tolerance in
Drosophila melanogaster. *Ecology and Evolution*, *6*(13), 4229–4242.
doi:10.1002/ece3.2185
Le Couteur, D. G., Solon-Biet, S., Cogger, V. C., Mitchell, S. J., Senior, A., De Cabo, R.,
… Simpson, S. J. (2016). The impact of low-protein high-carbohydrate diets
on aging and lifespan. *Cellular and Molecular Life Sciences*, *73*(6), 1237–
1252. doi:10.1007/s00018-015-2120-y
Lee, K. ., Cory, J. ., Wilson, K., Raubenheimer, D., & Simpson, S. . (2006). Flexible
diet choice offsets protein costs of pathogen resistance in a caterpillar.
Proceedings of the Royal Society B: Biological Sciences, *273*(1588), 823–829.
doi:10.1098/rspb.2005.3385
Lee, K. P., Simpson, S. J., & Wilson, K. (2008). Dietary protein-quality influences
melanization and immune function in an insect. *Functional Ecology*, *22*(6),
1052–1061. doi:10.1111/j.1365-2435.2008.01459.x
Leitão-Gonçalves, R., Carvalho-Santos, Z., Francisco, A. P., Fioreze, G. T., Anjos, M.,
Baltazar, C., … Ribeiro, C. (2017). Commensal bacteria and essential amino
acids control food choice behavior and reproduction. *PLOS Biology*, *15*(4),
e2000862. doi:10.1371/journal.pbio.2000862
Lesperance, D. N. A., & Broderick, N. A. (2020). Meta-analysis of Diets Used in
Drosophila Microbiome Research and Introduction of the Drosophila
Dietary Composition Calculator (DDCC). *G3: Genes, Genomes, Genetics*,
g3.401235.2020. doi:10.1534/g3.120.401235
Lochmiller, R. L., & Deerenberg, C. (2000). Trade-offs in evolutionary
immunology: Just what is the cost of immunity? *Oikos*, *88*(1), 87–98.
doi:10.1034/j.1600-0706.2000.880110.x
Longdon, B., Cao, C., Martinez, J., & Jiggins, F. M. (2013). Previous Exposure to an
RNA Virus Does Not Protect against Subsequent Infection in Drosophila
melanogaster. *PLoS ONE*, *8*(9), e73833. doi:10.1371/journal.pone.0073833
Longdon, B., Day, J. P., Alves, J. M., Smith, S. C. L., Houslay, T. M., McGonigle, J. E., …
Jiggins, F. M. (2018). Host shifts result in parallel genetic changes when
viruses evolve in closely related species. *PLOS Pathogens*, *14*(4), e1006951.
doi:10.1371/journal.ppat.1006951
Longdon, B., Hadfield, J. D., Day, J. P., Smith, S. C. L., McGonigle, J. E., Cogni, R., …
Jiggins, F. M. (2015). The causes and consequences of changes in virulence
following pathogen host shifts. *PLoS Pathogens*, *11*(3), e1004728.
doi:10.1371/journal.ppat.1004728
Longdon, B., Hadfield, J. D., Webster, C. L., Obbard, D. J., & Jiggins, F. M. (2011).
Host phylogeny determines viral persistence and replication in novel hosts.
PLoS Pathogens, *7*(9), e1002260. doi:10.1371/journal.ppat.1002260
Maklakov, A. A., Hall, M. D., Simpson, S. J., Dessmann, J., Clissold, F. J., Zajitschek,
F., … Brooks, R. C. (2009). Sex differences in nutrient-dependent
reproductive ageing. *Aging Cell*, *8*(3), 324–330. doi:10.1111/j.1474-
9726.2009.00479.x
Manokaran, G., Flores, H. A., Dickson, C. T., Narayana, V. K., Kanojia, K., Dayalan,

S., ... Simmons, C. P. (2020). Modulation of acyl-carnitines, the broad
mechanism behind Wolbachia -mediated inhibition of medically important
flaviviruses in Aedes aegypti . *Proceedings of the National Academy of*
Sciences, *117*(39), 24475–24483. doi:10.1073/pnas.1914814117
McKean, K. A., Yourth, C. P., Lazzaro, B. P., & Clark, A. G. (2008). The evolutionary
costs of immunological maintenance and deployment. *BMC Evolutionary*
Biology, *8*(1), 76. doi:10.1186/1471-2148-8-76
Merkling, S. H., Riahi, H., Overheul, G. J., Schenck, A., & van Rij, R. P. (2019).
Peroxisome-associated Sgroppino links fat metabolism with survival after
RNA virus infection in Drosophila. *Scientific Reports*, *9*(1), 2065.
doi:10.1038/s41598-019-38559-x
Moatt, J. P., Savola, E., Regan, J. C., Nussey, D. H., & Walling, C. A. (2020). Lifespan
Extension Via Dietary Restriction: Time to Reconsider the Evolutionary
Mechanisms? *BioEssays*, 1900241. doi:10.1002/bies.201900241
Mondotte, J. A., & Saleh, M.-C. (2017). Antiviral Immune Response and the Route
of Infection in Drosophila melanogaster. doi:10.1016/bs.aivir.2017.10.006
Nakagawa, S., Lagisz, M., Hector, K. L., & Spencer, H. G. (2012). Comparative and
meta-analytic insights into life extension via dietary restriction. *Aging Cell*,
11(3), 401–409. doi:10.1111/j.1474-9726.2012.00798.x
Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*,
401(6756), 877–884. doi:10.1038/44766
Peck, M. D., Babcock, G. F., & Alexander, J. W. (1992). The role of protein and
calorie restriction in outcome from Salmonella infection in mice. *Journal of*
Parenteral and Enteral Nutrition, *16*(6), 561–565.
doi:10.1177/0148607192016006561
Pike, V. L., Lythgoe, K. A., & King, K. C. (2019). On the diverse and opposing effects
of nutrition on pathogen virulence. *Proceedings of the Royal Society B:*
Biological Sciences, *286*(1906), 20191220. doi:10.1098/rspb.2019.1220
Piper, M. D. (2017). Using artificial diets to understand the nutritional physiology
of Drosophila melanogaster. *Current Opinion in Insect Science*, *23*, 104–111.
doi:10.1016/j.cois.2017.07.014
Ponton, F., Lalubin, F., Fromont, C., Wilson, K., Behm, C., & Simpson, S. J. (2011).
Hosts use altered macronutrient intake to circumvent parasite-induced
reduction in fecundity. *International Journal for Parasitology*, *41*(1), 43–50.
doi:10.1016/j.ijpara.2010.06.007
Ponton, F., Morimoto, J., Robinson, K., Kumar, S. S., Cotter, S. C., Wilson, K., &
Simpson, S. J. (2020). Macronutrients modulate survival to infection and
immunity in Drosophila. *Journal of Animal Ecology*, *89*(2), 460–470.
doi:10.1111/1365-2656.13126
Ponton, F., Wilson, K., Cotter, S. C., Raubenheimer, D., & Simpson, S. J. (2011).
Nutritional immunology: A multi-dimensional approach. *PLoS Pathogens*,
7(12), e1002223. doi:10.1371/journal.ppat.1002223
Ponton, F., Wilson, K., Holmes, A. J., Cotter, S. C., Raubenheimer, D., & Simpson, S.
J. (2013). Integrating nutrition and immunology: a new frontier. *Journal of*
Insect Physiology, *59*(2), 130–7. doi:10.1016/j.jinsphys.2012.10.011
Povey, S., Cotter, S. C., Simpson, S. J., Lee, K. P., & Wilson, K. (2009). Can the
protein costs of bacterial resistance be offset by altered feeding behaviour?
Journal of Animal Ecology, *78*(2), 437–446. doi:10.1111/j.1365-
2656.2008.01499.x

R Development Core Team. (2005). *R: A Language and Environment for Statistical*
Computing. Vienna, Austria: R Foundation for Statistical Computing.
Råberg, L., Graham, A. L., & Read, A. F. (2009). Decomposing health: tolerance
and resistance to parasites in animals. *Philosophical Transactions of the*
Royal Society B: Biological Sciences, *364*(1513), 37–49.
doi:10.1098/rstb.2008.0184
Rambaut, A. (2006). FigTree v1.4.4. Retrieved from
http://tree.bio.ed.ac.uk/software/ figtree/.
Rambaut, A., Suchard, M., Xie, D., & Drummond, A. (2014). Tracer v1.6. Retrieved
from http://tree.bio.ed.ac.uk/software/tracer/
Rao, S., Schieber, A. M. P., O'Connor, C. P., Leblanc, M., Michel, D., & Ayres, J. S.
(2017). Pathogen-Mediated Inhibition of Anorexia Promotes Host Survival
and Transmission. *Cell*, *168*(3), 503-516.e12. doi:10.1016/j.cell.2017.01.006
Raubenheimer, D., & Simpson, S. J. (1999). Integrating nutrition: a geometrical
approach. *Entomologia Experimentalis et Applicata*, *91*(1), 67–82.
doi:10.1046/j.1570-7458.1999.00467.x
Roberts, K. E., Hadfield, J. D., Sharma, M. D., & Longdon, B. (2018). Changes in
temperature alter the potential outcomes of virus host shifts. *PLOS*
Pathogens, *14*(10), e1007185. doi:10.1371/journal.ppat.1007185
Roy, B. A., & Kirchner, J. W. (2000). Evolutionary Dynamics of Pathogen
Resistance and Tolerance. *Evolution*, *54*(1), 51–63. doi:10.1111/j.0014-
3820.2000.tb00007.x
Savola, E., Montgomery, C., Waldron, F. M., Monteith, K., Vale, P., & Walling, C. A.
(2020). Testing evolutionary explanations for the lifespan benefit of dietary
restriction in Drosophila melanogaster. *BioRxiv*.
doi:https://doi.org/10.1101/2020.06.18.159731
Savola, E., Vale, P., & Walling, C. A. (2020). Larval diet affects adult reproduction
but not survival regardless of injury and infection stress in Drosophila
melanogaster. *BioRxiv*, 1–29.
doi:https://doi.org/10.1101/2020.10.16.342618 .
Sheldon, B. C., & Verhulst, S. (1996). Ecological immunology: costly parasite
defences and trade offs in evolutionary ecology. *Trends in Ecology &*
Evolution, *11*(8), 317–321.
Simpson, S. J., & Raubenheimer, D. (1995, July 1). The geometric analysis of
feeding and nutrition: a user's guide. *Journal of Insect Physiology*. Pergamon.
doi:10.1016/0022-1910(95)00006-G
Simpson, Stephen J., & Raubenheimer, D. (2011). The nature of nutrition: a
unifying framework. *Australian Journal of Zoology*, *59*(6), 350.
doi:10.1071/ZO11068
Skorupa, D. A., Dervisefendic, A., Zwiener, J., & Pletcher, S. D. (2008). Dietary
composition specifies consumption, obesity, and lifespan in *Drosophila*
melanogaster. *Aging Cell*, *7*(4), 478–490. doi:10.1111/j.1474-
9726.2008.00400.x
Sullivan, W., Ashburner, M., & Hawley, R. S. (2000). Drosophila protocols.
Drosophila Protocols.
Thomas-Orillard, M. (1984). Modifications of Mean Ovariole Number, Fresh
Weight of Adult Females and Developmental Time in Drosophila
melanogaster Induced by Drosophila C Virus. *Genetics*, *107*(4), 635–63544.
Traniello, J. F. A., Rosengaus, R. B., & Savoie, K. (2002). The development of

immunity in a social insect : Evidence for the group facilitation of disease
resistance. *Proceedings of the National Academy of Sciences*, *99*(10), 6838–
6842. doi:doi: 10.1073/pnas.102176599
Weindruch, R., & Walford, R. L. (1982). Dietary restriction in mice beginning at 1
year of age: Effect on life-span and spontaneous cancer incidence. *Science*,
215(4538), 1415–1418. doi:10.1126/science.7063854
Wilson, K., Holdbrook, R., Reavey, C. E., Randall, J. L., Tummala, Y., Ponton, F., …
Cotter, S. C. (2020). Osmolality as a Novel Mechanism Explaining Diet Effects
on the Outcome of Infection with a Blood Parasite. *Current Biology*, *30*(13),
2459–2467. doi:10.1016/j.cub.2020.04.058
Xu, J., & Cherry, S. (2014). Viruses and antiviral immunity in Drosophila.
Developmental and Comparative Immunology, *42*(1), 67–84.
doi:10.1016/j.dci.2013.05.002
828