Thermally tolerant symbionts may explain Caribbean octocoral resilience to heat stress

Jessie Pelosi[1,#,*], Katherine M. Eaton[1,^,*], Samantha Mychajliw[2], Casey P. terHorst[3], and Mary

Alice Coffroth[4,+]

[1]Department of Biological Sciences, University at Buffalo, Buffalo, NY 14260, [2]Department of

Environment and Sustainability, University at Buffalo, Buffalo, NY 14260, [3]Department of

Biology, California State University, Northridge, Northridge, CA 91330, [4]Department of

Geology, University at Buffalo, Buffalo, NY 14260

* These authors contributed equally to this work

Current address:

[#]Department of Biology, University of Florida, Gainesville, FL 32608

[^]Department of Biological Sciences, Auburn University, Auburn, AL, 36849

[+]Corresponding author: coffroth@buffalo.edu

Keywords: symbiosis, Symbiodiniaceae, octocorals, thermal tolerance

Abstract

Coral reef ecosystems are under threat from the frequent and severe impacts of anthropogenic climate change, particularly rising sea surface temperatures. The effects of thermal stress may be ameliorated by adaptation and/or acclimation of the host, symbiont, or holobiont (host + symbiont) to increased temperatures. We examined the role of the symbiont in promoting thermal tolerance of the holobiont, using *Antillogorgia bipinnata* (octocoral host) and *Breviolum antillogorgium* (symbiont) as a model system. We identified five distinct genotypes of *B. antillogorgium* from symbiont populations isolated from *A. bipinnata* colonies. Three symbiont genotypes were maintained at 26°C (ambient historical temperature) and two were maintained at 30°C (elevated historical temperature) for two years. We analyzed the *in vitro* growth rate and carrying capacity of each genotype at both ambient and elevated temperatures. All genotypes grew well at both temperature treatments, indicating thermal tolerance among these *B. antillogorgium* genotypes. We also inoculated juvenile *A. bipinnata* polyps with each of the five symbiont genotypes, and reared these polyps at both ambient and elevated temperatures. All genotypes were able to infect polyps at both temperature treatments. Survivorship of polyps at 30°C was significantly lower than survivorship at 26°C, but all treatments had surviving polyps at 56 days post-infection, suggestive of broad-scale thermal tolerance in this system. The widespread thermal tolerance observed in *B. antillogorgium* may play a part in the increased resilience of Caribbean octocorals during heat stress events.

Introduction

In recent years, anthropogenic climate change has severely impacted coral reefs, which

comprise some of the most biodiverse and economically important ecosystems on Earth (Hoegh-

Guldberg 2011; Hoegh-Guldberg et al. 2017). The success of reef ecosystems in oligotrophic

tropical waters is due to their obligate endosymbioses with dinoflagellate algae in the family

Symbiodiniaceae (Muscatine and Porter 1977). In this mutualistic relationship, the coral host

relies on photosynthate produced by its intracellular symbionts to thrive (Muscatine and

Cernichiari 1969; Muscatine and Porter 1977). In exchange for these metabolites, the symbiotic

algae recover host nitrogenous waste. This relationship, which is vital to the survival of the reef

ecosystem, is currently threatened by anthropogenic environmental changes.

Coral bleaching, or the loss of algal symbionts, occurs due to a variety of environmental

stressors including changes in salinity, pH, light intensity and, most commonly, water

temperature (Hoegh-Guldberg 1999). Most reef ecosystems now routinely encounter

temperatures at or above their thermal maximum each year (Hoegh-Guldberg et al. 2007; Eakin

et al. 2009) with bleaching events becoming more frequent and severe (Heron et al. 2016; van

Hooidonk et al. 2016; Hughes et al. 2018; Oliver et al. 2018). As ocean temperatures continue to

increase, it is essential to understand if and how corals and their symbionts can respond to the

present threat.

The holobiont (host + symbiont) can respond to rising seawater temperatures via changes

in the symbiont community within a single host, with the basic premise that hosts associating

with more thermally tolerant symbionts will be more likely to survive bleaching events (i.e.,

shuffling and/or switching, *sensu* Baker 2003). Although most coral species host a single

dominant symbiont type (Goulet 2006; LaJeunesse 2020), many hosts also harbor background or

cryptic symbionts that are often undetected by traditional techniques (Mieog et al. 2007; Correa

et al. 2009; Silverstein et al. 2012). During thermal stress, a more thermally tolerant background

symbiont can become the dominant symbiont type, potentially conferring thermal tolerance to

the holobiont (Berkelmans and van Oppen 2006; Jones et al. 2008; LaJeunesse et al. 2010;

Keshavmurthy et al. 2012; Silverstein et al. 2015). In the aforementioned studies, the background

symbionts often differed at the genus level and were typically replaced by the previously

dominant symbiont type once the stress had subsided (Berkelmans and van Oppen 2006; Jones et

al. 2008; LaJeunesse et al. 2010; Keshavmurthy et al. 2012; LaJeunesse 2020).

Thermally tolerant symbionts may arise through adaptation of symbiont populations,

even in the absence of evolutionary changes in the host. Large population sizes, short generation

times, and high mutation rates could greatly augment the standing genetic variation within

symbiont populations (van Oppen et al. 2011). This increased variation may allow for selection

of particular genotypes in response to elevated sea surface temperatures (van Oppen et al. 2011;

Torda et al. 2017; Quigley et al. 2018; Buerger et al. 2020). Numerous studies have established

that more thermally tolerant symbionts can be selected from within a single symbiont genus or

species (Howells et al. 2011; Chakravarti et al. 2017; Buerger et al. 2020). By sampling

symbionts in the genus *Cladocopium* (ITS type C1) from reefs with different temperature

regimes, Howells et al. (2011) showed that distinct symbiont genotypes had different thermal

tolerances. Symbionts from the warmer reef displayed increased performance and survival at

elevated temperatures compared to those from the cooler environment. Furthermore, juvenile

hosts inoculated with symbionts from the warmer reef performed better (grew faster, had less

bleaching, and had lower mortality) than those inoculated with symbionts from the cooler

region.

Chakravarti et al. (2017) and Buerger et al. (2020) presented evidence for selection

within a species by examining the adaptive capabilities of the symbionts within a monoclonal

culture of *Cladocopium goreaui*. Symbionts selected for thermal tolerance (i.e., those maintained

at 31°C for 2.5 to 4 years) performed better at 31°C than the original monoclonal culture that had

not undergone this selection treatment. Juvenile polyps infected with symbiont cells from the

elevated temperature grew at 27°C and 31°C, while polyps infected with the original culture

showed negative growth at 31°C. Infection with some strains of thermally selected cells,

however, did not reduce the rate or severity of bleaching compared to those infected with the

original culture (Buerger et al. 2020). These studies establish that selection for thermal tolerance

can occur over ecologically relevant time scales within at least some symbiont populations.

Given sufficient standing genetic variation, this is likely the case for other species within

Symbiodiniaceae and requires further inquiry (Chakravarti et al. 2017).

By starting with a single monoclonal culture, all genetic variation in the symbionts used

by Chakravarti et al. (2017) and Buerger et al. (2020) likely arose from *de novo* mutation.

Intraspecific markers have revealed numerous instances where host colonies harbor multiple

genotypes within a symbiont species (Santos et al. 2003, 2017; Howells et al. 2009; Kirk et al.

2009; Pettay and LaJeunesse 2007), suggesting the possibility for selection on pre-existing

genetic variation within *in hospite* populations, resulting from both mutation and recombination

(Baums et al. 2014; Chi et al. 2014; Wilkinson et al. 2015; Thornhill et al. 2017). Indeed, many

studies have found variation in thermal tolerance within a single symbiont species (Karim et al.

2015; Díaz-Almeyda et al. 2017; Grégoire et al. 2017; Bayliss et al. 2019). For example, Bayliss

et al. (2019) identified genetic and trait variation within and among *Breviolum* species isolated

from the octocoral *Antillogorgia bipinnata*. Here we ask if there are thermally tolerant genotypes

present in a symbiont species within a single host colony. We utilized the octocoral, *A. bipinnata*

and its host-specialist endosymbiotic algae, *Breviolum antillogorgium* as a model to study the

role of symbiont genetic variation in the thermal response of the holobiont. We first selected for

thermally tolerant symbionts from *in hospite* populations to capitalize on pre-existing genetic

variation. We then examined symbiont growth at two temperatures with an *in vitro* experiment

using these cultured isolates of *B. antillogorgium*. Subsequently, we examined whether thermal

tolerance in the symbiont could be conferred to the host using an *in hospite* experiment. The

presence of thermally tolerant symbionts within a population contained in a single host colony

expands the potential for the holobiont to ameliorate the consequences of increased temperatures,

potentially slowing the loss of these invaluable ecosystems.

126
Methods

Symbiont Isolation

Although culturing of Symbiodiniaceae is highly selective and rarely recovers the

symbionts dominant within the host (Santos et al. 2001; LaJeunesse 2002), *Breviolum*

antillogorgium (ITS2-type B1) has been successfully cultured and associates specifically with

octocorals in the genus *Antillogorgia* (Santos et al. 2004; Parkinson et al. 2015)*.* Symbiont cells

were isolated from sixteen adult colonies of *A. bipinnata* collected from 12-18 m depth at sites in

Elbow and Pickles Reefs, Florida Keys in September 2016 (Table 1). An approximately 2 cm

piece was sampled from each colony and homogenized in 2 mL of filtered seawater (FSW). Each

homogenate was passed through a series of meshes (250, 120, 70 and 20 μm), followed by

several 1 mL 0.22 μm FSW rinses for a final volume of 3-6 mL. Homogenates were spun at 800

rpm for five minutes. Supernatants were decanted, and the pellets resuspended in 10 mL of FSW.

This was repeated three additional times, followed by resuspending each pellet in 1 mL of f/2

media (Guillard and Ryther 1962). For each colony, 50 µL of the resultant cell suspension was

added to six flasks containing 30 mL of f/2 media. Three flasks of the symbionts from each

colony were maintained at 26°C, mimicking typical thermal conditions, and three were

maintained at 30°C to select for thermally tolerant strains from symbiont populations within a

single colony. Initial growth, observed as a small dot at the bottom of the flask, was immediately

transferred into a new flask with fresh media. This was repeated each time new growth was

observed over the first month, resulting in 131 cultures. All cultures were transferred to fresh

media monthly.

Genotype Identification

After two years (~500 asexual generations, assuming an average doubling time of ~1.5

days, based on growth rates from this study), total genomic DNA from 106 cultures and the host

source colonies was extracted using the 2X CTAB method (Coffroth et al. 1992), diluted to 5-10

ng µL^{-1} and the flanking region of the B7Sym15 microsatellite (Pettay and Lajeunesse 2007) was

sequenced (TACGEN, Richmond, California, USA). Sequences were compared to available

sequences on NCBI using BLASTn for symbiont species-level identification.

Cultures identified as *B. antillogorgium* (n=24) were amplified with six microsatellite

loci following published protocols: CA 6.38 (Santos et al. 2003), GV_1C (Andras et al. 2009),

B7Sym4, B7Sym8, B7Sym34, and B7Sym36 (Pettay and Lajeunesse 2007). Amplicons were

labeled with fluorescently tagged primers and visualized on a 6.5% polyacrylamide gel with the

LI-COR NEN Global IR2 DNA Sequencer (LI-COR Biotechnology Division) following Santos

et al. (2003). Allele sizes were scored by eye relative to size standards between 50 and 350 bp.

Based on the microsatellite results, five genetically distinct *B. antillogorgium* cultures (three

reared at 26°C and two reared at 30°C) were selected for use in the experiments outlined below

(Table 1).

In vitro Growth Experiments

Six replicates of each of the five cultures identified above were started by inoculating 18

mL of f/2 media with 10,000 cells mL^{-1}. Three of the replicates for each culture were

subsequently grown at 25.53 ± 0.89°C (mean ± SD; referred to as "26°C") and three were grown

at 29.84 ± 0.48°C (mean ±SD; referred to as "30°C") on a 12:12 h light:dark cycle. Light levels

were maintained at 120.58 ± 11.14 µmols m^{-2}s^{-1} (mean ±SD) (Philips Energy Advantage T5 HO

49W) and 103.79 ± 37.3 µmols m^{-2}s^{-1} (mean ±SD) (Philips T5 HO 24W/840) for the 26°C and

30°C incubators, respectively (Welch's Two Sample t-test, P = 0.13). Cell densities were

measured using six replicate hemocytometer cell counts every three days. Experiments were

continued until cell density peaked and exhibited a consistent decline in abundance. We used the

'growthrates' package ver. 0.8.1 (Petzoldt 2019) in R ver. 4.0.2 (R Core Team 2020) to fit

logistic growth curves to the cell counts over time, up to the maximum cell count for each

culture. We extracted values of r and K from the growthrates package and used these as

estimates of maximum population growth rate and carrying capacity, respectively. Samples of

each replicate were then preserved in 95% ethanol, from which DNA was extracted and

amplified following the methods detailed above to verify symbiont genotype at the end of the

experiment.

In hospite Experiments

Branches from adult colonies of _A. bipinnata,_ a gonochoric surface brooder (Kahng et al.

2011), were collected from Tennessee Reef, Florida Keys in November 2018 (N 24° 45.150' W

81° 45.275'). These branches were transported to flow-through seawater tanks at Keys Marine

Laboratory, where they were monitored daily for release of embryos. Brooded larvae were

collected from the colony surface on 16, 17, 22, 23, 25, and 26 November, washed three times in

0.22 μm filtered instant ocean (FIO), and transferred to Buffalo, NY. Following transport, larvae

(4-14 days old) were again washed in FIO to remove environmental contaminants and groups of

25 larvae were transferred to pre-conditioned polypropylene containers (volume = approx. 115

mL) with 50 mL of FIO. For pre-conditioning, the bottom and sides of each container were

roughened with sandpaper and then soaked for four weeks in 1.2 μm filtered seawater (FSW) to

establish a biofilm. Once larvae were added, water in the containers (FIO) was changed daily for

one week, then every other day until larvae settled and metamorphosed into polyps

(approximately 4 weeks). Containers of newly settled polyps (approximately 1-2 weeks post-

settlement) were then divided into five treatments (G1 through G5, Table 2) and each treatment

was inoculated with 1000 cells mL^{-1} of one of the five genotypes of *B. antillogorgium* utilized in

the *in vitro* experiments (Table 1). For each genotype, twelve containers were placed at 26ºC

(25.51 ± 1.10°C; mean ±SD) and eight at 30ºC (29.79 ± 0.28°C; mean ±SD) and maintained in a

12:12 h light:dark cycle. Initial inoculations occurred twice over a period of one week. Polyp

locations in the containers were mapped and survival for each individual polyp was recorded

every 3-4 days. As visual infection, indicated by a light brown coloration in the polyp, was not

observed within the first month following initial introduction of symbionts, inoculations were

then repeated with each water change and combined with feeding with Reefs-Roids (Polyplab,

Quebec, Canada) (2 mL of Reef-Roids solution at a concentration of 1 mg/mL FIO) to induce

symbiont uptake. Inoculations continued for an additional month, after which the experiment was

terminated due to overgrowth by an unidentified green alga which led to extensive polyp

mortality at that time. Following the termination of the experiment, surviving polyps were

collected, symbiont cell densities were calculated (as detailed above), and all collected polyps

were preserved in 95% ethanol. Symbiont DNA was extracted from collected polyps following

quantification of cell density and symbiont genotype was verified using the six microsatellite loci

described above.

Statistical Analyses

We analyzed the effects of temperature treatment and genotype on *in vitro* symbiont

maximum population growth rates and carrying capacity using separate two-way ANOVA,

including the interaction term between these factors. Data were log-transformed to meet

assumptions of homoscedasticity. In the *in hospite* experiment, we quantified polyp survivorship

as the proportion of surviving polyps at the end of one and two months post initial inoculation.

We analyzed one and two month survivorship as above; month one survivorship data were

arcsine transformed to meet assumptions of normality. We analyzed treatment and genotype

effects on symbiont cell densities as above; data were square-root transformed to meet

assumptions of normality. Assumptions of normality and homogeneity were checked using

residuals and Q-Q probability plots. All post-hoc pairwise comparisons were carried out using

Tukey's HSD (honest significance test) (Tukey 1949). *In vitro* statistical analyses were

conducted in R ver. 4.0.2 and *in hospite* analyses in R ver. 3.6.1 (R Core Team 2020). R scripts

are available at https://github.com/jessiepelosi/symbiont_tolerance and raw data are available at

https://www.bco-dmo.org/award/658940.

Results

Sequencing and Genotyping

Following two years of growth at either 26°C or 30°C, there were 131 viable cultures. As

noted above, culturing is notoriously selective and often not representative of the dominant

symbiont within the host, resulting in the recovery of background or cryptic symbionts (Santos et

al. 2001; LaJeunesse 2002; LaJeunesse et al. 2012; Parkinson et al. 2015). This was observed

here where the majority of cultures either could not be classified or aligned most closely with *B.*

minutum, a species commonly recovered from a range of host species where it is undetected by

normal techniques (LaJeunesse et al 2012; Parkinson et al 2015). Twenty-four of the cultures

were identified as *B. antillogorgium* based on a comparison of B7Sym15 microsatellite flanking

region sequences to available sequences on NCBI. Using microsatellite data, we found that 21 of

the 24 *B. antillogorgium* cultures (10 reared at 26°C and 11 reared at 30°C) contained a single

genotype. Among these 21 cultures, we identified five distinct *B. antillogorgium* genotypes

(three reared at 26°C and two reared at 30°C). These five genotypes were used in the *in vitro* and

in hospite experiments.

***In vitro* growth experiments**

Cultures reached their maximum cell density in 28-40 days and maintained steady state

growth for at least several days afterward (Fig 2a, Online Resource 1). The effect of temperature

on maximum population growth rate was dependent on the symbiont genotype

(Temperature*Genotype: $F_{4,20} = 30.2$, $p < 0.001$). At 30°C the mean growth rate for four of the

five genotypes decreased from that at 26°C, but to varying magnitudes; the mean growth was

slightly higher at 30°C for one genotype (G5) (Fig. 2a). The effect of temperature on carrying

capacity also depended on the symbiont genotype (Genotype*Temperature: $F_{4,20} = 17.90$, $p <$

0.001). Genotype responses to temperature ranged from a decrease in carrying capacity (G1,

G3), to little response (G2, G4), to an increase in carrying capacity (G5) (Fig. 2b). The time for

populations to reach their peak abundance was dependent on the interaction between temperature

and symbiont genotype (Temperature*Genotype: $F_{4,20}$ = 6.57, p = 0.002).

***In hospite* polyp experiments**

We found a significant interaction between the effects of temperature and genotype on

the symbiont cell densities in polyps at the end of the experiment ($F_{4,264}$ = 6.342, p < 0.001).

Average symbiont densities in polyps increased at higher temperature in some genotypes but not

others (Table 3, Fig. 3). At one month post-infection, polyp survivorship was not significantly

affected by symbiont genotype ($F_{4,85}$ = 0.64, p = 0.636), nor by treatment temperature ($F_{1,85}$ =

2.61, p = 0.110), and there was not a significant interaction term ($F_{4,85}$ = 0.673, p = 0.613) (Fig.

4a, b). At two months post-infection, survivorship decreased dramatically in both temperature

treatments (Fig. 4a, c), with those polyps at 30°C having significantly lower survivorship than

those reared at 26°C ($F_{1,72}$ = 26.90, p < 0.001). There was also a significant effect of symbiont

genotype on survivorship at the two-month mark ($F_{4,72}$ = 2.95, p = 0.026). There was not a

significant interaction between temperature and genotype on two-month survival ($F_{4,72}$ = 0.56, p

= 0.692).

Discussion

Cultured symbionts isolated from standing genetic variation within host colonies revealed

the presence of multiple genotypes capable of growth at 30°C, with three of the five unique

genotypes used here isolated from a single host colony (Table 1). Genotypes varied in their

response to thermal stress (i.e., growth rate and carrying capacity, Fig. 2) and there was no clear

evidence of selection for increased thermal tolerance over time in cultures reared at 30°C for

over 500 generations (2 years) (Figs. 2-4). However, the ability of all tested genotypes to grow

positively at 30°C suggests that *B. antillogorgium,* as a species, is thermally tolerant.

***In vitro* growth experiments**

By isolating genotypes that were able to grow at elevated temperatures from standing

genetic variation among *in hospite* populations, we established that thermally tolerant symbiont

genotypes are present within the host. We maintained these cultures at 30°C for two years to

select for long-term growth at elevated temperatures, anticipating that these cultures would adapt

and outperform other genotypes at elevated temperature. Unlike previous studies that found

evidence for thermal adaptation in Symbiodiniaceae after just one year of exposure to increased

temperatures (*Cladocopium* - Chakravarti et al. 2017; *Fugacium, Gerakladium, Symbiodinium* -

Chakravarti and van Oppen 2018), long-term growth at 30°C did not yield better performance for

B. antillogorgium at 30°C (as compared to 26°C). Our results suggest that local adaptation to this

temperature did not occur over this time period. Interestingly, growth rates were often counter to

our expectations, with G3 (historical thermal regime of 26°C) showing little change in growth

rate under increased temperature and G4 (historical thermal regime of 30°C) showing a decrease

in growth rate at 30°C compared to growth at 26°C (Fig. 2).

While growth rate serves as a good proxy for fitness of symbiont populations, carrying

capacity is likely a proxy for efficiency of resource use or metabolite buildup. We found

variation among genotypes in carrying capacity, but interestingly, also found variation in the

response of carrying capacity to increased temperature. Two of the five genotypes (G2, G5)

showed an increase in mean carrying capacity at higher temperature, but one genotype (G3)

showed a marked decrease in carrying capacity at higher temperature (Fig. 2b). These

differences were unrelated to the temperature at which cultures were historically grown, and the

three genotypes isolated from the same host showed the most variation in response (G3, G4, G5).

The metabolic theory of ecology and at least one experiment on phytoplankton predict decreases

in carrying capacity with increasing temperature (Savage et al. 2004; Bernhardt et al. 2018), but

only two of our five genotypes matched this prediction. However, higher metabolic rates should

result in increased growth rates. There was a weak positive correlation between r and K (t = 2.91,

df = 28, P = 0.007, r = 0.482) and these two variables combined to influence the time to reach

peak density. Three genotypes (G1, G4, G5) took longer to reach peak density at higher

temperature, but the other two genotypes reached peak density more quickly at higher

temperature. Although one genotype (G5) with long-term growth at 30°C showed increases in

growth rate and carrying capacity at the increased temperature, perhaps reflective of adaptation

to increased temperature, this result should be interpreted with caution, as the other genotype

with long-term growth at 30°C (G4) did not show the same response (Fig. 2b). The ability of all

cultures to survive and grow positively at 30°C for several months, regardless of historical

temperature, suggests that these genotypes of *B. antillogorgium* possess pre-existing, broad-scale

thermal tolerance, possibly due to plastic acclimatory responses.

Prior studies examining the potential for symbionts to adapt to elevated temperatures

have focused on a single genotype (Chakravarti et al. 2017; Chakravarti and van Oppen 2018),

but in the present study we used several genotypes within the same species, selected from pre-

existing genetic variation within *in hospite* populations. Variability in response to elevated

temperatures observed among the genotypes used in our experiments highlights the potential role

of intraspecific genetic variation in responding to thermal stress and further underscores the

importance of examining both intrageneric and intraspecific variation in thermal responses.

Other *in vitro* studies (Chakravarti et al. 2017; Chakravarti and van Oppen 2018; Buerger et al.

2020) have explored how dinoflagellate symbionts in other genera respond to temperatures

higher (31°C-34°C) than used in our experiments. It is possible that the rapid adaptation seen in

previous studies might also occur in *B. antillogorgium* if these genotypes were grown at higher

temperatures. Further work should explore the limits of thermal tolerance in *B. antillogorgium* to

temperatures greater than 30°C as these increasingly elevated temperatures may soon become

commonplace on coral reefs (Heron et al. 2016; van Hooidonk et al. 2016; Hughes et al. 2018;

Oliver et al. 2018).

***In hospite* polyp experiments**

Antillogorgia bipinnata polyps were infected by each symbiont genotype at both

experimental temperatures. The fact that polyps took up all genotypes is perhaps not surprising,

since *B. antillogorgium* is a host-specialist for *Antillogorgia* species (Parkinson et al. 2015), and

the symbiont cultures used in this study were initially isolated from *Antillogorgia* host colonies.

While Newkirk et al. (2020) found that certain strains of *Breviolum* were unable to establish a

stable symbiosis at elevated temperatures with the model scyphozoan *Cassiopeia xamachana*, by

focusing our study on a known symbiont of *A. bipinnata*, we demonstrate that the successful

establishment of the symbiosis may not always be limited by environmental temperature.

Although there were no consistent differences between *in hospite* symbiont densities,

final symbiont cell density varied with genotype and temperature, indicating that symbionts were

either taken up or grew at different rates within hosts. The lack of a consistent effect of

temperature on symbiont density within polyps suggests that the establishment of the symbiosis

at 30°C may be as stable as those at ambient temperatures (26°C). Given the significant

interaction between treatment temperature and symbiont genotype on symbiont density within

polyps, the ability of the symbiont to infect and thrive in the host likely depends on multiple

factors (Fig. 3), with potential for variation in the establishment and maintenance of the

symbiosis (Davy et al. 2012; Hawkins et al. 2016).

Our work highlights that *in vitro* studies of symbiont success under heat stress should not

be used as direct predictors of holobiont success. Despite this caveat, the overall thermal

tolerance recorded *in vitro* was observed in the polyp experiment, even though performance *in*

vitro was not directly mirrored *in hospite*. All treatments had relatively high polyp survival at

one month, with a more rapid decline in survivorship over the second month (Fig. 4a, b). Polyps

showed sustained survivorship over an extended period of heat exposure, with some surviving

polyps in all treatments at 56 days post-inoculation in spite of the algal overgrowth. This is a

substantially longer exposure to heat stress than has been previously reported for *in hospite*

experiments (i.e., Chakravarti et al. 2017: 28 days; Buerger et al. 2020: 7 days). Our results

suggest that the observed thermal tolerance exhibited by our symbiont types may have

contributed to or enhanced the thermal tolerance of the holobiont, although the long-term value

of this has yet to be determined.

In this study, the host-symbiont pairs exhibited thermal tolerance. *Breviolum* cultures

grown at 30°C did not show enhanced performance at 30°C (as compared to 26°C). This

indicates that selection for thermal tolerance of certain genotypes within the colonies likely

occurred prior to the isolation of these symbionts. It is possible that thermal tolerance has been

selected for in *Breviolum* over much longer time periods than those in this experiment in

response to historical thermal events. Studies of both long- and short-term responses to thermal

stress show that these organisms can successfully function at elevated temperatures, suggesting

that adaptation or acclimatization may ameliorate holobiont responses to thermal stress.

Holobionts may have adapted to variation in temperature regimes over thousands of years

(Smith-Keune and van Oppen 2006; Howells et al. 2013, 2016; Hume et al. 2015; Kenkel and

Matz 2016). For example, corals found in naturally high-temperature environments (e.g.,

Persian/Arabian Gulf, warm water pools) show increased thermal tolerance compared to

conspecifics from cooler habitats (Howells et al. 2013, 2016; Bay and Palumbi 2014; Hume et al.

2015). Rapid adaptation or acclimatization may increase thermal tolerance over shorter time

scales, exemplified by reduced levels of bleaching in the years following extreme bleaching

events (Maynard et al. 2008; Guest et al. 2012; Pratchett et al. 2013; Logan et al. 2014;

McClanahan et al. 2014; Brown et al. 2015; Coles et al. 2018).

Ultimately, our results indicate that there exists standing genetic variation within

symbiont populations, which in this case, contained genotypes that were thermally tolerant.

These symbionts may be vital for the holobiont's survival, although the basis for this thermal

tolerance requires further investigation. Various other factors including host-symbiont GxG

interactions and alternative selection pressures may influence holobiont success at elevated

temperatures (Baird et al. 2009; Chakravarti et al. 2017; Gabay et al. 2019). Additionally,

interactions between the entire coral microbiome (including endosymbiotic dinoflagellates,

bacteria, viruses and fungi) may be key in determining thermal stress responses of the holobiont

(Vega Thurber et al. 2009; Littman et al. 2011; Bourne et al. 2016).

Role of the symbiont and implications for Caribbean reefs

Most previous studies of the capacity for rapid thermal adaptation in symbionts have

focused on Pacific strains of Symbiodiniaceae. Our results shed new light on the existence of

thermally tolerant Caribbean coral endosymbionts in genera in addition to those thermally

tolerant species within the genus *Durusdinium* (LaJeunesse et al. 2014). We provide key

evidence that even in the absence of rapid adaptation to elevated temperatures, thermal tolerance

may be widespread in some *Breviolum* species. It is likely that broad thermal tolerance is an

inherent trait among these genotypes and perhaps among other species within *Breviolum*,

particularly those within the B1 (ITS2-type) lineage (Pleistocene (B1) radiation, *sensu*

Lajeunesse 2005; Parkinson et al. 2015) such as *B. antillogorgium*. During thermal stress, for

example, the octocoral *Briareum asbestinum* lost its dominant *Breviolum* ITS2 type B19

symbiont (Poland et al. 2013), which was replaced by a *Breviolum* ITS2 type B1 (Lewis and

Coffroth 2004; Poland and Coffroth 2019). Transitioning from a dominant symbiont species with

lower thermal tolerance to a more thermally tolerant cryptic species during bleaching events may

buffer Caribbean octocorals from increasing temperatures.

Coral reefs are in decline worldwide and some regions, such as the Caribbean, are

experiencing a phase shift from scleractinian-dominated to octocoral-dominated ecosystems

(Ruzicka et al. 2013; Lenz et al. 2015; Lasker et al. 2020). This pattern has been attributed, in

part, to high levels of octocoral recruitment (Lasker et al. 2020). However, bleaching in

octocorals is rarely observed during mass bleaching events (Lasker et al. 1984; Lasker 2003;

Prada et al. 2010) and if it is observed, most species show a high level of resilience (Prada et al.

2010). The fact that species within *Breviolum* are the dominant symbionts in Caribbean

octocorals (van Oppen et al. 2005; Goulet et al. 2008; Franklin et al. 2012), and that some

symbiont species within this genus appear to be thermally tolerant, as shown here, may

contribute to this resilience. Additional studies are needed to determine the prevalence of these

thermally tolerant symbionts, that tend to exist at background level, and their role in mitigating

the effects of increasing temperatures on reef cnidarians. The survival of Caribbean reefs may be

reliant on evolutionary rescue (Gomulkiewicz and Holt 1995) perpetrated by selection for these

more thermally tolerant symbiont genotypes.

Acknowledgements

We would like to thank H. Franklin, undergraduate volunteers, staff of the KML, the FKNMS and the FWC (Permit FKNMS-2014-018-A2 and SAL-18-2052-SR, respectively). This work was supported by NSF-OCE-1559286 (MAC) and NSF-OCE-1559105 (CPt), a UB Honors Research & Creativity Grant (JAP) and a UB CURCA Award (KME). We are grateful to H. Lasker, A. Martinez, and C. Wells for comments on previous drafts of this manuscript.

Conflict of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

Andras JP, Kirk NL, Coffroth MA, Harvell CD (2009) Isolation and characterization of microsatellite loci in Symbiodinium B1/B184, the dinoflagellate symbiont of the Caribbean sea fan coral, *Gorgonia ventalina*. Mol. Ecol. Resour. 9:989–993

Baird AH, Bhagooli R, Ralph PJ, Takahashi S (2009) Coral bleaching: the role of the host. Trends Ecol Evol (Amst) 24:16–20

Baker AC (2003) Flexibility and Specificity in Coral-Algal Symbiosis: Diversity, Ecology, and Biogeography of *Symbiodinium*. Annu. Rev. Ecol. Evol. Syst. 34:661–689

Baums IB, Devlin-Durante MK, LaJeunesse TC (2014) New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population genetic studies. Mol. Ecol. 23:4203–4215

Bayliss SLJ, Scott ZR, Coffroth MA, terHorst CP (2019) Genetic variation in *Breviolum antillogorgium*, a coral reef symbiont, in response to temperature and nutrients. Ecol. Evol. 9:2803–2813

Bay RA, Palumbi SR (2014) Multilocus adaptation associated with heat resistance in reef-building corals. Curr. Biol. 24:2952–2956

Berkelmans R, van Oppen MJH (2006) The role of zooxanthellae in the thermal tolerance of corals: a "nugget of hope" for coral reefs in an era of climate change. Proc. Biol. Sci. 273:2305–2312

Bernhardt JR, Sunday JM, O'Connor MI (2018) Metabolic Theory and the Temperature-Size Rule Explain the Temperature Dependence of Population Carrying Capacity. Am. Nat. 192:687–697

Bourne DG, Morrow KM, Webster NS (2016) Insights into the Coral Microbiome: Underpinning
the Health and Resilience of Reef Ecosystems. Annu. Rev. Microbiol. 70:317–340

Brown BE, Dunne RP, Edwards AJ, Sweet MJ, Phongsuwan N (2015) Decadal environmental
'memory' in a reef coral? Mar. Biol. 162:479–483

Buerger P, Alvarez-Roa C, Coppin CW, Pearce SL, Chakravarti LJ, Oakeshott JG, Edwards OR,
van Oppen MJH (2020) Heat-evolved microalgal symbionts increase coral bleaching
tolerance. Sci. Adv. 6:eaba2498

Chakravarti LJ, Beltran VH, van Oppen MJH (2017) Rapid thermal adaptation in
photosymbionts of reef-building corals. Glob. Chang. Biol. 23:4675–4688

Chakravarti LJ, van Oppen MJH (2018) Experimental evolution in coral photosymbionts as a
tool to increase thermal tolerance. Front. Mar. Sci. 5:

Chi J, Parrow MW, Dunthorn M (2014) Cryptic sex in *Symbiodinium* (Alveolata, Dinoflagellata)
is supported by an inventory of meiotic genes. J. Eukaryot. Microbiol. 61:322–327

Coffroth MA, Lasker HR, Diamond ME, Bruenn JA, Bermingham E (1992) DNA fingerprints of
a gorgonian coral: a method for detecting clonal structure in a vegetative species. Mar. Biol.
114:317–325

Coles SL, Bahr KD, Rodgers KS, May SL, McGowan AE, Tsang A, Bumgarner J, Han JH
(2018) Evidence of acclimatization or adaptation in Hawaiian corals to higher ocean
temperatures. PeerJ 6:e5347

Correa AMS, McDonald MD, Baker AC (2009) Development of clade-specific *Symbiodinium*
primers for quantitative PCR (qPCR) and their application to detecting clade D symbionts
in Caribbean corals. Mar. Biol. 156:2403–2411

Davy SK, Allemand D, Weis VM (2012) Cell biology of cnidarian-dinoflagellate symbiosis.
Microbiol. Mol. Biol. Rev. 76:229–261

Díaz-Almeyda EM, Prada C, Ohdera AH, Moran H, Civitello DJ, Iglesias-Prieto R, Carlo TA,
LaJeunesse TC, Medina M (2017) Intraspecific and interspecific variation in
thermotolerance and photoacclimation in *Symbiodinium* dinoflagellates. Proc. Biol. Sci.
284:

Eakin CM, Lough JM, Heron SF (2009) Climate variability and change: monitoring data and
evidence for increased coral bleaching stress. In: van Oppen M.J.H., Lough J.M. (eds) Coral
Bleaching. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 41–67

Franklin EC, Stat M, Pochon X, Putnam HM, Gates RD (2012) GeoSymbio: a hybrid, cloud-
based web application of global geospatial bioinformatics and ecoinformatics for
Symbiodinium-host symbioses. Mol. Ecol. Resour. 12:369–373

Gabay Y, Parkinson JE, Wilkinson SP, Weis VM, Davy SK (2019) Inter-partner specificity
limits the acquisition of thermotolerant symbionts in a model cnidarian-dinoflagellate
symbiosis. ISME J. 13:2489–2499

Gomulkiewicz R, Holt RD (1995) When does evolution by natural selection prevent extinction?
Evolution 49:201

Goulet TL (2006) Most corals may not change their symbionts. Mar. Ecol. Prog. Ser. 321:1–7

Goulet TL, Simmons C, Goulet D (2008) Worldwide biogeography of *Symbiodinium* in tropical octocorals. Mar. Ecol. Prog. Ser. 355:45–58

Grégoire V, Schmacka F, Coffroth MA, Karsten U (2017) Photophysiological and thermal tolerance of various genotypes of the coral endosymbiont *Symbiodinium* sp. (Dinophyceae). J. Appl. Phycol. 29:1893–1905

Guest JR, Baird AH, Maynard JA, Muttaqin E, Edwards AJ, Campbell SJ, Yewdall K, Affendi YA, Chou LM (2012) Contrasting patterns of coral bleaching susceptibility in 2010 suggest an adaptive response to thermal stress. PLoS ONE 7:e33353

Guillard RR, Ryther JH (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. Can. J. Microbiol. 8:229–239

Hawkins TD, Hagemeyer JCG, Warner ME (2016) Temperature moderates the infectiousness of two conspecific *Symbiodinium* strains isolated from the same host population. Environ. Microbiol. 18:5204–5217

Heron SF, Maynard JA, van Hooidonk R, Eakin CM (2016) Warming Trends and Bleaching Stress of the World's Coral Reefs 1985-2012. Sci. Rep. 6:38402

Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral reefs. Mar. Freshwater Res. 50:839

Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatziolos ME (2007) Coral reefs under rapid climate change and ocean acidification. Science 318:1737–1742

Hoegh-Guldberg O (2011) Coral reef ecosystems and anthropogenic climate change. Reg. Environ. Change 11:215–227

Hoegh-Guldberg O, Poloczanska ES, Skirving W, Dove S (2017) Coral Reef Ecosystems under Climate Change and Ocean Acidification. Front. Mar. Sci. 4:

Howells EJ, van Oppen MJH, Willis BL (2009) High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*. Coral Reefs 28:215–225

Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH (2011) Coral thermal tolerance shaped by local adaptation of photosymbionts. Nat. Clim. Chang. 2:116–120

Howells EJ, Berkelmans R, van Oppen MJH, Willis BL, Bay LK (2013) Historical thermal regimes define limits to coral acclimatization. Ecology 94:1078–1088

Howells EJ, Abrego D, Meyer E, Kirk NL, Burt JA (2016) Host adaptation and unexpected symbiont partners enable reef-building corals to tolerate extreme temperatures. Glob. Chang. Biol. 22:2702–2714

Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Wilson SK (2018)

Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. Science
359:80–83

Hume BCC, D'Angelo C, Smith EG, Stevens JR, Burt J, Wiedenmann J (2015) *Symbiodinium*
thermophilum sp. nov., a thermotolerant symbiotic alga prevalent in corals of the world's
hottest sea, the Persian/Arabian Gulf. Sci. Rep. 5:8562

Jones AM, Berkelmans R, van Oppen MJH, Mieog JC, Sinclair W (2008) A community change
in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field
evidence of acclimatization. Proc. Biol. Sci. 275:1359–1365

Kahng SE, Benayahu Y, Lasker HR (2011) Sexual reproduction in octocorals. Mar. Ecol. Prog.
Ser. 443:265–283

Karim W, Nakaema S, Hidaka M (2015) Temperature effects on the growth rates and
photosynthetic activities of *Symbiodinium* cells. J. Mar. Sci. Eng. 3:368–381

Kenkel CD, Matz MV (2016) Gene expression plasticity as a mechanism of coral adaptation to a
variable environment. Nat. Ecol. Evol. 1:14

Keshavmurthy S, Hsu C-M, Kuo C-Y, Meng P-J, Wang J-T, Chen CA (2012) Symbiont
communities and host genetic structure of the brain coral *Platygyra verweyi*, at the outlet of
a nuclear power plant and adjacent areas. Mol. Ecol. 21:4393–4407

Kirk NL, Andras JP, Harvell CD, Santos SR, Coffroth MA (2009) Population structure of
Symbiodinium sp. associated with the common sea fan, *Gorgonia ventalina*, in the Florida
Keys across distance, depth, and time. Mar. Biol. 156:1609–1623

LaJeunesse T (2002) Diversity and community structure of symbiotic dinoflagellates from
Caribbean coral reefs. Mar. Biol. 141:387–400

LaJeunesse TC (2005) "Species" radiations of symbiotic dinoflagellates in the Atlantic and Indo-
Pacific since the Miocene-Pliocene transition. Mol. Biol. Evol. 22:570–581

LaJeunesse TC, Smith R, Walther M, Pinzón J, Pettay DT, McGinley M, Aschaffenburg M,
Medina-Rosas P, Cupul-Magaña AL, Pérez AL, Reyes-Bonilla H, Warner ME (2010) Host-
symbiont recombination versus natural selection in the response of coral-dinoflagellate
symbioses to environmental disturbance. Proc. Biol. Sci. 277:2925–2934

LaJeunesse TC, Parkinson JE, Reimer JD (2012) A genetics-based description of *Symbiodinium*
minutum sp. nov. and *S. psygmophilum* sp. nov. (Dinophyceae), two dinoflagellates
symbiotic with cnidaria. J. Phycol. 48:1380–1391

LaJeunesse TC, Wham DC, Pettay DT, Parkinson JE, Keshavmurthy S, Chen CA (2014)
Ecologically differentiated stress-tolerant endosymbionts in the dinoflagellate
genus*Symbiodinium* (Dinophyceae) Clade D are different species. Phycologia 53:305–319

LaJeunesse T (2020) Zooxanthellae. Curr Biol 30:R1110–R1113

Lasker HR, Peters EC, Coffroth MA (1984) Bleaching of reef coelenterates in the San Blas
Islands, Panama. Coral Reefs 3:183–190

Lasker HR (2003) Zooxanthella densities within a Caribbean octocoral during bleaching and
non-bleaching years. Coral Reefs 22:23–26

Lasker HR, Martínez-Quintana Á, Bramanti L, Edmunds PJ (2020) Resilience of octocoral
forests to catastrophic storms. Sci. Rep. 10:4286

Lenz EA, Bramanti L, Lasker HR, Edmunds PJ (2015) Long-term variation of octocoral
populations in St. John, US Virgin Islands. Coral Reefs 34:1099–1109

Lewis CL, Coffroth MA (2004) The acquisition of exogenous algal symbionts by an octocoral
after bleaching. Science 304:1490–1492

Littman R, Willis BL, Bourne DG (2011) Metagenomic analysis of the coral holobiont during a
natural bleaching event on the Great Barrier Reef. Environ. Microbiol. Rep. 3:651–660

Logan CA, Dunne JP, Eakin CM, Donner SD (2014) Incorporating adaptive responses into
future projections of coral bleaching. Glob. Chang. Biol. 20:125–139

Maynard JA, Anthony KRN, Marshall PA, Masiri I (2008) Major bleaching events can lead to
increased thermal tolerance in corals. Mar. Biol. 155:173–182

McClanahan TR, Ateweberhan M, Darling ES, Graham NAJ, Muthiga NA (2014) Biogeography
and change among regional coral communities across the Western Indian Ocean. PLoS
ONE 9:e93385

Mieog JC, van Oppen MJH, Cantin NE, Stam WT, Olsen JL (2007) Real-time PCR reveals a
high incidence of *Symbiodinium* clade D at low levels in four scleractinian corals across the
Great Barrier Reef: implications for symbiont shuffling. Coral Reefs 26:449–457

Muscatine L, Cernichiari E (1969) Assimilation of photosynthetic products of zooxanthellae by a
reef coral. Biol. Bull. 137:506–523

Muscatine L, Porter JW (1977) Reef Corals: Mutualistic Symbioses Adapted to Nutrient-Poor
Environments. Bioscience 27:454–460

Newkirk CR, Frazer TK, Martindale MQ, Schnitzler CE (2020) Adaptation to bleaching: are
thermotolerant symbiodiniaceae strains more successful than other strains under elevated
temperatures in a model symbiotic cnidarian? Front. Microbiol. 11:822

Oliver JK, Berkelmans R, Eakin CM (2018) Coral bleaching in space and time. Coral Bleaching.
Springer, pp 27–49

Parkinson JE, Coffroth MA, LaJeunesse TC (2015) New species of Clade B *Symbiodinium*
(Dinophyceae) from the greater Caribbean belong to different functional guilds:
S. aenigmaticum sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and
S. pseudominutum sp. nov. J. Phycol. 51:850–858

Pettay DT, Lajeunesse TC (2007) Microsatellites from clade B *Symbiodinium* spp. specialized
for Caribbean corals in the genus *Madracis*. Mol. Ecol. Notes 7:1271–1274

Petzoldt T (2019) growthrates. Available at: https://github.com/tpetzoldt/growthrates

Poland DM, Mansfield JM, Hannes AR, Lewis CLF, Shearer TL, Connelly SJ, Kirk NL,
Coffroth MA (2013) Variation in Symbiodinium communities in juvenile Briareum
asbestinum (Cnidaria: Octocorallia) over temporal and spatial scales. Mar. Ecol. Prog. Ser.
476:23–37

Poland DM, Coffroth MA (2017) Trans-generational specificity within a cnidarian–algal
symbiosis. Coral Reefs 36:119–129

Poland DM, Coffroth MA (2019) Host growth and survivorship varies with endosymbiotic algal
partner in developing cnidarians. Mar. Ecol. Prog. Ser. 612:87–100

Prada C, Weil E, Yoshioka PM (2010) Octocoral bleaching during unusual thermal stress. Coral
Reefs 29:41–45

Pratchett MS, McCowan D, Maynard JA, Heron SF (2013) Changes in bleaching susceptibility
among corals subject to ocean warming and recurrent bleaching in Moorea, French
Polynesia. PLoS ONE 8:e70443

Quigley K.M., Baker A.C., Coffroth M.A., Willis B.L., van Oppen M.J.H. (2018) Bleaching
Resistance and the Role of Algal Endosymbionts. Ch. 6 pp. 111-151. In: Coral Bleaching:
Patterns, Processes, Causes and Consequences eds; J. Lough and M. van Oppen. Springer
Ecological Series. Cham, Switzerland

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Astria. URL https://www.R-project.org/.

Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS,
Macaulay KA, Bartlett LA, Meyers MK, Colee J (2013) Temporal changes in benthic
assemblages on Florida Keys reefs 11 years after the 1997/1998 El Niño. Mar. Ecol. Prog.
Ser. 489:125–141

Santos SR, Taylor DJ, Coffroth MA (2001) Genetic comparisons of freshly isolated versus
cultured symbiotic dinoflagellates: implications for extrapolating to the intact symbiosis. J.
Phycol. 37:900–912

Santos SR, Guti   rrez-Rodr   guez C, Lasker HR, Coffroth MA (2003) *Symbiodinium* sp.
associations in the gorgonian *Pseudopterogorgia elisabethae* in the Bahamas: high levels of
genetic variability and population structure in symbiotic dinoflagellates. Mar. Biol.
143:111–120

Santos SR, Shearer TL, Hannes AR, Coffroth MA (2004) Fine-scale diversity and specificity in
the most prevalent lineage of symbiotic dinoflagellates (*Symbiodinium*, Dinophyceae) of the
Caribbean. Mol. Ecol. 13:459–469

Savage VM, Gillooly JF, Brown JH, Charnov EL (2004) Effects of body size and temperature on
population growth. Am. Nat. 163:429–441

Silverstein RN, Correa AMS, Baker AC (2012) Specificity is rarely absolute in coral-algal
symbiosis: implications for coral response to climate change. Proc. Biol. Sci. 279:2609–
2618

Silverstein RN, Cunning R, Baker AC (2015) Change in algal symbiont communities after
bleaching, not prior heat exposure, increases heat tolerance of reef corals. Glob. Chang.
Biol. 21:236–249

Smith-Keune C, van Oppen M (2006) Genetic structure of a reef-building coral from thermally
distinct environments on the Great Barrier Reef. Coral Reefs 25:493–502

Thornhill DJ, Howells EJ, Wham DC, Steury TD, Santos SR (2017) Population genetics of reef
coral endosymbionts (*Symbiodinium*, Dinophyceae). Mol. Ecol. 26:2640–2659

Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N,
Foret S, Matz M, Miller DJ, Moya A, Putnam HM, Ravasi T, van Oppen MJH, Thurber RV,
Vidal-Dupiol J, Voolstra CR, Watson S-A, Whitelaw E, Munday PL (2017) Rapid adaptive
responses to climate change in corals. Nat. Clim. Chang. 7:627–636

Tukey JW (1949) Comparing individual means in the analysis of variance. Biometrics 5:99–
114.

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadia G, Raymundo L, Williams G,
Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of
the Paris Agreement. Sci. Rep. 6:39666

van Oppen MJH, Mieog JC, Sánchez CA, Fabricius KE (2005) Diversity of algal endosymbionts
(zooxanthellae) in octocorals: the roles of geography and host relationships. Mol. Ecol.
14:2403–2417

van Oppen MJH, Souter P, Howells EJ, Heyward A, Berkelmans R (2011) Novel genetic
diversity through somatic mutations: fuel for adaptation of reef corals? Diversity (Basel)
3:405–423

Vega Thurber R, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA, Angly F,
Dinsdale E, Kelly L, Rohwer F (2009) Metagenomic analysis of stressed coral holobionts.
Environ. Microbiol. 11:2148–2163

Wilkinson SP, Fisher PL, van Oppen MJH, Davy SK (2015) Intra-genomic variation in
symbiotic dinoflagellates: recent divergence or recombination between lineages? BMC
Evol. Biol. 15:325

Tables

Table 1. Five genotypes of *Breviolum antillogorgium* selected for *in vitro* and *in hospite* experiments. Letter color refers to selection temperature, blue indicating cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the start of the experiment. Cultures were isolated from colonies collected at Elbow Reef, Florida Keys.

Culture	Genotype	Selection Tempt	Source Colony	Reef	Depth	Location	Accession No.
16-0590F	G-1	26 C	7	Elbow	18 m	N 25 07.925 W 80 15.717	XXX
16-0875	G-2	26 C	5	Elbow	11 m	N 25 07.956 W 80 15.810	XXX
16-1631	G-3	26 C	8	Elbow	18 m	N 25 07.925 W 80 15.717	XXX
16-0587	G-4	30 C	8	Elbow	18 m	N 25 07.925 W 80 15.717	XXX
16-0763	G-5	30 C	8	Elbow	18 m	N 25 07.925 W 80 15.717	XXX

Table 2. Experimental set up for the *in hospite* study. Letter color refers to selection temperature,

blue indicating cultures reared at 26 C and red indicating cultures reared at 30 C two years prior

to the start of the experiment.

Treatment	Mean # Polyps/container	# polyps/treatment	Containers/ Treatment
G-1 @ 26	13	160	12
G-2 @ 26	13	159	12
G-3 @ 26	13	163	12
G-4 @ 26	13	166	12
G-5 @ 26	14	169	12
G-1 @ 30	8	55	8
G-2 @ 30	8	67	8
G-3 @ 30	8	58	8
G-4 @ 30	8	56	8
G-5 @ 30	8	58	8

710 **Table 3.** P-values for significant pairwise comparisons from Tukey's HSD post-hoc tests of

711 symbiont cell densities in polyps (*in hospite* experiment). No other pairwise comparisons were

712 significant.

Genotype at Temperature	P-values
G1 at 30°C > G3 at 30°C	0.0039
G3 at 26° C >G5 at 26°C	0.0229
G1 at 30°C > G2 at 26°C	0.031
G1 at 30°C >G4 at 30°C	0.045
G1 at 30°C > G5 at 26°C	0.003
G4 at 26°C > G3 at 30°C	0.039

713
714

Figures

 Figure 1. Experimental design for a.) *in vitro* growth experiment and b.) *in hospite* experiment.

 Figure colors refer to temperature treatment (blue, maintained at 26°C; red, maintained at 30°C),

 with tip figures colored based on treatment (not historical temperature).

Figure 2. a.) Mean logistic growth fits to *in vitro* cell densities of each genotype of *Breviolum*

antillogorgium over time. Blue represents growth at 26°C and red at 30°C. b.) Estimated mean

(+/- SE) maximum symbiont population growth rates (r) and C) estimated mean (+/- SE)

735 carrying capacities (K). Genotype colors refer to the historical temperature at which they were

736 grown, blue indicating cultures maintained at 26°C and red indicating cultures maintained at

737 30°C two years prior to the start of the experiment. Bars represent medians, boxes the

738 interquartile range, and whiskers the range of the data.

Figure 3. Symbiont densities within *Antillogorgia bipinnata* polyps inoculated with five

symbiont genotypes at 26°C and 30°C. Statistically significant differences for these measures in

Table 3. Genotype color refers to historical temperature, blue indicating cultures maintained at

26°C and red indicating cultures maintained at 30°C two years prior to the start of the

experiment. Boxes represent interquartile range and whiskers the range of the data. Points

indicate outliers.

Figure 4. a.) Mean polyp survivorship over time for *A. bipinnata* polyps inoculated with five

symbiont genotypes at 26°C and 30°C. Error bars are ±SE. Blue lines refer to treatments at 26°C

and red lines refer to treatments at 30°C. Differences in average survivorship at the end of the b.)

760   first month and c.) second month. Genotype color refers to historical temperature, blue indicating

761   cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the

762   start of the experiment. Boxes represent interquartile range and whiskers the range of the data.

763   Points indicate outliers.