Thermally tolerant symbionts may explain Caribbean octocoral resilience to heat stress

Jessie Pelosi[1, #,*], Katherine M. Eaton[1,^,*], Samantha Mychajliw[2], Casey P. terHorst[3], and Mary

Alice Coffroth[4,+]

[1]Department of Biological Sciences, University at Buffalo, Buffalo, NY 14260, [2]Department of

Environment and Sustainability, University at Buffalo, Buffalo, NY 14260, [3]Department of

Biology, California State University, Northridge, Northridge, CA 91330, [4]Department of

Geology, University at Buffalo, Buffalo, NY 14260

*These authors contributed equally to this work.

Current addresses:

[#]Department of Biology, University of Florida, Gainesville, FL 32608

[^]Department of Biological Sciences, Auburn University, Auburn, AL 36849

[+]Corresponding author: coffroth@buffalo.edu

Keywords: symbiosis, Symbiodiniaceae, octocorals, thermal tolerance

Abstract

Coral reef ecosystems are under threat from the frequent and severe impacts of anthropogenic climate change, particularly rising sea surface temperatures. The effects of thermal stress may be ameliorated by adaptation and/or acclimation of the host, symbiont, or holobiont (host + symbiont) to increased temperatures. We examined the role of the symbiont in promoting thermal tolerance of the holobiont, using *Antillogorgia bipinnata* (octocoral host) and *Breviolum antillogorgium* (symbiont) as a model system. We identified five distinct genotypes of *B. antillogorgium* from symbiont populations isolated from *Antillogorgia* colonies in the Florida Keys. Three symbiont genotypes were cultured and maintained at 26°C (ambient historical temperature) and two were cultured and maintained at 30°C (elevated historical temperature) for two years. Following culturing, we analyzed the growth rate and carrying capacity of each symbiont genotype at both ambient and elevated temperatures in culture (*in vitro*). All genotypes grew well at both temperatures, indicating thermal tolerance among these *B. antillogorgium* cultures. Prior culturing at the elevated temperature, however, did not result in increased thermal tolerance. We then inoculated juvenile *A. bipinnata* polyps with each of the five symbiont genotypes, and reared these polyps at both ambient and elevated temperatures (*in hospite* experiment). All genotypes were able to establish symbioses with polyps in both temperature treatments. Survivorship of polyps at 30°C was significantly lower than survivorship at 26°C, but all treatments had surviving polyps at 56 days post-infection, suggestive of broad thermal tolerance in *B. antillogorgium,* which may play a part in the increased resilience of Caribbean octocorals during heat stress events.

Introduction

In recent years, anthropogenic climate change has severely impacted coral reefs, which

comprise some of the most biodiverse and economically important ecosystems on Earth (Hoegh-

Guldberg 2011; Hoegh-Guldberg et al. 2017). The success of reef ecosystems in oligotrophic

tropical waters is due to their obligate endosymbioses with dinoflagellate algae in the family

Symbiodiniaceae (Muscatine and Porter 1977). In this mutualistic relationship, the coral host

relies on photosynthate produced by its intracellular symbionts to thrive (Muscatine and

Cernichiari 1969; Muscatine and Porter 1977). In exchange for these metabolites, the symbiotic

algae recover host nitrogenous waste. This relationship, which is vital to the survival of the reef

ecosystem, is currently threatened by anthropogenic environmental changes.

Coral bleaching, or the loss of algal symbionts, occurs due to a variety of environmental

stressors including changes in salinity, pH, light intensity, and, most commonly, water

temperature (Hoegh-Guldberg 1999). Most reef ecosystems now routinely encounter

temperatures at or above their thermal maximum each year (Hoegh-Guldberg et al. 2007; Eakin

et al. 2009), with bleaching events becoming more frequent and severe (Heron et al. 2016; van

Hooidonk et al. 2016; Hughes et al. 2018; Oliver et al. 2018). As ocean temperatures continue to

increase, it is essential to understand if and how corals and their symbionts can respond to the

present threat.

The holobiont (host + symbiont) can respond to rising seawater temperatures via changes

in the symbiont community within a single host, with the basic premise that hosts associating

with more thermally tolerant symbionts will be more likely to survive bleaching events (i.e.,

shuffling and/or switching, *sensu* Baker 2003). Although most coral species host a single

dominant symbiont type (Goulet 2006; LaJeunesse 2020), many hosts also harbor background or

cryptic symbionts that are often undetected by traditional techniques (Mieog et al. 2007; Correa

et al. 2009; Silverstein et al. 2012). During thermal stress, a more thermally tolerant background

symbiont can become the dominant symbiont type, potentially conferring thermal tolerance to

the holobiont (Berkelmans and van Oppen 2006; Jones et al. 2008; LaJeunesse et al. 2009;

Keshavmurthy et al. 2012; Silverstein et al. 2015). In the aforementioned studies, the background

symbionts often differed at the genus level and were typically replaced by the previously

dominant symbiont type once the stress had subsided.

Thermally tolerant holobionts may arise through adaptation of symbiont populations,

even in the absence of evolutionary changes in the host. Large population sizes, short generation

times, and high mutation rates could greatly augment the standing genetic variation within

symbiont populations (van Oppen et al. 2011). With greater standing variation, natural selection

is more likely to produce populations with greater thermal tolerance (van Oppen et al. 2011;

Torda et al. 2017; Quigley et al. 2018; Buerger et al. 2020). Numerous studies have established

that more thermally tolerant symbionts can be selected from within a single symbiont genus or

species (Howells et al. 2011; Chakravarti et al. 2017; Buerger et al. 2020). By sampling

symbionts in the genus *Cladocopium* (ITS type C1) from reefs with different temperature

regimes, Howells et al. (2011) showed that distinct symbiont genotypes had different thermal

tolerances. Symbionts from the warmer reef displayed increased performance and survival at

elevated temperatures as compared to those from the cooler environment. Furthermore, juvenile

hosts inoculated with symbionts from the warmer reef performed better (grew faster, had less

bleaching, and had lower mortality) than those inoculated with symbionts from the cooler region.

Chakravarti et al. (2017) and Buerger et al. (2020) presented evidence for selection

within a species by examining the adaptive capabilities of the symbionts within a monoclonal

culture of *Cladocopium goreaui*. Symbionts selected for thermal tolerance (i.e., those maintained

at 31°C for 2.5 to 4 years) performed better at 31°C than the original monoclonal culture that had

not undergo this selection treatment. Juvenile polyps infected with symbiont cells from the

elevated temperature grew at 27°C and 31°C, while polyps infected with the original culture

showed negative growth at 31°C. Infection with some strains of thermally selected cells,

however, did not reduce the rate or severity of bleaching compared to those infected with the

original culture (Buerger et al. 2020). These studies establish that selection for thermal tolerance

can occur over ecologically relevant time scales within at least some symbiont populations.

Given sufficient standing genetic variation, this is likely the case for other species within

Symbiodiniaceae and requires further inquiry (Chakravarti et al. 2017).

By starting with a single monoclonal culture, all genetic variation in the symbionts used

by Chakravarti et al. (2017) and Buerger et al. (2020) likely arose from *de novo* mutation.

Intraspecific markers have revealed numerous instances where host colonies harbor multiple

genotypes within a symbiont species (Santos et al. 2003; Pettay and LaJeunesse 2007; Howells et

al. 2009; Kirk et al. 2009), suggesting the possibility for selection on pre-existing genetic

variation within *in hospite* populations, resulting from both mutation and recombination (Baums

et al. 2014; Chi et al. 2014; Wilkinson et al. 2015; Thornhill et al. 2017). Indeed, many studies

have found variation in thermal tolerance within a single symbiont species (Parkinson and

Baums 2014; Karim et al. 2015; Díaz-Almeyda et al. 2017; Grégoire et al. 2017; Bayliss et al.

2019). For example, Bayliss et al. (2019) identified genetic and trait variation within and among

Breviolum species isolated from the octocoral *Antillogorgia bipinnata*.

Here we ask if there are thermally tolerant genotypes present in a symbiont species

within a single host colony. We utilized the octocoral *A. bipinnata* and its host-specialist

endosymbiotic algae, *Breviolum antillogorgium*, as a model to study the role of symbiont genetic

variation in the thermal response of the holobiont. We first selected for thermally tolerant

symbionts from *in hospite* populations to capitalize on pre-existing genetic variation. We then

examined symbiont growth at two temperatures with an *in vitro* experiment using these cultured

isolates of *B. antillogorgium*. Subsequently, we examined whether thermal tolerance in the

symbiont could be conferred to the host using an *in hospite* experiment. The presence of

thermally tolerant symbionts within a population contained in a single host colony expands the

potential for the holobiont to ameliorate the consequences of increased temperatures, potentially

slowing the loss of these invaluable ecosystems.

Methods

Symbiont isolation

Although culturing of Symbiodiniaceae is highly selective and rarely recovers the

symbionts dominant within the host (Santos et al. 2001; LaJeunesse 2002), *Breviolum*

antillogorgium (ITS2-type B1) has been successfully cultured and associates specifically with

octocorals in the genus *Antillogorgia* (Santos et al. 2004; Parkinson et al. 2015). Symbiont cells

were isolated from sixteen adult *Antillogorgia* colonies collected from 12-18 m depth at sites in

Elbow and Pickles Reefs, Florida Keys in September 2016 (Table 1). A sample of each host

colony was preserved in 95% ethanol for comparison of dominant host symbionts with those

isolated. An approximately 2 cm piece was sampled from each colony and homogenized in 2 mL

of filtered seawater (FSW). Each homogenate was passed through a series of mesh filters (250,

120, 70 and 20 µm), followed by several 1 mL 0.22 µm FSW rinses for a final volume of 3-6

mL. Homogenates were spun at 800 rpm for five minutes. Supernatants were decanted, and the

pellets were resuspended in 10 mL of FSW. This was repeated three additional times, followed

by resuspending each pellet in 1 mL of f/2 media (Guillard and Ryther 1962). For each colony,

50 µL of the resultant cell suspension was added to six flasks containing 30 mL of f/2 media.

Three flasks of the symbionts from each colony were maintained at 26°C, mimicking typical

thermal conditions, and three were immediately placed at 30°C to select for thermally tolerant

genotypes that can survive at an elevated temperature. An elevated temperature of 30°C was

chosen based on the historical thermal regime of the collection site during November/December,

when newly settled *A. bipinnata* polyps acquire symbionts (Williams and Miller 2015; Online

Resource 1). Initial growth, observed as a small dot at bottom of the flask, was immediately

transferred into a new flask with fresh media. This was repeated each time new growth was

observed in any flask over the first month, resulting in 131 cultures. All cultures were transferred

to fresh media monthly.

Genotype identification

After two years (approximately 735-1084 asexual generations, assuming a doubling time

of 1.01-1.97 days, based on growth rates from this study), total symbiont genomic DNA from

106 cultures and the host source colonies was extracted using the 2X CTAB method (Coffroth et

al. 1992) and diluted to 5-10 ng µL^{-1}, and the flanking region of the B7Sym15 microsatellite

(Pettay and LaJeunesse 2007) was sequenced (TACGEN, Richmond, California, USA).

Sequences were compared to available sequences on NCBI using BLASTn for symbiont species-

level identification.

Cultures identified as *B. antillogorgium* (n=24) and samples from the source colonies of

these cultures were amplified with five microsatellite loci following published protocols: CA

6.38 (Santos et al. 2003), GV_1C (Andras et al. 2009), B7Sym8, B7Sym34, and B7Sym36

(Pettay and LaJeunesse 2007). Amplicons were labeled with fluorescently tagged primers and

visualized on a 6.5% polyacrylamide gel with the LI-COR NEN Global IR2 DNA Sequencer

(LI-COR Biotechnology Division) following Santos et al. (2003). Allele sizes were scored by

eye relative to size standards between 50 and 350 bp. Based on the microsatellite results, 21 *B.*

antillogorgium cultures were unialgal and contained one of five genotypes. Ultimately, this

resulted in the occurrence of three distinct genotypes (referred to here as G1, G2, and G3) from

one or more of the cultures reared at 26°C and two genotypes (referred to here as G4 and G5)

from one or more of the cultures reared at 30°C (see Results). One of each of these five

genotypes were used in the experiments outlined below (Table 1).

***In vitro* growth experiments**

Six replicates of each of the five cultures identified above were started by inoculating 18

mL of f/2 media with 10,000 cells mL^{-1}. Three of the replicates for each culture were

subsequently grown at 25.53 ± 0.89°C (mean ± SD; referred to as "26°C") and three were grown

at 29.84 ± 0.48°C (mean ±SD; referred to as "30°C") on a 12:12 h light:dark cycle. Light levels

were maintained at 120.58 ± 11.14 µmols m^{-2}s^{-1} (mean ± SD) (Philips Energy Advantage T5 HO

49W) and 103.79 ± 37.3 µmols m^{-2}s^{-1} (mean ± SD) (Philips T5 HO 24W/840) for the 26°C and

30°C incubators, respectively (Welch's Two Sample t-test, p = 0.13). Cell densities were

measured using six replicate hemocytometer cell counts every three days. Experiments were

continued until cell density peaked and exhibited a consistent decline in abundance. We used the

'growthrates' package ver. 0.8.1 (Petzoldt 2019) in R ver. 4.0.2 (R Core Team 2020) to fit

logistic growth curves to the cell counts over time, up to the maximum cell count for each

culture. We extracted values of r and K from the 'growthrates' package and used these as

estimates of maximum population growth rate and carrying capacity, respectively. Samples of

each replicate were then preserved in 95% ethanol, from which DNA was extracted and

amplified following the methods detailed above to verify symbiont genotype at the end of the

experiment.

***In hospite* polyp experiments**

Branches from adult colonies of *A. bipinnata,* a gonochoric surface brooder (Kahng et al.

2011), were collected from Tennessee Reef, Florida Keys in November 2018 (N 24° 45.150' W

81° 45.275'). These branches were transported to flow-through seawater tanks at Keys Marine

Laboratory, where they were monitored daily for release of embryos. Aposymbiotic brooded

larvae were collected from the colony surface on 16, 17, 22, 23, 25, and 26 November, washed

three times in 0.22 μm filtered instant ocean (FIO), and transferred to Buffalo, NY. Following

transport, larvae (4-14 days old) were again washed in FIO to remove environmental

contaminants and groups of 25 larvae were transferred to pre-conditioned polypropylene

containers (volume = approximately 115 mL) with 50 mL of FIO. For pre-conditioning, the

bottom and sides of each container were roughened with sandpaper and then soaked for four

weeks in 1.2 μm filtered seawater (FSW) to establish a biofilm. Once larvae were added, water in

the containers (FIO) was changed daily for one week, then every other day until larvae settled

and metamorphosed into polyps (approximately 4 weeks). Containers of newly settled polyps

(approximately 1-2 weeks post-settlement) were then divided into five treatments (G1 through

G5, Online Resource 2) and each treatment was inoculated with 1000 cells mL^{-1} of one of the

five genotypes of *B. antillogorgium* utilized in the *in vitro* experiments (Table 1). For each

genotype, 9-10 containers were placed at 26ºC (25.51 ± 1.10°C; mean ± SD) and 6-8 at 30ºC

(29.79 ± 0.28°C; mean ± SD) and maintained in a 12:12 h light:dark cycle. Initial inoculations

occurred twice over a period of one week. Polyp locations in the containers were mapped and

survival for each individual polyp was recorded every 3-4 days with water changes. Visual

infection, indicated by a light brown coloration in the polyp, was limited within the first month

following initial introduction of symbionts. To increase the likelihood of the establishment of the

symbiosis, inoculations were repeated with each water change during the second month and

combined with feeding with Reefs-Roids (Polyplab, Quebec, Canada) (2 mL of Reef-Roids

solution at a concentration of 1 mg/mL FIO) to induce symbiont uptake. Inoculations continued

throughout the second month, and establishment of the symbiosis through visual observation was

recorded throughout the experiment. An unidentified green alga, first observed in three

containers on Day 42, but subsequently found in several additional containers on Days 56-59, led

to extensive polyp mortality over time. After 69 days, all surviving polyps were collected,

symbiont cell density was determined (as detailed above), and tissue was preserved in 95%

ethanol. Symbiont DNA was extracted from preserved tissue and symbiont genotype was

verified using the five microsatellite loci described above.

Statistical analyses

We analyzed the effects of temperature treatment and genotype on *in vitro* symbiont

maximum population growth rates and carrying capacity using separate two-way ANOVAs for

each dependent variable (r and K), including the interaction term between these factors

(temperature and genotype). Data were log-transformed to meet assumptions of

homoscedasticity for the *in vitro* analyses. For the *in hospite* experiment, we quantified polyp

survivorship as the proportion of surviving polyps after 31 and 52 days post-initial inoculation.

These time points were prior to the observation of widespread contamination by the unidentified

green alga, to avoid the possibility that this affected survivorship. Containers in which the

contaminating algae was observed prior to Day 52 (n=5) were excluded from survivorship

analyses beginning at the first observation of contamination. We analyzed survivorship over the

first 31 days and then from Day 32 to Day 52 as described above; with survivorship for Days 1-

31 arcsine transformed to meet assumptions of normality (survivorship data for Days 32-52 were

not transformed as the raw data fit the assumptions of the model without transformation). We

analyzed temperature and genotype effects on visually observed infection at Days 31 and 52 as

well as on symbiont cell densities at the end of the experiment (Day 69) as above. Symbiont cell

densities were square-root transformed to meet assumptions of the model. Assumptions of

normality and homogeneity were checked using residuals and Q-Q probability plots. All post-

hoc pairwise comparisons were carried out using Tukey's HSD (honest significance test) (Tukey

1949). *In vitro* statistical analyses were conducted in R ver. 4.0.2 and *in hospite* analyses in R

ver. 3.6.1 (R Core Team 2020). R scripts are available at

www.github.com/jessiepelosi/symbiont_tolerance and raw data are available at https://www.bco-

dmo.org/award/658940.

Results

Sequencing and genotyping

Following two years of growth at either 26ºC or 30ºC, there were 131 viable cultures.

The majority of cultures (106 of 131) either could not be classified or aligned most closely with

B. minutum, a species commonly recovered from a range of species where it is undetected by

normal techniques (LaJeunesse et al. 2012; Parkinson et al. 2015). Twenty-four of the cultures

were identified as *B. antillogorgium* based on a comparison of B7Sym15 microsatellite flanking

region sequences to available sequences on NCBI. Three of these cultures contained multiple

genotypes. Each of the remaining 21 *B. antillogorgium* cultures were unialgal and contained one

of five genotypes. Ten *B. antillogorgium* cultures reared at 26ºC were either genotype G1, G2 or

G3, while 11 *B. antillogorgium* cultures reared at 30ºC were genotype G4 or G5. These five

genotypes were used in the *in vitro* and *in hospite* experiments. Based on microsatellite

genotyping, one of the five genotypes (G3) was the same as the dominant genotype in the source

colony; the other four genotypes represented background symbionts as is common in culturing

from host tissue (Santos et al. 2001; LaJeunesse 2002; LaJeunesse et al. 2012; Parkinson et al.

2015).

***In vitro* growth experiments**

Cultures reached their maximum cell density in 28-40 days and maintained this density

for at least several days afterward (Fig 2a, Online Resource 3). The effect of temperature on

maximum population growth rate was dependent on the symbiont genotype

(Temperature*Genotype: $F_{4,20} = 30.20$, $p < 0.001$). At 30ºC the mean growth rate for two of the

five genotypes (G1 and G2) was significantly lower than that at 26ºC; there was no significant

effect of temperature for the remaining three genotypes (Fig. 2a, b). The effect of temperature on

carrying capacity also depended on the symbiont genotype (Temperature*Genotype: $F_{4,20} =$

17.90, $p < 0.001$). One genotype (G3) showed a decrease in carrying capacity at the elevated

temperature, but the remaining genotypes showed non-significant responses to temperature (Fig.

2c). The time for populations to reach their peak abundance was also dependent on the

interaction between temperature and symbiont genotype (Temperature*Genotype: $F_{4,20} = 6.57$, p

$= 0.002$).

In hospite **polyp experiments**

For the first 31 days, we did not find a significant impact of genotype ($F_{4,73}$ = 1.84, p =

0.130), temperature ($F_{1,73}$ = 0.03, p = 0.861), nor Temperature*Genotype ($F_{4,73}$ = 0.83, p = 0.508)

on the proportion of visually infected polyps (Fig. 3a). Similarly, between 32 and 52 days, we

did not find a significant impact of genotype ($F_{4,56}$ = 0.66, p = 0.623), temperature ($F_{1,56}$ = 3.77, p

= 0.057), nor Temperature*Genotype ($F_{4,56}$ = 1.22, p = 0.312) on the proportion of visually

infected polyps (Fig. 3b). We found a significant interaction between the effects of temperature

and genotype on the symbiont cell densities in polyps at the end of the experiment, when the vast

majority of polyps (range 66-100% of polyps per treatment) were infected (69 days post first

inoculation, $F_{4,264}$ = 6.34, p < 0.001). Average symbiont densities in polyps varied between

genotypes, depending on temperature at Day 69 (termination of the experiment, Fig. 3c). Polyp

survivorship at the end of the first 31 days was not significantly affected by symbiont genotype

($F_{4,74}$ = 0.81, p = 0.522), nor by treatment temperature ($F_{1,74}$ = 3.05, p = 0.085), and there was not

a significant interaction term ($F_{4,74}$ = 0.76, p = 0.574) (Fig. 4a, b). Polyp survivorship at the end

of 52 days decreased dramatically in both temperature treatments (Fig. 4a, c), with those polyps

at 30°C having significantly lower survivorship than those reared at 26°C ($F_{1,68}$ = 15.64, p <

0.001). There was not a significant effect of genotype ($F_{4,68}$ = 2.11, p = 0.089) nor the interaction

between temperature and genotype on survival for the second period (Temperature*Genotype,

$F_{4,68}$ = 0.13, p = 0.970). Including the five containers removed after Day 42 (due to algal

contamination) did not change the results. Although survivorship results are only reported prior

to the appearance of an unidentified green alga, all treatments were maintained for 69 days. At

that time, all treatments except G4 at 30ºC had surviving polyps (range 2-70 polyps per

treatment).

Discussion

Cultured symbionts isolated from standing genetic variation within host colonies revealed the presence of multiple genotypes capable of growth at 30ºC, with three of the five unique genotypes used here isolated from a single host colony (Table 1). However, there was no clear evidence of selection for increased thermal tolerance in cultures reared at 30ºC for over 700 generations (2 years) compared to those reared at 26ºC over this time (Figs. 2-4). Genotypic variability was clear as genotypes differed in their response to thermal stress (i.e., growth rate and carrying capacity, Fig. 2). Regardless of this variability, all tested genotypes showed growth at 30ºC, suggesting that *Breviolum antillogorgium,* as a species, is thermally tolerant.

In vitro growth experiments

By isolating genotypes that were able to grow at elevated temperatures from standing genetic variation among *in hospite* populations, we established that thermally tolerant symbiont genotypes are present within the host. We maintained these cultures at 30°C for two years to select for long-term growth at elevated temperatures, anticipating that these cultures would adapt and outperform other genotypes at elevated temperatures. Unlike previous studies that found evidence for thermal adaptation in Symbiodiniaceae after one to 2.5 years of exposure to increased temperatures (*Cladocopium* - Chakravarti et al. 2017; *Fugacium, Gerakladium, Symbiodinium* - Chakravarti and van Oppen 2018), a history of long-term growth at 30°C did not yield better performance for *B. antillogorgium* at 30°C (as compared to 26°C). Our results suggest that local adaptation to this temperature did not occur over this time period. It is possible that all examined genotypes possessed pre-existing adaptations to temperatures at or above 30ºC, as over the past decade and a half, if not longer, this threshold has been frequently exceeded during the summer months in the Florida Keys (Online Resource 1).

Growth rate serves as a good proxy for fitness of symbiont populations. Carrying

capacity may serve as a proxy for efficiency of resource use because it suggests that more

organisms can be sustained at the same level of resource. We found variation among genotypes

in carrying capacity, but interestingly, also found variation in the response of carrying capacity to

increased temperature. One genotype (G3) showed a significant decrease in carrying capacity at

higher temperature (Fig. 2c), but the remaining four genotypes showed non-significant responses

to temperature. The ability of all cultures to survive and grow positively at 30°C for several

months, regardless of historical temperature, suggests that these genotypes of *B. antillogorgium*

possess pre-existing, broad thermal tolerance, possibly due to plastic acclimatory responses.

Prior studies examining the potential for symbionts to adapt to elevated temperatures

have focused on a single genotype (Chakravarti et al. 2017; Chakravarti and van Oppen 2018),

but in the present study we used several genotypes within the same species, selected from pre-

existing genetic variation within *in hospite* populations. Variability in response to elevated

temperatures observed among the genotypes used in our experiments highlights the potential role

of intraspecific genetic variation in responding to thermal stress and further underscores the

importance of examining both intrageneric and intraspecific variation in thermal responses.

Other *in vitro* studies (Chakravarti et al. 2017; Chakravarti and van Oppen 2018; Buerger et al.

2020) have explored how dinoflagellate symbionts in other genera respond to temperatures

higher (31°C-34°C) than used in our experiments. It is possible that the rapid adaptation seen in

previous studies might also occur in *B. antillogorgium* if these genotypes were grown at higher

temperatures. It is clear that elevated temperatures in excess of 30°C are becoming frequent

during the summer months in the Florida Keys (Online Resource 1) and further work should

explore the limits of thermal tolerance in *B. antillogorgium* to temperatures greater than 30°C.

***In hospite* polyp experiments**

There were no significant differences in the percent of visually infected polyps between

genotypes and temperature treatments through Day 52. In general, the percent of infected polyps

on Day 52 was largely congruent with the average number of symbiont cells per polyp at the end

of the experiment; those genotypes that had a greater proportion of visibly infected polyps also

had a greater average number of symbiont cells per polyp (Fig. 3b, c). Final symbiont cell

density varied with genotype and temperature, indicating that symbionts were either taken up or

grew at different rates within hosts. The lack of a consistent effect of temperature on symbiont

density within polyps suggests that the establishment of the symbiosis at 30°C may be as stable

as those at ambient temperatures (26°C). Given the significant interaction between treatment

temperature and symbiont genotype on symbiont density within polyps at the end of the

experiment (Day 69), the ability of the symbiont to infect and thrive in the host likely depends on

multiple factors, with potential for variation in the establishment and maintenance of the

symbiosis (Davy et al. 2012; Hawkins et al. 2016). It is clear that biotic interactions (e.g.,

recognition and phagocytosis of algal symbionts, Davy et al. 2012) drive the establishment,

while abiotic factors, such as temperature, may dictate the maintenance of the symbiosis (e.g.,

Rädecker et al. 2021).

The overall thermal tolerance recorded *in vitro* was observed in the polyp experiment,

even though performance *in vitro* was not directly mirrored *in hospite* (i.e., those symbiont

genotypes that had the highest growth rates *in vitro* did not necessarily confer the highest

survivorship under the same conditions *in hospite*). All treatments had relatively high polyp

survival early in the experiment, with a more rapid decline in survivorship towards the end (Fig.

4). Polyps showed sustained survivorship over an extended period of heat exposure, with some

surviving polyps in all treatments at 56 days post-inoculation. Furthermore, after 69 days, all

treatments except G4 at 30ºC had surviving polyps. This is a substantially longer exposure to

heat stress than has been previously reported for *in hospite* experiments (i.e., Chakravarti et al.

2017: 28 days; Buerger et al. 2020: 7 days). Our results suggest that the observed thermal

tolerance exhibited by our symbiont types may have contributed to or enhanced the thermal

tolerance of the holobiont, although the long-term value of this has yet to be determined.

365   In this study, the host-symbiont pairs exhibited thermal tolerance. At the elevated

temperature, however, hosts infected with *Breviolum* cultures that had been reared at 30°C did

not consistently outperform their counterparts (i.e., hosts infected with cultures reared at 26°C).

This indicates that selection for thermal tolerance of certain genotypes likely occurred prior to

the isolation of these symbionts. Increasingly elevated temperatures are becoming more

commonplace on coral reefs (Heron et al. 2016; van Hooidonk et al. 2016; Hughes et al. 2018;

Oliver et al. 2018), and repeated exposure to these elevated temperatures may have selected for

symbiont genotypes capable of surviving such extremes. Ultimately, our results indicate that

there exists standing genetic variation within symbiont populations, which in this case, contained

genotypes that were thermally tolerant. These symbionts may be vital for the holobiont's

survival, although the basis for this thermal tolerance requires further investigation.

Role of the symbiont and implications for Caribbean reefs

377   Most previous studies of the capacity for rapid thermal adaptation in symbionts have

focused on Pacific strains of Symbiodiniaceae. Our results shed new light on the existence of

thermally tolerant Caribbean coral endosymbionts in genera in addition to those thermally

tolerant species within the genus *Durusdinium* (LaJeunesse et al. 2014). We provide key

evidence that even in the absence of rapid adaptation to elevated temperatures, thermal tolerance

may be widespread in some *Breviolum* species. It is likely that broad thermal tolerance is an

inherent trait among these genotypes and perhaps among other species within *Breviolum*,

particularly those within the B1 (ITS2-type) lineage (Pleistocene (B1) radiation, *sensu*

LaJeunesse 2005; Parkinson et al. 2015) such as *B. antillogorgium*. Under induced bleaching

and/or thermal stress, for example, the octocoral *Briareum asbestinum* lost its dominant

Breviolum ITS2 type B19 symbiont, which was replaced by a *Breviolum* ITS2 type B1 species

(Lewis and Coffroth 2004; Poland and Coffroth 2019). Transitioning from a dominant symbiont

species with lower thermal tolerance to a more thermally tolerant cryptic species during

bleaching events may buffer Caribbean octocorals from increasing temperatures.

Coral reefs are in decline worldwide and some regions, such as the Caribbean, are

experiencing a phase shift from scleractinian-dominated to octocoral-dominated ecosystems

(Ruzicka et al. 2013; Lenz et al. 2015; Lasker et al. 2020). This pattern has been attributed, in

part, to high levels of octocoral recruitment (Lasker et al. 2020). However, bleaching in

octocorals is rarely observed during mass bleaching events (Lasker et al. 1984; Lasker 2003;

Prada et al. 2010) and if it is observed, most species show a high level of resilience (Prada et al.

2010). The fact that species within *Breviolum* are the dominant symbionts in Caribbean

octocorals (LaJeunesse 2002; van Oppen et al. 2005; Goulet et al. 2008), and that some symbiont

species within this genus appear to be thermally tolerant, as shown here, may contribute to this

resilience. Additional studies are needed to determine the prevalence of these thermally tolerant

symbionts, which may exist at background level, and their role in mitigating the effects of

increasing temperatures on reef cnidarians. The survival of Caribbean reefs may be reliant on

evolutionary rescue (Gomulkiewicz and Holt 1995) perpetrated by selection for these more

thermally tolerant symbiont genotypes within a given species; a process that could be

particularly important for specialist hosts that are not able to harbor different species of

symbionts.

Acknowledgements

We would like to thank H. Franklin, undergraduate volunteers, staff of the KML, the FKNMS

and the FWC (Permit FKNMS-2014-018-A2 and SAL-18-2052-SR, respectively). This work

was supported by NSF-OCE-1559286 (MAC) and NSF-OCE-1559105 (CPt), a UB Honors

Research & Creativity Grant (JAP), and a UB CURCA Award (KME). We are grateful to D.

Williams for providing the NOAA temperature data and to H. Lasker, A. Martinez, C. Wells, and

two anonymous reviewers for helpful and insightful comments on previous drafts of this

manuscript.

Conflict of Interest

416 On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

Andras JP, Kirk NL, Coffroth MA, Harvell CD (2009) Isolation and characterization of
microsatellite loci in *Symbiodinium* B1/B184, the dinoflagellate symbiont of the Caribbean
sea fan coral, *Gorgonia ventalina*. Mol. Ecol. Resour. 9:989–993

Baker AC (2003) Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and
biogeography of *Symbiodinium*. Annu. Rev. Ecol. Evol. Syst. 34:661–689

Baums IB, Devlin-Durante MK, LaJeunesse TC (2014) New insights into the dynamics between
reef corals and their associated dinoflagellate endosymbionts from population genetic
studies. Mol. Ecol. 23:4203–4215

Bayliss SLJ, Scott ZR, Coffroth MA, terHorst CP (2019) Genetic variation in *Breviolum*
antillogorgium, a coral reef symbiont, in response to temperature and nutrients. Ecol. Evol.
9:2803–2813

Berkelmans R, van Oppen MJH (2006) The role of zooxanthellae in the thermal tolerance of

corals: a "nugget of hope" for coral reefs in an era of climate change. Proc. Biol. Sci.
273:2305–2312

Buerger P, Alvarez-Roa C, Coppin CW, Pearce SL, Chakravarti LJ, Oakeshott JG, Edwards OR,
van Oppen MJH (2020) Heat-evolved microalgal symbionts increase coral bleaching
tolerance. Sci. Adv. 6:eaba2498

Chakravarti LJ, Beltran VH, van Oppen MJH (2017) Rapid thermal adaptation in
photosymbionts of reef-building corals. Glob. Chang. Biol. 23:4675–4688

Chakravarti LJ, van Oppen MJH (2018) Experimental evolution in coral photosymbionts as a
tool to increase thermal tolerance. Front. Mar. Sci. 5:227

Chi J, Parrow MW, Dunthorn M (2014) Cryptic sex in *Symbiodinium* (Alveolata, Dinoflagellata)
is supported by an inventory of meiotic genes. J. Eukaryot. Microbiol. 61:322–327

Coffroth MA, Lasker HR, Diamond ME, Bruenn JA, Bermingham E (1992) DNA fingerprints of
a gorgonian coral: a method for detecting clonal structure in a vegetative species. Mar. Biol.
114:317–325

Correa AMS, McDonald MD, Baker AC (2009) Development of clade-specific *Symbiodinium*
primers for quantitative PCR (qPCR) and their application to detecting clade D symbionts
in Caribbean corals. Mar. Biol. 156:2403–2411

Davy SK, Allemand D, Weis VM (2012) Cell biology of cnidarian-dinoflagellate symbiosis.
Microbiol. Mol. Biol. Rev. 76:229–261

Díaz-Almeyda EM, Prada C, Ohdera AH, Moran H, Civitello DJ, Iglesias-Prieto R, Carlo TA,
LaJeunesse TC, Medina M (2017) Intraspecific and interspecific variation in
thermotolerance and photoacclimation in *Symbiodinium* dinoflagellates. Proc. Biol. Sci.
284:20171767

Eakin CM, Lough JM, Heron SF (2009) Climate variability and change: monitoring data and
evidence for increased coral bleaching stress. In: van Oppen M.J.H., Lough J.M. (eds) Coral
Bleaching. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 41–67

Gomulkiewicz R, Holt RD (1995) When does evolution by natural selection prevent extinction?
Evolution 49:201

Goulet TL, Simmons C, Goulet D (2008) Worldwide biogeography of *Symbiodinium* in tropical
octocorals. Mar. Ecol. Prog. Ser. 355:45–58

Goulet TL (2006) Most corals may not change their symbionts. Mar. Ecol. Prog. Ser. 321:1–7

Grégoire V, Schmacka F, Coffroth MA, Karsten U (2017) Photophysiological and thermal
tolerance of various genotypes of the coral endosymbiont *Symbiodinium* sp. (Dinophyceae).
J. Appl. Phycol. 29:1893–1905

Guillard RR, Ryther JH (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt,

and *Detonula confervacea* (Cleve) Gran. Can. J. Microbiol. 8:229–239

Hawkins TD, Hagemeyer JCG, Warner ME (2016) Temperature moderates the infectiousness of two conspecific *Symbiodinium* strains isolated from the same host population. Environ. Microbiol. 18:5204–5217

Heron SF, Maynard JA, van Hooidonk R, Eakin CM (2016) Warming trends and bleaching stress of the world's coral reefs 1985-2012. Sci. Rep. 6:38402

Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatziolos ME (2007) Coral reefs under rapid climate change and ocean acidification. Science 318:1737–1742

Hoegh-Guldberg O, Poloczanska ES, Skirving W, Dove S (2017) Coral reef ecosystems under climate change and ocean acidification. Front. Mar. Sci. 4:158

Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral reefs. Mar. Freshwater Res. 50:839

Hoegh-Guldberg O (2011) Coral reef ecosystems and anthropogenic climate change. Reg. Environ. Change 11:215–227

Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH (2011) Coral thermal tolerance shaped by local adaptation of photosymbionts. Nat. Clim. Chang. 2:116–120

Howells EJ, van Oppen MJH, Willis BL (2009) High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*. Coral Reefs 28:215–225

Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Wilson SK (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. Science 359:80–83

Jones AM, Berkelmans R, van Oppen MJH, Mieog JC, Sinclair W (2008) A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. Proc. Biol. Sci. 275:1359–1365

Kahng SE, Benayahu Y, Lasker HR (2011) Sexual reproduction in octocorals. Mar. Ecol. Prog. Ser. 443:265–283

Karim W, Nakaema S, Hidaka M (2015) Temperature effects on the growth rates and photosynthetic activities of *Symbiodinium* cells. J. Mar. Sci. Eng. 3:368–381

Keshavmurthy S, Hsu C-M, Kuo C-Y, Meng P-J, Wang J-T, Chen CA (2012) Symbiont communities and host genetic structure of the brain coral *Platygyra verweyi*, at the outlet of a nuclear power plant and adjacent areas. Mol. Ecol. 21:4393–4407

Kirk NL, Andras JP, Harvell CD, Santos SR, Coffroth MA (2009) Population structure of *Symbiodinium* sp. associated with the common sea fan, *Gorgonia ventalina*, in the Florida Keys across distance, depth, and time. Mar. Biol. 156:1609–1623

LaJeunesse TC, Parkinson JE, Reimer JD (2012) A genetics-based description of *Symbiodinium minutum* sp. nov. and *S. psygmophilum* sp. nov. (Dinophyceae), two dinoflagellates symbiotic with cnidaria. J. Phycol. 48:1380–1391

LaJeunesse TC, Smith RT, Finney J, Oxenford H (2009) Outbreak and persistence of opportunistic symbiotic dinoflagellates during the 2005 Caribbean mass coral "bleaching" event. Proc. Biol. Sci. 276:4139–4148

LaJeunesse TC, Wham DC, Pettay DT, Parkinson JE, Keshavmurthy S, Chen CA (2014) Ecologically differentiated stress-tolerant endosymbionts in the dinoflagellate genus*Symbiodinium* (Dinophyceae) Clade D are different species. Phycologia 53:305–319

LaJeunesse TC (2005) "Species" radiations of symbiotic dinoflagellates in the Atlantic and Indo-Pacific since the Miocene-Pliocene transition. Mol. Biol. Evol. 22:570–581

LaJeunesse T (2002) Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. Mar. Biol. 141:387–400

LaJeunesse T (2020) Zooxanthellae. Curr Biol 30:R1110–R1113

Lasker HR (2003) Zooxanthella densities within a Caribbean octocoral during bleaching and non-bleaching years. Coral Reefs 22:23–26

Lasker HR, Martínez-Quintana Á, Bramanti L, Edmunds PJ (2020) Resilience of octocoral forests to catastrophic storms. Sci. Rep. 10:4286

Lasker HR, Peters EC, Coffroth MA (1984) Bleaching of reef coelenterates in the San Blas Islands, Panama. Coral Reefs 3:183–190

Lenz EA, Bramanti L, Lasker HR, Edmunds PJ (2015) Long-term variation of octocoral populations in St. John, US Virgin Islands. Coral Reefs 34:1099–1109

Lewis CL, Coffroth MA (2004) The acquisition of exogenous algal symbionts by an octocoral after bleaching. Science 304:1490–1492

Mieog JC, van Oppen MJH, Cantin NE, Stam WT, Olsen JL (2007) Real-time PCR reveals a high incidence of *Symbiodinium* clade D at low levels in four scleractinian corals across the Great Barrier Reef: implications for symbiont shuffling. Coral Reefs 26:449–457

Muscatine L, Cernichiari E (1969) Assimilation of photosynthetic products of zooxanthellae by a reef coral. Biol. Bull. 137:506–523

Muscatine L, Porter JW (1977) Reef corals: mutualistic symbioses adapted to nutrient-poor environments. Bioscience 27:454–460

Oliver JK, Berkelmans R, Eakin CM (2018) Coral bleaching in space and time. Coral Bleaching.

Springer, pp 27–49

Parkinson JE, Baums IB (2014) The extended phenotypes of marine symbioses: ecological and
evolutionary consequences of intraspecific genetic diversity in coral-algal associations.
Front. Microbiol. 5:445

Parkinson JE, Coffroth MA, LaJeunesse TC (2015) New species of Clade B *Symbiodinium*
(Dinophyceae) from the greater Caribbean belong to different functional guilds:
S. aenigmaticum sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and
S. pseudominutum sp. nov. J. Phycol. 51:850–858

Pettay DT, LaJeunesse TC (2007) Microsatellites from clade B *Symbiodinium* spp. specialized
for Caribbean corals in the genus *Madracis*. Mol. Ecol. Notes 7:1271–1274

Petzoldt T (2019) growthrates. URL: https://github.com/tpetzoldt/growthrates.

Poland DM, Coffroth MA (2019) Host growth and survivorship varies with endosymbiotic algal
partner in developing cnidarians. Mar. Ecol. Prog. Ser. 612:87–100

Prada C, Weil E, Yoshioka PM (2010) Octocoral bleaching during unusual thermal stress. Coral
Reefs 29:41–45

Quigley KM, Baker AC, Coffroth MA, Willis BL, van Oppen MJH (2018) Bleaching resistance
and the role of algal endosymbionts. Ch. 6 pp. 111-151. In: Coral Bleaching: Patterns,
processes, causes and consequences eds; J. Lough and M. van Oppen. Springer Ecological
Series. Cham, Switzerland

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.

Rädecker N, Pogoreutz C, Gegner HM, Cárdenas A, Roth F, Bougoure J, Guagliardo P, Wild C,
Pernice M, Raina J-B, Meibom A, Voolstra CR (2021) Heat stress destabilizes symbiotic
nutrient cycling in corals. Proc Natl Acad Sci USA 118(5):e2022653118

Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS,
Macaulay KA, Bartlett LA, Meyers MK, Colee J (2013) Temporal changes in benthic
assemblages on Florida Keys reefs 11 years after the 1997/1998 El Niño. Mar. Ecol. Prog.
Ser. 489:125–141

Santos SR, Gutiérrez-Rodríguez C, Lasker HR, Coffroth MA (2003) *Symbiodinium* sp.
associations in the gorgonian *Pseudopterogorgia elisabethae* in the Bahamas: high levels of
genetic variability and population structure in symbiotic dinoflagellates. Mar. Biol.
143:111–120

Santos SR, Shearer TL, Hannes AR, Coffroth MA (2004) Fine-scale diversity and specificity in
the most prevalent lineage of symbiotic dinoflagellates (*Symbiodinium*, Dinophyceae) of the
Caribbean. Mol. Ecol. 13:459–469

Santos SR, Taylor DJ, Coffroth MA (2001) Genetic comparisons of freshly isolated versus

cultured symbiotic dinoflagellates: implications for extrapolating to the intact symbiosis. J. Phycol. 37:900–912

Silverstein RN, Correa AMS, Baker AC (2012) Specificity is rarely absolute in coral-algal symbiosis: implications for coral response to climate change. Proc. Biol. Sci. 279:2609–2618

Silverstein RN, Cunning R, Baker AC (2015) Change in algal symbiont communities after bleaching, not prior heat exposure, increases heat tolerance of reef corals. Glob. Chang. Biol. 21:236–249

Thornhill DJ, Howells EJ, Wham DC, Steury TD, Santos SR (2017) Population genetics of reef coral endosymbionts (*Symbiodinium*, Dinophyceae). Mol. Ecol. 26:2640–2659

Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N, Foret S, Matz M, Miller DJ, Moya A, Putnam HM, Ravasi T, van Oppen MJH, Thurber RV, Vidal-Dupiol J, Voolstra CR, Watson S-A, Whitelaw E, Munday PL (2017) Rapid adaptive responses to climate change in corals. Nat. Clim. Chang. 7:627–636

Tukey JW (1949) Comparing individual means in the analysis of variance. Biometrics 5:99–114

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadia G, Raymundo L, Williams G, Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of the Paris Agreement. Sci. Rep. 6:39666

van Oppen MJH, Mieog JC, Sánchez CA, Fabricius KE (2005) Diversity of algal endosymbionts (zooxanthellae) in octocorals: the roles of geography and host relationships. Mol. Ecol. 14:2403–2417

van Oppen MJH, Souter P, Howells EJ, Heyward A, Berkelmans R (2011) Novel genetic diversity through somatic mutations: fuel for adaptation of reef corals? Diversity (Basel) 3:405–423

Wilkinson SP, Fisher PL, van Oppen MJH, Davy SK (2015) Intra-genomic variation in symbiotic dinoflagellates: recent divergence or recombination between lineages? BMC Evol. Biol. 15:325

Williams DE, Miller MW (2015) Water temperature data from reef sites off the upper Florida Keys from 2003-09-18 to 2019-11-25 (NCEI Accession 0126994). [Water temperature data from reef sites off the upper Florida Keys from 2003-09-18 to 2019-11-25 (NCEI Accession 0126994)]. NOAA National Centers for Environmental Information. Dataset. https://www.ncei.noaa.gov/archive/accession/0126994. Accessed [7 Jan 2021.]

Figure Captions

Figure 1. Experimental design for a) *in vitro* growth experiment and b) *in hospite* experiment. Figure colors refer to historical temperature treatment (blue, maintained at 26°C; red, maintained at 30°C), with tip figures colored based on experimental treatment (not historical temperature).

Figure 2. a) Mean logistic growth fits to *in vitro* cell densities of each genotype of *Breviolum antillogorgium* over time. Blue represents growth at 26°C and red at 30°C. b) Estimated mean (+/- SE) maximum symbiont population growth rates (r) and c) estimated mean (+/- SE) carrying capacities (K). Colors for genotype names (x-axis) refer to the historical temperature at which they were grown, blue indicating cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the start of the experiment. Letters above bars represent differences among groups based on Tukey post-hoc tests.

Figure 3. Mean percent of visually infected polyps at the end of a) Day 31 and b) Day 52. c) Symbiont cell densities within *Antillogorgia bipinnata* polyps inoculated with five symbiont genotypes at 26ºC and 30ºC at the end of the experiment (Day 69). Colors for genotype names (x-axis) refers to historical temperature, blue indicating cultures maintained at 26ºC and red indicating cultures maintained at 30ºC two years prior to the start of the experiment. Color of boxes refers to experimental temperature (blue- 26ºC, red- 30ºC). Boxes represent interquartile range and whiskers the range of the data. Points indicate outliers. Letters above boxes represent differences among groups based on Tukey post-hoc tests.

Figure 4. a) Mean polyp survivorship over time for *A. bipinnata* polyps inoculated with five

symbiont genotypes at 26ºC and 30ºC. Error bars are ±SE. Blue lines refer to treatments at 26 ºC

and red lines refer to treatments at 30ºC. Differences in average survivorship at b) Day 31 and c)

Day 52. Colors for genotype names (x-axis) refer to historical temperature, blue indicating

cultures maintained at 26ºC and red indicating cultures maintained at 30ºC two years prior to the

start of the experiment. Color of boxes refers to experimental temperature (blue- 26ºC, red-

30ºC). Boxes represent interquartile range and whiskers the range of the data. Points indicate

outliers. Letters above boxes represent differences among groups based on Tukey post-hoc tests.

Online Resource 1. Average water temperature at Elbow Reef A at 3 m depth from 2005 to

2015. Data from Williams and Miller (2015). Horizontal lines are at 30ºC (red) and 26ºC (blue)

for reference.

Online Resource 2. Experimental set up for the *in hospite* study. Letter color refers to historical

temperature, blue indicating cultures reared at 26ºC and red indicating cultures reared at 30ºC

two years prior to the start of the experiment.

Online Resource 3. Mean logistic growth fits to *in vitro* cell densities of each genotype of

Breviolum antillogorgium over time for each replicate for each genotype and temperature

treatment.

a

Isolate
symbionts from
host

30°C   26°C

Grow symbiont
populations in
vitro for 2 years

30°C   26°C   30°C   26°C

in vitro experiment

b

Isolate
symbionts from
host

30°C   26°C

Grow symbiont
populations in
vitro for 2 years

30°C   26°C   30°C   26°C

in hospite experiment