Parabacteroides distasonis **enhances Type 1 Diabetes autoimmunity via molecular mimicry**

Qian Huang[1], I-Ting Chow[2], Claudia Brady[1], Amol Raisingani[1], Danmeng Li[3], David A. Ostrov[3], Mark A. Atkinson[3,4], William W. Kwok[2], C. Ronald Kahn[5], Emrah Altindis[1]*

[1] Boston College Biology Department, Higgins Hall, 140 Commonwealth Avenue Chestnut Hill, MA 02467, USA

[2] Benaroya Research Institute at Virginia Mason, Seattle, WA 98101, USA

[3] Department of Pathology, Immunology and Laboratory Medicine, College of Medicine, University of Florida Diabetes Institute, Gainesville, FL 32610-3633, USA

[4] Department of Pediatrics, College of Medicine, University of Florida Diabetes Institute, Gainesville, FL 32610-3633, USA

[5] Joslin Diabetes Center, Harvard Medical School, Boston, MA, 02215, USA

*Corresponding Author: Emrah Altindis, Boston College Biology Department, Higgins Hall, 140 Commonwealth Avenue Chestnut Hill, MA 02467. E-mail: altindis@bc.edu

One Sentence Summary: The human gut commensal bacterium, *Parabacteroides distasonis*, accelerates type 1 diabetes in the NOD mouse model of the disease and involves expression of an insulin B:9-23 epitope mimic, supporting a potential disease mechanism involving molecular mimicry.

ABSTRACT

Type 1 Diabetes (T1D) is an autoimmune disease characterized by destruction of pancreatic β-cells. Focusing on the main insulin epitope, insulin B-chain 9-23 (insB:9-23), we explored whether a microbial insB:9-23 mimic could modulate T1D. We now demonstrate that a microbial insB:9-23 mimic of *Parabacteroides distasonis*, a human gut commensal, exclusively stimulates non-obese diabetic (NOD) mouse T cells specific to insB:9-23. Indeed, immunization of NOD mice with either the bacterial mimic peptide or insB:9-23 further verified the cross-reactivity in vivo. Modeling *P. distasonis* peptide revealed a potential pathogenic register 3 binding. *P. distasonis* colonization of the female NOD mice gut accelerated T1D onset. In addition, adoptive transfer of splenocytes from NOD mice colonized with *P. distasonis* to NOD.SCID recipients conferred the enhanced disease phenotype. Integration analysis of published infant T1D gut microbiome data revealed that *P. distasonis* peptide is not present in the gut microbiota in the first year of life of infants that eventually develop T1D. Furthermore, *P. distasonis* peptide can stimulate human T cell clones specific to insB:9-23 and T1D patients demonstrated a strong humoral immune response to *P. distasonis* than controls. Taken together, our studies define a potential molecular mimicry link between T1D pathogenesis and the gut microbiota.

INTRODUCTION

Type 1 diabetes (T1D) is an autoimmune disease characterized by selective destruction of pancreatic β-cells by autoreactive T cells (1). The incidence rate of T1D in children is rising on an annual basis both in Europe (3.4 %) and the USA (1.4 %) (2, 3). Genome-wide association studies (GWAS) have identified ~50 genetic regions that influence the risk of developing T1D, but genetics alone cannot account for the increasing incidence rates of T1D. For example, incidence rates of T1D in two neighboring regions, Russian Karelia and Finnish Karelia are six-fold different; despite a very similar genetic background of the inhabitants (4). Furthermore, even among identical twins, the concordance of T1D is only 40-60% (5, 6). Various environmental factors have been studied including diet, birth mode, infections and antibiotics; however, the roles of these factors and cause of T1D largely remain unknown (7-9). This said, recent gut microbiome studies have observed an altered microbial and metabolite composition, and increased intestinal permeability in subjects with T1D (10-14). In addition to gut microbes, viral infections have also been suggested to play a role in T1D, yet no studies have identified a direct causal link (15). Taken together, these data suggest that genetics alone cannot explain the risk of T1D autoimmunity.

One of the earliest markers of T1D autoimmunity is the development of islet autoantibodies against insulin (IAA), glutamic acid decarboxylase (GADA), insulinoma-associated autoantigen 2 (IA-2A), zinc transporter 8 (ZnT8A) and islet specific glucose-6-phosphatase catalytic subunit related protein (IGRP, recently named G6PC2) (16, 17). Among these, insulin is the only autoantigen exclusively expressed by β-cells, with IAA usually the first to be detected (18, 19). In humans, IAA develop months to years before the onset of overt diabetes (20), and, there is a significant correlation between IAA concentration and rate of progression to T1D (21). In addition to humoral immune responses, insulin is a target of CD4[+] T cells in both humans and murine

autoimmune diabetes. In theNOD mouse model of T1D (22), over 90% of the anti-insulin CD4$^+$ T

cell clones target amino acids 9-23 of the insulin B chain (insB:9-23) (23). Importantly, InsB:9-23

specific T-cells have also been identified in islets obtained from human organ donors with T1D

(24, 25) as well as in peripheral blood lymphocytes of living T1D patients (19, 26, 27). Clearly

lacking in this field are studies that have the ability to pull together such findings and, bring

together a central hypothesis for testing of a strong, well supported, central hypothesis.

In this regard, the concept of molecular mimicry has long been noted as a potential trigger

for autoimmunity; one when the immune response cannot discriminate a foreign antigen from a

self-protein (8, 28). In this study, we hypothesized that exposure to a microbial insulin epitope

mimic could stimulate an autoimmune response initiating T1D onset. To address this question, we

screened for microbial mimics of insB:9-23 and subsequently evaluated one such mimic from a

gut commensal, *Parabacteroides distasonis*; this, for the notion of stimulating insB:9-23-specific

human and NOD T cells ex vivo as well as to modulate T1D progression in vivo. We also leveraged

existing human gut microbiota datasets from the BABYDIET and DIABIMMUNE studies to

explore microbial colonization and expression of insB:9-23 mimics during the first years of life in

children who later developed T1D. Taken together, our studies demonstrated that insulin epitope

mimics in the gut microbiota may play an important role in T1D onset ,via a molecular mimicry

mechanism, particularly when exposed after the neonatal period of immune development.

RESULTS

***Parabacteroides distasonis* insB:9-23 like peptide cross-reacts with NOD mouse T cell hybridomas specific to insB:9-23**

To test our aforementioned molecular mimicry hypothesis, we focused on the dominant T-cell epitope, insB:9-23. We first identified microbial insB:9-23 peptide mimics, searching the NCBI databases for viral (taxid:10239), bacterial (taxid:2) and fungal (taxid:4751) proteomes using the insB:9-23 sequence (SHLVEALYLVCGERG) as the query sequence. We identified 47 microbial peptides with over eight residues identical with the insulin peptide (**Table S1**). Among these microbial mimics, 17 bacterial and viral mimics contained the previously identified residues in the insB:9-23 critical for binding to the NOD MHC I-A^{g7} and human HLA-DQ2-8 molecule (**Table 1**). To test whether microbial insB:9-23 mimics can stimulate NOD T cells, we used IIT3 and 8F10 T cell hybridomas. IIT3 recognizes presented insulin protein as well as the insB:9-23 peptide, whereas 8F10 recognizes only the insB:9-23 peptide (29). We also used a C3g7 cell line expressing a large amount of I-A^{g7}, as antigen presenting cells (APCs). Among the 17 microbial peptides tested, only one stimulated IIT-3 hybridomas (**Figure 1A**). This peptide contained in the N-terminal of the hypoxanthine phosphoribosyltransferase (hpt) protein of *P. distasonis 33B* formerly known as *Bacteroides sp. 2_1_33B (30)* (**Figure S1A**). Yang et al. recently demonstrated that residues from insB:15-20 are essential to stimulate the human T cell clones via register 3 (27). Indeed, the *P. distasonis* peptide is the only tested microbial mimic that has all six residues conserved in this region (**Table 1, S1**). Notably, *P. distasonis* 33B and D13 strains are the only organisms that possess this unique insB:9-23 mimic sequence in their genomes in the NCBI dataset (as of September 12, 2020, **Figure S1B-C, Supplementary Document 1-2**). Taken together, as a

member of the human gut microbiome, *P. distasonis* makes an interesting candidate for T1D

autoimmunity.

With these results, we repeated the T cell hybridoma stimulation experiment using *P.*

distasonis lysate or growth culture instead of the 15-mer peptide, but did not observe a robust

response (**Figure S3A-B**). However, it is well known that some bacterial proteins are not expressed

in vitro and require specific conditions mimicking in vivo (31). Indeed, using RT-qPCR, we

observed extremely low gene expression of Hpt in our *P. distasonis* culture (**Figure S3C**).

Moreover, we performed an SDS-PAGE gel of the bacterial lysate and analyzed all bands between

20 and 50 kDa. *P. distasonis* Hpt protein, which is predicted to be 27.3 kDa, was not identified

among the 464 proteins detected by liquid chromatography-mass spectrometry (LC-MS) MS

analysis (**Table S2**).

Structural modeling of *P. distasonis* insB:9-23-like peptide suggests potential binding to

MHC in a specific register.

Previous studies have demonstrated the importance of specific insB:9-23 peptide positions in

anchoring the peptide to the MHC binding cleft in a manner that determines the binding register

(32). To understand the molecular mechanism underlying such cross-reactivity, we developed an

in silico model of the *P. distasonis* peptide compared to the native insB:9-23 epitope (**Figure 1B,**

S2). Glutamic acid at position 9 (P9) of insB:9-23 confers binding in "register 2", whereas non-

glutamic acid side chains, such as arginine, impose peptide binding in "register 3"(32). Variants

of insB:9-23 presented in register 3 elicit stronger autoreactive T cell responses compared to

insB:9-23 presented in register 2. In the context of register 3, the *P. distasonis* mimic and insB:9-

23 peptide share a high degree of sequence homology except at P9. The *P. distasonis* mimic

peptide has a tyrosine (Y) at P9 versus serine (S) in the native insulin epitope, whereas most of the

other microbial mimics have an arginine (R) at P9, which is not expected to shift the register. The

binding model of insB:9-23 and *P. distasonis* peptide revealed that the *P. distasonis* mimic peptide

has the potential to bind MHC in register 3, thus resulting in presentation of peptide side chains

that are identical between insB:9-23 and *P. distasonis* (**Figure 1B, S2A-B**). These data support

our observation that T cells responding to the register 3 bound insulin peptide have the ability to

cross react with the *P. distasonis* peptide in register 3 (**Figure 1A**).

Immunization of NOD mice with *P. distasonis* peptide stimulates cross-reactive T-cells

To further explore this peptide cross-reactivity, we immunized NOD mice with either (i) insB:9-

23 peptide or, (ii) *P. distasonis* mimic peptide with Complete Freund's Adjuvant (CFA). After 7

days, we collected the popliteal lymph nodes and examined T cell activity by enzyme-linked

immunospot (ELISpot) for interferon-gamma (IFNg) and IL-2 secretion; this, following *in vitro*

stimulation with either the insB9-23 or microbial mimic peptide. Immunization of mice with the

P. distasonis peptide generated a cross-reactive immune response to insB9-23 (**Figure 1C and**

1E). Likewise, immunization of mice with insB:9-23 stimulated a cross-reactive response to the

P. distasonis peptide, which was even stronger than the stimulation with the insulin peptide

(**Figure 1D and 1F**).

P. distasonis **colonization accelerates diabetes onset in female NOD mice**

Based on our findings on cross-reactivity, we hypothesized that *P. distasonis* colonization and

exposure to the microbial insB:9-23 mimic can trigger the immune response and stimulate

autoimmunity in NOD mice. To test this hypothesis, we orally gavaged both male and female

NOD mice with *P. distasonis* (10^9 CFU *P. distasonis*/mouse/day) or saline daily after weaning,

starting at 3 weeks of age (**Figure 2A**, n=40/group). Neither male nor female NOD mice carried

P. distasonis in their gut under specific pathogen free (SPF) housing conditions (**Figure 2B**),

providing an ideal control to determine the effect of the *P. distasonis* colonization on T1D onset.

After 30 days of oral gavage, we detected colonization of the gut with *P. distasonis* (**Figure 2B**).

At 12 weeks of age (pre-diabetic), we randomly separated five mice from each group and collected

the pancreas for immunostaining. Interestingly, the *P. distasonis* colonized female NOD mice

showed significantly higher insulitis scores as compared to controls (**Fig 2C, Fig S4A**

representative figures). Moreover, *P. distasonis* colonization significantly accelerated T1D onset

in female NOD mice (survival: 42% versus 19.5%) (**Figure 2D**). This effect was specific to the

female mice as no differences in insulitis or T1D incidence were observed in male mice

administered *P. distasonis* versus saline (**Figure 2C-D**).

***P. distasonis* stimulates a specific humoral immune response in colonized NOD mice.**

To determine whether *P. distasonis* colonization can stimulate IAA seroconversion, we measured

IAA levels with ELISA at 12 weeks of age in study animals. Only a small fraction of the mice had

circulating IAA at this early point, with no significant differences between study groups detected

(**Fig. S4B**). Western blot analysis using plasma samples obtained from the same NOD mice (12

weeks) revealed that *P. distasonis* colonization stimulated a very strong humoral immune response

to the bacterium in both female and male NOD mice (**Figure 2E-F**). To evaluate the specificity of

this humoral response, we performed a similar Western blot using the plasma samples to detect

lysates from another well studied gut commensal, *Bacteroides fragilis,* that was shown to have

immunomodulatory effects in NOD mice *(33).* As expected, we detected a weak humoral immune

response to *B. fragilis* lysates in both the saline and *P. distasonis* treated groups (**Figure S4C-D**).

Serum LPS levels were not significantly different between the groups indicating that this *P.*

distasonis specific humoral response did not result from gut barrier dysfunction (leaky gut)

stimulated by *P. distasonis* colonization (**Figure 2G**). We measured multiple (n=32) serum

cytokines and chemokines using a multiplex Luminex assay (Millipore) to determine whether *P.*

distasonis colonization of the female NOD mice alters the systemic immune response but did not

detect any differences between the two groups (**Table S3**). These results suggest that *P. distasonis*

stimulates a specific immune response but not IAA or elevated proinflammatory cytokines.

Adoptive transfer of splenocytes enhance T1D onset in immunodeficient NOD.SCID mice

We next performed an adoptive transfer experiment (**Figure 3A**) using immunodeficient

NOD.SCID mice, which lack functional B or T cells, as recipients (34). We orally gavaged a new

cohort of the NOD mice with *P. distasonis* or saline (n=16-20/group) as described above and

transferred $5x10^7$ splenocytes from individual NOD donor mice to 6 week old NOD.SCID

recipients (1:1 ratio, sex-matched, n=16-20 per group) (**Figure 3B**). Consistent with the

spontaneous T1D experiment (**Figure 2D**), recipient female NOD.SCID mice that received

splenocytes from the *P. distasonis* treated group developed T1D at a significantly higher rate

(**Figure 3C**). Thus, an adaptive immune response stimulated by *P. distasonis* exposure in female

NOD mice is sufficient to transfer and accelerate T1D.

Subcutaneous injection of *P. distasonis* peptide is protective in female NOD mice

Subcutaneous injection of mouse insulin 2, B:9-23 into female NOD mice for 25 weeks was shown

to be protective in female NOD mice, with this result explained by a T helper 2 (Th2) cell shift in

immune response (35).   To determine whether a different route of *P. distasonis* peptide

administration is protective, we injected female (n=20/peptide) and male (n=20/peptide) NOD

mice with i) insB:9-23 insulin 2, ii) the *P. distasonis* B:9-23 mimic, iii) saline (vector), or iv)

tetanus toxoid (TT) control peptide weekly for 20 weeks beginning at 4 weeks of age. Consistent

with the oral gavage experiments (**Figures 2D and 3C**), we did not observe any effect of *P.*

distasonis B:9-23 peptide on male NOD mice (**Figure 3D**) while insB:9-23 and the *P. distasonis*

peptide significantly delayed T1D onset in female NOD mice (**Figure 3E**). This further confirms

the cross reactivity between insB:9-23 and the *P. distasonis* peptide mimic, but the route of

exposure (i.e., subcutaneous versus mucosal) appears to impact the ability to modulate T1D onset

(**Figure 2D and 3E**).

P. distasonis **B:9-23 like peptide is not present in the gut microbiota of children developing**

T1D in the first year of life.

To investigate a potential role for the *P. distasonis* insB:9-23 mimic in human T1D, we analyzed

published human gut microbiome data emanating from the BABYDIET study, where 16S Illumina

sequencing was used to analyze stool samples collected from 22 children who developed islet

autoantibodies, with 10 developing diabetes after a median follow-up time of 5 years, as well as

from 22 healthy subjects from birth to 3 years of age (36). We searched for the operational

taxonomic units (OTUs) identified as *P. distasonis* and classified the data according to sample

collection time (0-1, 1-2 and 2-3 years of age) and subject disease status at the end of the study

(control, autoantibody (Ab) positive, or T1D). *P. distasonis* was present in control children at all

three time points, but was only identified in the second year of age and later in the children who

developed T1D. Moreover, compared to controls, fewer *P. distasonis* OTUs were identified for

the first year in the 12 children who developed autoantibodies but did not progress to T1D during

follow-up (**Figure 4A**). A key limitation of this 16S rRNA gene sequencing data is the inability to

determine whether the identified *P. distasonis* OTUs carry the insulin B9-23 mimic peptide.

To address this, we next analyzed published DIABIMMUNE metagenomic data obtained

from three countries: Russia, Finland and Estonia (11). In this study, researchers performed

shotgun metagenomics sequencing on stool samples collected annually from genetically

predisposed subjects from birth to age 3 years in Finland, Russia and Estonia (n=74/country). To

identify optimized alignments, we first assembled the contigs and searched for DNA sequences

capable of encoding the *P. distasonis* peptide. Consistent with the BABYDIET results (**Figure**

4A), we were unable to identify the *P. distasonis* epitope mimic in the first year of life for subjects

who later developed autoantibodies (seroconverted, Ab positive **Figure 4B**). We next analyzed the

data stratified by country. While no subjects were exposed to the *P. distasonis* peptide in the first

year of age, the peptide was identified in all subjects who develop autoantibodies in Estonia after

age one and in Russia after age two. In contrast, the children who developed autoantibodies in

Finland were never exposed to this peptide in the first three years of age; suggesting a potential

later time point of exposure or a different trigger (**Fig. 4C**). In sum, these results suggest that the

late exposure to the *P. distasonis* insB:9-23 mimic in years 2-3 of childhood might trigger an

immune response and stimulate autoimmunity. Indeed, these observations are consistent with our

findings in NOD mice. Colonization of the gut microbiota with to *P. distasonis* after weaning, at

a late time point relative to immune development, accelarated the disease in female NOD mice.

P. distasonis **peptide can stimulate insB:9-23 specific human T-cell clones**

To further characterize the immune response against *P. distasonis*, we investigated the humoral response in human T1D patients to determine their exposure. Specifically, we tested antibody responses using serum obtained from female subjects with T1D and age and ethnicity matched controls (n=8/group, median age 16.2, median duration 4.3 years). Indeed, four of eight female subjects with T1D demonstrated an elevated immune response to *P. distasonis* lysates compared to the eight controls and other T1D four samples (**Figure 4D Table S4**). Because T1D is T-cell mediated, we also tested whether the *P. distasonis* insB:9-23 mimic or 17 other insB:9-23 like peptides (**Table 1**) could stimulate insB:11-23-specific T cell clones isolated from participant T1D#3 in (27). The insB:9-23R^{22E} peptide was previously shown to be more potent than insB:9-23 in stimulating insB:9-23 hybridomas from NOD mice (32) and human T-cell clones (27); hence we used both insB:9-23R^{22E} and insB:9-23 as positive controls and a scrambled peptide as a negative control (**Table 1**). Consistent with our findings with NOD mice hybridomas (**Figure 1**), the *P. distasonis* peptide (peptide 1) was the only microbial peptide among 17 tested that could stimulate human T cells to produce IFN-g (**Figure 4E, F**).

DISCUSSION

There have been tremendous efforts to understand the triggers of T1D autoimmunity, however, the etiology of the disease is complex and remains incompletely characterized. In this study, we investigated molecular mimicry as a potential mechanism linking microbial flora to T1D onset. Molecular mimicry mechanisms are based on the degeneracy of T cell recognition (37, 38) and can be either pathogenic (39) or protective (40). While the molecular mimicry hypothesis was popular in T1D research in the 1990s (41-43), progress in this area is very limited in the last 20 years (44). In the current study, taking advantage of these growing genome databases, we identified candidate microbial mimic peptides for the T1D target epitope, insB:9-23. We further show that stimulation of the immune system with a gut commensal's, *P. distasonis,* insB:9-23 like peptide can stimulate an immune response to the endogenous insB:9-23. Moreover, exposure to *P. distasonis* during gut microbiota development caused increased infiltration to the islets and accelerated diabetes onset in female NOD mice. This pathogenic effect was mediated by splenocytes as evidenced by adoptive transfer to female NOD.SCID mice.

A plethora of human gut microbiome studies demonstrated that the composition of the gut microbiota of patients with autoimmune diseases including multiple sclerosis (45), systemic lupus erythematosus, anti-phospholipid syndrome (46), Crohn's disease (47-49), ulcerative colitis (50), inflammatory bowel diseases (51, 52), and coeliac disease (53) are significantly different from healthy controls. Although the methodologies and conclusions differ in microbiota studies from subjects with and at-risk for T1D (11-13, 54), there is a common pattern. Specifically, the diversity of T1D patients' gut microbiome is decreased while there is an increased prevalence of *Bacteriodetes* taxa and an altered metabolomic profile compared to healthy controls (14). Most of these studies are observational and do not define any mechanism related to T1D onset. One

exception is the DIABIMMUNE study (13) wherein authors concluded that the higher T1D rates

in Finnish Karelia and Estonia compared to Russian Karelia might be related to the gut microbiota

composition. They suggest that while Russian infants are exposed to *Escherichia coli* LPS, the

Finnish and Estonian infants are exposed to *Bacteroides* species' LPS, which inhibits endotoxin

tolerance. They also showed that injection of *E. coli* LPS protected NOD mice from developing

T1D; however, they did not report any data validating the LPS hypothesis for human T1D.

As the largest human gut microbiome study collecting 12,005 samples from 903 children

in four countries in seven locations, The Environmental Determinants of Diabetes in the Young

(TEDDY) trial identified *Parabacteroides* as the most significantly associated genus with T1D

(55). This is consistent with our findings related to *P. distasonis* mimic analysis from Russia and

Estonia wherein 100% of the children developing T1D had the *P. distasonis* peptide in their gut

microbiota in years 2-3 of life. The TEDDY study also recently showed that human gut microbiota

development is divided in three phases: a developmental phase (months 3-14), a transitional phase

(months 15-30), and a stable phase (months 31-46) (55). Using the DIABIMMUNE sequencing

data, we showed that the *P. distasonis* insB:9-23 mimic peptide is not present in the gut microbiota

during the developmental phase (i.e., the first year of life) in the children who go on to develop

T1D. Our data suggest that the absence of *P. distasonis* in the developmental phase may be critical

in T1D pathogenesis. A recent study showed that in addition to thymic education of the immune

system for self-tolerance, the microbiota provides a post-thymic education to provide tolerance to

commensal microbiota (56). Thus, the presence of the *P. distasonis* insB:9-23 mimic in this early

stage might promote tolerance to the insulin epitope. Hence, there might be a window of

opportunity for exposure during the developmental phase of gut microbiota development during

which the *P. distasonis* peptide might stimulate regulatory T cells (Tregs) and immunotolerance

for insB:9-23. Although further studies are needed to investigate the direct link between human

T1D and *P. distasonis*, the cross-reactivity of the insB:9-23 mimic with human T cell clones

represents a clear potential mechanism. Furthermore, the strong humoral immune response to *P.*

distasonis lysate clearly shows the exposure and humoral immune response in a fraction of T1D

patients.

In addition to their critical role in NOD mouse diabetes onset (57), insB:9-23 specific CD4[+]

T cells were identified in human islets (24, 25) and in peripheral blood of T1D patients (19, 26,

27). While our study is insB:9-23 centric, there are other epitopes recognized by human islet-

infiltrating or circulating T-cells in insulin B-chain (58), A-chain (59, 60) or C-peptides (24, 25).

A recent discovery showed that hybrid insulin peptides (HIPs) are novel epitopes formed with

post-translational modifications formed in the beta cell granules (61-63). Likewise, defective

ribosomal products (Drips) of insulin were shown to lead of aberrant insulin polypeptides

rendering beta cells immunogenic (64). Babon et al. recently showed that CD4[+] and CD8[+] T cell

lines or clones isolated from the islets of nine donors with T1D are stimulated by several islet

antigens including GAD, insulin, IA-2 and HIPs (65). Given the enormous number of microbial

peptides produced by the gut microbiota, we expect that others may exist with the potential to

mimic these additional epitopes and trigger cognate autoantigen reactive T cells. Indeed, Tai et al.

identified a microbial peptide mimic produced by *Fusobacteria* that can stimulate IGRP-specific

mouse T cells and promote diabetes development in a new TLR-deficient (TLR$^{-/-}$) and

MyD88$^{-/-}$ NY8.3 NOD mouse model (66).

Although gut microbiota has a critical role in modulating T1D onset in NOD mice (67),

there are very few studies focusing on the effects of a single bacterium. For example, the presence

of segmented filamentous bacteria correlated with protection in NOD mice (68). Likewise,

Hanninen et al. showed that oral gavage of female NOD mice with *Akkermensia muciniphila* from

weaning until 10 weeks of age delayed diabetes (69). Using a chemically induced enhanced gut

permeability condition, Sofi et al. showed that oral administration of heat-killed (HK) *B. fragilis*

suppressed autoimmunity in NOD mice *(33).* On the other hand, intravenous injection of NOD

mice with HK *B. fragilis* accelerated disease progression. Indeed, the importance of immunization

route is well appreciated in vaccinology (70) and highlighted by our experiments where oral

gavage of female NOD mice with *P. distasonis* stimulated a pathogenic response via the mucosal

immune system whereas subcutaneous immunization with the *P. distasonis* insB:9-23 mimic

stimulated protection. This latter finding corroborates a previous report demonstrating that

immunization with the native insB:9-23 peptide reduces T1D incidence in female NOD mice (35).

To our knowledge, *P. distasonis* is the first bacterium that can accelerate the disease onset

in the NOD mice. We showed that *P. distasonis* is not naturally present in the NOD mice gut

microbiome in our SPF facility. Although this provides a good control to determine the effects of

P. distasonis on T1D onset, it has limitations. Because *P. distasonis* is not previously used for

genetic mutations and do not have established protocols, we failed to make a mutant of *P.*

distasonis that has a deletion or mutations of the mimic peptide. Further *in vivo* studies are needed

to establish the direct link between the mimic peptide and the effects of the colonization.

Today, we have databases with enormous amounts of microbial and microbiome

sequencing data that can be levereaged to address the role of molecular mimicry in autoimmunity.

Indeed, we see the return of the molecular mimicry hypothesis in the autoimmunity field with

recent studies on T1D (66), lupus (71), inflammatory cardiomyopathy (72), and multiple sclerosis

(73). These studies, including ours, constitute a new link to gut antigens and autoimmune diseases

with the potential to ultimately provide new tools including vaccines, antibiotics or probiotics for

prevention and treatment of autoimmune disease. The uniqueness of the *P. distasonis* insB:9-23

mimic peptide in NCBI databases supports our ongoing efforts to characterize its effect based on

Koch's postulates.

In conclusion, our data show that the *P. distasonis* insB:9-23 like peptide stimulates human

and NOD mice T cells specific to native insB:9-23. Intestinal colonization of female NOD mice

with *P. distasonis* accelerates autoimmune diabetes. This phenotype can be adoptively transferred

to NOD.SCID mice demonstrating the direct role of immune cells stimulated by *P. distasonis.*

More importantly, human data show that delayed exposure to the *P. distasonis* insB:9-23 mimic

might be pathogenic in the first three years of life.

MATERIAL AND METHODS

Bioinformatics

We performed a comprehensive bioinformatics search for the presence of the microbial peptide sequences with homology to human insB:9-23 sequence at the National Center for Biotechnology Information (NCBI). A bioinformatics search was performed by NCBI BLASTp using insB:9-23 sequence as query against viral (NCBI taxonomic ID: 10239), bacterial (NCBI taxonomic ID: 2) and fungal (NCBI taxonomic ID: 10239) proteomes. The whole peptide sequence of each significant hit was compared with insB:9-23 using a multiple sequence alignment program (Clustal Omega) to determine the number of the identical and conserved residues. This yielded the data in Tables 1 and S1. An additional bioinformatics search was performed to determine whether P. distasonis peptide is unique. We used P. distasonis peptide (15 aa) or P. distasonis hypoxanthine phosphoribosyltransferase protein sequences as queries and searched against BLASTp using non redundant protein sequences. This yielded the data in Figure S1 and Supplementary documents 1-2.

Mouse procedures

NOD/ShiLtJ (NOD) and NOD. Cg-*Pcrkdc scid*/J (NOD.Scid) mice were purchased from Jackson Laboratory. Mice were maintained and bred in Boston College Animal Care Facility. All of the mice were maintained under specific pathogen-free conditions in a 12-h dark/light cycle with autoclaved food, water and bedding. All experiments complied with regulations and ethics guidelines of the National Institute of Health and were approved by the IACUC of Boston College (no B2019-003, 2019-004).

Human Plasma Samples

Peripheral blood was collected from living subjects following the provision of written informed consent (and assent in the case of minors) in accordance with IRB approved protocols (University of Florida IRB# 201400703) and the Declaration of Helsinki.

Human T-Cell Clone Stimulation.

Twenty 15-mer peptides (**Table 1**) used in human and mouse T-cell stimulation experiments were chemically synthesized by Genscript (TFA removal, >85% purity). The peptides were dissolved in DMSO at 20 mg/mL (~12 mM). Human T-cell clone stimulation assays were performed as previously described (74). Briefly, insulin B: 11-23-specific T-cell clones were stimulated in 96-well round bottom plates with irrelevant, specific, or microbial mimotope peptides in the presence of irradiated DQ8-transdimer-transfected HEK293 cells as APCs. 50 µl of supernatants from cultures of T cell clones were collected after 48 h stimulation and added to each well of 96-well round bottom plates precoated with IFN-γ (clone MD-1) capturing antibodies (BioLegend). After overnight incubation, bound cytokines were detected by biotinylated anti-IFN-γ (clone 4s.B3) and quantified using a Victor2 D time resolved fluorometer (Perkin Elmer). Experiments were performed in the presence of 1 µg/ml of anti-CD28 antibodies. Unless otherwise stated, peptide concentrations were 2.5 µM.

Binding Model for *P. distasonis* peptide

Coordinates for the insB:9-23 peptide bound to HLA-DQ8 were superimposed on I-Ag7 complexed to GAD peptide. The insB:9-23 was extracted and merged with I-Ag7, and energy minimized to model peptide binding in register 2. Side chains of the insB:9-23 peptide were mutated to accommodate the register 3 shift, followed by energy minimization. *P. distasonis*

403 peptide and insB:9-23 peptide sequences were aligned and the coordinates mutated to the *P.*

404 *distasonis* peptide, and the resulting model of peptide complexed to I-Ag7 was energy minimized.

405 The MHCI binding predictions were made on 11/30/2019 using the IEDB analysis resource

406 Consensus tool (75) which combines predictions from ANN aka NetMHC (4.0) (76-78), SMM

407 (79) and Comblib (80) .

NOD mice T-cell stimulation and antigen presentation assay

We performed the experiments as described previously (29). The peptides were ordered from

Genscript (TFA removal, >85% purity) and dissolved in a base buffer consisting of 50mM NaCl,

10mM HEPES, pH6.8 with 200uM TCEP. Because *P. distasonis* peptide has a neutral charge, we

included 5% DMSO to dissolve it (final DMSO concentration is less than 1%). The C3g7 cell line,

which express an abundance of MHC-II I-Ag7, was used as APC and were treated with 1/2log

dilutions of peptide (starting from 10uM). These cells were then cultured with the IIT-3 or 8F10 T

cell hybridomas for 18 h. Culture supernatants were assayed for IL-2 by incubation with the CTLL-

2 cell line. CTLL-2 cells are responsive to IL-2 and only actively divide in the presence of IL-2.

We then determined the 3H uptake (CPM) to measure how many CTLL cells are proliferating

using a scintillation counter.

Immunization of the NOD mice

The experiments were performed as described previously (81). Briefly, 13 week old male NOD

mice (n=2 mice per group) were immunized in the footpad with the *P. distasonis* peptide or insulin

B:9-23 peptide (10nmoles/mouse). After 7 days, the draining lymph nodes were removed and

pooled for examination by ELISpot. In the ELISpot assay, node cells were recalled with the various

peptides to elicit either an IL-2 or IFN-g response. Spots were analyzed by Immunospot 5.0

(C.T.L.)

Reanalysis of the published metagenomics data

BABYDIET 16S sequencing data (36) (22 cases and 22 matched controls) were provided by Dr. David Endesfelder (Scientific Computing Research Unit, Helmholtz Zentrum München, Munich, Germany). To determine the abundance of the OTUs corresponding to each group, we calculated the normalized average number of *P. distasonis* OTUs (Figure 4A) for each group based on the collection time of the stool samples, e.g. 0-1, 1-2 and 2-3 years. Data is shown with median OTU number/group normalized to the means of each sample. We downloaded the DIABIMMUNE data with the help of Dr. Alex Kostic. Because the reads were not ideal for a peptide search, we first assembled the reads using SPAdes (11) and they are available at https://github.com/ablab/spades. We used these assembled contigs for a TBLASTN search to identify "RILVELLYLVCSEYL" encoding sequences in these samples. We then classified the data based on the collection time of the stool samples, e.g. 0-1, 1-2 and 2-3 years (Figure 4B) and country (Figure 4C).

Bacterial culture

Parabacteroides distasonis D 13 strain was purchased from Dr. Emma Allen-Vercoe's laboratory at University of Guelph. Dr. Allen-Vercoe's group isolated this bacterium from the colon of a ulcerative colitis patient. *P. distasonis D13* strain and *Bacteroides fragilis* (ATCC® 25285) were cultured in anaerobic culture broth (Tryptic Soy Broth supplemented with 5ug/ml Hemin (BD Biosciences) and 1ug/ml Vitamin K1 (VWR)) at 37°C in anaerobic chamber (Coy Laboratory Product).

Oral gavage of *Parabacteroides distasonis* into NOD mice

After *P. distasonis* were cultured from frozen stock at 37°C overnight (OD=1.5), *P. distasonis* were collected by centrifuging at 4000 rpm for 10 min and washed 3 times with sterile saline

(Baxter). After the last wash, *P. distasonis* were re-suspended with saline and oral gavaged using

22ga plastic feeding tube (Instech Laboratories) (100ul/mouse, 10^9 CFU/mouse). 3-week-old NOD

mice were oral gavaged with *Parabacteroides distasonis D13* daily for four weeks right after

weaning for 4 weeks (n=40/group/sex). Control groups were gavaged with sterile saline.

Fecal bacterial DNA extraction

Mouse fecal samples were collected as fecal pellets in Eppendorf tubes and stored at -80°C right

after collection. The DNA were contracted from 100 mg mouse fecal samples using QIAamp

PowerFecal Pro DNA kit (Qiagen) following manufacturer's instruction.

Quantification of *Parabacteroides distasonis* by qPCR

Bacterial DNA was extracted from NOD mice fecal samples (9-week old) 2 weeks after oral

gavage was completed as described in Fecal bacteria DNA extraction section. qPCR was

conducted using QuantStudio 3 Real-Time PCR System (Applied Biosystems) at 95°C for 10 mins

followed by 40 cycles of 15 sec at 95°C and 1 min at 60°C with Power SYBR Green Master Mix

(Applied Biosystems) following manufacturer's instruction. The primers were described

previously and listed in Table S5 (82, 83). The relevant abundance of *Parabacteroides distasonis*

was determined by normalized with *Eubacteria.*

HPT expression in *Parabacteroides distasonis* by RT-qPCR

Bacterial RNA was extracted from Parabacteroides distasonis using ZymoBIOMICS DNA/RNA

kit (Zymo Research). RNA reverse-transcription and first strand synthesis were performed using

QuantStudio 3 (Applied Biosystems) following manufacturer's instruction. cDNAs were applied

for quantitative PCR through the same system mentioned above All sample reactions running in

triplicate. The primers were designed to reverse transcribe 200 bp fragments of HPT using primer

blast. The primers target all bacteria 16S and *Parabacteroides distasonis* 16S rRNA were

described previously and listed in Table S5(84). Data obtained were normalized using 16S rRNA

expression levels as reference and relative expression was determined by calculating the $\Delta\Delta$Cq.

Immunization of NOD mice with *P. distasonis* peptide

The peptides were synthesized by GenScript USA Inc. 15-mer peptides, Insulin epitope peptide

(LVELLYLVCSEYLNH), insulin 2 peptide (SHLVEALYLCGERG), or 14-mer tetanus toxin

peptide (QYIKANSKFIGIFE) (Purity>85%; 10ug/mouse/week) in saline were subcutaneously

injected into NOD mice (n=20) weekly from 4 weeks old until 25 weeks old as described

previously (35). Tail vein blood glucose was monitored weekly and mice with >250mg/dl for two

consecutive days as diabetic.

Immune cells transplantation

Splenocytes were isolated from spleen and pancreatic lymph nodes of 15 weeks donor NOD mice

by filtering through a sterile nylon mesh followed by red blood cell lysis with ACK (Lonza). The

isolated splenocytes were intravenously injected through lateral tail vein into recipient NOD SCID

mice (9 weeks old) (5×10^7 cells/mouse). Donor NOD mice and recipient NOD.SCID mice were

sex matched. Recipient mice were monitored for diabetes by checking tail vein blood glucose level

(>250mg/dl) twice per week. The experiment terminated 11 weeks after the transplantation unless

the mice were diagnosed with diabetes. Donor mice were 15-week male or female NOD mice

colonized with either *P. distasonis* or their sex matched saline control (n=16).

Insulitis score

Pancreas were isolated from *P. distasonis* colonized 12 weeks NOD mice and control mice (n=5)

followed by fixation with 4% paraformaldehyde overnight at 4°C, embedding with paraffin,

sectioning into 5µm thickness and stained with hematoxylin-eosin (Harvard University, BIDMC

Histology Core). Slides were analyzed using EVOS XL Core microscope with a LPlan PH2

10×/0.25 and a LPlan PH2 20×/0.40 objective. The pancreatic insulitis was screened and evaluated

as described previously (85).

Western Blot analysis

Bacteria lysis protein were extracted with CelLytic B Cell Lysis Reagent (Sigma) following

manufacturer's instruction. Protein concentration were determined by BCA assay (Thermo

Scientific). 20 µg bacterial protein was loaded for well for the gel. Standard western blots were

performed by using tank blotting (Bio-rad) as membrane transfer equipment for 1h at 4°C.

Membrane were then blocked with 5% non fat milk+PBST. After washing, each membrane was

incubated with mouse serum (1: 2,500) or human serum (1: 10,000) overnight at 4°C followed by

washing and Sheep anti-mouse (1:3000, Millipore) or goat anti-human antibody (1:3000, Santa

Cruz) incubation for 1h. After washing, the blots were developed with chemiluminescent substrate

ECL (Thermo Fisher)

Endotoxin and IAA quantification and ELISA

507   12-week NOD Mouse serum was collected and stored at −80 °C until use. Serum endotoxin level

508   was measured by using a Pierce Chromogenic Endotoxin Quant Kit (Thermo Scientific) following

509   the manufacturers' instructions. We used mouse IAA ELISA Kit (MyBioSource, MBS26087) to

510   measure IAA levels. The experiment was completed according manufacturer's instructions.

Luminex

MCYTMAG-70K-PX32 Milliplex Mouse Cytokine/Chemokine MAGNETIC BEAD Premixed

32 Plex Kit (**MilliporeSigma**) was used to determine circulating cytokines and chemokines in

mice plasma. The experiment was performed according to manufacturer's instructions and the

plates were read using Luminex® 200™ (Thermo Fisher) at Multiplex Core of Forsyth Institute.

Statistics

Data are presented as the mean ± SEM. Survival curves were analyzed by log-rank (Mantel-cox)

test. Statistical significance was evaluated using unpaired two-tailed Student's *t*-test for two-group

comparison or ANOVA, followed by a Tukey-Kramer post-hoc, Bonferroni's post-hoc where

appropriate. A P value of less than 0.05 was considered significant (*).

Proteomics Analysis. Gel slices were submitted for mass-spectrometry analysis and protein

identification, which were performed at the Taplin Mass Spectrometry Facility at Harvard Medical

School as previously described (86)

REFERENCES

1. Pugliese A. Autoreactive T cells in type 1 diabetes. *J Clin Invest.* 2017;127(8):2881-91.

2. Patterson CC, Harjutsalo V, Rosenbauer J, Neu A, Cinek O, Skrivarhaug T, et al. Trends and cyclical variation in the incidence of childhood type 1 diabetes in 26 European centres in the 25 year period 1989-2013: a multicentre prospective registration study. *Diabetologia.* 2019;62(3):408-17.

3. Mayer-Davis EJ, Lawrence JM, Dabelea D, Divers J, Isom S, Dolan L, et al. Incidence Trends of Type 1 and Type 2 Diabetes among Youths, 2002-2012. *N Engl J Med.* 2017;376(15):1419-29.

4. Kondrashova A, Reunanen A, Romanov A, Karvonen A, Viskari H, Vesikari T, et al. A six-fold gradient in the incidence of type 1 diabetes at the eastern border of Finland. *Ann Med.* 2005;37(1):67-72.

5. Redondo MJ, Jeffrey J, Fain PR, Eisenbarth GS, and Orban T. Concordance for islet autoimmunity among monozygotic twins. *N Engl J Med.* 2008;359(26):2849-50.

6. Zhao G, Vatanen T, Droit L, Park A, Kostic AD, Poon TW, et al. Intestinal virome changes precede autoimmunity in type I diabetes-susceptible children. *Proc Natl Acad Sci U S A.* 2017;114(30):E6166-E75.

7. Op de Beeck A, and Eizirik DL. Viral infections in type 1 diabetes mellitus--why the beta cells? *Nat Rev Endocrinol.* 2016;12(5):263-73.

8. Christen U, Hintermann E, Holdener M, and von Herrath MG. Viral triggers for autoimmunity: is the 'glass of molecular mimicry' half full or half empty? *J Autoimmun.* 2010;34(1):38-44.

9. Rewers M, and Ludvigsson J. Environmental risk factors for type 1 diabetes. *Lancet.* 2016;387(10035):2340-8.

10. Davis-Richardson AG, Ardissone AN, Dias R, Simell V, Leonard MT, Kemppainen KM, et al. Bacteroides dorei dominates gut microbiome prior to autoimmunity in Finnish children at high risk for type 1 diabetes. *Frontiers in microbiology.* 2014;5:678.

11. Vatanen T, Franzosa EA, Schwager R, Tripathi S, Arthur TD, Vehik K, et al. The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature.* 2018;562(7728):589-94.

554   12.   Kostic AD, Gevers D, Siljander H, Vatanen T, Hyotylainen T, Hamalainen AM, et al. The
555   dynamics of the human infant gut microbiome in development and in progression toward
556   type 1 diabetes. *Cell Host Microbe.* 2015;17(2):260-73.

557   13.   Vatanen T, Kostic AD, d'Hennezel E, Siljander H, Franzosa EA, Yassour M, et al.
558   Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans.
559   *Cell.* 2016;165(4):842-53.

560   14.   Dedrick S, Sundaresh B, Huang Q, Brady C, Yoo T, Cronin C, et al. The Role of Gut
561   Microbiota and Environmental Factors in Type 1 Diabetes Pathogenesis. *Front Endocrinol*
562   *(Lausanne).* 2020;11:78.

563   15.   Coppieters KT, Boettler T, and von Herrath M. Virus infections in type 1 diabetes. *Cold*
564   *Spring Harb Perspect Med.* 2012;2(1):a007682.

565   16.   Atkinson MA, Eisenbarth GS, and Michels AW. Type 1 diabetes. *Lancet.*
566   2014;383(9911):69-82.

567   17.   Peters L, Posgai A, and Brusko TM. Islet-immune interactions in type 1 diabetes: the nexus
568   of beta cell destruction. *Clin Exp Immunol.* 2019;198(3):326-40.

569   18.   Zhang L, Nakayama M, and Eisenbarth GS. Insulin as an autoantigen in NOD/human
570   diabetes. *Curr Opin Immunol.* 2008;20(1):111-8.

571   19.   Alleva DG, Crowe PD, Jin L, Kwok WW, Ling N, Gottschalk M, et al. A disease-
572   associated cellular immune response in type 1 diabetics to an immunodominant epitope of
573   insulin. *J Clin Invest.* 2001;107(2):173-80.

574   20.   Palmer JP, Asplin CM, Clemons P, Lyen K, Tatpati O, Raghu PK, et al. Insulin antibodies
575   in insulin-dependent diabetics before insulin treatment. *Science.* 1983;222(4630):1337-9.

576   21.   Steck AK, Johnson K, Barriga KJ, Miao D, Yu L, Hutton JC, et al. Age of islet
577   autoantibody appearance and mean levels of insulin, but not GAD or IA-2 autoantibodies,
578   predict age of diagnosis of type 1 diabetes: diabetes autoimmunity study in the young.
579   *Diabetes Care.* 2011;34(6):1397-9.

580   22.   Unanue ER. Antigen presentation in the autoimmune diabetes of the NOD mouse. *Annu*
581   *Rev Immunol.* 2014;32:579-608.

582   23.   Daniel D, Gill RG, Schloot N, and Wegmann D. Epitope specificity, cytokine production
583   profile and diabetogenic activity of insulin-specific T cell clones isolated from NOD mice.
584   *Eur J Immunol.* 1995;25(4):1056-62.

585   24.   Pathiraja V, Kuehlich JP, Campbell PD, Krishnamurthy B, Loudovaris T, Coates PT, et al.
586   Proinsulin-specific, HLA-DQ8, and HLA-DQ8-transdimer-restricted CD4+ T cells
587   infiltrate islets in type 1 diabetes. *Diabetes.* 2015;64(1):172-82.
588   25.   Michels AW, Landry LG, McDaniel KA, Yu L, Campbell-Thompson M, Kwok WW, et
589   al. Islet-Derived CD4 T Cells Targeting Proinsulin in Human Autoimmune Diabetes.
590   *Diabetes.* 2017;66(3):722-34.
591   26.   Nakayama M, McDaniel K, Fitzgerald-Miller L, Kiekhaefer C, Snell-Bergeon JK,
592   Davidson HW, et al. Regulatory vs. inflammatory cytokine T-cell responses to mutated
593   insulin peptides in healthy and type 1 diabetic subjects. *Proc Natl Acad Sci U S A.*
594   2015;112(14):4429-34.
595   27.   Yang J, Chow IT, Sosinowski T, Torres-Chinn N, Greenbaum CJ, James EA, et al.
596   Autoreactive T cells specific for insulin B:11-23 recognize a low-affinity peptide register
597   in human subjects with autoimmune diabetes. *Proc Natl Acad Sci U S A.*
598   2014;111(41):14840-5.
599   28.   Cusick MF, Libbey JE, and Fujinami RS. Molecular mimicry as a mechanism of
600   autoimmune disease. *Clin Rev Allergy Immunol.* 2012;42(1):102-11.
601   29.   Wan X, Zinselmeyer BH, Zakharov PN, Vomund AN, Taniguchi R, Santambrogio L, et
602   al. Pancreatic islets communicate with lymphoid tissues via exocytosis of insulin peptides.
603   *Nature.* 2018;560(7716):107-11.
604   30.   Sakamoto M, and Benno Y. Reclassification of Bacteroides distasonis, Bacteroides
605   goldsteinii and Bacteroides merdae as Parabacteroides distasonis gen. nov., comb. nov.,
606   Parabacteroides goldsteinii comb. nov. and Parabacteroides merdae comb. nov. *Int J Syst*
607   *Evol Microbiol.* 2006;56(Pt 7):1599-605.
608   31.   Abuaita BH, and Withey JH. Bicarbonate Induces Vibrio cholerae virulence gene
609   expression by enhancing ToxT activity. *Infect Immun.* 2009;77(9):4111-20.
610   32.   Stadinski BD, Zhang L, Crawford F, Marrack P, Eisenbarth GS, and Kappler JW.
611   Diabetogenic T cells recognize insulin bound to IAg7 in an unexpected, weakly binding
612   register. *Proc Natl Acad Sci U S A.* 2010;107(24):10978-83.
613   33.   Sofi MH, Johnson BM, Gudi RR, Jolly A, Gaudreau MC, and Vasu C. Polysaccharide A-
614   Dependent Opposing Effects of Mucosal and Systemic Exposures to Human Gut
615   Commensal Bacteroides fragilis in Type 1 Diabetes. *Diabetes.* 2019;68(10):1975-89.

34.   Fuchtenbusch M, Larger E, Thebault K, and Boitard C. Transfer of diabetes from
prediabetic NOD mice to NOD-SCID/SCID mice: association with pancreatic insulin
content. *Horm Metab Res.* 2005;37(2):63-7.

35.   Devendra D, Paronen J, Moriyama H, Miao D, Eisenbarth GS, and Liu E. Differential
immune response to B:9-23 insulin 1 and insulin 2 peptides in animal models of type 1
diabetes. *J Autoimmun.* 2004;23(1):17-26.

36.   Endesfelder D, zu Castell W, Ardissone A, Davis-Richardson AG, Achenbach P, Hagen
M, et al. Compromised gut microbiota networks in children with anti-islet cell
autoimmunity. *Diabetes.* 2014;63(6):2006-14.

37.   Mason D. A very high level of crossreactivity is an essential feature of the T-cell receptor.
Immunol Today. 1998;19(9):395-404.

38.   Calis JJ, de Boer RJ, and Kesmir C. Degenerate T-cell recognition of peptides on MHC
molecules creates large holes in the T-cell repertoire. *PLoS Comput Biol.*
2012;8(3):e1002412.

39.   Wucherpfennig KW, and Strominger JL. Molecular mimicry in T cell-mediated
autoimmunity: viral peptides activate human T cell clones specific for myelin basic protein.
Cell. 1995;80(5):695-705.

40.   Mana P, Goodyear M, Bernard C, Tomioka R, Freire-Garabal M, and Linares D. Tolerance
induction by molecular mimicry: prevention and suppression of experimental autoimmune
encephalomyelitis with the milk protein butyrophilin. *Int Immunol.* 2004;16(3):489-99.

41.   Atkinson MA, Bowman MA, Campbell L, Darrow BL, Kaufman DL, and Maclaren NK.
Cellular immunity to a determinant common to glutamate decarboxylase and coxsackie
virus in insulin-dependent diabetes. *J Clin Invest.* 1994;94(5):2125-9.

42.   Schloot NC, Willemen SJ, Duinkerken G, Drijfhout JW, de Vries RR, and Roep BO.
Molecular mimicry in type 1 diabetes mellitus revisited: T-cell clones to GAD65 peptides
with sequence homology to Coxsackie or proinsulin peptides do not crossreact with
homologous counterpart. *Hum Immunol.* 2001;62(4):299-309.

43.   Tian J, Lehmann PV, and Kaufman DL. T cell cross-reactivity between coxsackievirus and
glutamate decarboxylase is associated with a murine diabetes susceptibility allele. *J Exp*
Med. 1994;180(5):1979-84.

44. Honeyman MC, Stone NL, Falk BA, Nepom G, and Harrison LC. Evidence for molecular
mimicry between human T cell epitopes in rotavirus and pancreatic islet autoantigens. *J*
Immunol. 2010;184(4):2204-10.

45. Maruvada P, Leone V, Kaplan LM, and Chang EB. The Human Microbiome and Obesity:
Moving beyond Associations. *Cell Host Microbe.* 2017;22(5):589-99.

46. Rinaldi M, Perricone R, Blank M, Perricone C, and Shoenfeld Y. Anti-Saccharomyces
cerevisiae autoantibodies in autoimmune diseases: from bread baking to autoimmunity.
Clin Rev Allergy Immunol. 2013;45(2):152-61.

47. Gevers D, Kugathasan S, Denson LA, Vazquez-Baeza Y, Van Treuren W, Ren B, et al.
The treatment-naive microbiome in new-onset Crohn's disease. *Cell Host Microbe.*
2014;15(3):382-92.

48. Lamps LW, Madhusudhan KT, Havens JM, Greenson JK, Bronner MP, Chiles MC, et al.
Pathogenic Yersinia DNA is detected in bowel and mesenteric lymph nodes from patients
with Crohn's disease. *The American journal of surgical pathology.* 2003;27(2):220-7.

49. Chassaing B, Rolhion N, de Vallee A, Salim SY, Prorok-Hamon M, Neut C, et al. Crohn
disease--associated adherent-invasive E. coli bacteria target mouse and human Peyer's
patches via long polar fimbriae. *J Clin Invest.* 2011;121(3):966-75.

50. Carding S, Verbeke K, Vipond DT, Corfe BM, and Owen LJ. Dysbiosis of the gut
microbiota in disease. *Microb Ecol Health Dis.* 2015;26:26191.

51. Frank DN, St Amand AL, Feldman RA, Boedeker EC, Harpaz N, and Pace NR. Molecular-
phylogenetic characterization of microbial community imbalances in human inflammatory
bowel diseases. *Proc Natl Acad Sci U S A.* 2007;104(34):13780-5.

52. Navaneethan U, Venkatesh PG, and Shen B. Clostridium difficile infection and
inflammatory bowel disease: understanding the evolving relationship. *World J*
Gastroenterol. 2010;16(39):4892-904.

53. Quagliariello A, Aloisio I, Bozzi Cionci N, Luiselli D, D'Auria G, Martinez-Priego L, et
al. Effect of Bifidobacterium breve on the Intestinal Microbiota of Coeliac Children on a
Gluten Free Diet: A Pilot Study. *Nutrients.* 2016;8(10).

54. Heintz-Buschart A, May P, Laczny CC, Lebrun LA, Bellora C, Krishna A, et al. Integrated
multi-omics of the human gut microbiome in a case study of familial type 1 diabetes. *Nat*
Microbiol. 2016;2:16180.

55.   Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal
development of the gut microbiome in early childhood from the TEDDY study. *Nature.*
2018;562(7728):583-8.

56.   Lathrop SK, Bloom SM, Rao SM, Nutsch K, Lio CW, Santacruz N, et al. Peripheral
education of the immune system by colonic commensal microbiota. *Nature.*
2011;478(7368):250-4.

57.   Nakayama M, Abiru N, Moriyama H, Babaya N, Liu E, Miao D, et al. Prime role for an
insulin epitope in the development of type 1 diabetes in NOD mice. *Nature.*
2005;435(7039):220-3.

58.   Higashide T, Kawamura T, Nagata M, Kotani R, Kimura K, Hirose M, et al. T cell epitope
mapping study with insulin overlapping peptides using ELISPOT assay in Japanese
children and adolescents with type 1 diabetes. *Pediatr Res.* 2006;59(3):445-50.

59.   Kent SC, Chen Y, Bregoli L, Clemmings SM, Kenyon NS, Ricordi C, et al. Expanded T
cells from pancreatic lymph nodes of type 1 diabetic subjects recognize an insulin epitope.
Nature. 2005;435(7039):224-8.

60.   Mannering SI, Harrison LC, Williamson NA, Morris JS, Thearle DJ, Jensen KP, et al. The
insulin A-chain epitope recognized by human T cells is posttranslationally modified. *J Exp*
Med. 2005;202(9):1191-7.

61.   Mannering SI, Di Carluccio AR, and Elso CM. Neoepitopes: a new take on beta cell
autoimmunity in type 1 diabetes. *Diabetologia.* 2019;62(3):351-6.

62.   Purcell AW, Sechi S, and DiLorenzo TP. The Evolving Landscape of Autoantigen
Discovery and Characterization in Type 1 Diabetes. *Diabetes.* 2019;68(5):879-86.

63.   Delong T, Wiles TA, Baker RL, Bradley B, Barbour G, Reisdorph R, et al. Pathogenic
CD4 T cells in type 1 diabetes recognize epitopes formed by peptide fusion. *Science.*
2016;351(6274):711-4.

64.   Kracht MJ, van Lummel M, Nikolic T, Joosten AM, Laban S, van der Slik AR, et al.
Autoimmunity against a defective ribosomal insulin gene product in type 1 diabetes. *Nat*
Med. 2017;23(4):501-7.

65.   Babon JA, DeNicola ME, Blodgett DM, Crevecoeur I, Buttrick TS, Maehr R, et al.
Analysis of self-antigen specificity of islet-infiltrating T cells from human donors with type
1 diabetes. *Nat Med.* 2016;22(12):1482-7.

66.   Tai N, Peng J, Liu F, Gulden E, Hu Y, Zhang X, et al. Microbial antigen mimics activate
709   diabetogenic CD8 T cells in NOD mice. *J Exp Med.* 2016;213(10):2129-46.

710   67.   Markle JG, Frank DN, Mortin-Toth S, Robertson CE, Feazel LM, Rolle-Kampczyk U, et
711   al. Sex differences in the gut microbiome drive hormone-dependent regulation of
712   autoimmunity. *Science.* 2013;339(6123):1084-8.

713   68.   Kriegel MA, Sefik E, Hill JA, Wu HJ, Benoist C, and Mathis D. Naturally transmitted
714   segmented filamentous bacteria segregate with diabetes protection in nonobese diabetic
715   mice. *Proc Natl Acad Sci U S A.* 2011;108(28):11548-53.

716   69.   Hanninen A, Toivonen R, Poysti S, Belzer C, Plovier H, Ouwerkerk JP, et al. Akkermansia
717   muciniphila induces gut microbiota remodelling and controls islet autoimmunity in NOD
718   mice. *Gut.* 2018;67(8):1445-53.

719   70.   Song L, Xiong D, Song H, Wu L, Zhang M, Kang X, et al. Mucosal and Systemic Immune
720   Responses to Influenza H7N9 Antigen HA1-2 Co-Delivered Intranasally with Flagellin or
721   Polyethyleneimine in Mice and Chickens. *Front Immunol.* 2017;8:326.

722   71.   Greiling TM, Dehner C, Chen X, Hughes K, Iniguez AJ, Boccitto M, et al. Commensal
723   orthologs of the human autoantigen Ro60 as triggers of autoimmunity in lupus. *Sci Transl*
724   *Med.* 2018;10(434).

725   72.   Gil-Cruz C, Perez-Shibayama C, De Martin A, Ronchi F, van der Borght K, Niederer R, et
726   al. Microbiota-derived peptide mimics drive lethal inflammatory cardiomyopathy. *Science.*
727   2019;366(6467):881-6.

728   73.   Harkiolaki M, Holmes SL, Svendsen P, Gregersen JW, Jensen LT, McMahon R, et al. T
729   cell-mediated autoimmune disease due to low-affinity crossreactivity to common microbial
730   peptides. *Immunity.* 2009;30(3):348-57.

731   74.   Chow IT, Gates TJ, Papadopoulos GK, Moustakas AK, Kolawole EM, Notturno RJ, et al.
732   Discriminative T cell recognition of cross-reactive islet-antigens is associated with HLA-
733   DQ8 transdimer-mediated autoimmune diabetes. *Sci Adv.* 2019;5(8):eaaw9336.

734   75.   Kim Y, Ponomarenko J, Zhu Z, Tamang D, Wang P, Greenbaum J, et al. Immune epitope
735   database analysis resource. *Nucleic Acids Res.* 2012;40(Web Server issue):W525-30.

736   76.   Nielsen M, Lundegaard C, Worning P, Lauemoller SL, Lamberth K, Buus S, et al. Reliable
737   prediction of T-cell epitopes using neural networks with novel sequence representations.
738   *Protein Sci.* 2003;12(5):1007-17.

77. Lundegaard C, Lamberth K, Harndahl M, Buus S, Lund O, and Nielsen M. NetMHC-3.0:
accurate web accessible predictions of human, mouse and monkey MHC class I affinities
for peptides of length 8-11. *Nucleic Acids Res.* 2008;36(Web Server issue):W509-12.

78. Andreatta M, and Nielsen M. Gapped sequence alignment using artificial neural networks:
application to the MHC class I system. *Bioinformatics.* 2016;32(4):511-7.

79. Peters B, and Sette A. Generating quantitative models describing the sequence specificity
of biological processes with the stabilized matrix method. *BMC Bioinformatics.*
2005;6:132.

80. Sidney J, Assarsson E, Moore C, Ngo S, Pinilla C, Sette A, et al. Quantitative peptide
binding motifs for 19 human and mouse MHC class I molecules derived using positional
scanning combinatorial peptide libraries. *Immunome Res.* 2008;4:2.

81. Mohan JF, Levisetti MG, Calderon B, Herzog JW, Petzold SJ, and Unanue ER. Unique
autoreactive T cells recognize insulin peptides generated within the islets of Langerhans in
autoimmune diabetes. *Nat Immunol.* 2010;11(4):350-4.

82. Dziarski R, Park SY, Kashyap DR, Dowd SE, and Gupta D. Pglyrp-Regulated Gut
Microflora Prevotella falsenii, Parabacteroides distasonis and Bacteroides eggerthii
Enhance and Alistipes finegoldii Attenuates Colitis in Mice. *PLoS One.*
2016;11(1):e0146162.

83. Croswell A, Amir E, Teggatz P, Barman M, and Salzman NH. Prolonged impact of
antibiotics on intestinal microbial ecology and susceptibility to enteric Salmonella
infection. *Infect Immun.* 2009;77(7):2741-53.

84. Srivastava S, Singh V, Kumar V, Verma PC, Srivastava R, Basu V, et al. Identification of
regulatory elements in 16S rRNA gene of Acinetobacter species isolated from water
sample. *Bioinformation.* 2008;3(4):173-6.

85. Flodstrom-Tullberg M, Yadav D, Hagerkvist R, Tsai D, Secrest P, Stotland A, et al. Target
cell expression of suppressor of cytokine signaling-1 prevents diabetes in the NOD mouse.
Diabetes. 2003;52(11):2696-700.

86. Pollock N, Dhiman R, Daifalla N, Farhat M, and Campos-Neto A. Discovery of a unique
Mycobacterium tuberculosis protein through proteomic analysis of urine from patients with
active tuberculosis. *Microbes Infect.* 2018;20(4):228-35.

Acknowledgements

We first want to thank Prof. Emil Unanue and Dr. Anthony Vomund (University of Washington, St. Louis) for testing microbial insulins on T cell hybridomas, immunization experiments and with their help to interpret the data. Thanks to Jonathan.Dreyfuss and Hui Pan (Joslin Bioinformatics Core) for bioinformatics analysis. Thanks to Babak Momeni (Boston College) for sharing his laboratory's anaerobic chamber. We acknowledge BC Biology Department undergraduate students Tu Tran, Ruixu Huang with the microscopy work and David Kim and Scott Hsu for their help with the animal experiments. We thank to Alex Kostic, Tao Xu, Thomas Serwold and Shio Kobayashi (Joslin Diabetes Center) for the useful Discussion. Thanks to Annette Ziegler and David Endesfelder for providing the BABYDIET 16S sequencing data for reanalysis. Thanks to Amanda Posgai for editing the manuscript.

Funding: This work was supported by National Institutes of Health Grants 1K01DK117967-01 (to E.A.) and R01DK031026 and R01DK033201 (to C.R.K.) and NIH P01 AI042288 to MAA. E.A. was supported by The G. Harold and Leila Y. Mathers Charitable Foundation Research Grant and an Iacocca Family Foundation Fellowship.

Author contributions: Q.H assisted with all NOD mice experiments including peptide immunization, oral gavage, splenocyte transfer and ELISA/Luminex experiments, qPCR and Western blot analysis using human and mice plasma. A.V and E.U assisted with the experiments using NOD mice T-cell hybridomas. I.C and W.K assisted with the experiments using NOD mice T-cell clones. C.B and A.R assisted with the peptide immunization and oral gavage experiments. D.O and D.L assisted with MHC-binding models. M.A provided the human plasma samples. E.A

791   completed bioinformatics analysis. E.A and C.R.K designed the research. All authors assisted with

792   the analysis of the data. E.A supervised the project and wrote the paper.

793   **Competing Interests:** The authors do not have any conflict of interest related to this study.

794   **Data and materials availability:** Because the reads were not ideal for a peptide search, we first

795   assembled the reads using SPAdes (11) and they are available at https://github.com/ablab/spades.

FIGURES

Figure 1. *P. distasonis* **insB:9-23peptide can stimulate insB:13-21 specific NOD mice T cell hybridomas and structurally mimic the epitope. (A)** CD4+ T cells recognize *P. distasonis* peptide. Response of the B:13–21-specific IIT-3 hybridomas (red) and B:12–20-specific8F10 (blue) to insulin B9-23 or *P. distasonis* peptides covalently linked to I-A^{g7} expressed on macrophages. **(B)** Binding model for *P. distasonis* mimic peptide compared to insB9-23. In 3B, I-A^{g7} α-chain in tan, I-A^{g7} β-chain in green. insB:9-23 peptide shown as yellow sticks, *P. distasonis* peptide shown as white sticks. Red sticks and spheres show positions identical in *P. distasonis* peptide and insB:9-23 peptide. **(C-F)** Mice (n=2/group) were immunized with either insB:9-23 peptide (**C-E**) or *P. distasonis* peptide (**D-F**) and draining lymph nodes were removed and pooled after 7 days. Cells were stimulated either by *P. distasonis* peptide, insB:9-23, insulin, ConA (positive control) or scrambled peptide (negative control). Secretion of IL-2 (**C-D**) and interferon gamma (**E-F**) were determined by EliSPOT.

Figure 2. P. distasonis colonization enhances disease onset in female NOD mice. **(A)** Schematic overview of the *P. distasonis* oral gavage experiments (n=40/group/sex). Blue arrow shows fecal sample collection for qPCR experiments and red arrow shows when 5 mice were removed from each group (week 12). **(B)** Relative abundance of *P. distasonis* in fecal samples determined by qPCR (week 12, n=12-14/male, n=13-17/female). **(C)** Insulitis scores obtained from mice at week 12 (n=5/group/sex). **(D)** Diabetes incidence in NOD mice (n=35/group/sex) after daily saline or *P. distasonis* oral gavage for 30 days after weaning *(P<0.05).* **(E-F)** Western blot analysis using plasma samples from (E) female and (F) male mice either oral gavaged with *P. distasonis* or saline (week 12, n=5/group/sex). **(G)** Plasma LPS levels (week 12, n=5/group/sex).

Figure 3. Adoptive transfer accelerated diabetes in female NOD.SCID mice. (A) Schematic overview of the adoptive transfer experiments from NOD mice to NOD.SCID mice. 5×10^{7} splenocytes/mouse were transferred from individual NOD mice to NOD.SCID mice at 6 weeks of age (1:1 ratio, same gender, n=16-20). **(B)** Diabetes incidence of donor NOD mice until adoptive transfer (n=20). **(C)** Diabetes incidence of the recipient NOD.SCID mice after adoptive transfer (n=16-20) *(P<0.05)*. **(D-E)** Diabetes incidence of (D) male and (E) female mice (n=20/group/sex) immunized with either *P. distasonis* peptide or ins2:B:9-23 peptide or Tetanus toxin peptide or vector *(P<0.05)*. Data are mean ± SEM. Survival curves were analyzed by log-rank (Mantel-cox) test.

Figure 4. Reanalysis of BABYDIET and DIABIMMUNE metagenome data for P. distasonis peptide and identification of *P. distasonis* specific immune response in humans. (**A**) BABYDIET data was analyzed for the presence of *P. distasonis* OTUs in three different groups, control, antibody-positive and T1D. (**B-C**) Reanalysis of DIABIMMUNE metagenome data for the presence of *P. distasonis* ins9:23 like peptide (**B**) in all three countries and (**C**) in individual countries. (**D**) Insulin B: 11-23-specific human T-cell clones were stimulated with each of selected 17 microbial peptides or 2 negative control (irrelevant and 20) peptides or 2 positive control (18-19) (peptide concentration, 2.5 µM) and IFN-γ secretion was measured *(P<0.05)*. The corresponding peptides for the numbers are listed in Table 1. (**E**) P. distasonis peptide produced a dose response in stimulating IFN-γ secretion. B:9-23 and B:9-23R22E are positive controls of the experiment. (**F**) Western blot analysis using plasma samples obtained from eight female subjects with T1D or eight healthy female subjects against *P. distasonis* lysate. Data are mean ± SEM. Survival curves were analyzed by log-rank (Mantel-cox) test.

848 **Table 1. The sequences of the microbial mimics of the insB:9-23 tested in this study. Red**

849 **indicate identical residues.**

No	Classification	Source	B:9-23 like peptides	Identity
1	**Positive**	Human insulin B:9-23 peptide	SHLVEALYLVCGERG	Control
2	**Control**	Hyperactive peptide	SHLVEALYLVCGEEG	Control
3	**Human Gut**	*Parabacteroides distasonis 33B*	RILVELLYLVCSEYL	9 of 15
4	**Microbiome**	*Bacteroides sp. CAG:144*	LDFKEALYLGCGDRT	8 of 15
5		*Ruminococcus gnavus CAG:126*	DPRRSALYLFCGKRC	7 of 15
6		*Coprococcus eutactus*	NHDKEALYIYCDETE	7 of 15
7		*Clostridiales bacterium VE202-14*	VRAGYALFLVCDEEK	7 of 15
8	**Human**	*Corynebacterium genitalium*	FVHEDALHLVCGERI	9 of 15
9	**Genital Microbiome**	*Lactobacillus vaginalis*	LQSMEIPYLVCGERE	8 of 15
10	**Viruses**	*Lymphocystis disease virus 1*	AHLVAALQRVCGNRG	10 of 15
11		*Cyprinid herpesvirus 1*	SHPNVFIALVCGERG	9 of 15
12		*Grouper iridovirus*	GELIDALTEHCGDRG	7 of 15
13	**Human pathogen**	*Burkholderia multivorans*	LHLARALYEMCGEFP	8 of 15
14	**Other microbes**	*Bradyrhizobium japonicum SEMIA 5079*	VSGKHALYLYCGERG	9 of 15
15		*Streptomyces griseus*	RDRVEALRLVCGEAM	9 of 15
16		*Tetrasphaera japonica T1-X7*	HWLVEIAYLVCGDRR	8 of 15
17		*Saccharomonospora halophila*	TAHGVAEYLVCGERR	8 of 15
18		*Brevundimonas sp. BAL3*	WVGFETLYLYCGERL	8 of 15
19		*Metarhizium robertsii*	DHWDEAGFLVCGERG	10 of 15
20	**Negative Control**	*Negative control peptide (random)*	LELYARVGVSECHGL	Control

A.

>EEY82194.1 hypoxanthine phosphoribosyltransferase [Bacteroides sp. 2_1_33B]
MLN**RILVELLYLVCSEYL**NHNSRCLGFYSAYKCITNFGMLLHFFVHRRKSCTFVPSKTERFMDRIRLKDKEFELFIPESDIQAAIAKMAVQIKADVEGKNPLFVGVLNGAFMFVAELMRE
LDVPYELTFARYSSYQGTSSTGILNEIMPVQADIRGRMVILLEDIIDTGFTMSYVMEKLRSEGAADVRLATMLFKPESLKCELTPDYVGLQIPADFIVGHGLDYDELGRSYKDIYKVVE

>EEU50637.1 hypoxanthine phosphoribosyltransferase [Parabacteroides sp. D13]
MLN**RILVELLYLVCSEYL**NHNSRCLGFYSAYKCITNFGMLLHFFVHRRKSCTFVPSKTERFMDRIRLKDKEFELFIPESDIQAAIAKMAVQIKADVEGKNPLFVGVLNGAFMFVAELMRE
LDVPYELTFARYSSYQGTSSTGILNEIMPVQADICGRMVILLEDIIDTGFTMSYVMEKLRSEGAADVRLATMLFKPESLKCELTPDYVGLQIPADFIVGHGLDYDELGRSYKDIYKVVE

B.   **C.**

A.

No insulitis

Peri-insulitis

Moderate insulitis

Severe insulitis

B.

NOD mice serum IAA

IAA concentration (nU/ml)

Male   Female

● Saline   ■ P.distasonis

C.

Female

Saline Mice   *P.distasonis* Mice

250 kDa
150 kDa
100 kDa
75 kDa
50 kDa -
37 kDa -
25 kDa -
15 kDa -
10 kDa -

No Serum   1 2 3 4 5 6 7 8 9 10

***B. fragilis* lysate**

Male

Saline Mice   *P.distasonis* Mice

250 kDa
150 kDa
100 kDa
75 kDa
50 kDa -
37 kDa -
25 kDa -
15 kDa -

1 2 3 4 5 6 7 8 9 10

B. *fragilis* lysis