1

A Gut Microbial Peptide and Molecular Mimicry in the Pathogenesis of Type 1 Diabetes

Khyati Girdhar[1], Qian Huang[1], I-Ting Chow[2], Claudia Brady[1], Amol Raisingani[1], Patrick

Autissier[1], Mark A. Atkinson[3,4], William W. Kwok[2], C. Ronald Kahn[5], Emrah Altindis[1]*

[1] Boston College Biology Department, Higgins Hall, 140 Commonwealth Avenue Chestnut Hill,

MA 02467, USA

[2] Benaroya Research Institute at Virginia Mason, Seattle, WA 98101, USA

[3] Department of Pathology, Immunology and Laboratory Medicine, College of Medicine,

University of Florida Diabetes Institute, Gainesville, FL 32610-3633, USA

[4] Department of Pediatrics, College of Medicine, University of Florida Diabetes Institute,

Gainesville, FL 32610-3633, USA

[5] Joslin Diabetes Center, Harvard Medical School, Boston, MA, 02215, USA

        *Corresponding Author: Emrah Altindis, Boston College Biology Department, Higgins

Hall, 140 Commonwealth Avenue Chestnut Hill, MA 02467. E-mail: altindis@bc.edu

ABSTRACT

Type 1 Diabetes (T1D) is an autoimmune disease characterized by the destruction of

pancreatic β-cells. One of the earliest aspects of this process is development of autoantibodies and

T-cells directed at an epitope in the B-chain of insulin (insB:9-23). Analysis of microbial protein

sequences with homology to insB:9-23 sequence revealed 17 peptides showing >50% identity to

insB:9-23. Of these, one peptide, found in the normal human gut commensal *Parabacteroides*

distasonis, activated both human T cell clones from T1D patients and T-cell hybridomas from non-

obese diabetic (NOD) mice specific to insB:9-23. Immunization of NOD mice with *P. distasonis*

insB:9-23 peptide mimic or insB:9-23 peptide verified immune cross-reactivity. Colonization of

female NOD mice with *P. distasonis* accelerated the development of T1D, increasing

macrophages, dendritic cells and destructive CD8+ T-cells, while decreasing FoxP3+ regulatory

T-cells. Western blot analysis identified *P. distasonis* reacting antibodies in sera of NOD mice

colonized with *P. distasonis* and human T1D patients. Furthermore, adoptive transfer of

splenocytes from *P. distasonis* treated mice to NOD/SCID mice enhanced disease phenotype in

the recipients. Finally, analysis of human infant gut microbiome data revealed that exposure of

infants to *P. distasonis* may modulate disease pathogenesis. Taken together, these data demonstrate

the potential role for an insB:9-23-mimimetic peptide from gut microbiota as a molecular trigger

or modifier of T1D pathogenesis.

SIGNIFICANCE STATEMENT

In Type 1 diabetes (T1D), immune cells destroy pancreatic β-cells. The trigger of this response, however, is unknown. Some sequences (epitopes) in the insulin molecule form a major target for this autoimmune response. We have identified a sequence in a human gut bacterium that can mimetic this insulin epitope. Immune cells specific to insulin cross-react with this bacterial mimetic. Further, this bacterium can accelerate diabetes onset in a mouse model of T1D, inducing destructive and decreasing protective immune cells. We found this mimetic in the gut of children developing T1D. Furthermore, T1D patients have a stronger immune response to this bacterium compared to healthy individuals. Taken together, this bacterial mimetic in human gut has the potential to trigger/modify T1D onset.

1. INTRODUCTION

Type 1 diabetes (T1D) is an autoimmune disease characterized by selective destruction of pancreatic β-cells by autoreactive T-cells[1]. Genome-wide association studies (GWAS) have identified 152 genetic regions that influence the risk of developing T1D[2]; however, multiple studies have shown that the incidence rate of T1D in children is rising at rates exceeding what can be explained on a genetic basis alone[3]. Indeed, even among identical twins, the concordance of T1D is only 65%[4]. Likewise, there is a six-fold difference in incidence of T1D in neighboring regions of Karelia in Russia and Finland, despite the very similar genetic background of the inhabitants[5].

Various environmental factors have been studied as potential modifiers or triggers of the autoimmune response in T1D including diet, birth mode, infection, and antibiotics. Viral infections have been suggested to play a role in T1D pathogenesis, but most of these viruses have been proposed to act by direct infection of the β-cell[6]. Recently, attention has been focused on the gut microbiome as a potential disease modifier through its effects on metabolite composition, intestinal permeability, and regulation of the immune response in subjects with T1D[7, 8]. However, the exact environmental modifiers and how they might affect T1D pathogenesis remain largely unknown[9].

One of the earliest markers of T1D is the development of islet autoantibodies[10]. These autoantibodies target several autoantigens including insulin, glutamic acid decarboxylase (GAD), insulinoma-associated autoantigen-2 (IA-2), zinc transporter-8 (ZnT8), and an islet-specific glucose-6-phosphatase catalytic subunit related protein (IGRP or G6PC2)[11]. Among these, insulin autoantibodies (IAA) are usually the first to be detected, and insulin is the only autoantigen exclusively expressed by β-cells[12]. In humans, IAA may develop years before the onset of overt

diabetes(13) and are especially prominent in early-onset T1D(14). They also show a significant

correlation with the rate of progression from prediabetes to overt disease(15). More importantly,

insulin or insulin-derived peptides are a target of disease pathogenetic T-cells in both humans with

T1D and in the most established murine model, the NOD mouse. In NOD mice, over 90% of the

anti-insulin T-cell clones target a single 15-amino acid peptide corresponding to the insulin B-

chain 9-23 sequence (insB:9-23)(15). InsB:9-23 specific T-cells have also been identified in

islets(16) and peripheral blood lymphocytes of T1D patients(17, 18).

In the present study, we hypothesized that exposure to a microbial peptide that resembles

the insulin epitope, insB:9-23, could stimulate or modify the autoimmune response initiating

T1D(19). To address this hypothesis, we analyzed bacterial, viral and fungal genome databases to

identify microbial proteins that have >50% sequence homology to human insB:9-23. Of these, 17

were synthesized and tested for ability to activate insB:9-23 specific T-cells. Herein, we

demonstrate that one of these peptides, a peptide from the gut commensal organism

Parabacteroides distasonis, could activate both human and NOD mouse insB:9-23-specific T-

cells *ex-vivo* to the same extent as the human insulin B:9-23 peptide. This bacterial peptide also

cross-reacts with the immune cells obtained from the mice immunized with the human insB:9-23

peptide. Furthermore, administration of *P. distasonis* bacteria by oral gavage accelerated T1D

progression in NOD mice *in vivo* by stimulating innate immune cells and CD8+ T-cells and

decreasing regulatory T-cells. Analysis of human gut microbiota datasets from the longitudinal

DIABIMMUNE study revealed the presence of this bacterial peptide in children who later

developed autoantibodies to insulin itself. Taken together, these studies demonstrate that an insulin

B:9-23-like epitope in normal gut microbiota may mimic the native insulin peptide and has

potential to play a role in onset or progression of T1D.

2. RESULTS

2.1 A *P. distasonis* insB:9-23-like peptide stimulates NOD mouse hybridomas and human T-cell clones specific to insB:9-23

To determine if any microbe might have DNA encoding proteins with sequences resembling the dominant T-cell epitope involved in the autoimmune response to the insulin B:9-23 peptide linked to T1D (SHLVEALYLVCGERG)(15-18), we used BlastP to search NCBI databases for predicted proteomes of all sequenced viruses (taxid:10239), bacteria (taxid:2), and fungi (taxid:4751). We identified 47 microbial peptides with over eight residues identical to the insulin peptide **(Supplementary Table S1)**. Among these predicted microbial peptides, we selected 17 bacterial and viral peptides that contained the largest number of previously identified residues in the insB:9-23 peptide critical for this interaction(18) to be synthesized (**Figure 1A**).

These 17 microbial insB:9-23-like peptides, as well as a negative control (irrelevant/scramble) and a positive control (insB:9-23), were then tested for their ability to stimulate an insB:11-23-specific human T-cell clone isolated from a T1D patient(18) (**Figure 1B**). In addition, we included a second positive control, a variant of the insB:9-23 peptide in which R at position 22 is substituted by E (insB:9-23R^{22E}) and which has been shown to be even more potent than native insB:9-23 in stimulating insB:9-23 hybridomas from NOD mice(20) and human T-cell clones(18). As expected, in the human T-cell assay, both positive controls stimulated interferon-gamma (IFN-γ) secretion with insB:9-23R^{22E} being more active than the wild type sequence. Among the 17 microbial peptides tested, only one could stimulate human T-cells to produce the IFN-γ (**Figure 1B**). This peptide (*hprt4-18*) represents amino acids 4-18 in the N-terminus of the hypoxanthine phosphoribosyltransferase (*hprt*) protein of *Parabacteriodes distasonis* 33B and D13 strains (formerly known as *Bacteroides sp. 2_1_*33B/ *Parabacteroides*

sp. D13)(21). Interestingly, *P. distasonis* 33B and D13 are the only organisms in the NCBI dataset

(as of October, 2021) that possess this insB:9-23 mimic sequence in their genomes. Next, we tested

the same set of microbial insB:9-23-like peptides and controls on NOD IIT-3 T-cell hybridomas,

previously shown to recognize the insB:9-23 epitope(22). In this assay, C3g7 cells that have high

expression of MHC-II I-A^{g7} were used as antigen-presenting cells (APCs) and treated with each

peptide at concentrations from 0.01 to 10μM. Co-culturing of these cells with IIT-3 T-cell

hybridomas stimulated interleukin-2 (IL-2) secretion only in presence of insB:9-23 and the *hprt4-*

18 peptide, and produced similar dose-response curves, with the bacterial peptide being only

slightly less potent **(Figure 1C).** Thus, both human T-cell clones and NOD mouse T-cell

hybridomas specific to insB:9-23 were reactive to *hprt4-18*, raising the possibility that *hprt4-18*

may have the potential to modulate the development of T1D.

2.2 *P. distasonis hprt4-18* peptide stimulates a T-cell response to insB:9-23 in vivo

To further explore the cross-reactivity of the microbial and insulin-derived peptide

sequences, NOD mice were immunized with either insB:9-23 peptide or the *hprt4-18* peptide in

Complete Freund's Adjuvant (CFA). After 7 days, lymphocytes were isolated from the popliteal

lymph nodes and stimulated with either insB:9-23, *hprt4-18* or control peptides. T-cell activation

was assessed using an enzyme-linked immunospot (ELISpot) assay. Consistent with the in-vitro

results above, lymphocytes from mice immunized with the *hprt4-18* peptide exhibited a strong

immune response to both *hprt4-18* peptide and insB:9-23 as measured by IFNγ and IL-2 secretion

(**Figure 1D &1E**). Conversely, lymphocytes from mice immunized with insB:9-23 peptide

showed a strong response to both insB:9-23 peptide and the microbial *hprt4-18* peptide (**Figure**

1F & 1G). Taken together with the assays using the hybridoma T-cell lines, these data indicate

that the microbial peptide *hprt4-18* strongly cross-react with the recognition and signaling

machinery for human insB:9-23 in stimulating an immune response.

2.2 *P. distasonis* colonization accelerates diabetes onset in NOD mice

Based on this cross-reactivity, we hypothesized that *P. distasonis* colonization and

potential exposure to the microbial *hprt4-18* peptide could trigger and/or modify the immune

response and stimulate autoimmunity in NOD mice. To test this hypothesis, both male and female

NOD mice were orally gavaged with a saline suspension of *P. distasonis* (10^8 CFU/mouse/day) or

saline itself daily, starting at 4 weeks of age (i.e., after weaning) and followed for 30 weeks (**Figure**

2A, n=40/group). Two weeks after the last oral gavage, qPCR using fecal DNA samples revealed

significant levels of *P. distasonis* in the feces of the treated mice, whereas neither male nor female

control NOD mice housed under specific pathogen-free (SPF) conditions had *P. distasonis* in their

gut microbiome (**Figure 2B**). At 12 weeks of age, when mice are in the pre-diabetic stage, five

mice from each group were randomly selected and examined histologically for insulitis. In the *P.*

distasonis colonized female NOD mice there was more than 2-fold increase in severe insulitis

scores as compared to controls (**representative images in 2C, Figure 2D**). Consistent with the

increase in insulitis, female NOD mice subjected to *P. distasonis* colonization showed significantly

accelerated onset of T1D with only 19.5% disease-free in the *P. distasonis* colonized group versus

42% disease-free in the control group (**Figure 2E**). This effect was specific to the female mice,

with no differences in insulitis scores or T1D incidence in male NOD mice with *P. distasonis*

colonization (**Supplementary Figure S1A-B)**.

To determine whether *P. distasonis* colonization stimulates an immune response against

the bacterium, we performed Western blot analysis in which *P. distasonis* cellular protein lysates

were resolved on SDS-PAGE, transferred to PVDF membranes, then probed using serum samples

from 12-week-old *P. distasonis* colonized or control NOD mice. *P. distasonis* colonization of the

NOD mice gut stimulated an antibody response to multiple bacterial proteins in both female and

male NOD mice (**Figure 2F, Supplementary Figure S1C**). This response was specific; Western

blots using protein lysates from *Bacteroides fragilis*, another gut microbe that has been shown to

have immunomodulatory effects in NOD mice(23), produced only a weak humoral immune

response in both the saline and *P. distasonis* treated groups **(Supplementary Figure S1D-E)**.

Importantly, serum LPS levels were not significantly different between the groups, indicating that

the *P. distasonis* humoral response was not the result from of general gut barrier dysfunction

stimulated by bacterial colonization **(Supplementary Figure S1F)**.

To determine whether there is a similar enrichment in the humoral immune responses

against *P. distasonis* occurs in humans with T1D, Western blot analysis was performed using

serum samples from 12 T1D patients (median age 17.5, median disease duration 8 years, males)

with age, sex and ethnicity matched controls. As shown in **Supplementary Figure S1G,** there was

a very weak humoral response against *P. distasonis* in healthy subjects, strong reactivity was

observed in seven out of 12 T1D patients. Thus, consistent with our findings in NOD mice, at least

a subset of T1D patients appear to have a humoral immune response to *P. distasonis* proteins.

2.3 Adoptive transfer of splenocytes enhances T1D onset in NOD/SCID mice

To determine if the enhanced development of T1D in the NOD receiving *P. distasonis*

colonized mice was T-cell mediated, we performed an adoptive transfer experiment using

immunodeficient NOD/SCID mice, which lack functional B and T cells, as recipients(24). To this

end, a new cohort of NOD mice was orally gavaged with *P. distasonis* or saline (n=15-20/group)

as described above. At week 15, these mice were sacrificed, and 5×10^6 splenocytes were

transferred from individual NOD diabetes-free donors to 9-week-old NOD/SCID recipients (1:1

ratio, sex-matched) after which the recipients were followed for 10 weeks (**Supplementary Figure**

S2A). Consistent with the effect of *P. distasonis* colonization on spontaneous T1D in NOD mice

(**Figure 2G**), female NOD/SCID mice that received splenocytes from the *P. distasonis* treated

group developed T1D at a higher rate than those receiving splenocytes from control NOD mice.

Thus, at 10 weeks following transfer, 62.5% of recipients of receiving control splenocytes

remained disease free, and this was decreased to 31% in mice receiving splenocytes from *P.*

distasonis colonized NOD donors (**Figure 2G**). Hence, the immune response stimulated by *P.*

distasonis in female NOD mice was sufficient to accelerate T1D in NOD/SCID mice.

2.5 *P. distasonis* colonization increases CD8+ T-cells and decreases Foxp3+ regulatory T cells

in the splenocytes of female NOD mice.

To determine the effects of *P. distasonis* treatment on NOD mice immune cell composition,

a new cohort of 12-week-of age NOD mice was orally gavaged with either *P. distasonis* or saline,

and the T-cell populations in splenocytes and pancreatic lymph nodes (PLNs) assessed by flow

cytometry. The gating strategy for different T-cell subsets and innate immune cells are described

in **Supplementary Figure S3A & B**. We found that in the spleens of *P. distasonis* colonized NOD

mice there was a significant 31% increase in CD8+ T-cells, leading to a 30 % decrease in the

CD4/CD8 ratio (**Figure 3A & 3B**). PLN cells isolated from the *P. distasonis* colonized mice

showed similar trends in CD4+ T-cells, CD8+ T-cells, and CD4/CD8 ratio, but these did not quite

reach statistical significance (**Supplementary Figure S4A & S4B**). Using FACS analysis, we also

determined the various subsets of cells including naive cells (CD44^{lo} CD62L^{hi}), effector memory

cells (T$_{EM}$, (CD44^{hi} CD62L^{lo}), and, central memory cells (T$_{CM}$, CD62L^{hi} CD44^{hi}) in the

TCRβ+/CD8+/CD4+ cell population in both the spleen and PLN (**Figure 3C-3F**). This revealed a

significant increase in the TCRβ+/CD8+/CD4+ naive T-cell population (**Figure 3D, 3E**) and a

decrease in T_{EM} CD4$^+$ T-cells in splenocytes (**Figure 3C-3F**) but no alterations in TCRβ+/CD4+

naive cells in PLNs of *P. distasonis* colonized mice **(Supplementary Figure S4C-S4F)**. There

was also a two-fold increase in the naive cell population in both CD4+ and CD8+ T-cells in spleen

(**Figure 3E & F**) and in the CD8+ naive cell population in PLNs **(Supplementary Figure S4F)**.

There were no differences in the central memory cells and effector memory cell populations in

TCRβ+/CD8+ cells in splenocytes (**Figure 3D & 3F**) or in TCRβ+/CD8+/CD4+ cells in PLNs

(Supplementary Figure S4D-S4F).

T-regulatory cells (Treg) play a key role in modulating T1D autoimmunity by suppressing

self-reactive T-cell proliferation(25). This is modulated by Forkhead box protein P3 (Foxp3)

expression and interleukin 10 (IL-10) secretion. It has previously been shown that CD4+ CD25+

CD44+ Treg cells positively correlate with Foxp3 expression and production of IL-10(26). We

found a 15 % decrease of CD4+CD25+CD44+ Treg cells in both splenocytes (**Figure 3G & 3H**)

and PLN cells **(Supplementary Figure S4G & S4H**) and a 1.6-fold increase in the total CD4+

CD25+ PLN cell population **(Supplementary Figure S4H)** of *P. distasonis* colonized mice. While

there were no differences in other cell subsets **(Figure 3H).** In the CD4+ T-cell population, the

percentage of Foxp3+ cells were significantly decreased in the splenocytes and PLNs of *P.*

distasonis colonized NOD mice by 21% and 17.8%, respectively (**Figures 4A, 4B & S5A,**

representative images). There was no difference in the pancreas and spleen weight or in the total

number of the spleen and PLN cells between *P. distasonis* colonized mice and the control mice

(**Supplementary Figure 5B).** Thus, *P. distasonis* colonization decreases anti-inflammatory CD4+

CD25+ CD44+ and Foxp3+ Tregs and increase inflammatory CD8+ T-cells in splenocytes and

PLNs, which could contribute to the enhancement of T1D in female NOD mice.

2.6 *P. distasonis* colonization in female NOD mice increases dendritic cells and macrophages

in the splenocytes

To further explore the mechanisms that lead to an increase in CD8+ T-cells in *P. distasonis*

colonized NOD female mice, we assessed the number of the APCs in spleens and PLNs. CD11c+

CD11b+ dendritic cells and F4/80+ macrophages play an essential role in accelerating diabetes in

NOD mice[27, 28]. FACS analysis revealed a 1.5-fold increase in CD11b+ CD11c+ dendritic cells

and a 1.6-fold increase in F4/80+ macrophages in the splenocytes of *P. distasonis* colonized mice

(**Figure 4C-F**), with no significant differences in dendritic cells in PLNs (**Supplementary Figure**

S5D-S5G). Moreover, CD11c+ CD11b- dendritic cells[29] increased 1.8-fold in the spleen of *P.*

distasonis colonized mice (**Figure 4D**). There was also a 1.3-fold increase in circulatory

macrophages ($SSC^{lo}Ly6C^{hi}$ cells), while there was no change in the number of residential

macrophages ($SSC^{lo}Ly6C^{lo}$) in this population (**Figure 4G**). However, there was 7-fold increase

in F4/80+ macrophage population in PLN in *P. distasonis* colonized mice (**Supplementary Figure**

S5G). These findings are consistent with previous studies demonstrating a role of CD11c+

CD11b+ dendritic cells and F4/80+ macrophages in accelerating diabetes in NOD mice[27, 28]

and demonstrated that *P. distasonis* colonization can stimulate both innate and adaptive immune

responses.

2.7 Critical timing of *P. distasonis hprt4-18* exposure in gut microbiome samples of children

developing T1D in early life.

To investigate the potential role for the *P. distasonis* insB:9-23 mimetic peptide in human T1D, we reanalyzed human gut microbiome data from the DIABIMMUNE study[57] using shotgun metagenomic sequencing performed on stool samples collected monthly from infants 0-3 years of age who were genetically predisposed to T1D living in Finland, Russia and Estonia and correlated this with development of islet autoantibodies, i.e., seroconversion(30). Using the metagenomic data of this study, we searched for the specific DNA sequences encoding the *P. distasonis hprt4-18* peptide. We found that in the children who remained negative for autoantibody markers for T1D, sequences encoding *P. distasonis hprt4-18* were detectable in 20-40% patients with no significant change between ages 0 to 3. By comparison, in children who later became autoantibody positive, *hprt4-18* encoding sequences were absent in the first year of life, began to appear by age 2 and then were found in 60% of the children by age 3 (**Figure 5A**).

The exact frequency and timing of appearance of the *hprt4-18* sequence in the microbiome varied by country. Thus, in subjects from Estonia (14 of 74 infants) and Russia (4 of 74 infants), for those who became seropositive, we did not find the *hprt4-18* sequence in their gut microbiota during the first year of life, but was found in 100% of subjects who developed autoantibodies by the age of 2 or 3 (**Figure 5B**). In contrast, this sequence was found in 15-40% of the seronegative infants throughout this time period. Interestingly, while children in Finland who were seronegative has similar prevalence of the microbial sequence as those in Estonia and Russia, infants who became seropositive in Finland (16 of 74 infants) had no detectable *hprt4-18* sequences during the first three years of life (**Figure 5B**). These results suggest that a delayed exposure to the *P. distasonis hprt4-18* peptide, i.e., after the first year of life, may be able to trigger or potentiate an autoimmune response that leads to development of T1D in genetically susceptible individuals.

284 This is also consistent with our findings in NOD mice where colonization of the female NOD mice

285 with *P. distasonis* after weaning accelerated diabetes development.

286 **3. DISCUSSION**

287  Despite major increases in our understanding of the role of autoimmunity in pathogenesis

288 of T1D, the triggering events which lead to disease development remain poorly understood. It is

289 well known that an important component of the early autoimmune response in individuals who

290 ultimately develop T1D is the development of autoantibodies and T-cell reactivity to islet proteins,

291 especially insulin. While there are multiple potential epitopes, the dominant sequence within the

292 insulin molecule to which reactivity occurs is a sequence in the B-chain involving amino acids 9-

293 23(14). This B:9-23 peptide (SHLVEALYLVCGERG) can bind to DQ8 molecules utilizing three

294 different registers, with the amino acid V at position 12, E at position 13 or A at position 14(31).

295 It has been shown that a R→E substitution at position 22 of the B:9-23 peptide creates an even

296 more potent agonist for activating B:9-23 T cell clones, implicating the A at position 14 as the p1

297 anchor(18).

298  In this study, we have investigated the potential role of molecular mimicry as a link

299 between microbial flora and this component of T1D development. Molecular mimicry mechanisms

300 are based on the degeneracy of T-cell recognition(32, 33) and can be either pathogenic(34) or

301 protective(35). While molecular mimicry has long been postulated as a potential factor in

302 autoimmune diseases, including T1D (36, 37), progress in this area has been limited due to a lack

303 of identification of microbial sequences which might trigger this response(38). In the current study,

304 taking advantage of the growing genome databases for microbes in the environment, we have

305 identified 47 microbial peptides with high sequence homology to insB:9-23. We demonstrate that

of these, a peptide with the sequence of *hprt4-18* from *P. distasonis,* can be recognized by and

activate both murine and human T-cells known to respond to insB:9-23 and can cross-stimulate an

immune response to the insB:9-23 peptide in mice. Moreover, exposure of NOD mice to *P.*

distasonis in the gut microbiome during early life results in increased insulitis and accelerates

diabetes onset. This pathogenic effect is mediated by CD8+ T splenocytes, as evidenced by the

ability to adoptively transfer this increase in diabetes risk to female NOD/SCID mice.

Microbiota and gut microbiota in particular have been shown to play a role in modulating

T1D onset in NOD mice(7, 39-41), but previous reports have identified primarily protective

effects, and most through somewhat non-specific mechanisms. For example, NOD mice reared in

germ-free environments have higher rates of development of diabetes than those raised in

conventional facilities(40). In terms of protective effects of some specific bacterial species, the

presence of segmented filamentous bacteria in the gut correlates with diabetes protection in NOD

female mice, which normally have a high incidence of disease(42). Likewise, female NOD mice

colonized at 3-10 weeks of age with *Akkermensia muciniphila* show delayed diabetes

development(43), and oral administration of heat-killed *B. fragilis* has been shown to suppress

autoimmunity in NOD mice when administered under conditions which induce increased gut

permeability(23). Since literally hundreds of treatments have been shown to decrease development

of diabetes in NOD mice (44, 45), however, the specificity of these effects has been

questioned(46). By contrast, in the present study we show that exposure to *P. distasonis* can

accelerate disease development in NOD mice and that a specific peptide encoded in their genome,

hprt4-18, can serve as a mimic of the major insulin auto-epitope at position B9-23. This peptide

and the insulin peptide can also stimulate a bidirectional cross-reactive immune response, proving

its nature as a molecular mimic.

In addition to the immunologic cross-reactivity, our study also provides some insight

into the mechanisms of immune cell regulation by this bacterium. Thus, colonization of NOD mice

with *P. distasonis* increases the CD8+ T-cell population and decreases the CD4/CD8 T-cell ratio.

A reduction in this ratio has been previously shown to be associated with increased T1D in

humans(47). G9C8 CD8+ T-cell clones originally isolated from the islets of young NOD mice

cells are activated by an insB:15-23 peptide, i.e., the C-terminal fragment of the insB:9-23 peptide,

and this accelerates diabetes when adoptively transferred in NOD/SCID mice, even in the absence

of CD4+ T-cells(48, 49). Wong *et al.* showed that cross presentation of insulin by dendritic cells

can stimulate G9C8 T-cell clones, and both insulin and the insB:15-23 peptide can stimulate

proliferation of the naive cell phenotype in NOD mice (48). The *P. distasonis* sequence is identical

to this peptide in 6 of its 9 residues. We also observed that *P. distasonis* colonization in NOD mice

decreased Foxp3+ Treg cells in spleen and PLNs. This effect on Foxp3+ Treg cells is particularly

interesting because Foxp3+ Treg cells are dysregulated in newly diagnosed T1D patients(50) and

increasing Foxp3+ Treg cells in NOD mice can delay the onset of diabetes in NOD mice(25, 51).

A plethora of human gut microbiome studies have demonstrated that the composition of

gut microbiota in patients with autoimmune diseases, including multiple sclerosis(52), systemic

lupus erythematosus, anti-phospholipid syndrome(53), Crohn's disease(54-56), ulcerative

colitis(57), inflammatory bowel diseases(58, 59), and celiac disease(60) are significantly different

from those in healthy controls. Although the methodologies and conclusions differ in studies of

microbiota from subjects with or at-risk for T1D(8, 30, 61), most show that the diversity of gut

microbiota is decreased in T1D patients with increased prevalence of *Bacteroidetes* species. This

is also associated with an altered serum metabolomic profile compared to healthy controls(7).

While most of these studies have not been able to define any mechanism by which this might relate

to disease development, data in the DIABIMMUNE study[57] has suggested that the higher T1D

rates in Finnish Karelia and Estonia compared to Russian Karelia may in part be related to an

action of different lipopolysaccharides (LPS) present in the gut microbiota on the immune

response.

The Environmental Determinants of Diabetes in the Young (TEDDY) study, which has

collected over 12,000 fecal samples from 903 children at risk for T1D in four countries, has

identified *Parabacteroides* as the genus most significantly associated with T1D(62). This is

consistent with our findings that a *P. distasonis* peptide can induce T-cells and antibodies that

cross-react with the major epitope in insulin involved in the autoimmune response. Reanalysis of

the DIABIMMUNE metagenomic data show that 100% of the children becoming seropositive for

autoantibodies related to T1D in Russia and Estonia have the DNA encoding the *P. distasonis*

hprt4-18 peptide in their gut microbiota in first 2-3 years of life compared to less than 40% of

children remaining seronegative. Interestingly, most of these seroconverters were negative for this

microbial sequence in the first year of life, suggesting the timing of exposure to this microbial

antigen may be important in disease development. The TEDDY study showed that human gut

microbiota development can be divided into three phases: a developmental phase (months 3-14),

a transitional phase (months 15-30), and a stable phase (months 31-46)(62). Our data suggest that

exposure to *P. distasonis* in the transitional phase may be critical in T1D pathogenesis. Further

studies will be needed to directly investigate the link between human T1D and *P. distasonis*, but

the cross-reactivity of the *P. distasonis hprt*4-18 peptide with insB:9-23 activated human T cell

clones represents a clear potential mechanism.

While we have focused only on insulin and the B:9-23 peptide as an antigenic determinant

of disease, there are clearly other epitopes recognized by both circulating and islet-infiltrating T-

cells in the insulin B-chain(63), A-chain(64, 65), or C-peptides(16, 66). The recent discovery that

hybrid insulin peptides (HIPs) may be novel epitopes formed by post-translational modifications

formed in the beta cell(67, 68) also opens the possibility of microbial sequences that may more

closely mimic these hybrid peptides. Defective ribosomal products (Drips) of insulin have also

been shown to lead to aberrant insulin polypeptides rendering beta cells immunogenic(69). We

must also keep in mind that insulin is not the only antigen to which antibodies or reactive T-cell

lines are formed in T1D. Other islet antigens include GAD, IA-2, and HIPs(70). Given the

enormous number of microbial peptides produced by the gut microbiota, we expect that other

microbial peptides may exist with the potential to mimic these epitopes and/or trigger related

autoantigen reactive T-cells. Indeed, Tai *et al.* identified a microbial peptide mimic produced by

Fusobacteria that can stimulate islet-specific glucose-6-phosphatase (IGRP) specific mouse T-

cells and promote diabetes development in a new Toll-like receptor (TLR) deficient (TLR$^{-/-}$) and

MyD88$^{-/-}$ NY8.3 NOD mouse model(71).

In summary, our data define a novel molecular mimicry mechanism in which a specific

sequence in a normal commensal gut microbe can mimic a sequence in insulin B-chain and trigger

or modify the immune response involved in development of T1D. This finding may provide a new

target for treat and a window of opportunity to prevent or delay T1D development. These data also

have implications for other diseases with an autoimmune component. Today, we have databases

with enormous amounts of microbial and microbiome sequence data, which can be leveraged to

address the role of molecular mimicry in the autoimmunity, not only of T1D, but also lupus

erythematosis(72), inflammatory cardiomyopathy(73), and multiple sclerosis(74). Our findings

demonstrate a new link to gut antigens and autoimmune diseases with the potential to ultimately

provide new tools, including vaccines, antibiotics, or probiotics for the prevention and treatment

of autoimmune diseases.

4. MATERIAL AND METHODS

4.1 Bioinformatics

A bioinformatics search was performed using NCBI BLASTp for the presence of the microbial peptide sequences with homology to human insB:9-23 sequence as query against viral (NCBI taxonomic ID: 10239), bacterial (NCBI taxonomic ID: 2), and fungal (NCBI taxonomic ID: 10239) proteomes. The whole peptide sequence of each significant hit was compared with insB:9-23 using a multiple sequence alignment program (Clustal Omega) to determine the number of identical and conserved residues. This yielded the data in **Supplementary Table S1**. An additional bioinformatics search was performed to determine whether *P. distasonis* peptide is unique. *P. distasonis* peptide (15aa) or *P. distasonis* hypoxanthine phosphoribosyltransferase *(hprt4-18)* protein sequences were used as queries and searched against BLASTp using non-redundant protein sequences.

4.2 Bacterial culture

Parabacteroides distasonis D13 strain was purchased from Dr. Emma Allen-Vercoe's laboratory at the University of Guelph. Dr. Allen-Vercoe's group isolated this bacterium from the colon of an ulcerative colitis patient. *P. distasonis* D13 strain and *Bacteroides fragilis* (ATCC® 25285) were cultured in anaerobic culture broth (Tryptic Soy Broth supplemented with 5 µg/ml Hemin (BD Biosciences) and 1µg/ml Vitamin K1 (VWR) at 37°C in an anaerobic chamber (Coy Laboratory Product).

4.3 Animals

NOD/ShiLtJ (NOD) and NOD. Cg-*Pcrkdc scid*/J (NOD/SCID) mice were purchased from Jackson Laboratory. Mice were maintained and bred in the Boston College Animal Care Facility.

The mice were maintained under specific pathogen-free conditions in a 12-hrs dark/light cycle

with autoclaved food, water, and bedding. All experiments complied with regulations and ethics

guidelines of the National Institute of Health and were approved by the IACUC of Boston College

(Protocol No.#B2019-003 and 2019-004). For colonization, *P. distasonis* bacteria were re-

suspended in saline at a concentration of 10^9 CFU/mL and oral gavaged using 22ga plastic feeding

tube (Instech Laboratories) (100µl/mouse, 10^8 CFU/mouse). 3-week-old NOD mice were orally

gavaged with *P. distasonis* D13 daily for four weeks right after weaning for 4 weeks. Control

groups were gavaged with sterile saline.

To determine colonization, the DNA was extracted from 100 mg mouse fecal samples (9-

week-old, 2 weeks after final oral gavage treatment) using QIAamp PowerFecal Pro DNA kit

(Qiagen) following the manufacturer's instruction. qPCR was conducted using QuantStudio 3

Real-Time PCR System (Applied Biosystems) with Power SYBR Green Master Mix (Applied

Biosystems) following manufacturer's instruction. The primers were described previously and

listed in Supplementary **Table S2**(75, 76). The relevant abundance of *P. distasonis* was determined

through being normalized with *Eubacteria.*

4.4 Human T-Cell Clone Stimulation.

Twenty 15-mer peptides **(Figure 1A)** used in human and mouse T-cell stimulation

experiments were chemically synthesized by Genscript (TFA removal, >85% purity). The peptides

were dissolved in DMSO at 20 mg/mL (~12 mM). Human T-cell clone stimulation assays were

performed as previously described(77). Briefly, insB:11-23-specific T-cell clones were stimulated

in 96-well round- bottom plates with irrelevant, specific, or microbial mimotope peptides in the

presence of irradiated DQ8cis- or DQ8trans-expressing HEK293 cells as APCs. 50 µl of

supernatants from cultures of T cell clones were collected after 48 hrs of stimulation and added to

each well of 96-well round-bottom plates precoated with IFN-γ (clone MD-1) capturing antibodies

(BioLegend). After overnight incubation, bound cytokines were detected by biotinylated anti-IFN-

γ (clone 4s.B3) and quantified using a Victor2 D time-resolved fluorometer (Perkin Elmer). The

% activity was calculated by dividing the SI of mutated peptides by the SI of the wild-type peptide.

Experiments were performed in the presence of 1 µg/ml of anti-CD28 antibodies. Unless otherwise

stated, peptide concentrations were 2.5 µM.

452 4.5 NOD mice T-cell stimulation and antigen presentation assay

This experiment was performed as described previously(22). Briefly, the peptides were

ordered from Genscript (TFA removal, >85% purity) and dissolved in a base buffer consisting of

50 mM NaCl, 10 mM HEPES, pH 6.8 with 200 µM TCEP. *P. distasonis hprt4-18* peptide -

(**RILVELLYLVCSEYL)** was dissolved in 5% DMSO (the final DMSO concentration is less than

1%). The C3g7 cell line, which expresses an abundance of MHC-II I-A^{g7}, was used as APCs and

was treated with 1/2log dilutions of peptide (starting from 10 µM). These cells were then cultured

with the IIT-3 T-cell hybridomas for 18 hrs. Culture supernatants were assayed for IL-2 by

incubation with the CTLL-2 cell line. CTLL-2 cells are responsive to IL-2 and only actively divide

in the presence of IL-2. Next, CTLL cell proliferation was determined by 3H uptake (CPM) using

a scintillation counter.

463 4.6 Immunization of the NOD mice

The experiments were performed as described previously(78). Briefly, 13 weeks old male

NOD mice (n=3 mice per group) were immunized in the footpad with the either *hprt4-18* peptide

or insB:9-23 peptide (10 nmol/mouse). After 7 days, the draining (popliteal) lymph nodes were

removed and pooled for examination by ELISpot. In the ELISpot assay, node cells were recalled

with the various peptides to elicit either an IL-2 or IFN-γ response. Spots were analyzed by

Immunospot 5.0 (C.T.L.)

4.7 Reanalysis of the published metagenomics data

DIABIMMUNE data was downloaded with the help of Dr. Alex Kostic. The reads were

assembled using SPAdes as the reads were not ideal for peptide search(30) and are now available

at https://github.com/ablab/spades. These assembled contigs were used for a tBLASTn search to

identify "RILVELLYLVCSEYL" encoding sequences in these samples. The data was classified

based on the country and collection time of the stool samples, e.g., 0-1, 1-2, and 2-3 years.

4.8 Splenocyte transplantation

Splenocytes were isolated from 15 weeks donor NOD mice by filtering through a sterile

70 µm nylon mesh. The isolated splenocytes were incubated for 3 mins in red blood cell lysis with

ACK (Lonza) followed by washing with media. The isolated splenocytes were intravenously

injected (5×10^7 cells/mouse) through lateral tail vein into recipient NOD/SCID mice (9 weeks old).

Donor NOD mice and recipient NOD/SCID mice were sex-matched. Recipient mice were

monitored for diabetes by checking tail vein blood glucose levels (>250 mg/dl) twice per week.

The experiment terminated 10 weeks after the transplantation or unless the mice were diagnosed

with diabetes. Donor mice were 15-week male or female NOD mice colonized with either *P.*

distasonis or their sex-matched saline control (n=15-19).

4.9 FACS and Flow Cytometry:

We collected spleen and Pancreatic Lymph Nodes (PLNs) cells from saline-treated   and

P. distasonis colonized female NOD mice at 12 weeks age. Single-cell suspension was obtained

by mechanical disruption of spleen and PLNs using an Ammonium-Chloride-Potassium (ACK)

lysis buffer for 5 minutes followed by filtration with a 70-µm filter. For staining of surface

markers, cells were incubated in fluorescently labeled antibodies **(Supplementary Table S4)** for

15 minutes in PBS containing 2% FBS at room temperature. Cells were then washed with 2 mL

of staining buffer and fixed with 1% Paraformaldehyde (PFA). For Foxp3 staining, fix and perm

permeabilization kit (Thermo fisher) was used as per manufacturer instruction. All the flow

cytometry antibodies were obtained from BioLegend, and Flow cytometry was performed using

BD FACSAria III sorter (BD biosciences) at the Boston College Core facility. The data was

analyzed using FlowJo 10.0 software. Gating strategy for the determination of different cells

subsets are described in supplementary **Figure S2.**

4.10 Insulitis score

Pancreata were isolated from *P. distasonis* colonized 12 weeks NOD mice and control mice

(n=5) followed by fixation with 4% paraformaldehyde overnight at 4°C. Pancreata were embedded

in paraffin blocks and were sectioned into 5µm thickness. To determine insulitis scores, pancreatic

sections were stained with hematoxylin-eosin at Harvard University BIDMC Histology Core.

Slides were analyzed using EVOS XL Core microscope with an LPlan PH2 10 × /0.25 and an

LPlan PH2 20×/0.40 objective. The pancreatic insulitis was screened and evaluated as described

previously(79).

4.11 Western Blot analysis

Protein lysate was extracted using CelLytic B Cell Lysis Reagent (Sigma) following the

manufacturer's instruction. Protein concentrations were determined using BCA assay

(ThermoFisher Scientific). 20 µg of total denatured bacterial protein was loaded into each well in

SDS gel. Wet transfer was performed by using PVDF membrane for 1 h at 4°C. The membrane

was then blocked with 5% non-fat milk in PBST. After washing, the membrane was incubated

with mouse serum (1: 2,500) or human serum (1: 10,000) overnight at 4°C followed by washing

three-time wash with PBST and incubation with secondary antibodies, sheep anti-mouse IgG

antibody (1:3000, Millipore) or goat anti-human IgG antibody (1:3000, SantaCruz) for 1 h at room

temperature. The membrane was washed and developed with chemiluminescent substrate ECL

(ThermoFisher Scientific).

4.12 Endotoxin ELISA

12-weeks old NOD mice serum was collected and stored at −80 °C until use. Serum

endotoxin levels were measured using a Pierce Chromogenic Endotoxin Quant Kit (ThermoFisher

Scientific) following the manufacturers' instructions.

4.13. **Human Plasma Samples**

Peripheral blood was collected from living subjects following the provision of written

informed consent (and assent in the case of minors) in accordance with IRB approved protocols

(University of Florida IRB# 201400703) and the Declaration of Helsinki. The patient

demographics outlined in **Supplementary Table 3**.

4.14 Statistics

Data are presented as the mean ± SEM. Survival curves were analyzed by a log-rank

(Mantel-cox) test. Statistical significance was evaluated using unpaired two-tailed Student's *t*-test

for two-group comparison or ANOVA, followed by Dunnet's or Šidák's multiple comparisons

tests whichever was recommended. A *p-value* of less than *0.05* was considered significant (*) at

the confidence interval of 95%.

Acknowledgements

We first want to thank Prof. Emil Unanue and Dr. Anthony Vomund (University of Washington, St. Louis) for testing microbial insulins on T-cell hybridomas, immunization experiments, and their help to interpret the data. Thanks to Jonathan Dreyfuss and Hui Pan (Joslin Bioinformatics Core) for bioinformatics analysis. Thanks to Babak Momeni (Boston College) for sharing his laboratory's anaerobic chamber. We thank Bret Judson and the Boston College Imaging Core for infrastructure and support. We acknowledge our undergraduate students Tu Tran, Ruixu Huang, David Kim, Scott Hsu, Kaan Sevgi and Maximilian Figura and Typhania Zanou for their help with the microscopy work and the animal experiments. Thanks to Nancy McGilloway and Todd Gaines for their support in the Boston College Animal Care Facility. We thank Alex Kostic, Tao Xu, Thomas Serwold, and Shio Kobayashi (Joslin Diabetes Center) for valuable Discussion.

Funding: This work was supported by (i) National Institutes of Health Grants 1K01DK117967-01 (to E.A.), R01DK031036 and R01121967 (to C.R.K.) and NIH P01 AI042288 (to MAA) and (ii) The G. Harold and Leila Y. Mathers Charitable Foundation Research Grant MF-1905-00311 (to E.A).

Author contributions: K.G assisted with NOD mice experiments, NOD mice colonization followed by FACS staining, data analysis and writing the manuscript. Q.H assisted with NOD mice experiments; NOD mice oral gavage, splenocytes transfer, qPCR, and Western blot analysis using mice serum. I.C and W.K assisted with the experiments using NOD mice T-cell clones. C.B and A.R assisted with oral gavage experiments and following the diabetes onset in NOD mice. P.A assisted with FACS facilities and instrument handling. CRK assisted with research design and data

analysis. All authors assisted with the analysis of the data. E.A completed bioinformatics analysis

for the mimic identification, supervised the project, analyzed the data and wrote the paper.

Competing Interests: The authors do not have any conflict of interest related to this study.

Data and materials availability: Because the DIABIMMUE study metagenomics sequencing

reads were not ideal for a peptide search, we first assembled the reads using SPAdes (11), and they

are available at https://github.com/ablab/spades.

REFERENCES:

1.   Pugliese A (2017) Autoreactive T cells in type 1 diabetes. *J Clin Invest* 127(8):2881-2891.
2.   Robertson CC*, et al.* (2021) Fine-mapping, trans-ancestral and genomic analyses identify causal variants, cells, genes and drug targets for type 1 diabetes. *Nat Genet* 53(7):962-971.
3.   Mayer-Davis EJ*, et al.* (2017) Incidence Trends of Type 1 and Type 2 Diabetes among Youths, 2002-2012. *N Engl J Med* 376(15):1419-1429.
4.   Redondo MJ, Jeffrey J, Fain PR, Eisenbarth GS, & Orban T (2008) Concordance for islet autoimmunity among monozygotic twins. *N Engl J Med* 359(26):2849-2850.
5.   Kondrashova A*, et al.* (2005) A six-fold gradient in the incidence of type 1 diabetes at the eastern border of Finland. *Ann Med* 37(1):67-72.
6.   Coppieters KT, Boettler T, & von Herrath M (2012) Virus infections in type 1 diabetes. *Cold Spring Harb Perspect Med* 2(1):a007682.
7.   Dedrick S*, et al.* (2020) The Role of Gut Microbiota and Environmental Factors in Type 1 Diabetes Pathogenesis. *Front Endocrinol (Lausanne)* 11:78.
8.   Kostic AD*, et al.* (2015) The dynamics of the human infant gut microbiome in development and in progression toward type 1 diabetes. *Cell Host Microbe* 17(2):260-273.
9.   Boljat A*, et al.* (2017) Environmental Risk Factors for Type 1 Diabetes Mellitus Development. *Exp Clin Endocrinol Diabetes* 125(8):563-570.
10.   Ziegler AG*, et al.* (2013) Seroconversion to multiple islet autoantibodies and risk of progression to diabetes in children. *JAMA* 309(23):2473-2479.
11.   Atkinson MA, Eisenbarth GS, & Michels AW (2014) Type 1 diabetes. *Lancet* 383(9911):69-82.
12.   Zhang L, Nakayama M, & Eisenbarth GS (2008) Insulin as an autoantigen in NOD/human diabetes. *Curr Opin Immunol* 20(1):111-118.
13.   Palmer JP, Asplin CM, Clemons P, & al. e (1983) Insulin antibodies in insulin-dependent diabetics before insulin treatment. *Science* 222:1337.
14.   Harrison LC (2021) The dark side of insulin: A primary autoantigen and instrument of self-destruction in type 1 diabetes. *Mol Metab*:101288.
15.   Steck AK*, et al.* (2011) Age of islet autoantibody appearance and mean levels of insulin, but not GAD or IA-2 autoantibodies, predict age of diagnosis of type 1 diabetes: diabetes autoimmunity study in the young. *Diabetes Care* 34(6):1397-1399.
16.   Michels AW*, et al.* (2017) Islet-Derived CD4 T Cells Targeting Proinsulin in Human Autoimmune Diabetes. *Diabetes* 66(3):722-734.
17.   Nakayama M*, et al.* (2015) Regulatory vs. inflammatory cytokine T-cell responses to mutated insulin peptides in healthy and type 1 diabetic subjects. *Proc Natl Acad Sci U S A* 112(14):4429-4434.
18.   Yang J*, et al.* (2014) Autoreactive T cells specific for insulin B:11-23 recognize a low-affinity peptide register in human subjects with autoimmune diabetes. *Proc Natl Acad Sci U S A* 111(41):14840-14845.
19.   Cusick MF, Libbey JE, & Fujinami RS (2012) Molecular mimicry as a mechanism of autoimmune disease. *Clin Rev Allergy Immunol* 42(1):102-111.
20.   Stadinski BD*, et al.* (2010) Diabetogenic T cells recognize insulin bound to IAg7 in an unexpected, weakly binding register. *Proc Natl Acad Sci U S A* 107(24):10978-10983.
21.   Sakamoto M & Benno Y (2006) Reclassification of Bacteroides distasonis, Bacteroides goldsteinii and Bacteroides merdae as Parabacteroides distasonis gen. nov., comb. nov.,

Parabacteroides goldsteinii comb. nov. and Parabacteroides merdae comb. nov. *Int J Syst Evol Microbiol* 56(Pt 7):1599-1605.

22.   Wan X*, et al.* (2018) Pancreatic islets communicate with lymphoid tissues via exocytosis of insulin peptides. *Nature* 560(7716):107-111.

23.   Sofi MH*, et al.* (2019) Polysaccharide A-Dependent Opposing Effects of Mucosal and Systemic Exposures to Human Gut Commensal Bacteroides fragilis in Type 1 Diabetes. *Diabetes* 68(10):1975-1989.

24.   Fuchtenbusch M, Larger E, Thebault K, & Boitard C (2005) Transfer of diabetes from prediabetic NOD mice to NOD-SCID/SCID mice: association with pancreatic insulin content. *Horm Metab Res* 37(2):63-67.

25.   Tang Q*, et al.* (2004) In vitro-expanded antigen-specific regulatory T cells suppress autoimmune diabetes. *J Exp Med* 199(11):1455-1465.

26.   Liu T, Soong L, Liu G, Konig R, & Chopra AK (2009) CD44 expression positively correlates with Foxp3 expression and suppressive function of CD4+ Treg cells. *Biol Direct* 4:40.

27.   Saxena V, Ondr JK, Magnusen AF, Munn DH, & Katz JD (2007) The countervailing actions of myeloid and plasmacytoid dendritic cells control autoimmune diabetes in the nonobese diabetic mouse. *J Immunol* 179(8):5041-5053.

28.   Carrero JA*, et al.* (2017) Resident macrophages of pancreatic islets have a seminal role in the initiation of autoimmune diabetes of NOD mice. *Proc Natl Acad Sci U S A* 114(48):E10418-E10427.

29.   Klementowicz JE*, et al.* (2017) Cutting Edge: Origins, Recruitment, and Regulation of CD11c(+) Cells in Inflamed Islets of Autoimmune Diabetes Mice. *J Immunol* 199(1):27-32.

30.   Vatanen T*, et al.* (2018) The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature* 562(7728):589-594.

31.   Wang Y*, et al.* (2018) C-terminal modification of the insulin B:11-23 peptide creates superagonists in mouse and human type 1 diabetes. *Proc Natl Acad Sci U S A* 115(1):162-167.

32.   Mason D (1998) A very high level of crossreactivity is an essential feature of the T-cell receptor. *Immunol Today* 19(9):395-404.

33.   Calis JJ, de Boer RJ, & Kesmir C (2012) Degenerate T-cell recognition of peptides on MHC molecules creates large holes in the T-cell repertoire. *PLoS Comput Biol* 8(3):e1002412.

34.   Wucherpfennig KW & Strominger JL (1995) Molecular mimicry in T cell-mediated autoimmunity: viral peptides activate human T cell clones specific for myelin basic protein. *Cell* 80(5):695-705.

35.   Mana P*, et al.* (2004) Tolerance induction by molecular mimicry: prevention and suppression of experimental autoimmune encephalomyelitis with the milk protein butyrophilin. *Int Immunol* 16(3):489-499.

36.   Tian J, Lehmann PV, & Kaufman DL (1994) T cell cross-reactivity between coxsackievirus and glutamate decarboxylase is associated with a murine diabetes susceptibility allele. *J Exp Med* 180(5):1979-1984.

37.   Atkinson MA*, et al.* (1994) Cellular immunity to a determinant common to glutamate decarboxylase and coxsackie virus in insulin-dependent diabetes. *J Clin Invest* 94(5):2125-2129.

38.   Honeyman MC, Stone NL, Falk BA, Nepom G, & Harrison LC (2010) Evidence for
molecular mimicry between human T cell epitopes in rotavirus and pancreatic islet
autoantigens. *J Immunol* 184(4):2204-2210.
39.   Wen L*, et al.* (2008) Innate immunity and intestinal microbiota in the development of Type
1 diabetes. *Nature* 455(7216):1109-1113.
40.   Markle JG*, et al.* (2013) Sex differences in the gut microbiome drive hormone-dependent
regulation of autoimmunity. *Science* 339(6123):1084-1088.
41.   Sun J*, et al.* (2015) Pancreatic beta-Cells Limit Autoimmune Diabetes via an
Immunoregulatory Antimicrobial Peptide Expressed under the Influence of the Gut
Microbiota. *Immunity* 43(2):304-317.
42.   Kriegel MA*, et al.* (2011) Naturally transmitted segmented filamentous bacteria segregate
with diabetes protection in nonobese diabetic mice. *Proc Natl Acad Sci U S A*
108(28):11548-11553.
43.   Hanninen A*, et al.* (2018) Akkermansia muciniphila induces gut microbiota remodelling
and controls islet autoimmunity in NOD mice. *Gut* 67(8):1445-1453.
44.   Atkinson MA, Roep BO, Posgai A, Wheeler DCS, & Peakman M (2019) The challenge of
modulating beta-cell autoimmunity in type 1 diabetes. *Lancet Diabetes Endocrinol*
7(1):52-64.
45.   Roep BO, Atkinson M, & von Herrath M (2004) Satisfaction (not) guaranteed: re-
evaluating the use of animal models of type 1 diabetes. *Nat Rev Immunol* 4(12):989-997.
46.   Skyler JS (2018) Hope vs hype: where are we in type 1 diabetes? *Diabetologia* 61(3):509-
516.
47.   Al-Sakkaf L*, et al.* (1989) Persistent reduction of CD4/CD8 lymphocyte ratio and cell
activation before the onset of type 1 (insulin-dependent) diabetes. *Diabetologia* 32(5):322-
325.
48.   Wong FS*, et al.* (2009) Activation of insulin-reactive CD8 T-cells for development of
autoimmune diabetes. *Diabetes* 58(5):1156-1164.
49.   Wong FS*, et al.* (1999) Identification of an MHC class I-restricted autoantigen in type 1
diabetes by screening an organ-specific cDNA library. *Nat Med* 5(9):1026-1031.
50.   Viisanen T*, et al.* (2019) FOXP3+ Regulatory T Cell Compartment Is Altered in Children
With Newly Diagnosed Type 1 Diabetes but Not in Autoantibody-Positive at-Risk
Children. *Front Immunol* 10:19.
51.   Grinberg-Bleyer Y*, et al.* (2010) IL-2 reverses established type 1 diabetes in NOD mice by
a local effect on pancreatic regulatory T cells. *J Exp Med* 207(9):1871-1878.
52.   Maruvada P, Leone V, Kaplan LM, & Chang EB (2017) The Human Microbiome and
Obesity: Moving beyond Associations. *Cell Host Microbe* 22(5):589-599.
53.   Rinaldi M, Perricone R, Blank M, Perricone C, & Shoenfeld Y (2013) Anti-Saccharomyces
cerevisiae autoantibodies in autoimmune diseases: from bread baking to autoimmunity.
Clin Rev Allergy Immunol 45(2):152-161.
54.   Gevers D*, et al.* (2014) The treatment-naive microbiome in new-onset Crohn's disease. *Cell*
Host Microbe 15(3):382-392.
55.   Lamps LW*, et al.* (2003) Pathogenic Yersinia DNA is detected in bowel and mesenteric
lymph nodes from patients with Crohn's disease. *Am J Surg Pathol* 27(2):220-227.
56.   Chassaing B*, et al.* (2011) Crohn disease--associated adherent-invasive E. coli bacteria
target mouse and human Peyer's patches via long polar fimbriae. *J Clin Invest* 121(3):966-
975.

57. Carding S, Verbeke K, Vipond DT, Corfe BM, & Owen LJ (2015) Dysbiosis of the gut
microbiota in disease. *Microb Ecol Health Dis* 26:26191.
58. Frank DN*, et al.* (2007) Molecular-phylogenetic characterization of microbial community
imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci U S A*
104(34):13780-13785.
59. Navaneethan U, Venkatesh PG, & Shen B (2010) Clostridium difficile infection and
inflammatory bowel disease: understanding the evolving relationship. *World J*
Gastroenterol 16(39):4892-4904.
60. Quagliariello A*, et al.* (2016) Effect of Bifidobacterium breve on the Intestinal Microbiota
of Coeliac Children on a Gluten Free Diet: A Pilot Study. *Nutrients* 8(10).
61. Vatanen T*, et al.* (2016) Variation in Microbiome LPS Immunogenicity Contributes to
Autoimmunity in Humans. *Cell* 165(6):1551.
62. Stewart CJ*, et al.* (2018) Temporal development of the gut microbiome in early childhood
from the TEDDY study. *Nature* 562(7728):583-588.
63. Higashide T*, et al.* (2006) T cell epitope mapping study with insulin overlapping peptides
using ELISPOT assay in Japanese children and adolescents with type 1 diabetes. *Pediatr*
Res 59(3):445-450.
64. Kent SC*, et al.* (2005) Expanded T cells from pancreatic lymph nodes of type 1 diabetic
subjects recognize an insulin epitope. *Nature* 435(7039):224-228.
65. Mannering SI*, et al.* (2005) The insulin A-chain epitope recognized by human T cells is
posttranslationally modified. *J Exp Med* 202(9):1191-1197.
66. Pathiraja V*, et al.* (2015) Proinsulin-specific, HLA-DQ8, and HLA-DQ8-transdimer-
restricted CD4+ T cells infiltrate islets in type 1 diabetes. *Diabetes* 64(1):172-182.
67. Mannering SI, Di Carluccio AR, & Elso CM (2019) Neoepitopes: a new take on beta cell
autoimmunity in type 1 diabetes. *Diabetologia* 62(3):351-356.
68. Delong T*, et al.* (2016) Pathogenic CD4 T cells in type 1 diabetes recognize epitopes
formed by peptide fusion. *Science* 351(6274):711-714.
69. Kracht MJ*, et al.* (2017) Autoimmunity against a defective ribosomal insulin gene product
in type 1 diabetes. *Nat Med* 23(4):501-507.
70. Babon JA*, et al.* (2016) Analysis of self-antigen specificity of islet-infiltrating T cells from
human donors with type 1 diabetes. *Nat Med* 22(12):1482-1487.
71. Tai N*, et al.* (2016) Microbial antigen mimics activate diabetogenic CD8 T cells in NOD
mice. *J Exp Med* 213(10):2129-2146.
72. Greiling TM*, et al.* (2018) Commensal orthologs of the human autoantigen Ro60 as triggers
of autoimmunity in lupus. *Sci Transl Med* 10(434).
73. Gil-Cruz C*, et al.* (2019) Microbiota-derived peptide mimics drive lethal inflammatory
cardiomyopathy. *Science* 366(6467):881-886.
74. Harkiolaki M*, et al.* (2009) T cell-mediated autoimmune disease due to low-affinity
crossreactivity to common microbial peptides. *Immunity* 30(3):348-357.
75. Dziarski R, Park SY, Kashyap DR, Dowd SE, & Gupta D (2016) Pglyrp-Regulated Gut
Microflora Prevotella falsenii, Parabacteroides distasonis and Bacteroides eggerthii
Enhance and Alistipes finegoldii Attenuates Colitis in Mice. *PLoS One* 11(1):e0146162.
76. Croswell A, Amir E, Teggatz P, Barman M, & Salzman NH (2009) Prolonged impact of
antibiotics on intestinal microbial ecology and susceptibility to enteric Salmonella
infection. *Infect Immun* 77(7):2741-2753.

77.   Chow IT, *et al.* (2019) Discriminative T cell recognition of cross-reactive islet-antigens is
associated with HLA-DQ8 transdimer-mediated autoimmune diabetes. *Sci Adv*
5(8):eaaw9336.
78.   Mohan JF, *et al.* (2010) Unique autoreactive T cells recognize insulin peptides generated
within the islets of Langerhans in autoimmune diabetes. *Nat Immunol* 11(4):350-354.
79.   Flodstrom-Tullberg M, *et al.* (2003) Target cell expression of suppressor of cytokine
signaling-1 prevents diabetes in the NOD mouse. *Diabetes* 52(11):2696-2700.

FIGURES:

Figure 1. *P. distasonis hprt4-18* peptide mimic can stimulate both human and NOD mice T

cells specific to insB:9-23. **(A)** Sequence alignment of 17 microbial peptides tested in this study.

The amino acids highlighted in red are identical to insB:9-23. **(B)** Treatment of insB:11-23-specific

human T-cell clones with each of selected 17 microbial peptides, where, negative control and

irrelevant peptides were used as the negative controls, ins B9-23 and insB:9-23 R^{22E} were used as

positive controls. ELISpot was used to determine IFN-γ secretion *(P<0.05)* (n=3). **(C)** Dose

response of IIT-3 hybridomas to insB:9-23 (black) or *hpt4-18* peptides to (red) where peptides

were presented to hybridomas as covalently linked to I-A^{g7} expressed on macrophages (C3g7

cells). Data were presented as IL-2 induced proliferation in CTLL-2 cells. **(D-G)** Mice (n=3/group)

were immunized with either *hprt4-18* peptide (**D-E**) or insB:9-23 peptide (**F-G**) after 7 days of

immunization, popliteal lymph nodes were stimulated either by *hprt4-18* peptide, insB:9-23, ConA

(positive control) or scrambled peptide (negative control). Secretion of IL-2 (**E & G**) and IFN-γ

(D & F) were determined by ELISpot. All samples in each panel are biologically independent.

Data were expressed as means ± SEM. **P<0.05, **P<0.01, ****P<0.0001*. Statistical analysis

was performed by one-way ANOVA using Dunnett's multiple comparison test.

Figure 2. *P. distasonis* colonization enhances disease onset in female NOD mice and

T1D patients have a stronger immune response to *P. distasonis* compared to healthy

individuals. (A) Schematic overview of the *P. distasonis* oral gavage experiments

(n=40/group/sex). Blue arrow shows the time-point (week 9) of fecal sample collection for qPCR

experiments for *P. distasonis* colonization and red arrow shows the time-point (week 12) for

pancreata collection for insulitis analysis (n = 5 mice/group/sex). **(B)** Relative abundance of *P.*

distasonis in fecal samples determined by qPCR (week 12, n=10-13/male, n=12-17/female). **(C)**

Representative images of islets insulitis score data where islets were scored as no insulitis, peri-

insulitis, moderate insulitis or having severe infiltration as shown in images. **(D)** Quantification of

insulitis scores obtained from *P. distasonis* colonized and saline gavaged female NOD mice at

week 12 (n=5/group/sex). **(E)** Diabetes incidence in female NOD mice (n=35/group/sex) after

daily oral gavage with either saline or *P. distasonis* for four weeks after weaning *(P<0.05)*. **(F)**

Western blot analysis performed using serum samples from female NOD mice either oral gavaged

with *P. distasonis* or saline (week 12, n=5/group/sex). **(G)** Diabetes incidence of the recipient

NOD/SCID mice after adoptive transfer of 5×10^7 splenocytes/mouse from individual female NOD

mice to female NOD/SCID mice at 6 weeks of age (1:1 ratio, same gender, n=15). All samples in

each panel are biologically independent. Log-rank (Mantel-cox) test was used for survival curves

and adoptive transfer experiment. Data were expressed as mean ± SEM. *P<0.05*, **P<0.01*,

***P<0.01*. Statistical analysis was performed by two-way ANOVA using Sídak multiple

comparison test or two-tailed unpaired Student's *t*-test.

791   **Figure 3: *P. distasonis* colonization increase CD8+ T-cells and Naïve cells phenotype**.

792 Mice were orally gavaged with either *P. distasonis* or saline for 4-weeks after weaning and

793 analyzed at 12 weeks of age. **(A)** Representative image from a single experimental cohort of FACS

794 analyses for splenic CD4+ and CD8+ T-cells from saline and *P. distasonis* gavaged mice. **(B)**

795 CD4+ and CD8+ cells as percent of TCR-β+ immune cell subsets and ratio of splenic CD4+ to

796 CD8+ T-cells. **(C)** CD44$^{lo/hi}$ and CD62L$^{lo/hi}$ T-cells in TCR-β+, CD4+ and CD8+ T-cells. **(D)**

797 Percent of CD44^{hi} CD62L^{lo} (T$_{EM}$)**,** CD44^{hi} CD62L^{hi} (T$_{CM}$)**,** CD44^{lo} CD62L^{hi} (Naive) in TCR-β^+

798 immune cell subsets, **(E)** Percent of CD44^{hi} CD62L^{lo} (T$_{EM}$)**,** CD44^{hi} CD62L^{hi} (T$_{CM}$)**,** CD44^{lo}

799 CD62L^{hi} (Naive) in CD4+ immune T-cell subsets; **(F)** Percent of CD44^{hi} CD62L^{lo} (T$_{EM}$)**,** CD44^{hi}

800 CD62L^{hi} (T$_{CM}$)**,** CD44^{lo} CD62L^{hi} (Naive) in CD8+ immune T-cell subsets **(G)** Representative

801 image of FACS analyses of CD4+, CD25+ Treg population in saline and *P. distasonis* gavaged

802 mice. **(H)** Percent of CD4+ CD25+ cells in CD4+ single cell subsets**,** CD44^{hi} CD62L^{lo} (T$_{EM}$) and

803 CD44^{lo} CD62L^{hi} (Naive) population in percent of CD4+ CD25+ single cell subsets. All samples

804 in each panel are biologically independent. Spleens were obtained from female NOD mice oral

805 gavaged with either *P. distasonis* (n=6) or saline (n=3). Data were expressed as means ± SEM. *,

806 *P<0.05*, **, *P<0.01*, ***, *P<0.001*. Statistical analysis was performed by two-tailed, unpaired

807 Student's t-test.

Figure 4: *P. distasonis* colonization decreases Foxp3+ cells and increases dendritic, and macrophage populations. **(A)** Representative image of FACS analyses of splenic Foxp3+ cells in saline and *P. distasonis* gavaged mice. **(B)** Percent of Foxp3+ cells in CD4+ T-cell subsets. **(C)** Representative image of FACS analyses of dendritic cells (CD11b-CD11c+ and CD11c+ CD11b+) in saline and *P. distasonis* gavaged mice. **(D)** Percent of CD11b+/-CD11c+/- population in splenic single cell subsets **(E)** Representative image of FACS analyses of macrophages

(F4/80+cells) in spleen of saline and *P. distasonis* gavaged mice. **(F)** Percent of F4/80+ cells in

CD11b+ cell subsets **(G)** Percent of F4/80+ cell subsets, where SSC^{lo}Ly6C^{lo} represents residential

macrophage and SSC^{lo}Ly6C^{hi} represents circulatory macrophages. All samples in each panel are

biologically independent. Spleens were obtained from female NOD mice oral gavaged with either

P. distasonis (n=6) or saline (n=3). Data were expressed as means ± SEM. *, *P<0.05*, **, *P<0.01*,

***, *P<0.001*. Statistical analysis was performed by two-tailed, unpaired Student's t-test.

Figure 5. Reanalysis of DIABIMMUNE gut microbiome data revealed the presence

of sequences encoding *P. distasonis hprt4-18* peptide in all children developing autoantibodies

in Estonia after age 1 and Russia after age 2 (A-B) Reanalysis of DIABIMMUNE gut

microbiome data for the presence of *hprt4-18* peptide **(A)** The percentage of subjects having

sequence encoding *hprt4-18* in the gut in all three countries (Finland, Estonia and Russia) and **(B)**

in individual countries at different phases of gut microbiota development (age 0 to 3). Blue

represents subjects that developed autoantibodies (seroconverted) and red represents subjects with

no autoantibodies (non-seroconverted). 16 of 74 subjects in Finland, 14 of 74 of subjects in

Estonia, 4 of 74 subjects in Russia developed autoantibodies during the study.