Cell group analysis reveals changes in upper-layer neurons associated with schizophrenia

Rujia Dai[1,2], Lulu Chen[3], Sihan Liu[1], Yu Chen[1], Yi Jiang[1,4], Jiacheng Dai[1], Qihang Wang[1], Liz Kuney[2], Richard Kopp[2], Guoqiang Yu[3], Yue Wang[3, *], Chao Chen[1, 5, 6, 7 *], Chunyu Liu[1, 2, 8, *]

[1]Center for Medical Genetics & Hunan Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, Changsha, China

[2]Department of Psychiatry, SUNY Upstate Medical University, NY, USA

[3]Department of Electrical and Computer Engineering, Virginia Polytechnic Institute and State University, VA, USA

[4]Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, USA

[5]National Clinical Research Center for Geriatric Disorders, Xiangya Hospital, Central South University, Changsha, China

[6]Hunan Key Laboratory of Animal Models for Human Diseases, Central South University, Changsha, China

[7]Hunan Key Laboratory of Molecular Precision Medicine, Central South University, Changsha, China

[8] School of Psychology, Shaanxi Normal University, Xi'an, China

* Corresponding author Contact:

Chunyu Liu, liuch@upstate.edu

Chao Chen, chenchao@sklmg.edu.cn

Yue Wang, yuewang@vt.edu

Keywords: schizophrenia; tissue deconvolution; gene expression; neural cell type;

prefrontal cortex; eQTL; TWAS

Abstract

Genome-wide association studies (GWAS) of schizophrenia (SCZ) have revealed over

100 risk loci. We investigated whether these SCZ-associated variants regulate gene

expression by cell type. Using a fully unsupervised deconvolution method, we

calculated gene expression by clusters of estimated cell types (cell-groups, CGs). Five

CGs emerged in the dorsolateral prefrontal cortices (DLPFC) of 341 donors with and

without SCZ. By mapping expression quantitative trait loci (eQTL) per CG, we

partitioned the CG-specific heritability of SCZ risk in GWAS. CG-specific expressions

and eQTLs were replicated in another independent deconvoluted bulk tissue data and

single-cell expression data. Further, we characterized CG-specific gene differential

expression and cell proportion changes in SCZ brains. We found upper-layer neurons

in the DLPFC to be associated with SCZ based on enrichment of risk heritability in

eQTLs, disease-related transcriptional signatures and cell type disproportion. Our

study suggests that neurons and related anomalous circuits in upper layers of the

DLPFC may have a major contribution to SCZ risk.

Introduction

Schizophrenia (SCZ) is a complex psychiatric disorder, with approximately 80% of phenotypic variation explained by genetics[1]. In recent years, a surge of genetic markers associated with SCZ has flowed from genome-wide association studies (GWAS)[2]. Nonetheless, interpreting the biological meaning of these GWAS signals remains a challenge. SCZ GWAS loci are located largely within non-coding regions of the genome, pointing to a mechanistic involvement in gene regulation[3]. Expression quantitative trait loci (eQTL) mapping provides an effective means for connecting and interpreting GWAS results with the underlying relevant expression regulation[4]. The PsychENCODE[5], CommonMind[6], Brain Cloud[7], and UK Brain Expression Consortium[8] projects produce large-scale eQTL resources. These data have been integrated with GWAS in the effort to pinpoint risk genes[9-11].

A primary pitfall of current brain eQTL analyses however, is the indifference to cell-type specificity[12]. All existing brain eQTL resources originate from homogenate bulk tissues in the brain, a mixture of many cell types or subtypes. Bulk tissue eQTL mapping offers only averaged effects across the composite cell types. It is therefore confounded by variations in proportion and gene expression. Reports of cell-type specificity of eQTLs from blood[13] provide the evidence that at least some common genetic variants shape gene expression by cell types[14]. However, a neural interpretation of cell-type-specificity of eQTL is not yet available.

To date, the majority of SCZ transcriptomic studies have similarly relied on data derived

from bulk tissue. These include differential analysis and co-expression analyses[15,16].

Such studies bear two major limitations: 1) it is unclear whether genes that are

differentially expressed between individuals with SCZ and controls are attributed to

differences in expression or to changes in cell proportion, and 2) it is unknown whether

changes in SCZ-related expression occur only within one specific cell type or within

multiple cell types.

Researchers investigate cell-type-specific expression using either of two approaches:

1) the sequencing individual or sorted cells using single-cell or single-nuclei

sequencing (scRNA-seq or snRNA-seq), and 2) the computational deconvolution of

bulk tissue transcriptomic data. To date, fewer than 50 human brain samples have

been sequenced and published using scRNA-seq or snRNA-seq[17,18]. For the time-

being, such methods are impractical for most genetic or case-control studies due to

the prohibitory costs of the large sample size needed to achieve adequate statistical

power.[19] Alternatively, the computational approach can deconvolute bulk tissue data

into cell-type-specific data with sufficient accuracy[20]. The computational deconvolution

of bulk tissue data can estimate cell type proportion and cell type-specific expression

in either a supervised or an unsupervised manner. Supervised deconvolution relies on

an accurate reference, for example, cell type-specific expression from the same

species and brain region. However, suitable references are often unavailable. Whether

reference developed from control samples can be used to deconvolute patient data is

another valid concern. In contrast, using unsupervised deconvolution sidesteps this

problem by estimating cell-type-specific expression and cell proportion without prior

information. Cell classification translates to a group of cells algorithmically

differentiated from other groups. Since these groups may not always reflect biologically

homogeneous cell types, we instead use the term cell groups (CGs). The resulting

CGs represent an estimated cluster of cell sub-types extractable from bulk tissue.

Using unsupervised deconvolution methods, like convex analysis of mixture

(CAM),[21,22]   extricates CGs based on genes expression patterns aggregated from all

tested individuals.

Some advanced deconvolution methods, can *infer* cell-type-specific expression per

individual, based upon the methylation[23] or gene expression[24] profile from the donor's

bulk tissue. Again, such supervised methods rely on suitable references. Ideally,

sample-wise and unsupervised deconvolution methods are needed to investigate CG-

specific expressions from data bulk tissue data in a cost-effective manner.

We hypothesized that GWAS variants associated with the risk for SCZ regulate gene

expression within one or more specific brain cell types. To achieve this goal, we

developed sample-wise convex analysis of mixtures (swCAM), an advanced fully

unsupervised deconvolution method[25]. We applied swCAM to RNA-seq data from 341

postmortem brain samples of donors with SCZ and healthy donors to define CGs

(Fig.1). We estimated CG-specific gene expression for each individual along with the

proportion of each individual CG to total CGs. An additional postmortem brain dataset

of 67 samples replicated the swCAM deconvolution results. Further, after generating

CG-specific eQTLs by integrating genotype data and deconvoluted expression, then

we tested whether the CG-specific eQTLs were enriched for SCZ GWAS signals. To

detect SCZ-related transcriptional changes per CG, we performed differential

expression analyses, followed by co-expression network and transcriptome-wide

association analyses of the deconvoluted expression data. We integrated results from

GWAS enrichment, cell proportion, and transcriptomic changes to target SCZ-

associated CGs. The results are accessible on our website

(http://lbpg.upstate.edu/module_search/).

Results

Sample-wise deconvolution of RNA-seq data from bulk brain tissue of donors

with and without SCZ

Five CGs emerged from swCAM deconvolution of bulk tissue data. Their identities

were annotated by enrichment testing of known cell-type marker genes. We calculated

the enrichment of marker genes of major brain cell types[26] (Supplemental table 1) in

the top cell-type differentially expressed genes (ctDEG) of each CG. Top ctDEGs were

identified by differential expression analysis of target CG versus the remainder of CGs

(log2FC>2, False Discovery Rate [FDR] q <0.05, Wilcoxon signed rank test). The five

annotated CGs represented four relatively homogeneous cell types and one mix of

several cell types (p value <0.05, Fisher's exact test). The relatively homogeneous

CGs included two neuronal CGs (CG2 and CG3), one astrocyte CG (CG1), and one

oligodendrocyte CG (CG5, Fig. 2A, Supplemental Fig. 1). CG4 contained a mixture of

astrocytes, endothelial cells, and microglia. The CG-specific expression of the four

relatively homogeneous CGs was correlated to the snRNA-seq data[27] of neurons,

astrocytes and oligodendrocytes from human frontal cortices (averaged rho_{neuron_CG2}

=0.40, rho_{neuron_CG3} =0.31, $rho_{astrocyte\_CG1}$=0.26, $rho_{oligodendrocyte\_CG5}$=0.58, all three p

values < 0.05, Supplemental Fig. 2).

To further differentiate the two neuronal clusters, marker gene-sets of standard

neuronal subtypes were applied. We initially suspected that CG2 and CG3 represented

inhibitory and excitatory neurons; however, we were dissuaded without convincing

differentiation by marker genes of inhibitory and excitatory neurons (Fisher's exact test

p>0.05, Supplemental Fig. 3). We then speculated that CG2 and CG3 may reflect the

origin of specific cortical layers. With only two detected neuronal CGs and six cortical

layers, we took a quantitative approach in allocating gene expression to gradient

cortical layers: 770 *upper-layer (upperL) genes* were expressed most abundantly in

the uppermost layer (layer 1) steadily decreasing by layer with the least expression in

the layer 6, while 875 *deep-layer (deepL) genes* were distributed in the opposite

direction (FDR<0.05, linear regression). *UpperL* and *deepL genes* corresponded to the

CG2 and CG3 clusters respectively (Fisher's exact test, $p_{upperL} < 2.2e-16$, $p_{deepL} = 7.90e$-

10, Fig. 2B).

The expression data of manually isolated neurons from each of the six layers of the

middle temporal cortex[28] was used to validate the identity of neuronal CGs. We

calculated the CG-specific difference for each gene by comparing the expression of

CG2 and CG3 individually to that of the other CGs. Layer-specific differences were

also calculated for each individual gene by comparing the neuronal expression of one

specific layer to that of the other layers. CG-specific difference and layer-specific

differences were then tested for correlation. CG3-specific differences were highly

correlated to layer-specific differences of neurons from layers 1 through 3, while CG3-

specific differences were poorly correlated to layer-specific differences of neurons from

layers 4 through 6. In contrast, correlations for CG2-specific differences ran in the

opposite direction (Supplemental Fig. 4). Based on this, we defined CG2 as

representing deep-layer neurons (deepN) and CG3 as upper-layer neurons (upperN).

At this juncture, the gene expression profiles for the five CGs were now garnered from

the bulk brain tissue of donors with and without SCZ, i.e., upperN, deepN, astrocytes

(ast), oligodendrocytes (oli), and a mixture CG of astrocytes, microglia, and endothelial

cells (mic_ast_endo) (Fig. 2C).

swCAM deconvolution results were replicated on a second dataset of 67 samples

(GEO: GSE12649). We detected five CGs in the replication dataset also

(Supplemental Fig. 5). Through marker gene enrichment tests as described earlier, we

found the five CGs in these data were upperN (p_{neuron}= 3.55e-25, p_{upperL}=8.30e-08),

deepN (p_{neuron}=3.43e-07, p_{deepL}= 0.0001), oli (p=4.16e-39), astrocyte (p=8.04e-11), and

a CG enriched for no specific cell type and a mixture of endothelial cells (p= 0.05). The

upperN, deepN and oli CGs were well re-captured while microglia CG were absent in

the second dataset.

Generation of eQTLs for CGs

eQTL mapping yielded examination of the genetic regulation of CG-specific expression.

We included 72 additional brain samples from individuals with bipolar disorder (BD) to

increase the sample size. Since we regress out the affection status, the inclusion of

BD patients will not interfere with the eQTL mapping results. The same five CGs were

still detected by swCAM deconvolution after adding the BD samples. After removing

known and hidden covariates in the CG-specific transcriptome, we focused on local

eQTLs (cis-eQTLs). assessing for any associations between individual expression of

genes with common variants within ±1 Mb of the gene body region. In total, we

identified 1,878,815 genome-wide significant eQTLs (FDR q < 0.05). After subtracting

shared eQTL across all CGs, we observed more eQTL signals in neuronal CGs (both

of upperN and deepN) than in other glial CGs (Fig. 3A). The number of single

nucleotide polymorphisms of eQTL (eSNPs) of neuronal CGs (n=302,608) was more

than that of eSNPs of oligodendrocyte CGs (n=80,229). No eQTL survived multiple

testing correction in astrocyte CG data. This was consistent with a previous result,

which reported that neurons express more genes and transcripts than do glia[18]. We

annotated the genomic regions for eSNPs with Ensembl annotations using ANNOVAR

software[29]. The majority of eSNPs were located in intergenic and intronic regions within

non-coding RNAs (Fig. 3C).

To examine the replication rate of eQTLs of the three relatively homogeneous CGs

(deepN, upperN, and oli), we compared our results to another published cell-type-

specific eQTL study based on sorted cells[30], which profiled the gene expression and

genotype of neurons (n of sample=42) and oligodendrocytes (n of sample=36). Using

the same eQTL mapping pipeline described earlier, we calculated the eQTLs using

these replication data. From this small sorted-cell dataset, we could identify 644 eQTLs

for neurons and 190 eQTLs for oligodendrocytes (FDR q<0.05). Having calculated π1

for the proportion of significant eQTLs from the sorted cells replicated in our

deconvoluted CGs, we found that upperN (π1=0.56) and deepN (π1=0.55) had greater

reproducibility than oli (π1=0.35). We also compared our results with eQTLs from bulk

brain tissue samples from BrainGVEX. CG-specific eQTLs showed good

reproducibility in the BrainGVEX dataset (π1 >0.97 in four CGs except for

mic_ast_endo by π1= 0.92, Fig. 3B).

Neuronal CGs explain risk heritability of SCZ

To associate brain cell type with SCZ, we used CG-specific eQTLs to assess the SCZ

GWAS risk heritability (Fig. 4). We hypothesized that if a CG is associated with SCZ,

their eQTLs will explain greater GWAS heritability. To test this hypothesis, we applied

stratified linkage disequilibrium (LD) score regression[31] (sLDSC) on GWAS summary

statistics and CG-specific eQTLs to partition risk heritability from summary statistics.

The proportion of risk heritability explained by upperN (h2=19.24%) was the highest

across all CGs and the second by the deepN (h2=19.03%). Combining upperN and

deepN could explain 22.80% of risk heritability. Combining all five CGs explained 23.67%

of risk heritability. However, bulk tissue explained slightly greater risk heritability

(29.78%). To control for the effect of eSNPs in each category, we calculated heritability

enrichment by the ratio measuring the proportion of heritability over the proportion of

eSNPs. After normalizing for this effect, the enrichment ratio per CG (1.32~1.51) was

higher than the enrichment levels found in bulk tissue (enrichment ratio=1.29).

Decreased proportion of upper-layer neuron CGs from donors with SCZ

correlated with genetic risk burden

To find out whether the proportion of each CG was altered in the brains of patients with

SCZ, we estimated the proportion of CGs for each sample using CAM. The proportion

of upperN decreased by 2.76% in patients than in controls, and the mixture CG of

mic_ast_endo increased by 0.8% in patients with SCZ (Wilcoxon signed rank test, p

value<0.05, Fig. 5A). Other CGs did not show significant change. Based on our testing

for genetic enrichment, we hypothesized that the changes in CG proportion could arise

from genetic risk variants. We used two strategies to test this hypothesis. First, we

performed QTL mapping using CG proportions as the phenotype but did not detect any

significant CG proportion-QTL signals. Second, we used polygenic risk score (PRS)

as a measure of genetic risk burden for each individual and tested the correlation

between PRS and CG proportions. We observed weak negative correlations between

SCZ PRS and the upperN proportion (r=-0.1, p value=0.04, Fig. 5B). High PRS is

associated with low upperN proportion.

CG-specific differential gene expression in SCZ

We further analyzed CG-specific differential expression comparing individuals with and

without SCZ using Wilcoxon signed rank test with significance by FDR q <0.05,

(Supplemental Table 2). We found substantial overlap in differentially expressed genes

(DEG) across five CGs with 330 upregulated genes and 344 downregulated genes

shared across five CGs (Supplemental Fig. 6). The numbers of shared DEGs across

all CGs was greater than the number of CG-specific DEGs, except for upregulated

genes in CG of mic_ast_endo having 367 genes. Approximately 33% of the genes

(n=14,865, Gencode v19) were affected in at least one CG in the SCZ cohort. The two

neuronal CGs possessed greater transcriptional changes than the three glial CGs,

(fold change: 0.20 in neuronal CGs versus 0.09 in glial related CGs, p value<2.2e-16,

Kolmogorov–Smirnov test).

We used Gene Ontology (GO) enrichment to annotate the function of DEGs (Table 1).

focusing on the functions of dysregulated genes in neuronal groups. A total of 1,537

downregulated DEGs were detected in the upperN. These were also enriched for

protein folding (FDR q= 3.20e-09) and antigen processing (FDR q= 8.01e-05). The

1,407 downregulated DEGs within deepN were enriched for positive regulation of cell

proliferation (FDR q= 2.24e-06) and trans-synaptic signaling (FDR q=4.10e-06).

Upregulated genes in both the upperN (n=229) and deepN (n=223) GCs were enriched

for carboxylic acid breakdown (FDR q_{deepN}=1.04e-06, FDR q_{upperN} =6.93e-08).

Transcriptome-wide association analysis identified SCZ-associated genes in all

CGs

To determine which genes had CG-specific dysregulated expression driven by

genetics, we performed a TWAS analysis using the CG-specific eQTLs. We identified

51 SCZ risk genes across 43 genetic loci (FDR<0.05, Supplemental Table 3). We

compared our TWAS results to the 193 SCZ risk genes identified in bulk tissue by

TWAS in Gandal et al[16]. Twelve genes in our results had been previously identified in

Gandal's results. Our analysis attributed these 12 genes to all 5 CGs: deepN

(*CYP17A1-AS1, AS3MT, CSPG4P11, EMB*), upperN (*CYP17A1-AS1, AS3MT,*

CSPG4P11, EMB, GATAD2A, NAGA, RP11-73M18.8, TYW5), ast (*RP11-73M18.8,*

CYP17A1-AS1, EMB), oli (*SNX19, AC011330.5, CSPG4P11, EMB*), and

mic_ast_endo (*BTN3A2, TOM1L2, CSPG4P11*). Additionally, 39 CG-specific TWAS

genes are novel to bulk tissue-based TWAS.

Out of the 51 TWAS genes, 17 genes are in deepN and 30 in upperN. Eight genes

existed in both subtypes (*CYP17A1-AS1, AS3MT, PIGB, GOLGA6L9, CSPG4P11,*

CNPPD1, EMB, ENDOG). Twenty-two genes were uniquely identified in upperN

(*DIABLO, C12orf65, RP11-73M18.8, STRC, AP3B2, WHAMM, SLX1B, ARL17B,*

GATAD2A, RPS5, ADAM15, RP1-130H16.16, WBP2NL, NAGA, ALMS1, TYW5, ITIH4,

TMEM110, SFMBT1, NDUFAF2, BTN2A1, DNPH1). Nine genes were identified in the

deepN only (*CWF19L2, MORN3, TIE1, RBBP5, CBR3, RFTN2, RP11-10L12.4,*

REEP2, STK4). Twenty-four TWAS genes are specific to glial CGs (9 from ast, 10 from

oli and 5 from mic_ast_endo).

Co-expression network analysis refined CG-specific biological processes

associated with SCZ

To investigate the coordinated expression changes in each CG, we performed

weighted gene correlation network[32] (WGCNA) on CG-specific expressions

(Supplemental Table 4). For this study, we identified *CG-specific modules,* a group of

genes showing concordant variation across samples in one specific CG. However,

modules of upperN and deepN were highly consistent. In total, we identified eight CG-

specific modules out of the 149 detected modules. Seven of the eight showed either

differential expression of eigengenes in SCZ or enrichment of SCZ GWAS signals.

These seven modules were defined as "SCZ-associated modules". (Table 2).

The neuronal CGs shared five of the seven SCZ-associated modules (Fig. 6). The

eigengenes of two modules showed upregulated expression in SCZ: neuron_ME28

(log2FC=0.02, FDR=6.8e-03) and neuron_ME30 (log2FC=0.03, FDR=6.3e-07). These

two were enriched in the pathways of neural stem cell maintenance and carbohydrate

transmembrane transport (FDR q<0.05). The other three modules showed only

enrichment of GWAS signals (neuron_ME32, neuron_ME27, and neuron_ME29).

They were enriched within pathways of regulation of actin filament polymerization,

regulation of dendritic spine morphogenesis, and SWI/SNF superfamily-type complex

respectively (FDR q<0.05).

The other two SCZ-associated modules were specific to glial CGs. In the CGs of

astrocytes, we identified ast _ME28, enriched for SCZ GWAS signals and involved with

the RNA splicing pathway (FDR q<0.05). In the oligodendrocyte CG-related module

(oli_ME14), the eigengene was downregulated in SCZ (log2FC=-0.01, FDR=0.04).

Member genes in this oli_ME14 were involved in neuron enseathment (FDR q<0.05).

Discussion

Combining CG-specific eQTL analysis and GWAS can reveal *how* (the regulation of

gene expression) and *where* (within which CGs) variants contribute to disease risk.

Recent studies have employed this approach identifying cell-type-specific eQTLs in

blood that have heightened risk heritability for immune-related diseases[33,34] However,

cell-type-specific eQTLs for human brains are not available and are difficult to obtain.

To bridge genetic findings of SCZ to brain cell types, we generated CG, a proxy of cell

type, using an unsupervised deconvolution method. We then mapped CG-specific

eQTLs using computationally partitioned gene expression specific to each CG per

individual. Based on GWAS data, we found distinct enrichment for risk heritability in

the eQTLs of upper-layer DLPFC neuronal CGs (upperN). Furthermore, we found that

the upperN CGs also displayed gene expression changes and altered cell proportions

in the individuals we studied with SCZ.

Our results are consistent with previous SCZ studies in implicating neurons in the risk

for SCZ. In a case-control study of bulk brain tissues, Gandal et al. showed the neuron-

specific co-expression module to be downregulated in SCZ[16]. Skene et al. calculated

the neuronal specificity of the human-homologous gene in mouse scRNA-seq data and

correlated this specificity to risk heritability enrichment within the genetic region[35]. They

found that four neuron subtypes in the mouse brain (i.e., medium spiny neurons,

pyramidal cells in hippocampal CA1, pyramidal cells in the somatosensory cortex, and

cortical interneurons) are associated with SCZ. Our study expands upon those findings,

confirming that a subtype of neurons, the group of upper-layer neurons in the DLPFC

are associated with SCZ.

Our study provides genetic support for an association between the upper-layer

neurons of the DLPFC and SCZ. This is supported by the enrichment of GWAS signals,

dysregulated gene expression and decreased proportions within the upperN CGs.

Previous research has shown upper-layer neurons of the DLPFC to be actively

engaged in the delayed activity of working memory[36], the impairment of which is a

major feature of SCZ. Accumulating evidence points to upper-layer neurons within the

DLPFC as associated with SCZ in terms of cell density[37], cell shape[38], dendritic

abnormalities[39,40], and mRNA expression[41]. However, no genetic evidence to date has

pinpointed upper-layer neurons as having a role in SCZ risk, that we are aware of. In

this study, we observed that GWAS risk variants contribute to SCZ by altering gene

expression within that upper-layer neurons. Our coexpression network analysis and

TWAS analysis further narrowed down specific genes and pathways. Therefore, we

suspect that previously reported phenotypic changes[37-41] within the upper-layer

neurons of patients with SCZ may be rooted in these genetic variants and their related

gene expression regulation.

Differential gene expression associated with SCZ appeared primarily within the upperN

CG. This included 1304 upregulated and 1407 downregulated genes. Yet, the effect

size of the upperN-specific DEGs equaled that of the deepN-specific DEGs, with both

surpassing that of other glial CGs. This finding demonstrates a profound transcriptional

change within the neuronal CGs of the donors with SCZ. The DEGs in upperN were

enriched in the pathways of cell proliferation and trans-synaptic signaling, which has

previously been suggested as involved in SCZ risk. For example, disrupted

proliferation of neurons has been reported in primate and murine models of SCZ[42].

Research shows an association between a group of 1,026 synaptic genes and the risk

for SCZ[43]. Even though these two studies are not specific to upper-layer neurons, our

discovery suggests that these important pathways may bolster our understanding of

the neuron-related mechanisms underlying SCZ.

We identified candidate SCZ risk genes in upperN associated with the pathways that

support neuronal function. Five TWAS genes in upperN are related to neuronal

development: *AS3MT, CSPG4P11, ADAM15, STRC, and WHAMM.* These genes have

been reported as involved in SCZ risk. For example, *AS3MT* participates in neural

stem cell differentiation and is a risk gene for SCZ[44]. *CSPG4P11* is also an eQTL gene

in fetal brains, and its expression has also been linked to SCZ risk[45,46]. *ADAM15, STRC,*

and *WHAMM* function in cell-matrix adhesion, which is linked to neurodevelopment

and SCZ[47,48]. Even though adult brain tissues comprised our database, our results

suggest a probable pathway in which SCZ risk variants dysregulate genes involved in

neuronal development at the earliest stages. Furthermore, such dysregulation

ultimately affects the functions of upper-layer neurons in the DLPFC at later

developmental stages. Other TWAS genes found within the upperN are involved

general cell functions such as the formation of synaptic vesicles (*AP3B2*), lysosomal

enzymes (*NAGA*), and mitochondrial function (*C12orf65, NDUFAF2, DIABLO,*

ENDOG). While our study could not prove the causality of CG-specific risk genes to

SCZ, further functional experiments, such as the manipulation of brain organoids,

could potentially confirm the function and mechanisms of these candidates.

We also asked whether the enrichment of genetic variants of SCZ risk and evidence

of transcriptional changes in upper-layer neurons are associated with proportional

changes of upperN CG to all CGs. We confirmed a reduced proportion of upperN to all

CGs in the DLPFC of patients with SCZ. This is consistent with an earlier finding that

there is a reduced density of neurons in layer 2 of the frontal cortex in individuals with

SCZ (~29%)[41]. Further testing of the correlation between the decreased proportion of

upperN and the polygenic risk burden of SCZ (i.e., PRS), revealed an inverse

correlation. This suggests that a reduction of upper-layer neurons in the brains of those

with SCZ is due to their increased genetic burden for SCZ risk. We hypothesize that

TWAS genes related to neurodevelopment may contribute to cell proportions. While

our findings do not directly confirm a causal relationship, we afforded a notable

hypothesis for future testing.

Although our major discussion centers on the upperN CG, alterations of glial CGs

contribute to the complex pathology of SCZ as well. While significant risk heritability

enrichment was not evident in the glial CGs, we did observe an increased proportion

of the mic_ast_endo CG within the SCZ cohort. It suggests cell proportion of

mic_ast_endo is increased with SCZ risk. Moreover, 27% of the genes have changed

expression in at least one glial CG. Network analysis also affirmed the association of

gene expression changes in the glial CGs to SCZ. Ubiquitous transcriptional

dysregulation in glial CGs and changes in their proportions are consistent with previous

findings involving glial cells in the onset of SCZ, as related to the involvement of

synaptic transduction[49], inflammation[50], and myelination[51]. Whether and how

alterations of glial cells interact with neurons and circuits would be an important

question for future studies.

The discovery of upper layer neuron is a good surprise from swCAM that we would not

expect using existing reference cell-type marker genes with supervised method. While

swCAM yielded reproducible unsupervised deconvolution results, particularly for

neuronal CGs, it also has its limitations. The CGs estimated by swCAM represent

groupings of cells with similar transcriptional profiles. Correspondingly, swCAM has a

relatively poor resolution in separating cell subtypes of cells. Glial subtypes with small

composition, such as microglia, have been particularly challenging. As single-cell data

are accumulating, a hybrid model integrating a suitable reference may boost swCAM's

resolution. Despite these constraints, swCAM enhances the merit of existing bulk-

tissue data by deriving expression for major CGs. With this method, it is possible to

determine whether disease-related gene expression is a feature of specific CGs or a

pattern of composite CGs. Furthermore, gene networks and genetic regulatory

relationships can emerge that may help to ferret out hidden clues to the molecular

mechanisms of complex psychiatric disorders.

Conclusion

Of five clusters of cell types estimated by gene expression *in silico*, the upper-layer

neurons within the DLPFC display distinctive enrichment of risk heritability and

transcriptional changes in the brain tissue of donors with SCZ. Furthermore, decreased

proportion of this cell group to the total cell groups is shown within the SCZ brain. The

uniquely dysregulated genes and pathways associated with upper-layer neurons,

particularly those related to neurodevelopment, should be targeted in future

experimental studies and quantitative analysis of SCZ in order to further understand

the molecular mechanism of SCZ.

Materials and methods

Brain tissues

440 The PsychENCODE/BrainGVEX project has 459 postmortem DLPFC samples,

441 including healthy controls, BD and SCZ patients. Brain samples originated from the

442 Stanley Medical Research Institute (SMRI) and the Banner Sun Health Research

443 Institute (BSHRI). Detailed information can be found at

444 https://www.synapse.org/#!Synapse:syn4921369/wiki/235539.

RNA sequencing

Total RNA was isolated from brain tissue at the University of Illinois at Chicago and at

the University of Chicago with the Qiagen miRNeasy mini kit. Sample with a RIN score

of 5.5 or greater were assayed by Agilent Bioanalyzer RNA 6000 Nano assay kit. All

total RNAs were processed into rRNA-depleted stranded libraries for sequencing in

Illumina HiSeq2000. Libraries were tri-plexed per lane to reach 40 million paired-end

reads per library. Details of library preparation and RNA sequencing can be found in

our previous study[16].

Data processing

Data processing included three steps: reads processing, quantification, and post-

quantification processing. Fastq files underwent adapter removal using cutadapt

(https://pypi.python.org/pypi/cutadapt), after which the resulting adapter-trimmed fastq

files   were   checked   for   quality   using   FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). A subset of 10,000 reads

estimated the insert mean size and standard deviation for use with Tophat. STAR was

aligned trimmed reads to the GENCODEv19 reference (modified to include artificial

ERCC RNA ExFold spike-in sequences). BAM files were sorted in samtools (v1.3).

Expression level was then calculated using RSEM (v1.2.29). Quality control metrics

were   calculated   using   RNA-SeQC (v1.1.8),   featureCounts (v1.5.1),   PicardTools

(v1.128), and Samtools (v1.3.1). The quantified fragments per kilobase of transcript

per million mapped reads (FPKM) were used for downstream analysis. Mitochondrial

genes, pseudoautosomal genes, and genes with FPKM fewer than 1 in more than 75%

of samples were dropped. We calculated the distance between samples and removed

samples with a z-score of normalized connectivity to other samples lower than -2. After

filtering, 341 samples and 14865 genes were retained for subsequent analyses. Linear

regression removed the effect of covariates including age, sex, RIN, PMI, brain bank,

batches, and principal components of sequencing statistics (seqPC). The seqPCs was

composed of the top 29 principal components analyzed from the sequencing statistics.

Covariates were selected by multivariate adaptive regression splines (MARs).

Genotype processing

Genotypes were called from three platforms, including 277 samples by whole genome

sequencing, 257 samples by Psych v1.1 beadchips, and 137 samples by Affymetrix

5.0 450K. Genotypes were imputed by Michigan Imputation Server using HRC (r1.1

2016) EUR samples as references from each platform. Variants with squared R less

than 0.3 or HWE (Hardy-Weinberg Equilibrium) p-value less than 1e-6 were removed.

We used DRAMS software[52] to correct mixed-up data IDs for the three platforms based

on relationships across various omics data. We have 251 samples genotyped on whole

genome sequencing and beadchip. The genotypes called from the two platforms were

highly consistent (r= 0.986). Genotypes of the three platforms were combined, with

discordant genotypes marked as missing. Polygenic risk scores were calculated using

PRSice (2.1.4) with GWAS summary data from Ripke et al[2] as base data set.

Sample-specific deconvolution

To isolate CGs *in-silico*, we developed a novel sample-wise deconvolution technique

(swCAM), applying it to the processed RNA-seq data. The original CAM framed the

deconvolution of mixture expression to solve the blind source separation problem

(sBSS), which was accomplished by nuclear norm regularization. The basic

assumption in the deconvolution was

X=AS+E (1)

which is the mixture sample being the weighted sum of the CG-specific expression S.

In the classic deconvolution method, S is a common matrix for all mixture samples.

However, each sample may have its sample-specific S_i:

$S_i = \bar{S} + \Delta S_i$ (i=1, …, M) (2)

The associated sample-specific BSS model is given by

$x_i = a_i(\bar{S} + \Delta S_i) + e_i$ (3)

If some samples have a group pattern in a particular source or some molecules'

expression are more highly correlated in one particular source, the associated samples

may share a similar pattern of ΔS_i, leading to the following swCAM objective function:

$\min \frac{1}{2}\sum_{i=1}^{M} ||x_i - a_i(\bar{S} + \Delta S_i)||_2^2 + \lambda \sum_{k=1}^{K} ||T_k||_*$ (4)

Where T_k consists of the kth column in all ΔS_i, representing between-sample variability

in source K; $||T_k||^*$ is the nuclear norm of T_k. λ is the regularization parameter of nuclear

norms. We solved equation (4) using quadratic programming and the alternating

direction method of multipliers (ADMM). Code of swCAM can be accessed at

https://github.com/Lululuella/swCAM.

Firstly, we used CAM to estimate the CG proportion in an unsupervised manner. Five

CGs were estimated (CG1-CG5) without cell annotation. Then the proportion of five

CGs and processed RNA-seq data were entered into swCAM to infer the CG-specific

expression for each sample. To obtain the top ctDEGs and thereby differentiate the

CGs, Wilcoxon signed rank test determined the difference of expression in each CG

(log2FC>2, FDR<0.05).

Processing of single-cell datasets

Three single-cell datasets provided further confirmation of CG identities[27,28,53]. We

used only frontal cortices. Raw counts data were downloaded. The Seurat package

(v2.0) was used for data processing[54]. We filtered out the following: genes expressing

in less than half of the smallest CGs, cells with unique feature counts (i.e., >5,000 and

<200), and cells with mitochondrial counts >5%. The data were then normalized by

library size and log-transformed. The cell identity from the original studies were used

for cell clustering. The averaged expression was calculated to represent gene

expressions for the specific cell type.

Processing the GSE12649 dataset for replication

The GEO website (GSE12649)[55] provided a replication dataset with 102 human

DLPFCs. Gene expression was measured by Affymetrix Human Genome U133A Array.

CEL files were processed in Affy package[56]. After quality control, the data retained

13,652 probes and 67 samples including 32 controls and 35 SCZ samples. Expression

was normalized by a robust multi-array average (RMA) function followed by quantile

normalization.

Cell group identification

We used two strategies to annotate the CGs, gene sets enrichment and correlation

with scRNA-seq data. In the identification of five general CGs, the marker genes of five

major cell types were collected from Zhang et al[26] and the marker genes for inhibitory

neurons and excitatory neurons were collected from Lake et al[57]. Fisher's exact test

was used to test the enrichment of marker genes in CGs. Meanwhile, spearman

correlation testing with scRNA-seq datasets[27,53] measured similarities between CG

expression and that of known cell types. To identify genes expressed in gradient

cortical layers, we downloaded RNA-seq data across human DLPFC layers from the

He et al. study[58]. Gene expression was set as a linear regression function for DLPFC

layers (from layer 1 to layer 6). Genes with coefficient >0 and FDR <0.05 were named

"deep-layer genes." Those with coefficient <0 and FDR <0.05 were named "upper-

layer genes". scRNA-seq data used for correlation testing with neuronal CGs was

downloaded   from   https://portal.brain-map.org/atlases-and-data/rnaseq,   which

provided neuronal expression for the multiple layers of human middle temporal gyri.

Spearman correlation test was performed on layered neuronal expression and that of

CG2 and CG3.

Pathway analysis

Gene ontology enrichment tests for biological processes, molecular function and

cellular components were performed using gProfileR[59], with *p*-values FDR-corrected.

CG-specific eQTL mapping

To identify CG-specific eQTLs in the brain, we performed cis-eQTL mapping using

package FastQTL[60]. A cis-window was defined for genes in each CG, within 1

megabase window up- and down-stream of the gene body. Expression was adjusted

for PEER hidden covariates with FDR-corrected p-values. eQTLs with FDR <0.05 were

retained. We compared and defined CG-specific eQTLs, finding significant eQTL in

one CG.

To compare CG-specific eQTLs with single-cell and bulk brain eQTLs, we used the

Storey's Qvalue package[61]. The proportion of true associations ($\pi 1$) was estimated by

parallels of significant CG-specific eQTLs in the single-cell eQTLs/BrainGVEX DLPFC

eQTLs. With the distribution of corresponding p values for the overlapping eQTLs we

calculated $\pi 0$, i.e., proportion of true null associations based on distribution. Then, $\pi 1$

= 1 - $\pi 0$ estimated the lowest bound for true-positive associations.

Heritability estimation with GWAS summary statistics

Partitioned heritability was measured through LD Score Regression v1.0.0[62],

identifying enrichment of GWAS summary statistics among CG-specific eQTLs by

accounting for LD. CG-specific eQTL annotation files were created by eSNPs detected

in each CG. Marking all HapMap3 SNPs that fell within the eQTL annotations produced

the annotation file. LD-scores were calculated for SNPs within the file (LD window of

1cM). The LD reference panel was downloaded from 1000 Genomes European Phase

Three. SNP LD-scores were interwoven in the computation of partitioned heritability

(proportion of h^2). Enrichment for each annotation file was calculated by the proportion

of heritability explained by each annotation file divided by the proportion of SNPs falling

in that annotation file. Enrichment p values were then corrected by multiple testing.

Transcriptome-wide association study

We performed TWAS analysis on the GWAS summary statistics[63] and the CG-specific

eQTLs using the TWAS/FUSION package[9]. To identify risk genes with evidence of

genetic control, we used genome-wide complex trait analysis to estimate cis-SNP

heritability h2g ($\pm$1MB window around gene body). We identified genes with significant

h2g (nominal p-value <0.05), which were used to calculate the SNP-based predictive

weights per gene. Using the FUSION package, we calculated five-fold cross-validation

of five models of expression prediction and evaluated the prediction models for

accuracy. The five models are best cis-eQTL, best linear unbiased predictor, Bayesian

sparse linear mixed model [BSLMM], and elastic-net regression, LASSO regression.

The model with the largest cross-validation R^2 was chosen for downstream association

analyses. TWAS statistics were calculated using different cell-group weights, LD SNP

correlations from the 1000 Genomes European Phase 3 reference panel. TWAS

association p values were FDR-corrected.

Gene co-expression analysis

We constructed the gene co-expression network for each CG using Weighted Gene

Co-Expression Network Analysis (WGCNA) on each CG-specific expression data. A

correlation matrix was calculated for the genes, which was then weighted after a scale-

free topology was approximated. The following weighted powers were chosen for the

respective CGs: 12 for ast and deepN, 14 for upperN, mic_ast_endo, and oli. The

signed adjacency, biweight midcorrelation, and blockwiseModules parameters were

selected to build the network. Other parameter settings included the following:

mergeCutHeight= 0.1, minModuleSize= 30, pamStage=FALSE, and deepSplit=4.

Associating the gene modules with SCZ

Two strategies were used to test the association between gene modules and SCZ.

First, we constructed a linear model in which dependent variable is the module

eigengene across all samples and independent variable is the disease state for the

samples (FDR-adjusted p value<0.05). Second, we tested the correlation between

GWAS significance and the module membership (kME), which is the expression

correlation between each module member gene and the module eigengene. GWAS

significance of each module member gene were calculated by MAGMA v1.06[64] using

the European subset of the 1000 Genomes as a reference panel for LD. An annotation

step was performed first in which GWAS significant SNPs were mapped to genes,

based on the presence of a GWAS significant SNP in the region between a gene's

start and stop sites +/- 10kb. A competitive gene-set analysis was then performed using

sets defined by gene co-expression modules. Resulting p values were FDR-corrected

for multiple comparisons. Modules meeting any of the two criteria above were defined

as "SCZ-associated modules".

Gene module preservation test

To test the similarity among the CG-specific modules, we tested preservation among

the modules detected per CG using $Z_{summary}$. The network-based preservation test

generated two types of statistics: 1) density-based preservation statistics to determine

whether the nodes in the reference network were still highly connected in the test

network; 2) connectivity-based preservation statistics to determine whether the

connectivity pattern in the reference network was preserved in test network. We

applied $Z_{summary}$ test by Peter Langfelder et al[65], in the formula

626
$$Z_{\text{summary}} = \frac{Z_{density} + Z_{connectivity}}{2}$$

$Z_{density}$ summarizes density preservation statistics, and $Z_{connectivity}$ summarizes

connectivity-based statistics.

Data and code availability

Access to the raw BrainGVEX data is controlled by the NIMH Repository and

Genomics Resources (NRGR), https://www.nimhgenetics.org/. Instructions can be

found   in   the   PsychENCODE   Knowledge   Portal:

https://www.synapse.org/#!Synapse:syn4921369. Scripts used in the data analysis for

this   manuscript   can   be   found   at   GitHub:

https://github.com/RujiaDai/CellSpecificAnalysis.

Acknowledgments

This work was supported by NIH grants 1U01 MH103340-01, 1R01ES024988, and New York State Empire Innovation Program (to C.L.) and National Natural Science Foundation of China (NSFC) grants 31970572 and 31871276, National Key Plan for Scientific Research and Development of China grant 2016YFC1306000, and Innovation-Driven Project of Central South University grants 2020CX003 (to C.C.). We thank the PsychENCODE consortium and other data contributors (Iwamoto et al, Habib et al, Zhang et al, Darmanis et al, Lake et al, Mendizabal et al, Hodge et al, He et al) for their data support. We thank James Liu for contributions in web site design and data visualization. We gratefully acknowledge the families of the brain donors, without whom this work would not have been possible.

Author contributions

Data processing: R.D., S.L., Y.J. Bioinformatics analysis: R.D., S.L., Y.C, J.D. Algorithm development: L.C., G.Y., Y.W. Web design: R.D., Q.W. The conception of study design: R.D., S.L, C.C., C.L. Writing of the manuscript: R.D. Substantive revisions of the manuscript: L.K, R.D, C.L., C.C., R.K.

Competing interests

The authors have no competing financial interests to declare.

654 **Additional information**

655 This work was honored with the Hugh Gurling Memorial Awards by the International

656 Society of Psychiatric Genetics at XXVI World Congress of Psychiatric Genetics.

Figures and tables

Fig. 1 Overview of the study. Using the sample-wise convex analysis of mixtures (swCAM)

deconvolution method, we extracted cell-group-specific (CG-specific) expression from RNA-

seq data from bulk brain tissue of 341 samples (SCZ=93, CTL=248). To annotate the identified

CGs, maker gene enrichment and correlation testing with scRNA-seq data were used. To

discover SCZ-associated cell types and genes, we performed CG-specific eQTL mapping,

GWAS heritability enrichment, CG-specific differential expression, CG-specific co-expression,

and transcriptome-wide association study. These analyses offer a framework for discerning cell

670   subtypes and underlying genes and pathways associated with SCZ. We found upper-layer

671   neuron CGs in human DLPFC to be associated with SCZ.

Fig. 2 Cell-group-specific expression for each individual. (A) Expression of brain cell-type

marker genes per individual estimated cell groups (CGs). (B) Gradients of gene expression

across cortical layers in deconvoluted neuronal CGs. The top row (red) denotes upper-layer

genes (upperL genes), which express in a decreasing gradient from layers 1 to 6, and the

bottom row (blue) denotes deep-layer genes (deepL genes), which express in an increasing

gradient from layers 1 to 6 (FDR<0.05, Spearman correlation). (C) tSNE plot of deconvoluted

groups, color-coded by the annotated CG. Each dot denotes one deconvoluted cell within a

sample. Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN: upper-layer neuron CG,

mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial cells, oli: oligodendrocyte

CG.

Fig. 3 Summary of cell-group-specific eQTLs. (A) The number of eQTLs and eSNPs

detected in each cell group (CG) (FDR corrected p value<0.05). (B) Similarity between CG-

specific eQTLs, bulk brain tissue eQTLs, and single-cell eQTLs, evaluated by π1 values. (C)

Genomic annotation of eSNPs. Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN:

upper-layer neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial cells,

oli: oligodendrocyte CG.

Fig. 4 Genetic risk heritability enrichment in cell groups. GWAS heritability is partitioned in

five estimated cell groups (CGs). The SNPs of CG-specific eQTLs (FDR corrected p value<0.05)

are cataloged in annotated files in sLDSC. Orange denotes a corrected p values <0.05 and

grey denotes corrected p-values >0.05. Ast: astrocyte CG, deepN: deep-layer neuron CG,

upperN: upper-layer neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and

endothelial cells, oli: oligodendrocyte CG.

Fig. 5 Proportion per cell group and correlation with genetic burden. (A) Proportions of

the five cell groups (CGs) for donors with and without SCZ, estimated with the CAM method.

"*" denotes corrected p-values of the Wilcoxon rank-sum test <0.05. (B) Spearman correlation

between polygenic risk score (PRS) and CG proportion. "*" denotes corrected p-values of

correlation test <0.05. Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN: upper-layer

neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial cells, oli:

oligodendrocyte CG.

748 **Fig. 6 Neuron-specific co-expression modules related to SCZ.** Top 10 genes ordered by

749 module membership (kME, i.e., correlation between gene expression and module eigengene)

750 are shown for each neuron-group-specific module.

762 **Table 1 Summary of cell group-specific differential gene expression**

Cell group	#Upreg-ulated gene	#Downreg-ulated gene	Mean (|log2FC|) ± sd	Representing functional pathways of upregulated genes	Representing functional pathway of downregulated genes
ast	950	1660	0.11±0.18	carboxylic acid catabolic process	defense response
deepN	1254	1297	0.22±0.26	organic acid metabolic process	positive regulation of cell proliferation
upperN	1304	1882	0.26±0.25	carboxylic acid catabolic process	protein folding
mic_ast_endo	1197	1083	0.11±0.18	regulation of cellular component size	positive regulation of immune system process
oli	747	1445	0.14±0.18	modulation of chemical synaptic transmission	positive regulation of immune system process
Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN: upper-layer neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial cells, oli: oligodendrocyte CG.					

771 **Table 2 Cell group-specific co-expression modules**

Cell group	Module name	Trait association	GWAS enrichment	Representative GO term (FDR<0.05)	Top 3 hub genes
ast	ast _ME28	-	SCZ2014, Education, IBD	RNA splicing	SEC62, PPIG, PNN
neuron	neuron_ ME28	SCZ↑	SCZ2014, Education	neuronal stem cell population maintenance	TSC22D1, ENTPD3, SYNE1
	neuron_ ME29	-	SCZ2014, Education	SWI/SNF superfamily-type complex	WDR48, STOML2, OS9
	neuron_ ME30	SCZ↑	SCZ2014, Education, Neuroticism	carhydrate transmembrane transport	FKBP5, SORT1, WIPF3
	neuron _ME32	-	SCZ2014, Education	negative regulation of actin filament polymerization	GRIA4, CUX2, EPHX4
	neuron _ME27	-	SCZ2014, Education, Neuroticism, BD2016	regulation of dendritic spine morphogenesis	RIMS3, NGEF, LZTS3
mic_ast _endo	mic_ast _endo_ ME25	-	-	Defense response to virus	RSAD2, DDX58, PSME2
oli	oli_ME14	SCZ↓	-	Ensheathment of neuron	RNASE1, GPR37, PLP1
Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN: upper-layer neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial cells, oli: oligodendrocyte CG.					

Supplemental materials

Supplemental Fig. 1 Cell group-enrichment of brain cell marker genes. Plot shows

enrichment of the top cell-type differentially expressed genes (ctDEGs) in estimated

cell groups (CGs) (log2FC>2, FDR <0.05) for human brain cell marker genes. Color

denotes log10-scaled p values of enrichment for significant associations and text in

cell denotes odds ratio (Fisher's exact test).

Supplemental Fig. 2 Correlations between estimated cell-group-specific (CG-specific)

expression and (A) single cell expression (Darmanis et al. 2015) and (B) single nuclei

expression (Habib et al. 2017). The Spearman correlation test was used. Size and the

color both denote the degree of correlation. Ast: astrocyte CG, deepN: deep-layer

neuron CG, upperN: upper-layer neuron CG, mic_ast_endo: mixture CG of microglia,

astrocyte, and endothelial cells, oli: oligodendrocyte CG.

Supplemental Fig. 3 Expression of marker genes of the inhibitory and excitatory

neuronal subtypes in deconvoluted neuronal cell groups (CGs). Ex: excitatory neurons;

In: inhibitory neurons.

Supplemental Fig. 4 Spearman correlation between expressions of estimated neuronal cell groups (CGs) and expressions of neurons from different layers of the middle temporal gyrus. Spearman correlation test was used.

Supplemental Fig. 5 Deconvolution of replication dataset GSE12649. (A) Marker gene

expression in estimated cell groups (CGs). (B) Expressions of upper-layer and deep-layer

genes in estimated CGs. (C) Enrichment test of marker genes in estimated CGs. The number

denotes odds ratio and depth of color denotes log10-transformed p-value. (D and E)

Correlations between estimated CG-specific expression and single cell expressions (Darmanis

et al. 2015) and between estimated CG-specific expression and single nuclei expressions

(Habib et al. 2017). Ast: astrocyte CG, deepN: deep-layer neuron CG, upperN: upper-

layer neuron CG, mic_ast_endo: mixture CG of microglia, astrocyte, and endothelial

cells, oli: oligodendrocyte CG; opc: oligodendrocyte precursor cells; nsc: neuronal

stem cells; SMC, smooth muscle cells; odc: oligodendrocytes; exPFC: glutamatergic

neurons from the PFC; GABA: GABAergic interneurons; exCA1/3: pyramidal neurons

from the hip CA region; exDG: granule neurons from the hip dentate gyrus region.

upregulated

downregulated

Supplemental Fig. 6 Overlap of differentially expressed genes detected in each cell

groups. The differential expression of genes in SCZ was calculated with Wilcoxon

signed rank test (pvalue <0.05) per cell group (CG). The left-side bars denote the

number of differentially expressed genes in each CG. The upper bars denote the

intersection size between sets of differentially expressed genes. Dark connected dots

on the bottom panel denote which substrates are considered for each intersection.

835 **Supplemental table 1** Collected brain cell marker genes

836 **Supplemental table 2** Differential expressed genes in SCZ per cell group

837 **Supplemental table 3** TWAS genes in SCZ per cell group

838 **Supplemental table 4** Coexpression-module member genes and their module

839 membership in each cell group

References

1   Sullivan, P. F., Kendler, K. S. & Neale, M. C. Schizophrenia as a complex trait: evidence from a

meta-analysis of twin studies. *Archives of general psychiatry* **60**, 1187-1192,

doi:10.1001/archpsyc.60.12.1187 (2003).

2   Schizophrenia Working Group of the Psychiatric Genomics, C. Biological insights from 108

schizophrenia-associated genetic loci. *Nature* **511**, 421-427, doi:10.1038/nature13595 (2014).

3   Richards, A. L. *et al.* Schizophrenia susceptibility alleles are enriched for alleles that affect gene

expression in adult human brain. *Molecular psychiatry* **17**, 193-201, doi:10.1038/mp.2011.11

(2012).

4   Myers, A. J. *et al.* A survey of genetic human cortical gene expression. *Nature genetics* **39**, 1494-

1499, doi:10.1038/ng.2007.16 (2007).

5   Wang, D. *et al.* Comprehensive functional genomic resource and integrative model for the

human brain. *Science* **362**, doi:10.1126/science.aat8464 (2018).

6   Fromer, M. *et al.* Gene expression elucidates functional impact of polygenic risk for

schizophrenia. *Nature neuroscience* **19**, 1442-1453, doi:10.1038/nn.4399 (2016).

7   Jaffe, A. E. *et al.* Developmental and genetic regulation of the human cortex transcriptome

illuminate schizophrenia pathogenesis. *Nature neuroscience* **21**, 1117-1125,

doi:10.1038/s41593-018-0197-y (2018).

8   Ramasamy, A. *et al.* Genetic variability in the regulation of gene expression in ten regions of

the human brain. *Nature neuroscience* **17**, 1418-1428, doi:10.1038/nn.3801 (2014).

9   Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies.

Nat Genet **48**, 245-252, doi:10.1038/ng.3506 (2016).

10   Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait

gene targets. *Nature genetics* **48**, 481-487, doi:10.1038/ng.3538 (2016).

11   Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference

transcriptome data. *Nat Genet* **47**, 1091-1098, doi:10.1038/ng.3367 (2015).

12   Brown, C. D., Mangravite, L. M. & Engelhardt, B. E. Integrative modeling of eQTLs and cis-

regulatory elements suggests mechanisms underlying cell type specificity of eQTLs. *PLoS Genet*

9, e1003649, doi:10.1371/journal.pgen.1003649 (2013).

13   van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs

and co-expression QTLs. *Nature genetics* **50**, 493-497, doi:10.1038/s41588-018-0089-9 (2018).

14   Dimas, A. S. *et al.* Common regulatory variation impacts gene expression in a cell type-

dependent manner. *Science* **325**, 1246-1250, doi:10.1126/science.1174148 (2009).

15   Gandal, M. J. *et al.* Shared molecular neuropathology across major psychiatric disorders

parallels polygenic overlap. *Science* **359**, 693-697, doi:10.1126/science.aad6469 (2018).

16   Gandal, M. J. *et al.* Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and

bipolar disorder. *Science* **362**, doi:10.1126/science.aat8127 (2018).

17   Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**, 332-

337, doi:10.1038/s41586-019-1195-2 (2019).

18   Velmeshev, D. *et al.* Single-cell genomics identifies cell type-specific molecular changes in

autism. *Science* **364**, 685-689, doi:10.1126/science.aav8130 (2019).

19   Huang, Q. Q., Ritchie, S. C., Brozynska, M. & Inouye, M. Power, false discovery rate and Winner's

Curse in eQTL studies. *Nucleic acids research* **46**, e133, doi:10.1093/nar/gky780 (2018).

20   Shen-Orr, S. S. & Gaujoux, R. Computational deconvolution: extracting cell type-specific

information from heterogeneous samples. *Current opinion in immunology* **25**, 571-578,

doi:10.1016/j.coi.2013.09.015 (2013).

21   Wang, N. *et al.* Mathematical modelling of transcriptional heterogeneity identifies novel

markers and subpopulations in complex tissues. *Scientific reports* **6**, 18909,

doi:10.1038/srep18909 (2016).

22   Chen, L. *et al.* debCAM: a bioconductor R package for fully unsupervised deconvolution of

complex tissues. *Bioinformatics* **36**, 3927-3929, doi:10.1093/bioinformatics/btaa205 (2020).

23   Rahmani, E. *et al.* Cell-type-specific resolution epigenetics without the need for cell sorting or

single-cell biology. *Nature communications* **10**, 3417, doi:10.1038/s41467-019-11052-9 (2019).

24   Wang, J., Devlin, B. & Roeder, K. Using multiple measurements of tissue to estimate subject-

and cell-type-specific gene expression. *Bioinformatics*, doi:10.1093/bioinformatics/btz619

(2019).

25   Chen, L. Mathematical Modeling and Deconvolution for Molecular Characterization of Tissue

Heterogeneity. *Doctoral dissertation*, doi:http://hdl.handle.net/10919/96553 (2020).

26   Zhang, Y. *et al.* Purification and Characterization of Progenitor and Mature Human Astrocytes

Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37-53,

doi:10.1016/j.neuron.2015.11.013 (2016).

27   Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nature methods* **14**,

955-958, doi:10.1038/nmeth.4407 (2017).

28   Hodge, R. D. *et al.* Conserved cell types with divergent features in human versus mouse cortex.

Nature **573**, 61-68, doi:10.1038/s41586-019-1506-7 (2019).

29   Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from

high-throughput sequencing data. *Nucleic acids research* **38**, e164, doi:10.1093/nar/gkq603

(2010).

30   Mendizabal, I. *et al.* Cell type-specific epigenetic links to schizophrenia risk in the brain.

Genome biology **20**, 135, doi:10.1186/s13059-019-1747-7 (2019).

31   Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nature genetics* **47**, 1228-1235, doi:10.1038/ng.3404 (2015).

32   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis.

BMC bioinformatics **9**, 559, doi:10.1186/1471-2105-9-559 (2008).

33   Schmiedel, B. J. *et al.* Impact of Genetic Polymorphisms on Human Immune Cell Gene

Expression. *Cell* **175**, 1701-1715 e1716, doi:10.1016/j.cell.2018.10.022 (2018).

34   Raj, T. *et al.* Polarization of the effects of autoimmune and neurodegenerative risk alleles in

leukocytes. *Science* **344**, 519-523, doi:10.1126/science.1249547 (2014).

35   Skene, N. G. *et al.* Genetic identification of brain cell types underlying schizophrenia. *Nature*

genetics **50**, 825-833, doi:10.1038/s41588-018-0129-5 (2018).

36   Finn, E. S., Huber, L., Jangraw, D. C., Molfese, P. J. & Bandettini, P. A. Layer-dependent activity

in human prefrontal cortex during working memory. *Nature neuroscience* **22**, 1687-1695,

doi:10.1038/s41593-019-0487-z (2019).

37   Benes, F. M., Davidson, J. & Bird, E. D. Quantitative cytoarchitectural studies of the cerebral

cortex of schizophrenics. *Archives of general psychiatry* **43**, 31-35,

doi:10.1001/archpsyc.1986.01800010033004 (1986).

38   Cullen, T. J. *et al.* Anomalies of asymmetry of pyramidal cell density and structure in

dorsolateral prefrontal cortex in schizophrenia. *The British journal of psychiatry : the journal of*

mental science **188**, 26-31, doi:10.1192/bjp.bp.104.008169 (2006).

39   Garey, L. J. *et al.* Reduced dendritic spine density on cerebral cortical pyramidal neurons in

schizophrenia. *Journal of neurology, neurosurgery, and psychiatry* **65**, 446-453,

doi:10.1136/jnnp.65.4.446 (1998).

40   Glantz, L. A. & Lewis, D. A. Decreased dendritic spine density on prefrontal cortical pyramidal

neurons in schizophrenia. *Archives of general psychiatry* **57**, 65-73,

doi:10.1001/archpsyc.57.1.65 (2000).

41   Joshi, D., Catts, V. S., Olaya, J. C. & Shannon Weickert, C. Relationship between somatostatin

and death receptor expression in the orbital frontal cortex in schizophrenia: a postmortem

brain mRNA study. *NPJ schizophrenia* **1**, 14004, doi:10.1038/npjschz.2014.4 (2015).

42   Selemon, L. D. & Zecevic, N. Schizophrenia: a tale of two critical periods for prefrontal cortical

development. *Translational psychiatry* **5**, e623, doi:10.1038/tp.2015.115 (2015).

43   Lips, E. S. *et al.* Functional gene group analysis identifies synaptic gene groups as risk factor for

schizophrenia. *Molecular psychiatry* **17**, 996-1006, doi:10.1038/mp.2011.117 (2012).

44   Li, M. *et al.* A human-specific AS3MT isoform and BORCS7 are molecular risk factors in the

10q24.32 schizophrenia-associated locus. *Nature medicine* **22**, 649-656, doi:10.1038/nm.4096

(2016).

45   O'Brien, H. E. *et al.* Expression quantitative trait loci in the developing human brain and their

enrichment in neuropsychiatric disorders. *Genome biology* **19**, 194, doi:10.1186/s13059-018-

1567-1 (2018).

46   Hauberg, M. E. *et al.* Large-Scale Identification of Common Trait and Disease Variants Affecting

Gene   Expression.   *American   journal   of   human   genetics*   **100**,   885-894,

doi:10.1016/j.ajhg.2017.04.016 (2017).

47   Hattori, T. *et al.* DISC1 regulates cell-cell adhesion, cell-matrix adhesion and neurite outgrowth.

Molecular psychiatry **15**, 778, 798-809, doi:10.1038/mp.2010.60 (2010).

48   Hirota, Y. & Nakajima, K. Control of Neuronal Migration and Aggregation by Reelin Signaling in

the Developing Cerebral Cortex. *Frontiers in cell and developmental biology* **5**, 40,

doi:10.3389/fcell.2017.00040 (2017).

49   Mei, Y. Y., Wu, D. C. & Zhou, N. Astrocytic Regulation of Glutamate Transmission in

Schizophrenia. *Frontiers in psychiatry* **9**, 544, doi:10.3389/fpsyt.2018.00544 (2018).

50   Monji, A., Kato, T. & Kanba, S. Cytokines and schizophrenia: Microglia hypothesis of

schizophrenia. *Psychiatry and clinical neurosciences* **63**, 257-265, doi:10.1111/j.1440-

1819.2009.01945.x (2009).

51   Dietz, A. G., Goldman, S. A. & Nedergaard, M. Glial cells in schizophrenia: a unified hypothesis.

The lancet. Psychiatry **7**, 272-281, doi:10.1016/S2215-0366(19)30302-5 (2020).

52   Jiang, Y. *et al.* DRAMS: A tool to detect and re-align mixed-up samples for integrative studies of

multi-omics   data.   *PLoS   computational   biology*   **16**,   e1007522,

doi:10.1371/journal.pcbi.1007522 (2020).

53   Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell level.

Proceedings of the National Academy of Sciences of the United States of America **112**, 7285-

7290, doi:10.1073/pnas.1507125112 (2015).

54   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic

data across different conditions, technologies, and species. *Nature biotechnology* **36**, 411-420,

doi:10.1038/nbt.4096 (2018).

55   Iwamoto, K., Bundo, M. & Kato, T. Altered expression of mitochondria-related genes in

postmortem brains of patients with bipolar disorder or schizophrenia, as revealed by large-

scale DNA microarray analysis. *Human molecular genetics* **14**, 241-253,

doi:10.1093/hmg/ddi022 (2005).

56   Gautier, L., Cope, L., Bolstad, B. M. & Irizarry, R. A. affy--analysis of Affymetrix GeneChip data

at the probe level. *Bioinformatics* **20**, 307-315, doi:10.1093/bioinformatics/btg405 (2004).

57   Lake, B. B. *et al.* Integrative single-cell analysis of transcriptional and epigenetic states in the

human adult brain. *Nature biotechnology* **36**, 70-80, doi:10.1038/nbt.4038 (2018).

58   He, Z. *et al.* Comprehensive transcriptome analysis of neocortical layers in humans,

chimpanzees and macaques. *Nature neuroscience* **20**, 886-895, doi:10.1038/nn.4548 (2017).

59   Reimand, J., Kull, M., Peterson, H., Hansen, J. & Vilo, J. g:Profiler--a web-based toolset for

functional profiling of gene lists from large-scale experiments. *Nucleic acids research* **35**,

W193-200, doi:10.1093/nar/gkm226 (2007).

60   Ongen, H., Buil, A., Brown, A. A., Dermitzakis, E. T. & Delaneau, O. Fast and efficient QTL mapper

for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479-1485,

doi:10.1093/bioinformatics/btv722 (2016).

61   Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proceedings of the*

National Academy of Sciences of the United States of America **100**, 9440-9445,

doi:10.1073/pnas.1530509100 (2003).

62   Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in

genome-wide association studies. *Nature genetics* **47**, 291-295, doi:10.1038/ng.3211 (2015).

63   Bipolar, D., Schizophrenia Working Group of the Psychiatric Genomics Consortium. Electronic

address, d. r. v. e., Bipolar, D. & Schizophrenia Working Group of the Psychiatric Genomics, C.

Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell*

173, 1705-1715 e1716, doi:10.1016/j.cell.2018.05.046 (2018).

64   de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set analysis

of GWAS data. *PLoS computational biology* **11**, e1004219, doi:10.1371/journal.pcbi.1004219

(2015).

65   Langfelder, P., Luo, R., Oldham, M. C. & Horvath, S. Is my network module preserved and

reproducible? *PLoS computational biology* **7**, e1001057, doi:10.1371/journal.pcbi.1001057

(2011).

1017