Single cell transcriptional profiling reveals helper, effector, and regulatory MAIT cell populations enriched during homeostasis and activation

Charles Kyriakos Vorkas[1,2†], Chirag Krishna[3†], Kelin Li[4], Jeffrey Aubé[4], Daniel W. Fitzgerald[1,5], Linas Mazutis[6], Christina S. Leslie[3*], Michael S. Glickman[2,7*]

1. Division of Infectious Diseases, Weill Cornell Medicine, Cornell University, NY, NY, USA
2. Immunology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, NY, NY, USA
3. Computational and Systems Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, NY, NY, USA
4. Division of Chemic Biology and Medicinal Chemistry, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
5. Center for Global Health, Weill Cornell Medicine, Cornell University, NY, NY, USA
6. Single Cell Research Initiative, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA
7. Division of Infectious Diseases, Memorial Sloan Kettering Cancer Center, NY, NY, USA

[†] These authors contributed equally to this manuscript.

* Correspondence: cleslie@cbio.mskcc.org (C.S.L.), glickmam@mskcc.org (M.S.G)

Abstract

Mucosal-associated invariant T (MAIT) cells are innate-like lymphocytes that recognize microbial

vitamin B metabolites and have emerging roles in infectious disease, autoimmunity, and cancer.

Although MAIT cells are identified by a semi-invariant T cell receptor, their phenotypic and

functional heterogeneity is not well understood. Here we present an integrated single cell

transcriptomic analysis of over 76,000 human MAIT cells during acute and chronic antigen-

specific activation with the MR1 ligand 5-OP-RU and non-specific TCR stimulation. We show

that MAIT cells span a broad range of homeostatic, effector, helper, tissue-infiltrating, regulatory,

and exhausted phenotypes, with distinct gene expression programs associated with CD4$^+$ or

CD8$^+$ co-expression. During acute activation, MAIT cells rapidly adopt a cytotoxic phenotype

characterized by high expression of *GZMB*, *IFNG* and *TNF*. In contrast, chronic stimulation

induces heterogeneous states defined by proliferation, cytotoxicity, immune modulation, and

exhaustion. These scRNAseq-defined MAIT cell subtypes were also detected in individuals

recently exposed to *Mycobacterium tuberculosis* infection, confirming their presence during

human infection. Our study provides the first comprehensive atlas of human MAIT cells in

activation conditions and defines substantial functional heterogeneity, suggesting complex roles

in health and disease.

Introduction

Mucosal-associated invariant T (MAIT) cells are a subset of re-circulating innate-like lymphocytes enriched in human liver, lung and gut with highly variable abundance in peripheral blood in healthy donors (<1%-18% in humans)(Godfrey et al., 2019; Li et al., 2018; Vorkas et al., 2018). MAIT cells express an evolutionarily conserved T cell receptor (TCR; TRAV1-2 in humans) that can recognize microbially-derived, non-peptide small molecule metabolites of the vitamin B pathway(Harriff et al., 2018; Treiner et al., 2003) presented by the oligomorphic MHC Ib-related molecule, MR1(Seshadri et al., 2017; Treiner et al., 2003). In contrast to conventional T cells that exit the thymus in a "naive" state and require peptide-specific antigen priming over days to weeks to proliferate, most mature MAIT cells enter the circulation as effector cells and are activated within hours of antigen recognition(Legoux et al., 2019). This innate-like property confers their capacity to be among the first responders to bacterial and fungal pathogens(Dias et al., 2017; Jahreis et al., 2018; Shaler et al., 2017; Vorkas et al., 2018) as well as to viral infections through TCR-independent cytokine stimulation (Lamichhane et al., 2019; Leng et al., 2019; van Wilgenburg et al., 2016).

Despite a growing body of literature implicating MAIT cells in both homeostasis and disease(Godfrey et al., 2019), their specialized functional states in humans are poorly characterized relative to conventional T cells. For example, whereas conventional T cells are broadly divided into two lineages defined by CD4 or CD8 co-receptor expression and stable functional phenotypes, whether these same lineages can be defined in MAIT cells remains unclear. CD4$^+$ MAIT cells are reported to express increased CD25 and IL2 along with decreased cytotoxic markers relative to CD8$^+$ MAIT cells (Gherardin et al., 2018b; Kurioka et al., 2017; Lamichhane et al., 2019; Vorkas et al., 2018). While CD8$^+$ and CD4$^-$CD8$^-$ (double-negative; DN) MAIT cells have similar functional

phenotypes(Kurioka et al., 2017), the DN subset is reported to have depressed cytotoxic function

with increased propensity to activation-induced apoptosis(Dias et al., 2018). Moreover, while

prior studies have suggested that MAIT cell functional states may depend on the quality and

duration of activating stimuli(Kelly et al., 2019; Lamichhane et al., 2019), the early and late effects

of TCR-stimulation on MAIT cell state remain largely unknown.

Here, we present a single cell transcriptomic atlas of human MAIT cells, comprising more than

76,000 cells from peripheral blood under acute and chronic activation with 5-OP-RU or

αCD3/CD28 in the presence of IL2 or IL2/TGFβ. Our study elucidates distinct transcriptional

programs of CD4$^+$ and CD8$^+$ MAIT cells and identifies novel MAIT cell functional clusters that we

validate in human subjects recently exposed to *Mycobacterium tuberculosis*. Our results

demonstrate broad functional heterogeneity of MAIT cells and suggest that these innate T cells

can participate in complex immune responses in addition to their role in direct killing of

pathogens.

Results

CD4 or CD8 expression on MAIT cells is associated with distinct transcriptional states

To simulate acute and chronic MAIT cell activation under antigen-specific and non-specific T cell

receptor stimulation, we employed an in vitro activation assay using human PBMCs incubated

with MR1 ligand (5-OP-RU) or αCD3/CD28 beads for 15 hours (D1; acute) or 7 days (D7; chronic)

compared to unstimulated control cells. A subset of conditions at D7 also included TGFβ to

further assess potential regulatory phenotypes(Bhattacharyya et al., 2018; Rouxel et al., 2017).

Cells were harvested at each time point for flow cytometric assays and flow-cytometry assisted

cell sorting (FACS) using MR1 tetramers (Fig. 1a, Supplementary Fig. 1, and Supplementary

Table 1). To define the immunologic landscape of MAIT cell states after acute and chronic

activation, we performed droplet microfluidics single cell RNA sequencing of these sorted MAIT

cells from the peripheral blood of three donors in the activation conditions defined above. 76,845

MAIT cells passed quality control and were used for downstream analysis.

Although prior studies have demonstrated that MAIT cells can express CD4 or CD8 co-

receptors(Gherardin et al., 2018b; Kurioka et al., 2017; Vorkas et al., 2018), it is unclear whether

these markers define functionally distinct MAIT cell populations. We first sought to resolve this

question using our single cell data by performing a differential expression analysis between

CD4$^+$CD8$^-$ and CD4$^-$CD8$^+$ MAIT cells at D1 (Fig. 1b,c) or D7 (Fig. 1d, e) in the absence of TCR

stimulation. At D1 (Fig. 1c) CD4$^+$ MAIT cells more highly express genes associated with

conventional naïve and central memory T cells including *LEF1*, *SELL*, and *CCR7*, but also

markers of activation and co-stimulation such as *CTLA4*, *IL2RA*, *TNFRSF4*, and *TNFRSF18* (Fig

1b, Supplementary Table 2). CD8$^+$ MAIT cells more highly express markers of cytotoxic activity

such as *NKG7*, *KLRD1*, *EOMES*, *GNLY*, and *GZMB* as well as the transcription factor *RORA*.

We confirmed the differential expression of CD25, OX40, GZMB, and EOMES at the protein level

within CD4$^+$, CD8$^+$, DN and CD4$^+$CD8$^+$ (double-positive; DP) MAIT cell subsets using flow

cytometry (Fig. 1c). At D7, CD4$^+$ MAIT cells upregulated *CTLA4*, *TNFRSF4*, *SELL*, *IL2RA* and

CCR7, whereas CD8$^+$ MAIT cells upregulated *NKG7*, *GZMA*, *GZMK*, *PRF1*, SLAMF7 as well as

several genes involved in Type I interferon signaling (*IFI44L*, *IFITM3*, *IFI6*, *IFITM1*, and *ISG15*)

(Fig. 1d, Supplementary Table 3). We confirmed the differential expression of OX40, CTLA4,

GZMA, and GZMK in each subset at the protein level (Fig. 1e). Our analysis demonstrates that

CD4$^+$ MAIT cells are associated with expression of co-stimulatory receptors, IL2 signaling and

memory markers, whereas CD8$^+$ MAIT cells are defined by granzyme-mediated cytotoxicity and

type 1 interferon signaling. These results suggest that CD4 and CD8 expression on MAIT cells

may define distinct functional roles.

Single cell transcriptional analysis defines heterogeneous MAIT cell activation states

To fully characterize the phenotypic states of tetramer positive MAIT cells, we performed

dimensionality reduction by principal component analysis followed by Louvain clustering on cells

from all donors, timepoints, and stimuli (Fig. 2a). Clusters were broadly partitioned by timepoint

(Supplementary Fig. 2). Visualization with uniform manifold approximation and projection

(UMAP) together with differential expression analysis displayed 12 clusters that could be

distinguished by homeostatic, cytotoxic, Th1-associated cytokine release, proliferating, tissue-

infiltrating, immune regulatory, and exhausted states (Fig. 2b-c; Supplementary Table 3).

Clusters 1-3 were enriched at D1 (acute activation). Cluster 1 (C1) had high expression of *IFNG*,

GZMB, *TNF*, *TBX21*, and notably the transcription factor *FOXP3*. Since this cluster expressed

genes associated with granzymes and Th1-associated cytokines, it was termed "MAIT1

cytotoxic effector" (Koay et al., 2019), consistent with a similar cluster detected in Koay *et*

al.(Koay et al., 2019) C2 demonstrated increased *CD69* and pronounced expression of heat

shock proteins, *NR4A1*, and the AP-1 subunits *JUN* and *FOS* and was named "Early activated".

C3 expressed effector/central memory markers *CCR7* and *C62L*, markers of MAIT cell

intrathymic development *CD4* and *LEF1*(Koay et al., 2019), as well as *RUNX3*. This cluster also

upregulated type 1 interferon signaling genes, e.g. *IFIT3* and *ISG15*. Thus, C3 was named "ISG⁺

memory" (ISG: interferon-stimulated genes). Collectively these clusters define the landscape of

MAIT cells during homeostasis and early activation. The rapid induction of cytotoxic effector

programs at this timepoint suggest that MAIT cells may be poised to respond earlier than

conventional T cells.

In contrast, clusters 4-12 were enriched with chronic activation and demonstrated marked

phenotypic heterogeneity. C4 and C5 ("Homeostatic effectors") displayed similar expression

patterns, with high levels of *IL7R*, *GZMM*, *GZMK*, *KLRG1* and *IFI44L* and decreased expression

of *CD69* and *CD25*. C4 was distinguished from C5 by expression of vascular endothelial growth

factors *VEGFA/B* and was named "Homeostatic effector VEGF." C6 ("Degranulating effector")

showed high expression of *LAMP1*, cytokines *TNF* and *TNFSF13B* (B-cell activating factor),

IFNGR1, as well as the homing molecule *CXCR3*. C7 ("MR1⁺ MAIT17") expressed *GZMA*,

ENTPD1, *RORA*(Koay et al., 2019) as well as *MR1* itself, suggesting that this population may

interact with other MAIT cells. C8 ("Tissue-infiltrating-like") highly expressed *ITGA1* and *ITGAE*

and was only present in activation conditions including TGFβ. C9 was termed "Innate-like" given

its high expression of *HOPX*, *KLRB1*, and *NCR3* together with *GZMA* and *GZMK*. Clusters 10-

12 were defined by a gradient of activation/exhaustion marker expression including *PDCD1*,

CTLA4, *PDCD1* and *LAG3*, with C12 ("Late activated/exhausted") resembling a terminally

exhausted population. C11 ("Proliferating") demonstrated a strong mitotic phenotype. C10

("Cytotoxic/Regulatory") upregulated *GZMA*, *GNLY* and *IL26* as well as immune modulating

genes *TGFβ* and *LAG3*. Taken together, these analyses define a highly diverse MAIT cell

transcriptional landscape shaped by chronic activation, with important parallels with

conventional CD4$^+$ and CD8$^+$ T cells.

Acute MAIT cell activation induces a cytotoxic phenotype marked by GZMB, IFNG, and

TNF

We next asked how the prevalence of these populations varied across stimuli and donors during

acute and chronic activation (Fig. 2d-e and Supplementary Fig. 2b). In particular, C1 "MAIT1

cytotoxic effector"—which expressed high levels of *IFNG*, *GZMB*, and *TNF*—was enriched by

TCR stimulation conditions with 5-OP-RU or αCD3/CD28 compared to cells incubated with IL2

alone (Fig. 3a, b). To validate and further explore these early transcriptional phenotypes, we

performed flow cytometric characterization of MAIT cells under the same acute activation

conditions.

First, we confirmed the differential expression of common T cell surface activation markers IL7R,

CD69, and CD25 (Fig. 3c). Unexpectedly, IL7R was significantly downregulated by 5-OP-RU, but

not αCD3/CD28. CD69 and CD25 were significantly upregulated by both stimulation conditions,

though CD69 was more highly expressed with aCD3/CD28, likely indicating the relative potency

of this non-specific T cell stimulus. These results demonstrate that MAIT cells rapidly adopt

distinct immune phenotypes after acute antigen-specific and non-specific TCR stimulation.

As *GZMB*, *IFNG*, and *TNF* were significantly upregulated in early activation clusters C1 and C2,

we next asked if these markers were co-expressed at the protein level. A representative gating

strategy of intracellular co-staining for GZMB, IFNγ and TNF is presented alongside cumulative

data from 30 healthy donors (Fig. 3d). These analyses confirmed the presence of a MAIT cell

population with a combined granzyme and Th1-associated cytokine phenotype enriched with

acute activation, consistent with "MAIT1 cytotoxic" C1 enriched in the αCD3/CD28 condition in

the single cell RNA-seq analysis. The primary acute activation response in MAIT cells was

upregulation of granzyme B alone, which was increased in αCD3/CD28 relative to 5-OP-RU

conditions (Fig. 3d). 5-OP-RU and αCD3/CD28 both induced significant co-expression of GZMB

and TNFα, though increased co-expression was observed with αCD3/CD28 (Fig. 3d). Only

αCD3/CD28 induced IFNγ and TNFα co-expression (Fig. 3d). These results are consistent with

increased expression of IFNγ and TNFα with non-specific TCR stimulation.

We next performed a differential expression analysis between MAIT cells stimulated with 5-OP-

RU and aCD3/CD28 at D1 to define the gene expression programs associated with antigen-

specific or non-specific TCR stimulation (Fig. 3e and Supplementary Table 4). Although both

acute TCR-stimulation conditions induced similar transcriptional states (Fig. 2d), we did detect

stimulus-specific gene expression patterns, with αCD3/CD28 robustly upregulating *GZMB*,

PRF1, *IFNG*, *IL2RA*, *BATF3*, *CTLA4*, and *LAG3*. In contrast, 5-OP-RU induced a strong type 1

interferon signaling signature with upregulation of *IFIT1*, *ISG15*, *IFI44L*, *IFIT3*, *IFITM1* and *IFI6*.

These findings are consistent with distinct MAIT cell transcriptional programs driven by acute

antigen-specific signals versus non-specific T cell stimulation.

MAIT cells upregulate FOXP3 during acute and chronic activation

Given differential expression of *FOXP3* observed in C2, we sought to further characterize

peripheral FOXP3 expressing MAIT cell subsets by flow cytometry. Although few FOXP3[+] MAIT

cells were detected in resting cells, FOXP3 protein was significantly upregulated with 5-OP-RU

and αCD3/CD28 during acute stimulation (Fig. 4a,b). In contrast, only 5-OP-RU significantly

upregulated FOXP3 during chronic stimulation (Fig. 4b).

As FOXP3 is a transcriptional regulator of CD4$^+$ regulatory T (Treg) cells and often co-expressed

with CD25(Liston et al., 2008; Yamaguchi et al., 2011), we hypothesized that MAIT cells could

share similar staining characteristics. First, we compared FOXP3 expression in CD4$^+$ and CD8$^+$

MAIT cells with non-MAIT CD4$^+$ T cells (Fig. 4c). While CD4$^+$ MAIT cells had increased FOXP3

expression at baseline, only chronic 5-OP-RU-activated CD8$^+$ MAIT cells stably upregulated

FOXP3 (Fig. 4c). The addition of TGFβ did not significantly affect MAIT cell FOXP3 expression,

though there was a trend for increased FOXP3$^+$ expression with TGFβ co-stimulation (Fig. 4c).

We next compared FOXP3/CD25 co-expression in MAIT cells with non-MAIT CD4$^+$ T cells and

found that 5-OP-RU-activated MAIT cells strongly upregulated FOXP3, but CD3/CD28

stimulation only upregulated FOXP3 in conventional CD4$^+$ T cells (Fig. 4d). In contrast to Treg

cells, the FOXP3$^+$CD25$^+$ MAIT cells were strongly induced in the CD8$^+$ MAIT cell subset after 5-

OP-RU activation (Fig. 4e).   As previously reported(Hoffmann et al., 2004), αCD3/CD28

stimulation alone resulted in significant Treg cell expansion (Fig. 4e). Analysis of absolute

numbers of FOXP3$^+$CD25$^+$ MAIT cells demonstrate that FOXP3$^+$CD25$^+$ MAIT cells were most

prevalent within the CD8$^+$ MAIT cell subset (Fig. 4f) and that the vast majority of FOXP3

expressing cells were detected among CD25$^+$IL7R$^-$ cells and rarely among CD25$^-$ cells (Fig. 4g).

Chronic activation of MAIT cells expands homeostatic subpopulations while also inducing

proliferative, cytotoxic, regulatory, and exhausted phenotypes

We next examined clusters enriched during chronic activation (Fig. 5a). These clusters were

broadly associated with expression of *MKI67*, *GZMA*, *GZMB*, *TNF*, *CTLA4*, *ICOS*, *LAG3*, *FAS*,

and *PDCD1* (Fig. 5a-b). Despite largely overlapping transcriptional states observed between

chronic 5-OP-RU and αCD3/CD28 conditions (Fig. 2e), 5-OP-RU significantly upregulated genes

associated with lymphoid development (*LTB*), self-renewal (*LST1*), maturation (*CD52*), motility

and proliferation (*STMN1*, *ACTG1*) and recruitment of immune cells (*CCL3*, *CCL4*)

(Supplementary Table 4). In contrast, αCD3/CD28 upregulated genes associated with

cytotoxicity (*GZMB*, *GZMK*), cell stress (*HILPDA*), and exhaustion (*CTLA4*, *LAG3*)

(Supplementary Table 4). This granzyme/exhaustion signature observed was largely driven by

C12 defined by cells from Donor 1 (Supplementary Fig. 2c).

The transcriptional differences between stimuli were further elucidated at the protein level, where

proliferation marker Ki67 was significantly upregulated in the 5-OP-RU but not the αCD3/CD28

condition (Fig. 5c,d). This was consistent with robust antigen-specific MAIT cell expansion not

observed with chronic αCD3/CD28 (Fig. 5e). Similar to acute activation, we again detected

significant downregulation of IL7R at the protein level only in the 5-OP-RU condition, which was

associated with significant upregulation of CD25 not observed with chronic αCD3/CD28

stimulation (Fig. 5f). Similar to acute activation, GZMB was the dominant MAIT cell effector

protein (Fig. 5g). Activation/exhaustion markers CTLA4, PD1, FAS, ICOS and LAG3 expression

were also enhanced with chronic 5-OP-RU stimulation at the protein level (Fig. 5g).

MAIT cell subtypes are present during human tuberculosis infection

Finally, we assessed whether the MAIT cell phenotypes observed during acute and chronic

activation could also be detected during natural human exposure to *Mycobacterium tuberculosis*

(*Mtb*). To this end, we analyzed MAIT cell phenotypes in PBMCs collected from household

contacts of active TB patients in Port-au-Prince, Haiti compared to healthy unexposed donors

from the same community.

First, we focused our analysis on a subset of markers that characterized CD4+ MAIT cells in our

single cell analyses. We observed a similar differential expression pattern of CD25, CCR7 and

OX40 at the protein level in CD4+ MAIT cells relative to other subsets (Fig. 6a-c). Moreover, each

of these markers was significantly upregulated in contacts recently exposed to *Mtb,* indicating

that the acute activation gene expression pattern characteristic of clusters 1-3 in vitro

correspond to in vivo MAIT cell subsets in the context of recent *Mtb* exposure. Consistent with

our previous work(Vorkas et al., 2018), these findings indicate that CD4+ MAIT cells respond to

acute *Mtb* exposure.

We next interrogated MAIT cell FOXP3 expression, which was predominant in the CD8+ subset

after chronic antigenic stimulation (Fig. 4e,f).   We found that recently exposed contacts

expressed increased FOXP3 relative to healthy controls (Fig. 6d). Consistent with the scRNA-

seq results, FOXP3+CD25+ MAIT cells were most prevalent in the CD8+ subset, and

FOXP3+CD25+ CD8+ MAIT cells were significantly enriched in contacts (Fig. 6e). When stratifying

by *Mtb*-specific IFNγ release assay (IGRA) status, which distinguishes contacts who were

exposed and latently infected (IGRA+) from those who remained uninfected (IGRA-), we

observed that FOXP3+CD25+ CD8+ MAIT cells were enriched in latently infected contacts (Fig.

6f), further confirming that this MAIT cell subtype is responding to *Mtb* infection. These MAIT

cell phenotypes detected in TB contacts closely resemble the CD8+FOXP3+CD25+ MAIT cells

detected with antigen specific stimulation residing in clusters 1-3 (Fig. 3 and 4) and indicate that

these cell states defined by single cell transcriptomics are present during acute human *Mtb*

exposure and infection.

Discussion

MAIT cells have been well-characterized as innate-like, rapid responders to microbial antigens

with the capacity to mount cytotoxic responses and secrete Th1/Th17-associated cytokines

within hours of antigen recognition(Dias et al., 2017; Dusseaux et al., 2011; Vorkas et al., 2018).

However, less is known about the transcriptional programs that define MAIT cell adaptation

during acute and chronic activation states. Here we provide the first single cell transcriptional

atlas of over 76,000 human MAIT cells in homeostatic and acute/chronic activation states

coupled with flow cytometric and functional characterization. Our analysis demonstrates distinct

transcriptional programs defined by incubation period and quality of stimulus, revealing novel

MAIT cell subpopulations not yet described by bulk RNA-seq or traditional flow cytometric

approaches.

First, our work contributes further evidence to support divergent transcriptional and functional

phenotypes associated with human MAIT cell CD4 or CD8 co-expression. Several prior studies

have used CD8 as a prerequisite for defining MAIT cells or did not distinguish between

subsets(Coulter et al., 2017; Estevez et al., 2020; Koay et al., 2019; Slichter et al., 2016).

Differential expression analysis revealed that CD4$^+$ MAIT cells upregulated *IL7R*, *CD25*, *CCR7*,

SELL along with TNF superfamily receptors *TNFRSF4* and *TNFRSF18*, suggesting that CD4$^+$

MAIT cells respond to distinct co-stimulatory signals from the CD8$^+$ subset. We hypothesize

that this CD4$^+$ MAIT cell signature may confer unique tissue-homing functions through *CCR7*

and *SELL*. Future studies will help to evaluate whether this CD4$^+$ MAIT cell signature confers the

capacity to exit the circulation and migrate into tissues.

Recent work in mice and humans shows that CD4 and the transcription factor, LEF1, are co-

expressed during MAIT cell intrathymic development with reciprocal downregulation of CD319

(SLAMF7)(Koay et al., 2019). CD4 is subsequently downregulated as MAIT cells mature with

reciprocal upregulation of CD8 and CD319 during thymic egress. In a parallel observation in

human peripheral blood MAIT cells, we observed that CD4$^+$ MAIT cells significantly upregulated

LEF1, while *SLAMF7* was significantly differentially expressed in CD8$^+$ MAIT cells. We

hypothesize that the peripheral CD4$^+$ MAIT cells may comprise newly generated cells emigrating

from the thymus(Fink and Hendricks, 2011). Our findings motivate further study of how MAIT

cell activation conditions can select for CD4 or CD8 co-expression in peripheral circulating

populations, and further, how these subsets differentially respond to major pathogens. Indeed,

our TB data demonstrate that CD25, CCR7 and OX40 are significantly upregulated in CD4$^+$ MAIT

cells of TB contacts, suggesting that this subset responds to acute *Mtb* exposure.

The second aim of our study was to distinguish between transcriptional phenotypes induced

during acute and chronic activation and to characterize functional programs associated with

MR1 ligand vs non-specific T cell stimulation. We observed that acutely activated MAIT cells

with either stimulus were driven towards similar states defined by a rapid upregulation of GZMB,

IFNγ and TNFα. At both the transcript and protein level, this signature was more potently

induced by αCD3/CD28, which is possibly explained by non-specific stimulation of all T cells

within PBMCs triggering cytokine release by MAIT and non-MAIT T cells alike(Trickett and Kwan,

2003). αCD3/CD28 may mimic other non-specific T cell stimuli derived from whole bacteria or

cytokines, which have previously been shown to significantly upregulate GZMB and IFNγ in MAIT

cells relative to 5-OP-RU alone(Lamichhane et al., 2019; Suliman et al., 2019). In contrast, acute

activation with 5-OP-RU induced a strong type I interferon-driven gene expression phenotype

along with T cell maturation marker CD52. Co-stimulation of MAIT cells with 5-OP-RU and type

1 interferons have previously been shown to enhance GZMB and IFNγ release(Lamichhane et

al., 2020). Our results demonstrate that acute antigen stimulation alone also enhances MAIT cell

type 1 interferon immunity, which may promote survival and proliferation observed during

chronic stimulation(Welsh et al., 2012) as well as modulate immunity against bacterial or viral

pathogens.(Trinchieri, 2010)

While previous studies have used chronic 5-OP-RU co-stimulation to expand MAIT cells for

downstream use during in vitro functional assays(Boulouis et al., 2020; Gherardin et al., 2018a),

this is the first investigation of the transcriptional landscape of chronic antigen-specific activation

during this proliferative phase. While we observed largely overlapping transcriptional signatures

between 5-OP-RU and αCD3/CD28 activation conditions, 5-OP-RU significantly upregulated

genes associated with T cell development, self-renewal, proliferation and recruitment of

inflammatory cells. In contrast, chronic αCD3/CD28 activation upregulated genes associated

with cytotoxicity, stress and exhaustion. These differences between antigen and mitogen

stimulation were recapitulated by flow cytometry. Taken together, these results demonstrate

that antigen-specific stimulation is required for MAIT cell survival, expansion and sustained

effector function during the chronic phase of stimulation.

Moreover, we describe a FOXP3$^+$ peripheral blood MAIT cell subset enriched in acute activation

cluster C1 with abundant FOXP3 detected at the protein level during chronic 5-OP-RU

stimulation. We observed that peripheral FOXP3$^+$ induction was detected almost exclusively in

the CD8$^+$CD25$^+$IL7R$^-$ subset, in contrast to a recent study(Ling et al., 2016). This surface co-

expression of FOXP3 and CD25 in CD8$^+$ MAIT cells parallels that of CD4$^+$ Treg cells and raises

the question of whether FOXP3 expression in MAIT cells may confer a similar immune regulatory

function(Bhattacharyya et al., 2018).   Importantly, we extend our findings of MAIT cell FOXP3

induction in vitro to immune responses of donors recently exposed to *Mtb*. We found that CD8$^+$

FOXP3$^+$CD25$^+$ MAIT cells accumulated after acute *Mtb* exposure in healthy household TB

contacts and were significantly more abundant in recently infected donors (IGRA+ contacts).

Our results suggest that CD8$^+$ FOXP3$^+$CD25$^+$ MAIT cells respond to acute *Mtb* infection.

In addition to FOXP3 expression, our analysis identifies several alternative immune regulatory

mechanisms mediated by MAIT cells. We observed that activation/exhaustion marker LAG3 was

significantly upregulated in chronically activated MAIT cell clusters (C7, C10, and C12) and was

significantly differentially expressed in CD8$^+$ and DP subsets. LAG3 can bind peptide-MHC II

complexes and inhibit CD4$^+$ T cell activation(Maruhashi et al., 2018). Additionally, MAIT cells

may target CD4$^+$ T cells via MR1 or through secretion of immunomodulatory cytokines like TGFβ.

Future studies will help to elucidate the relative contribution of these immune modulation

pathways on MAIT cell regulatory activity.

Our study also highlights that MAIT cell activation rapidly drives phenotypic and functional

exhaustion. This was evident in both antigen and non-specific stimulation conditions at the

transcriptional and protein level. These findings mirror evidence from chronic disease states

such as TB, viral hepatitis, and malignancy where increased MAIT cell PD1 expression is

associated with peripheral MAIT cell depletion and dysfunction and directly correlates with

severity of disease in some studies (Duan et al., 2019; Hengst et al., 2016; Jiang et al., 2020;

Leeansyah et al., 2013; Saeidi et al., 2015; Shaler et al., 2017; Yong et al., 2018). While immune

checkpoint molecules can be transiently upregulated after T cell activation, we observed

sustained expression in chronic activation clusters. At the protein level, these exhaustion

markers were significantly upregulated in the 5-OP-RU condition, suggesting that chronic

antigen-specific activation strongly induces negative feedback mechanisms to suppress further

activation and proliferation. Taken together, these data imply that checkpoint blockade may

enhance MAIT cell activity against infectious or malignant disease (Boulouis et al., 2020;

Leeansyah et al., 2015; Venken et al., 2018) or contribute to autoimmune complications of

immunotherapy(Sasson et al., 2020).

We acknowledge certain limitations to our study. The foremost of these is the small number of

individuals in our single cell cohort and donor-to-donor variability. However, the functional flow

cytometric validation in multiple donors, including a genetically distinct population from Haiti,

strengthens our confidence that the MAIT cell states observed can be applied to other

populations. Future studies will test the reproducibility of this transcriptional atlas in additional

healthy donors or in specific disease states.

In sum, we present an integrated transcriptional and immune landscape of human MAIT cell

activation at single cell resolution. Our results provide a detailed analysis of novel MAIT cell

clusters defined by acute and chronic stimulation, including distinct CD4$^+$ and CD8$^+$ MAIT cell

signatures. We also show that these phenotypes are recapitulated in the setting of acute *Mtb*

exposure and infection. This functional map of MAIT cell activation biology will help inform future

investigations into how MAIT cells specialize in healthy and disease states.

Acknowledgments

We would like to thank Ojasvi Chaudhury for his assistance in single-cell RNA-seq libraries

preparation. We acknowledge the Immune Monitoring, Flow Cytometry, and Integrated

Genomics Operation Core Facilities at the Sloan Kettering Institute, Memorial Sloan Kettering

Cancer, NY, NY for their expert consultation and services rendered during this project. We also

acknowledge the NIH Tetramer Core (Emory University, Atlanta, GA) for providing MR1

tetramers. This work was supported by the Ludwig Center for Cancer Immunotherapy, the Tri-I

TBRU, part of the TBRU Network (U19 AI111143), and P30 CA008748. L.M. acknowledges

support by The Alan and Sandra Gerry Metastasis and Tumor Ecosystems Center. C.K.V. was

supported by NIH NIAID K08 AI132739.

Conflicts of interest:

M.S.G. reports consulting fees and equity in Vedanta Biosciences, Inc., consulting fees from

Takeda, and is on the SAB of PRL-NYC.

Author Contributions

C.K.V. and M.S.G. conceived of the study. C.K.V. conducted the experiments. J.A. and K.L.

synthesized 5-A-RU. L.M. conducted inDROP scRNA-seq. D.F.W. recruited the Haitian TB

contact cohort. C.K. and C.S.L. analyzed single-cell RNA sequencing data. C.K.V., C.K., C.S.L.

and M.S.G. analyzed wrote the manuscript with input from all authors.

Methods

PBMC collection

Healthy donor PBMCs were obtained with written informed consent at Sloan Kettering Institute,

MSKCC, Rockefeller University or ATCC. De-identified blood samples were processed

according to the protocols approved by the Institutional Review Board of MSKCC and

Rockefeller University. TB household contacts and unexposed community controls were

recruited at GHESKIO centers, Port-au-Prince, Haiti (U01AI069421). PBMCs were isolated from

peripheral blood using Ficoll-Prep (GE Healthcare) or cell preservation tubes (CPT; Becton

Dickinson) according to the manufacturer's instructions and cryopreserved using 90% fetal

bovine serum (Gibco) 10% DMSO (Sigma Aldrich) and stored in liquid N_2 prior to thawing for in

vitro assays.

PBMC culture

PBMCs were cultured for 15 hrs or 7 days in RPMI (GIBCO) supplemented with 10% (v/v) heat

inactivated FBS, penicillin/streptomycin (100 U/mL), L-glutamine (2 mM), sodium pyruvate (1

mM), nonessential amino acids (0.1 mM), HEPES buffer (10 mM) and 2-mercaptoethanol (50 μM)

(from GIBCO or Sigma Aldrich) at 37°C 5% CO2 in 96 well plates in the presence of IL2 (GIBCO

100 IU/mL or Roche 1 ng/mL). Additional conditions included 1 ng/mL TGFβ (GIBCO). 2 μM 5-

A-RU was added directly to 50 μM methylglyoxal to form 5-OP-RU for in vitro stimulation assays.

5-A-RU was synthesized as previously described(Li et al., 2018), resuspended in 200 μM

aliquots, cryopreserved, and thawed as needed. αCD3/CD28 beads (Gibco) were incubated

with PBMCs at a 1:2 bead:cell ratio.

Flow cytometry

Cells were washed once in FACS buffer then incubated with Fc Receptor antibodies

(eBioscience; Fisher catalog # 14-1961-73) for 15 minutes at room temperature prior to staining.

Extracellular staining was performed at room temperature for 15 minutes in 50 μL FACS buffer

(eBioscience) while intracellular staining was performed after 2 hours of incubation in Brefeldin

A at 37°C followed by 45 minutes of incubation in 50 μL Permeabilization/Fixation buffer

(eBioscience), for 1 hour in 50 μl Permeabilization Buffer (eBioscience) at 4°C

The flow cytometric gating strategy to identify MAIT cells using MR1-5-OP-RU tetramers

generated by the NIH tetramer core facility(Corbett et al., 2014; Li et al., 2018) is summarized in

Supplementary Fig. 1. The % of MAIT cells of total CD3$^+$ cells and the absolute number of cells

sorted per condition are presented in Supplementary Tables 1a and b.

Cell staining was performed using the following fluorophore-conjugated antibodies (purchased

from BD, eBioscience or Biolegend; clones in parentheses): GZMB (GB11), ki67 (20Raj1), FoxP3

(PHC101), IFNγ (4S.BS3), TNFα (MAb11), CD25 (BC96), OX40 (ACT35), CD161 (DX12), CD3

(UCHT1), CD8 (SK1), CD4 (SK3), EOMES (WD1928), CTLA4 (BNI3), GZMA (CB9), GZMK

(GM26E7), IL7R (HIL-7R-M21), CD69 (FN50), ICOS (ISA-3), FAS (DX2), LAG3 (11C3C65) and PD1

(EH12.2H7), CCR7 (3D12).

Flow-assisted cell sorting and flow cytometric analysis

Stained cells were sorted on a BD Aria sorter or analyzed on a BD Fortessa analyzer. Flow

cytometry plots and analysis were performed using FCS Express version 7 (De Novo Software,

Pasadena, CA). Graphs and statistical analysis of flow cytometry data were generated using

Prism version 11 software (Graphpad Software, San Diego, CA).

Cell barcoding and scRNA-seq library preparation

The FACS-sorted live-cell suspensions were encapsulated in microfluidic droplets using a

Chromium instrument (10X genomics) and reagents provided in Single Cell 3' Reagent Kit (v3).

The barcoded-cDNA libraries were prepared according to the manual (CG00183 Rev B). The

viability of cells (stained with 0.2% (w/v) Trypan Blue) prior to loading onto the Chromium

instrument was 80-98%. Each sample was encapsulated at a final dilution of ~100 cells/µl.

Following the reverse transcription step the emulsion droplets were broken, barcoded-cDNA

purified with DynaBeads and amplified by 12-cycles of PCR: 98℃ for 180 s, 12x (98℃ for 15 s,

67˚C for 20 s, 72℃ for 60 s), and 72℃ for 60 s. The 50 ng of PCR-amplified barcoded-cDNA was

fragmented with the reagents provided in the kit, purified with SPRI beads and resulting DNA

library was ligated to the sequencing adapter followed by indexing PCR: 98℃ for 45 s; 14x (98℃

for 20 s, 54℃ for 30 s, 72℃ for 20 s), and 72℃ for 60 s. The final DNA library was double-size

purified (0.6–0.8X) with SPRI beads and sequenced on Illumina Nova-Seq platform (R1 – 26

cycles, i7 – 8 cycles, R2 – 70 cycles or higher) at depth of 220 million reads per sample.

Single cell RNA-seq analysis

Pre-processing of scRNA-seq fastq files was conducted using cellranger v3.0.2 (10x genomics).

scRNA-seq reads were aligned to the hg19 reference genome (ref-version 3.0.0), and the count

matrix of cell barcodes by genes used for downstream analysis was generated using the

cellranger count function with parameter –expect-cells=3000. The raw count matrix for each

sample was obtained from the cellranger count filter_matrix output. Briefly, for each sample,

cellranger plots the total UMI count against the cell barcode rank in decreasing order of total

counts, and filters cell barcodes out of the resulting count matrix based on the inflection point of

the plot. This step minimizes the number of empty droplets that are included in downstream

analyses. We created an initial count matrix combining all samples from all patients. Following

count matrix generation, cells with > 20% of transcripts derived from mitochondrial genes were

considered apoptotic, and were thus excluded. Following this step, all mitochondrial genes were

filtered out of the count matrix. Ribosomal genes and the noncoding RNAs NEAT1 and MALAT1

were excluded due to prior reports of strong influence on downstream clustering. Genes with

mean raw count < 3.0 were removed from the analysis, and we restricted to KLRB1+ cells,

resulting in a final count matrix of 76845 cells and xxx genes for downstream analysis. We used

Seurat v2.3.4 to perform standard library size and log normalization. The mean library size was

6680 transcripts per cell.

To examine potential batch effects in the data, we computed the prevalence of each cluster out

of all cells in each sample, followed by a centered log-ratio (CLR) transformation for

compositional data and PCA. This analysis demonstrated that samples clustered by timepoint,

not batch (i.e. donor). Differential expression analyses were conducted using the FindMarkers()

function in Seurat using a Wilcoxon test. Specifically, we compared each cluster to all other

clusters (one vs all), and genes were considered signature genes for a cluster if the log fold

change was greater than 0 and FDR $P <= 0.05$.

References

Bhattacharyya, A., Hanafi, L.A., Sheih, A., Golob, J.L., Srinivasan, S., Boeckh, M.J., Pergam, S.A., Mahmood, S., Baker, K.K., Gooley, T.A*., et al.* (2018). Graft-Derived Reconstitution of Mucosal-Associated Invariant T Cells after Allogeneic Hematopoietic Cell Transplantation. Biol Blood Marrow Transplant *24*, 242-251.

Boulouis, C., Sia, W.R., Gulam, M.Y., Teo, J.Q.M., Png, Y.T., Phan, T.K., Mak, J.Y.W., Fairlie, D.P., Poon, I.K.H., Koh, T.H*., et al.* (2020). Human MAIT cell cytolytic effector proteins synergize to overcome carbapenem resistance in Escherichia coli. PLoS Biol *18*, e3000644.

Corbett, A.J., Eckle, S.B., Birkinshaw, R.W., Liu, L., Patel, O., Mahony, J., Chen, Z., Reantragoon, R., Meehan, B., Cao, H*., et al.* (2014). T-cell activation by transitory neo-antigens derived from distinct microbial pathways. Nature *509*, 361-365.

Coulter, F., Parrish, A., Manning, D., Kampmann, B., Mendy, J., Garand, M., Lewinsohn, D.M., Riley, E.M., and Sutherland, J.S. (2017). IL-17 Production from T Helper 17, Mucosal-Associated Invariant T, and gammadelta Cells in Tuberculosis Infection and Disease. Front Immunol *8*, 1252.

Dias, J., Boulouis, C., Gorin, J.B., van den Biggelaar, R., Lal, K.G., Gibbs, A., Loh, L., Gulam, M.Y., Sia, W.R., Bari, S*., et al.* (2018). The CD4(-)CD8(-) MAIT cell subpopulation is a functionally distinct subset developmentally related to the main CD8(+) MAIT cell pool. Proc Natl Acad Sci U S A *115*, E11513-E11522.

Dias, J., Leeansyah, E., and Sandberg, J.K. (2017). Multiple layers of heterogeneity and subset diversity in human MAIT cell responses to distinct microorganisms and to innate cytokines. Proc Natl Acad Sci U S A *114*, E5434-E5443.

Duan, M., Goswami, S., Shi, J.Y., Wu, L.J., Wang, X.Y., Ma, J.Q., Zhang, Z., Shi, Y., Ma, L.J., Zhang, S*., et al.* (2019). Activated and Exhausted MAIT Cells Foster Disease Progression and Indicate Poor Outcome in Hepatocellular Carcinoma. Clin Cancer Res *25*, 3304-3316.

Dusseaux, M., Martin, E., Serriari, N., Peguillet, I., Premel, V., Louis, D., Milder, M., Le Bourhis, L., Soudais, C., Treiner, E., and Lantz, O. (2011). Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17-secreting T cells. Blood *117*, 1250-1259.

Estevez, O., Anibarro, L., Garet, E., Martinez, A., Pena, A., Barcia, L., Peleteiro, M., and Gonzalez-Fernandez, A. (2020). Multi-parameter flow cytometry immunophenotyping distinguishes different stages of tuberculosis infection. J Infect.

Fink, P.J., and Hendricks, D.W. (2011). Post-thymic maturation: young T cells assert their individuality. Nat Rev Immunol *11*, 544-549.

Gherardin, N.A., Loh, L., Admojo, L., Davenport, A.J., Richardson, K., Rogers, A., Darcy, P.K., Jenkins, M.R., Prince, H.M., Harrison, S.J*., et al.* (2018a). Enumeration, functional responses and cytotoxic capacity of MAIT cells in newly diagnosed and relapsed multiple myeloma. Sci Rep *8*, 4159.

Gherardin, N.A., Souter, M.N., Koay, H.F., Mangas, K.M., Seemann, T., Stinear, T.P., Eckle, S.B., Berzins, S.P., d'Udekem, Y., Konstantinov, I.E*., et al.* (2018b). Human blood MAIT cell subsets defined using MR1 tetramers. Immunol Cell Biol *96*, 507-525.

Godfrey, D.I., Koay, H.F., McCluskey, J., and Gherardin, N.A. (2019). The biology and functional importance of MAIT cells. Nat Immunol.

Harriff, M.J., McMurtrey, C., Froyd, C.A., Jin, H., Cansler, M., Null, M., Worley, A., Meermeier,
E.W., Swarbrick, G., Nilsen, A., *et al.* (2018). MR1 displays the microbial metabolome driving
selective MR1-restricted T cell receptor usage. Sci Immunol *3*.
Hengst, J., Strunz, B., Deterding, K., Ljunggren, H.G., Leeansyah, E., Manns, M.P., Cornberg,
M., Sandberg, J.K., Wedemeyer, H., and Bjorkstrom, N.K. (2016). Nonreversible MAIT cell-
dysfunction in chronic hepatitis C virus infection despite successful interferon-free therapy. Eur J
Immunol *46*, 2204-2210.
Hoffmann, P., Eder, R., Kunz-Schughart, L.A., Andreesen, R., and Edinger, M. (2004). Large-
scale in vitro expansion of polyclonal human CD4(+)CD25high regulatory T cells. Blood *104*,
895-903.
Jahreis, S., Bottcher, S., Hartung, S., Rachow, T., Rummler, S., Dietl, A.M., Haas, H., Walther,
G., Hochhaus, A., and von Lilienfeld-Toal, M. (2018). Human MAIT cells are rapidly activated by
Aspergillus spp. in an APC-dependent manner. Eur J Immunol *48*, 1698-1706.
Jiang, J., Cao, Z., Qu, J., Liu, H., Han, H., and Cheng, X. (2020). PD-1-expressing MAIT cells
from patients with tuberculosis exhibit elevated production of CXCL13. Scand J Immunol *91*,
e12858.
Kelly, J., Minoda, Y., Meredith, T., Cameron, G., Philipp, M.S., Pellicci, D.G., Corbett, A.J.,
Kurts, C., Gray, D.H., Godfrey, D.I., *et al.* (2019). Chronically stimulated human MAIT cells are
unexpectedly potent IL-13 producers. Immunol Cell Biol *97*, 689-699.
Koay, H.F., Su, S., Amann-Zalcenstein, D., Daley, S.R., Comerford, I., Miosge, L., Whyte, C.E.,
Konstantinov, I.E., d'Udekem, Y., Baldwin, T., *et al.* (2019). A divergent transcriptional
landscape underpins the development and functional branching of MAIT cells. Sci Immunol *4*.
Kurioka, A., Jahun, A.S., Hannaway, R.F., Walker, L.J., Fergusson, J.R., Sverremark-Ekstrom,
E., Corbett, A.J., Ussher, J.E., Willberg, C.B., and Klenerman, P. (2017). Shared and Distinct
Phenotypes and Functions of Human CD161++ Valpha7.2+ T Cell Subsets. Front Immunol *8*,
1031.
Lamichhane, R., Galvin, H., Hannaway, R.F., de la Harpe, S.M., Munro, F., Tyndall, J.D.,
Vernall, A.J., McCall, J.L., Husain, M., and Ussher, J.E. (2020). Type I interferons are important
co-stimulatory signals during T cell receptor mediated human MAIT cell activation. Eur J
Immunol *50*, 178-191.
Lamichhane, R., Schneider, M., de la Harpe, S.M., Harrop, T.W.R., Hannaway, R.F., Dearden,
P.K., Kirman, J.R., Tyndall, J.D.A., Vernall, A.J., and Ussher, J.E. (2019). TCR- or Cytokine-
Activated CD8(+) Mucosal-Associated Invariant T Cells Are Rapid Polyfunctional Effectors That
Can Coordinate Immune Responses. Cell Rep *28*, 3061-3076 e3065.
Leeansyah, E., Ganesh, A., Quigley, M.F., Sonnerborg, A., Andersson, J., Hunt, P.W.,
Somsouk, M., Deeks, S.G., Martin, J.N., Moll, M., *et al.* (2013). Activation, exhaustion, and
persistent decline of the antimicrobial MR1-restricted MAIT-cell population in chronic HIV-1
infection. Blood *121*, 1124-1135.
Leeansyah, E., Svard, J., Dias, J., Buggert, M., Nystrom, J., Quigley, M.F., Moll, M.,
Sonnerborg, A., Nowak, P., and Sandberg, J.K. (2015). Arming of MAIT Cell Cytolytic
Antimicrobial Activity Is Induced by IL-7 and Defective in HIV-1 Infection. PLoS Pathog *11*,
e1005072.
Legoux, F., Bellet, D., Daviaud, C., El Morr, Y., Darbois, A., Niort, K., Procopio, E., Salou, M.,
Gilet, J., Ryffel, B., *et al.* (2019). Microbial metabolites control the thymic development of
mucosal-associated invariant T cells. Science *366*, 494-499.
Leng, T., Akther, H.D., Hackstein, C.P., Powell, K., King, T., Friedrich, M., Christoforidou, Z.,
McCuaig, S., Neyazi, M., Arancibia-Carcamo, C.V., *et al.* (2019). TCR and Inflammatory Signals
Tune Human MAIT Cells to Exert Specific Tissue Repair and Effector Functions. Cell Rep *28*,
3077-3091 e3075.

Li, K., Vorkas, C.K., Chaudhry, A., Bell, D.L., Willis, R.A., Rudensky, A., Altman, J.D., Glickman,
M.S., and Aube, J. (2018). Synthesis, stabilization, and characterization of the MR1 ligand
precursor 5-amino-6-D-ribitylaminouracil (5-A-RU). PLoS One *13*, e0191837.
Ling, L., Lin, Y., Zheng, W., Hong, S., Tang, X., Zhao, P., Li, M., Ni, J., Li, C., Wang, L., and
Jiang, Y. (2016). Circulating and tumor-infiltrating mucosal associated invariant T (MAIT) cells in
colorectal cancer patients. Sci Rep *6*, 20358.
Liston, A., Nutsch, K.M., Farr, A.G., Lund, J.M., Rasmussen, J.P., Koni, P.A., and Rudensky,
A.Y. (2008). Differentiation of regulatory Foxp3+ T cells in the thymic cortex. Proc Natl Acad Sci
U S A *105*, 11903-11908.
Maruhashi, T., Okazaki, I.M., Sugiura, D., Takahashi, S., Maeda, T.K., Shimizu, K., and
Okazaki, T. (2018). LAG-3 inhibits the activation of CD4(+) T cells that recognize stable pMHCII
through its conformation-dependent recognition of pMHCII. Nat Immunol *19*, 1415-1426.
Rouxel, O., Da Silva, J., Beaudoin, L., Nel, I., Tard, C., Cagninacci, L., Kiaf, B., Oshima, M.,
Diedisheim, M., Salou, M.*, et al.* (2017). Cytotoxic and regulatory roles of mucosal-associated
invariant T cells in type 1 diabetes. Nat Immunol *18*, 1321-1331.
Saeidi, A., Tien Tien, V.L., Al-Batran, R., Al-Darraji, H.A., Tan, H.Y., Yong, Y.K.,
Ponnampalavanar, S., Barathan, M., Rukumani, D.V., Ansari, A.W.*, et al.* (2015). Attrition of
TCR Va7.2+CD161++ MAIT cells in HIV-tuberculosis co-infection is associated with elevated
levels of PD-1 expression. PLoS ONE *10*, 1-14.
Sasson, S.C., Zaunders, J.J., Nahar, K., Munier, C.M.L., Fairfax, B.P., Olsson-Brown, A., Jolly,
C., Read, S.A., Ahlenstiel, G., Palendira, U.*, et al.* (2020). Mucosal-associated invariant T
(MAIT) cells are activated in the gastrointestinal tissue of patients with combination ipilimumab
and nivolumab therapy-related colitis in a pathology distinct from ulcerative colitis. Clin Exp
Immunol.
Seshadri, C., Thuong, N.T., Mai, N.T., Bang, N.D., Chau, T.T., Lewinsohn, D.M., Thwaites,
G.E., Dunstan, S.J., and Hawn, T.R. (2017). A polymorphism in human MR1 is associated with
mRNA expression and susceptibility to tuberculosis. Genes Immun *18*, 8-14.
Shaler, C.R., Choi, J., Rudak, P.T., Memarnejadian, A., Szabo, P.A., Tun-Abraham, M.E.,
Rossjohn, J., Corbett, A.J., McCluskey, J., McCormick, J.K.*, et al.* (2017). MAIT cells launch a
rapid, robust and distinct hyperinflammatory response to bacterial superantigens and quickly
acquire an anergic phenotype that impedes their cognate antimicrobial function: Defining a
novel mechanism of superantigen-induced immunopathology and immunosuppression. PLoS
Biol *15*, e2001930.
Slichter, C.K., McDavid, A., Miller, H.W., Finak, G., Seymour, B.J., McNevin, J.P., Diaz, G.,
Czartoski, J.L., McElrath, M.J., Gottardo, R., and Prlic, M. (2016). Distinct activation thresholds
of human conventional and innate-like memory T cells. JCI Insight *1*.
Suliman, S., Murphy, M., Musvosvi, M., Gela, A., Meermeier, E.W., Geldenhuys, H., Hopley, C.,
Toefy, A., Bilek, N., Veldsman, A.*, et al.* (2019). MR1-Independent Activation of Human
Mucosal-Associated Invariant T Cells by Mycobacteria. J Immunol *203*, 2917-2927.
Treiner, E., Duban, L., Bahram, S., Radosavljevic, M., Wanner, V., Tilloy, F., Affaticati, P.,
Gilfillan, S., and Lantz, O. (2003). Selection of evolutionarily conserved mucosal-associated
invariant T cells by MR1. Nature *422*, 164-169.
Trickett, A., and Kwan, Y.L. (2003). T cell stimulation and expansion using anti-CD3/CD28
beads. J Immunol Methods *275*, 251-255.
Trinchieri, G. (2010). Type I interferon: friend or foe? J Exp Med *207*, 2053-2063.
van Wilgenburg, B., Scherwitzl, I., Hutchinson, E.C., Leng, T., Kurioka, A., Kulicke, C., de Lara,
C., Cole, S., Vasanawathana, S., Limpitikul, W.*, et al.* (2016). MAIT cells are activated during
human viral infections. Nat Commun *7*, 11653.

Venken, K., Favreau, M., Gaublomme, D., Menu, E., Vanderkerken, K., and Elewaut, D. (2018).
Checkpoint inhibition in the treatment of multiple myeloma: A way to boost innate-like T cell anti-
tumor function? Mol Immunol *101*, 521-526.
Vorkas, C.K., Wipperman, M.F., Li, K., Bean, J., Bhattarai, S.K., Adamow, M., Wong, P., Aube,
J., Juste, M.A.J., Bucci, V*., et al.* (2018). Mucosal-associated invariant and gammadelta T cell
subsets respond to initial Mycobacterium tuberculosis infection. JCI Insight *3*.
Welsh, R.M., Bahl, K., Marshall, H.D., and Urban, S.L. (2012). Type 1 interferons and antiviral
CD8 T-cell responses. PLoS Pathog *8*, e1002352.
Yamaguchi, T., Wing, J.B., and Sakaguchi, S. (2011). Two modes of immune suppression by
Foxp3(+) regulatory T cells under inflammatory or non-inflammatory conditions. Semin Immunol
23, 424-430.
Yong, Y.K., Saeidi, A., Tan, H.Y., Rosmawati, M., Enstrom, P.F., Batran, R.A., Vasuki, V.,
Chattopadhyay, I., Murugesan, A., Vignesh, R*., et al.* (2018). Hyper-Expression of PD-1 Is
Associated with the Levels of Exhausted and Dysfunctional Phenotypes of Circulating
CD161(++)TCR iValpha7.2(+) Mucosal-Associated Invariant T Cells in Chronic Hepatitis B Virus
Infection. Front Immunol *9*, 472.
671
672

Fig. 1

Figure 1. MAIT cell CD4 or CD8 expression are associated with distinct transcriptional signatures

(a) Experimental schematic of the MAIT cell in vitro activation assay used for single cell RNA sequencing (scRNA-seq) and flow cytometric analyses. A total of 30 healthy donor PBMCs were assayed by flow cytometry after 15 hrs (D1; acute activation) or 7 days (D7; chronic activation) of incubation under resting, 5OPRU or αCD3/CD28 conditions. A subset of three of these donors underwent flow cytometry-assisted cell sorting (FACS) for scRNA-seq under the same conditions. All conditions included IL2 and a subset of D7 conditions also included TGFβ. **(b)** Differentially expressed genes in CD8$^+$ (+ log FC; log fold change) or CD4$^+$ MAIT cells (- log FC) after 15 hours of incubation in the resting condition. y axis: log FDR-adjusted p value with a significance level of FDR < 0.05. **(c)** Mean% +/- SEM of CD25, OX40, GZMB, and EOMES staining in CD4$^+$, CD8$^+$, DN or DP MAIT cell subsets measured by flow cytometry in the same condition as (b). Data representative of 3 independent experiments. n=5-30 donors. **(d)** Differentially expressed genes in CD8$^+$ or CD4$^+$ MAIT cells after 7 days of incubation in the resting condition. Same axes as (b). **(e)** Mean% +/- SEM of OX40, CTLA4, GZMA, and GZMK staining in MAIT cell subsets measured by flow cytometry in the same condition as (d). Data representative of 2 independent experiments. n=5-10 donors. Flow cytometric statistical comparisons were made by unpaired t-test. *p<0.05 **p<0.005 *** p<0.0005 ****p<0.0001; GZMB: granzyme B; GZMA: granzyme A; GZMK: granzyme K; DN: double-negative (CD4$^-$CD8); DP: double-positive (CD4$^+$CD8$^+$)

Fig. 2

Figure 2. Single cell transcriptional analysis defines heterogeneous MAIT cell activation states.

(a) Uniform manifold approximation and projection (UMAP) plot and Louvain clustering of cells from all donors, timepoints and stimuli with ascribed names based on differentially expressed gene sets in each cluster. **(b)** Heat map showing top 15 differentially expressed genes per cluster after unsupervised hierarchical clustering. **(c)** UMAP plots of selected MAIT cell markers demonstrating expression level across all clusters (blue=low expression to yellow=high expression). **(d, e)** Dot plots demonstrating relative frequency of cells per condition in all donors.

Fig. 3

b IL7R KLRB1 CD69 IL2RA

GZMB IFNG TNF FOXP3

c
- Resting
- 5OPRU
- αCD3/CD28

%marker of MAIT

IL7R⁺   CD69⁺⁺   CD25⁺

d Gated on MAIT cells

REST   5OPRU   αCD3/CD28

GZMB-FITC / IFNG-PE

REST: 0.20% / 0.00% / 99.66% / 0.14%
5OPRU: 11.59% / 0.97% / 83.57% / 3.86%
αCD3/CD28: 38.83% / 5.74% / 52.40% / 3.03%

- REST
- 5OPRU
- aCD3/CD28

%subset of MAIT
GZMB⁺IFNG⁻ GZMB⁺IFNG⁺ GZMB⁻IFNG⁺

GZMB-FITC / TNF-BV605

REST: 0.17% / 0.00% / 99.78% / 0.06%
5OPRU: 10.63% / 0.00% / 86.96% / 2.42%
αCD3/CD28: 37.47% / 5.74% / 53.13% / 3.65%

%subset of MAIT
GZMB⁺TNF⁻ GZMB⁺TNF⁺ GZMB⁻TNF⁺

IFNG-PE / TNF-BV605

REST: 0.08% / 0.00% / 99.83% / 0.08%
5OPRU: 3.38% / 0.48% / 91.79% / 4.35%
αCD3/CD28: 5.64% / 1.98% / 83.30% / 9.08%

%subset of MAIT
IFNG⁺TNF⁻ IFNG⁺TNF⁺ IFNG⁻TNF⁺

e
-log10(p adj)

IFIT1 BATF3 GZMB IFNG
IL2RA
ISG15 PRF1
IFI44L
OAS1
IFIT3
IFIT2 MX1 ICOS
IL7R
IFITM1 CTLA4
IFI6
LY6E LAG3
GZMA

5OPRU   logFC   αCD3/CD28

Figure 3. Acute MAIT cell activation induces a cytotoxic phenotype marked by GZMB,

IFNG, and TNF

(a) UMAP plot highlighting clusters 1-3 enriched in day 1 (D1) incubation conditions. **(b)** UMAP

plots of selected MAIT cell marker expression levels at D1 (blue=low expression to yellow=high

expression). **(c)** Mean% +/- SEM of MAIT cell extracellular staining of activation markers

detected by flow cytometry after 15 hrs of incubation in three conditions: resting (blue), 5OPRU

(red), αCD3/CD28 (brown). Results are representative of 4 independent experiments: n=3-30

donors. **(d)** Representative flow cytometry density plots in one donor demonstrating GZMB,

TNFα and IFNγ co-staining (left) alongside cumulative plots (right) of mean% +/- SEM for each

co-staining condition in multiple donors. Results are representative of 4 independent

experiments. n=30 donors. **(e)** Differentially expressed genes in the αCD3/CD28 (+ log fold

change; log FC) or 5OPRU (- log FC) conditions at D1. y axis: log FDR-adjusted p value at

significance level of FDR $P < 0.05$. Flow cytometric statistical comparisons were made by

unpaired t-test. $^*p<0.05$ $^{**}p<0.005$ *** $p<0.0005$ $^{****}p<0.0001$; GZMB: granzyme B

Fig. 4

Figure 4. MAIT cells upregulate FoxP3 during acute and chronic activation

(a) Representative flow cytometry contour plots of MAIT cell FOXP3 intracellular staining in

resting or 5OPRU conditions after 1, 3, or 7 days of stimulation (black=resting; red=5OPRU). **(b)**

Mean% +/- SEM of intracellular MAIT cell FOXP3 staining after 1 (D1; acute) or 7 (D7; chronic)

days of incubation under resting (blue), 5OPRU (red), or αCD3/CD28 (brown) conditions. Data

representative of 3 independent experiments.   n=10-26 donors.   **(c)** Mean% +/- SEM of

intracellular FOXP3 staining in CD4$^+$ and CD8$^+$ MAIT cells compared to non-MAIT CD4$^+$ T cells

after 7 days of incubation in the same conditions as (b). **(d)** Representative flow cytometry

density plots from one donor demonstrating FOXP3$^+$CD25$^+$ MAIT cells (top row) or non-MAIT

CD4$^+$ T cells (bottom row) after 7 days of incubation in the same conditions as (b). **(e)** Mean%

+/- SEM of FOXP3$^+$CD25$^+$ CD4$^+$ or CD8$^+$ MAIT cells compared to non-MAIT CD4$^+$ T cells after 7

days in the same conditions as (b).   Cytokine co-incubation conditions included IL2 (left) or

IL2/TGFβ (right).   Representative of two independent experiments.   n=10 donors.   **(f)** Mean

absolute number +/- SEM of FOXP3$^+$CD25$^+$ CD4$^+$ or CD8$^+$ MAIT cells in the same incubation

conditions as (e). Representative of 1 independent experiment. n=5 donors. **(g)** Mean absolute

number +/- SEM of FOXP3$^+$ MAIT cells among CD25$^+$IL7R$^-$ MAIT cells or CD25$^-$ MAIT cells after

7 days in the same incubation conditions as (e).

Fig. 5

Figure 5. Chronic activation of MAIT cells expands homeostatic subpopulations while also

inducing proliferative, cytotoxic, immune regulatory and exhaustion phenotypes

(a) UMAP highlighting clusters 4-12 defined by day 7 (D7) incubation conditions. **(b)** UMAP plots

of selected MAIT cell marker expression levels at D7 (blue=low expression to yellow= high

expression). **(c)** Representative flow cytometry contour plots (left) of intracellular ki67 staining

after 15 hours (D1; acute) or 7 days (D7; chronic) of incubation at rest (black) or with 5OPRU (red)

accompanied by cumulative data of mean% +/- SEM of MAIT cell ki67 staining after 7 days of

incubation under resting (blue), 5OPRU (red) or αCD3/CD28 (brown) conditions (right). Data is

representative of one independent experiment. n=3 donors. **(d)** Mean% +/- SEM of MAIT cells

among total T cells after 7 days of incubation in the same conditions as (c). Data is representative

of two independent experiments. n=10 donors. **(e)** Mean% +/- SEM of MAIT cell surface

activation markers after 7 days of incubation in the same conditions as (c). Data is representative

of two independent experiments. n=3-8 donors. **(f)** GZMB, IFNγ, and TNFα intracellular co-

staining demonstrating mean% +/- SEM of each subset after 7 days of incubation under the

same conditions as (c). Data is representative of two independent experiments. n=8 donors. **(g)**

Mean% +/- SEM of activation/exhaustion marker staining after 7 days of incubation under the

same conditions as (c). Data is representative of one independent experiment. n=5 donors.

Data representative of one independent experiment. n=3 donors. Flow cytometric statistical

comparisons were made by unpaired t-test. $*p<0.05$ $**p<0.005$ $***$ $p<0.0005$ $****p<0.0001$;

GZMB: granzyme B

Fig. 6

Figure 6. MAIT cell subsets differentially respond to acute and chronic activation by

Mycobacterium tuberculosis

(a-c) Mean% +/- SEM of CD25 (a), CCR7 (b), and OX40 (c) staining of CD4$^+$, CD8$^+$ or DN MAIT

cell subsets in healthy household contacts of TB patients (red triangles) compared to healthy

unexposed community controls (blue circles). Same legend also applies to panels (d) and (e).

n=18-41 donors. **(d)** Mean% +/- SEM of FOXP3$^+$ (left) or FOXP3$^+$CD25$^+$ (right) MAIT cells. n=36-

42 donors. **(e)** Mean% +/- SEM of FOXP3$^+$CD25$^+$ CD4$^+$ (left) or CD8$^+$ MAIT cells (right) in the

same donors as (d). **(f)** Mean% +/- SEM of FOXP3$^+$CD25$^+$ CD4$^+$ or C8$^+$ MAIT cells stratified by

IGRA status in the same donors as (d). Flow cytometric statistical comparisons were made by

unpaired t-test. *p<0.05 **p<0.005 *** p<0.0005 ****p<0.0001; IGRA: *Mtb*-specific IFNγ release

assay.

Supplementary Figure 1

763 **Supplementary Figure 1. Flow cytometric gating strategy for MAIT cell identification.**

765 **(a)** Flow cytometric gating strategy to identify MAIT cells as live, CD3$^+$, MR1-5OPRU tetramer$^+$,

766 CD161^{++} cells after 15 hrs of overnight rest from thaw in one donor. **(b)** MAIT cell expansion

767 within the CD3$^+$ T cell subset after 7 days of 5OPRU incubation.

a

MAIT cells sorted per condition

Condition	Cytokine	Donor 1	Donor 2	Donor 3
D1_REST	IL2	11,351	26,751	37,295
D1_5OPRU	IL2	11,500	25,023	13,160
D1_αCD23/CD28	IL2	7,050	15,163	33,642
D7_REST	IL2	7,236	21,201	100,000
D7_5OPRU	IL2	53,097	100,000	100,000
D7_αCD23/CD28	IL2	4,087	20,967	100,000
D7_REST	IL2/TGFβ	12,020	19,292	100,000
D7_5OPRU	IL2/TGFβ	30,900	100,000	100,000
D7_αCD23/CD28	IL2/TGFβ	4,056	15,268	75,172

b

%MAIT cells of T cells per condition

Condition	Cytokine	Donor 1	Donor 2	Donor 3
D1_REST	IL2	0.4	0.4	2.8
D1_5OPRU	IL2	0.2	0.3	0.3
D1_αCD23/CD28	IL2	0.4	0.3	1.1
D7_REST	IL2	1.0	0.5	2.6
D7_5OPRU	IL2	5.1	19.2	22.5
D7_αCD23/CD28	IL2	0.4	0.5	3.8
D7_REST	IL2/TGFβ	1.2	0.6	2.9
D7_5OPRU	IL2/TGFβ	2.7	9.1	10.9
D7_αCD23/CD28	IL2/TGFβ	0.3	0.5	2.9

774 **Supplementary Table 1. Table of MAIT cell absolute numbers and % of CD3$^+$ T cells sorted.**

775 Sorted MAIT cells per condition expressed as **(a)** absolute number and **(b)** % of total CD3$^+$ T

776 cells.

Supplemental Figure 2

a

b

c

769 **Supplementary Figure 2. Single cell cluster prevalence per donor and timepoint**

771 **(a)** PCA of CLR-transformed cluster prevalence. Data show broad clustering by timepoint. **(b)**

772 Cluster prevalence per donor and stimulation.

Supplementary Table 2. Differential expression gene analysis of CD4⁺ versus CD8⁺ MAIT

cells in the resting condition at D1 and D7, with all donors pooled together and for

individual donors

Supplementary Table 3. Differentially expressed genes for each cluster presented in

Figure 2.

Supplementary Table 4. Differential expression gene analysis of MAIT cells stimulated with

5OPRU or aCD3/CD28 at D1 and D7, with all donors pooled together and for individual

donors.