Alleloscope: Integrative single cell analysis of allele-specific copy number alterations and chromatin accessibility in cancer

Chi-Yun Wu[1,2], Billy T. Lau[3,4], Heon Seok Kim[3], Anuja Sathe[3], Susan M. Grimes[4], *Hanlee P. Ji[3,4], *Nancy R. Zhang[1,2]

Institutions

[1]Graduate Group in Genomics and Computational Biology, University of Pennsylvania, Philadelphia, PA

[2]Department of Statistics, University of Pennsylvania, Philadelphia, PA

[3]Division of Oncology, Department of Medicine, Stanford University School of Medicine, Stanford, CA

[4]Stanford Genome Technology Center, Stanford University, Palo Alto, CA

*Corresponding author

Corresponding authors

Hanlee P. Ji

Email: genomics_ji@stanford.edu

Nancy R. Zhang

Email: nzh@wharton.upenn.edu

Abstract

Cancer progression is driven by both somatic copy number aberrations (CNAs) and chromatin remodeling, yet little is known about the interplay between these two classes of events in shaping the clonal diversity of cancers. We present Alleloscope, a method for allele-specific copy number estimation that can be applied to single cell DNA and ATAC sequencing data, either separately or in combination. This approach allows for integrative multi-omic analysis of allele-specific copy number and chromatin accessibility on the same cell. On scDNA-seq data from gastric, colorectal, and breast cancer samples, with extensive validation using matched linked-read sequencing, Alleloscope finds pervasive occurrence of highly complex, multi-allelic copy number aberrations, where cells that carry varying allelic configurations adding to the same total copy number co-evolve within a tumor. The contributions of such allele-specific events to intratumor heterogeneity have been under-reported and under-studied due to the lack of methods for their detection. On scATAC-seq from two basal cell carcinoma samples and a gastric cancer cell line, Alleloscope detects multi-allelic copy number events and copy neutral loss-of-heterozygosity, enabling the dissection of the contributions of chromosomal instability and chromatin remodeling in tumor evolution.

Introduction

Cancer is a disease caused by genetic alterations and epigenetic modifications which, in combination, shape the dysregulated transcriptional programming of tumor cells[1, 2]. These somatic genomic events lead to a diverse cellular population from which clones with advantageous alterations proliferate and eventually metastasize[3]. The comprehensive study of cancer requires the integrative profiling of genetic and epigenetic changes at the resolution of single cells. We combined the analysis of two such genomic dimensions – DNA copy number and chromatin accessibility – through massively parallel single cell sequencing assays.

First, consider copy number aberrations (CNAs), through which we have derived much of our current understanding of the relationship between genome instability and tumor evolution[4]. Total copy number profiling, which estimates the sum of the copy numbers of the two homologous chromosomes, is inadequate to characterize some types of cancer genomic aberrations. Such events include the pervasive copy-neutral losses of heterozygosity (LOH)[5-8] and the intriguing "mirrored events"[9, 10], where cells carrying amplification of one haplotype are intermingled with cells carrying amplification of the other haplotype. While the importance of allele-specific copy number profiling has been emphasized in bulk DNA sequencing analysis[5-8, 11], most single-cell studies profile only total copy number due to the difficulty of reliable allele-specific estimation at low per-cell coverage[12-19]. Recently, a major advance was made by Zaccaria and Raphael in the development of CHISEL[10], a method for single-cell allele-specific copy number analysis. By default, CHISEL relies on externally phased haplotypes derived from available reference cohorts for accurate estimation of allele-specific CNAs, and found evidence for

the existence of mirrored amplifications in a deeply sequenced breast tumor. This was

the first time that allele-specificity of CNA events were reported at single-cell resolution,

but the requirements for high sequencing depth and external phasing limit the general

applicability of CHISEL. Thus, despite these recent advances, the prevalence of mirrored

amplification events and the general complexity of CNA regions at single-cell allele-

specific resolution are still largely unexplored.

Epigenetic modifications are also an important genomic feature of cancer. One type of

epigenetic profiling is the measurement of chromatin openness, for which a variety of

methods have been developed[20] including transposase-accessible chromatin sequencing

(ATAC-seq)[21, 22], which can be performed at the bulk-tissue or single-cell level. Analysis

of chromatin structure has shown that epigenetic remodeling modulates the plasticity of

cells in cancer[23-27], leads to stem-like properties[28-30] and generates therapeutic

resistance[31-34]. Since copy number alterations involve large gains and losses of chromatin,

we expect the chromatin accessibility of a region to be influenced by the changes in

underlying copy number. Current scATAC-seq studies estimate total copy number profiles

by smoothing the read coverage and comparing position-wise against a control cell

population[24, 26]. Yet this appropriate control is difficult to identify, as even between normal

cells of distinct lineages, broad chromatin remodeling can lead to chromosome-level shifts

in coverage (examples of this will be given in Results). There is yet no method for reliable

total or allele-specific copy number profiling in scATAC-seq data, and thus, the estimation

of CNA and chromatin accessibility are confounded in current analysis pipelines.

Addressing these challenges, we present Alleloscope, a method for **allel**e-**s**pecific **cop**y

number **e**stimation and multiomic profiling in single cells. Alleloscope does not rely on

external phasing information, and can be applied to low coverage scDNA-seq data or to

scATAC-seq data with sample-matched bulk DNA sequencing data. To investigate the

single cell landscape of allele-specific CNA, we first apply Alleloscope on scDNAseq data

from four gastric cancer samples, four colorectal cancer samples, and two datasets from

different regions of a breast cancer sample[10, 12, 35]. For five of the gastrointestinal cancer

samples, results are extensively validated by 10x linked-read sequencing which provides

accurate phasing information[36-38]. In these datasets, Alleloscope accurately identifies

LOH and mirrored-subclonal amplification events, and furthermore, finds pervasive

occurrence of highly complex multi-allelic loci where cells that carry varying allelic

configurations adding to the same total copy number co-evolve within a tumor. The

ubiquity of such events in all three cancer types analyzed reveal that they may be an

important overlooked source of intratumor genetic heterogeneity.

Having characterized the complexity of allele-specific CNA events at single cell resolution,

we turn to scATAC-seq data from two basal cell carcinoma samples with paired bulk

whole exome sequencing data[26] and a complex polyclonal gastric cancer cell line that we

also analyzed by scDNA-seq. In these samples, we evaluate the accuracy of Alleloscope

in genotyping and clone assignment for scATAC-seq data and demonstrate its application

to the integrative analysis of CNA and chromatin accessibility.

Results

Overview of Alleloscope model and algorithm

First, we briefly overview Alleloscope's method for allele-specific copy number estimation

(Fig. 1). Clone assignment and integration with peak signals in scATAC-seq data will be

described later. Alleloscope relies on two types of data features: coverage, derived from

all reads that map to a given region, and allelic imbalance, derived from allele-informative

reads that cover heterozygous loci in the region. We start with some essential definitions.

For a given single nucleotide polymorphism (SNP) site, we refer to its mean coverage

across cells as *bulk coverage* and its mean variant allele frequency (VAF = ratio of

alternative allele read count to total read count) across cells as its *bulk VAF*. Between the

two parental haplotypes, we define the term "major haplotype" as the haplotype with

higher mean count across cells. Note that a haplotype may be the "major haplotype" of a

sample, but have lesser copy number than the other haplotype within some cells. For

each cell i, in any given CNA region r, we define two key parameters: (1) the major

haplotype proportion (θ_{ir}), defined as the count of the copies of the major haplotype

divided by the total copy number for the region in cell i and (2) total copy fold change

(ρ_{ir}), defined as the ratio of the total copy number of the region in cell i relative to that in

normal cells.

The algorithm starts by segmenting the genome into regions of homogeneous allele-

specific copy number using both the bulk coverage and bulk VAF profiles (Fig.1 Step 2).

This can be achieved by multiple existing algorithms, which may be combined to increase

detection sensitivity, see Methods for details. In our analyses of scATAC-seq data, the

segmentation was performed on the matched bulk or single cell DNA-seq data, which

ensures that the CNA regions passed to the characterization step are supported by

evidence from DNA and thus are not simply due to the broad chromatin remodeling that

occur in cancer.

Now consider each putative CNA region, indexed by r. An expectation-maximization (EM)

based algorithm is used to iteratively phase each SNP and estimate the major haplotype

proportion (θ_{ir}) for each cell (Fig. 1, step 3). For each SNP j lying in the region, let $I_j \in$

{0,1} be the indicator of whether the reference allele of SNP j is carried by the major

haplotype (please see the toy example in Fig. 1). An initial estimate $\hat{I}_j^{(0)}$ is first derived

from the bulk VAF profile. Then, in iteration t, Alleloscope computes $\hat{\theta}_{ir}^{(t)}$ by pooling

counts across sites within the region r, weighted by the current phasing $\hat{I}_j^{(t)}$, then updates

the estimate of I_j based on $\hat{\theta}_{ir}^{(t)}$ by pooling counts across cells. The estimates of θ_{ir} and

I_j usually converge within a few iterations as described in the Methods. If matched

scDNA-seq data are available for a sample sequenced by scATAC-seq, I_j values can be

estimated from scDNA-seq and then used to compute θ_{ir} for each cell in the scATAC-

seq data, enabling more information sharing between the two data types.

The estimated major haplotype proportions ($\hat{\theta}_{ir}$'s), along with a preliminarily normalized

coverage statistic ($\tilde{\rho}_{ir}$'s, see Methods for details), are then used to identify a set of normal

cells and diploid regions (Fig. 1, Step 4). This information is then used to estimate an

improved relative coverage fold-change ($\hat{\rho}_{ir}$) for each cell within each CNA region. If cell

i's true allele-specific copy numbers are homogeneous within the given region r, then its

true value of (θ_{ir}, ρ_{ir}) should belong to a set of canonical points displayed in Step 5 of

Fig. 1. Thus, the estimated values ($\hat{\rho}_{ir}, \hat{\theta}_{ir}$) are clustered across cells and associated with

one of the canonical values to yield the cell-level haplotype profiles for the CNA region.

These cell- and region-specific haplotype profiles serve as the basis for clone assignment

and subsequent integration with peak signals in scATAC-seq data. (Fig. 4b).

As we will show next in benchmark experiments, the above method allows accurate CNA

state characterization in a wide range of scenarios. However, there is one scenario that

is extremely difficult: balanced mirrored subclones at small clonal frequencies. To

increase sensitivity for such events, Alleloscope allows an additional refinement step that

starts with the $(\hat{\rho}_{ir}, \hat{\theta}_{ir})$ estimates and uses thresholding on $\hat{\rho}_{ir}$ to improve the estimation

of $\hat{\theta}_{ir}$. Please see Supplementary Methods for details.

Whole genome haplotypes validate Alleloscope in scDNA-seq allele-specific copy

number estimation

First, we explore the single cell landscape of allele-specific CNAs using scDNA-seq data.

The scDNA-seq samples included in this study are given in Supplementary Table 1. To

benchmark the phasing and allele-specific copy number state estimation accuracy, we

performed matched linked-read whole-genome sequencing data on five gastrointestinal

tumor samples representing different levels of chromosomal instability: P5931, P6335,

P6198, P5915, and P6461. Linked-read sequencing, in which one derives reads from

individual high molecular weight DNA molecules, provides variants that can be phased

into extended haplotypes covering megabases in length[36-38]. To evaluate the accuracy of

phasing, we compared the haplotypes estimated by Alleloscope to the haplotypes

obtained from linked-read WGS. Additionally, we used the linked-read WGS haplotype

with Alleloscope to derive gold-standard allele-specific copy number state assignments

for each cell, which are used to assess the impact of phasing errors on accuracy in allele-

specific copy number state estimation (Fig. 2a). This allows us to compute the sensitivity

(recall rate of CNA regions detected in gold-standard) and specificity of a method, as

defined in Supplementary Methods.

We first use the gastric cancer sample P5931 to demonstrate Alleloscope and illustrate

this benchmark design. The heatmap in Fig. 2b gives a bird's-eye-view of Alleloscope's

allele-specific copy number estimates for this sample, showing it to have a relatively

simple genome with few CNAs, which is expected given its MSI subtype. However,

detailed inspection of the four chromosomes carrying clear CNA events, chr7, chr8, chr20,

and chr21, reveals complexity at the allelic level. For each event, the scatter plots of

$\left(\hat{\theta}_{ir}, \hat{\rho}_{ir}\right)$ estimated by Alleloscope and colored by haplotype-specific CNA state

assignment are shown in Fig. 2c. Note that the $\left(\hat{\theta}_{ir}, \hat{\rho}_{ir}\right)$ clusters fall almost directly on

top of the expected canonical values (e.g. (1/2, 1) for diploid, (2/3, 1.5) for 1 copy gain of

major haplotype). Interestingly, chromosomes 7, 8, and 21 each show subclonal clusters

with differing allelic ratios adding up to the same total copy number, which would not have

been detectable without allele-specific estimation. We denote the major haplotype of a

region by "M", and the minor haplotype by "m". The chromosome 7 amplification exhibits

two tumor subclones with mirrored-subclonal CNAs (MMm and Mmm), each subclone

amplifying a different haplotype. Such a mirrored-subclonal CNA configuration is also

observed for the deletion on chromosome 21 (M- and m-). The chromosome 8

amplification exhibits four tumor subclones differentiated by their haplotype profiles—

MMm, Mmm, MMmm, and MMMm.

Comparing against the whole genome haplotypes derived from linked-reads, the phasing

accuracy is 98% for the chr21 deletion, ~90% for the two clonal amplifications (on chr8

and chr20), and 79% for the subclonal chr7 amplification (shown in the titles of the scatter

plots of Fig. 2c). Fig. 2d shows scatterplots of $\hat{\rho}_{ir}$ against major haplotype proportion

computed using haplotypes derived from linked-read sequencing ($\tilde{\theta}_{ir}$), with cells colored

by their state assignment given by Alleloscope without using the haplotype information

(i.e. each cell has the same color as in Fig. 2c). Comparing the scatterplots in Fig. 2d to

their counterparts in Fig. 2c reveals that Alleloscope's estimated cell haplotype profiles

are highly concordant with those derived directly with the haplotypes from linked-read

WGS. Specifically, the concordance is ~100% across all four events (the concordance for

each event is labeled in the scatter plots of Fig. 2d). This shows that the allele-specific

copy number estimation by Alleloscope is robust to errors in phasing (e.g. for chr7). We

also used two extreme cases—chr2 and chr3 of P5931 to demonstrate the refinement

step (Supplementary Fig. 1). For these two regions, linked-read sequencing revealed a

mirrored amplification event distinguishing two subclones each comprised of <10 out of

700 cells. The differing major haplotype proportions of these two subclones, at $\frac{1}{3}$ and $\frac{2}{3}$,

was missed by the original estimation, but accurately detected with the refinement step

of Alleloscope.

So far, CHISEL is the only other comparable method for allele-specific copy number

estimation with scDNA-seq data. Thus, we benchmarked Alleloscope against CHISEL by

comparing each method's estimates under default settings to the same method's

estimates using the true haplotype phasing information provided by linked-read WGS.

Thus, each method's performance is assessed using its own "gold standard" computed

with linked-read WGS haplotypes. We verified that CHISEL's gold standard is highly

similar to Alleloscope's gold standard, and thus, with the help of linked-read WGS

haplotypes both methods can achieve accurate estimation that is close to the truth. The

sensitivity and specificity of each method on the five samples are given in Table 1. For

CHISEL, there is an extra "correction step" where the inferred clones are used to generate

consensus allele-specific copy number profiles for all cells within each clone. We show

CHISEL's results with and without this correction step, as for most samples, it increases

sensitivity (but decreases specificity), but for (P5931) it makes the results worse. The

results obtained using different external phasing datasets and different block sizes for

CHISEL also suggest that CHISEL's accuracy depends on choices for these inputs

(Supplementary Table 2). Detailed heatmaps comparing Alleloscope and CHISEL using

P5931 as an example are given in Supplementary Fig. 2-3, with more discussion given in

Supplementary Results. Overall, Alleloscope maintains high accuracy (both sensitivity

and specificity) across the samples analyzed, substantially improving upon the current

state-of-the-art.

Further performance assessment via data downsampling and simulations

We also compared Alleloscope and CHISEL on the breast cancer sample that was

analyzed in detail by Zaccaria et al[10]. This sample does not have true phasing information,

but was sequenced at much higher coverage, and thus, we downsampled the reads to

50% and 25% of the original depth and compared the estimates obtained from the

downsampled datasets to those obtained from the original dataset. Heatmaps of the

genome-wide profiles estimated by CHISEL and Alleloscope on the original dataset, the

50% subsampled dataset and the 25% subsampled dataset are shown in Supplementary

Fig. 4. For CHISEL, the clone-corrected outputs are shown since they are more accurate

than the uncorrected ones. The accuracy of Alleloscope and CHISEL are shown in Table

1. We see that at 25% of the original depth, Alleloscope has a specificity of 82% as

compared to CHISEL's 65%, and sensitivity of 95% as compared to CHISEL's 64%.

Finally, we used simulations to investigate the performance of Alleloscope over a grid of

experimental parameters (number of cells, total per cell coverage, and total coverage at

heterozygous SNP sites, Supplementary Fig. 5). Across experimental design settings,

Alleloscope attains higher accuracy for deletions than amplifications, which is expected

due to the larger change in both coverage ratio and major haplotype proportion ($\hat{\theta}$) that

accompanies deletions. Phasing for deletions (and other LOH events) is accurate across

the board, but for amplifications it is low at 10% and 5% clonal frequencies. However, the

CNA state assignment accuracy is, to a certain extent, robust to phasing error, increasing

steadily with coverage, number of heterozygous SNPs, and number of cells.

Alleloscope finds pervasive occurrence of polyclonal CNA regions differentiated

by haplotype ratios

We start by considering copy neutral LOH events, which are well-known drivers of cancer

evolution that can only be identified through allele-specific copy number analysis. We use

the colorectal adenocarcinoma sample P6198 as illustration. The copy number heatmap

in cellranger (Fig. 3a) reveals that most of the genome is amplified for this sample, but

with few regions of normal copy number. One of these regions, chromosome 5, is shown

by cellranger to be copy neutral for its entire length. The bulk VAF clearly separates this

chromosome into two main regions, a normal region followed by a copy-neutral LOH (Fig.

3b). Concordantly, Alleloscope reveals a cluster centered at $(\rho, \theta) = (1,1)$ corresponding

to copy-neutral LOH only for the region on the right (Fig. 3b). Similarly, alleloscope also

revealed copy neutral LOH events in this sample on chr11 and chr16. Using haplotypes

from matched linked-read WGS, all of these LOH events are validated and the per-cell

LOH state assignment accuracy of Alleloscope is nearly 100% (Supplementary Fig. 6).

Next, we survey the overall landscape of single cell allele-specific copy number profiles

among the samples analyzed. Detailed segmentation plots and heatmaps for all samples,

listed in Supplementary Table 1, are in Supplementary Fig. 7-13. Across most of the

samples, we observe a high prevalence of complex subclonal CNAs indicated by multiple

clusters differentiated by allelic ratios for the same genomic region. Prototypical examples

from P6198, SNU601, P6335 and BC10X are shown in Fig. 3c. In some regions, such as

chromosome 9 of SNU601, 3q of SNU601, 2q and 16p of BC10x, we see as many as

seven subclonal clusters. In many cases there are multiple clusters corresponding to the

same total copy number but varying in allelic dosage. Minor subclones carrying deletion

of one haplotype can be easily masked by dominant subclones carrying amplifications of

the other haplotype in a bulk analysis or a single cell analysis that relies on total coverage.

Overall, the high subclonal diversity in these genomic regions reveal an aspect of

intratumor heterogeneity that was previously underreported.

Recurrent chromosomal instability events, affecting both haplotypes and producing

gradients in haplotype dosage, is a common theme across all samples analyzed.

Consider, for example, the region on chromosome 9 of P6198 (Fig. 3c), which reveals 7

subpopulations of cells: besides the normal cell cluster and the dominant tumor cell

cluster with the haplotype profile MMm, there is a small cluster of cells with copy neutral

LOH, two small subclones at four chromosome copies and two more at five chromosome

copies. This produces major haplotype ratios of $\{\frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{4}{5}, 1\}$ in different cells. Another

example of such complexity is chromosome 3q of SNU 601 and chromosome 2q of BC10x,

which share a similar pattern: two mirrored-subclonal CNAs (MMm, mmM) at total copy

number of 3, as well as mirrored-subclonal CNAs (MMMm, MMmm, mMMM) at total copy

number of 4, producing a gradient of haplotype ratios $\{\frac{1}{4}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}\}$. Interrogating the

evolutionary route by which such diversity was achieved, Alleloscope reveals that a whole

genome doubling event is likely to have taken place early in the development of BC10x

and P6198, but not in the development of SNU601 (see Supplementary Fig. 4&6). Thus,

the subclones at 2q in BC10x and 3q in SNU601 must have evolved through different

evolutionary routes: In BC10x, the early whole-genome doubling produced the cluster

MMmm, from which the other clusters of different haplotype profiles were most likely

derived through successive loss and gene conversion events. On the contrary, the

clusters on 3q of SNU601 were most likely a result of successive amplification events

starting from the normal haplotype profile Mm. The observation of such similar clonal

configurations in allele-specific copy number, in two samples coming from different cancer

types (breast and gastric), suggests that convergent evolution or similar evolutionary

pressures, where such haplotype dosage gradients serve as an important substrate for

selection in tumor evolution, may be at play. Large studies involving more samples would

be needed to characterize the evolutionary significance of such complex clonal patterns.

Juxtaposition of single cell copy number and chromatin remodeling events by

integrative scATAC-seq analysis

To demonstrate the integrative analysis of scATAC-seq data with Alleloscope, we first

consider two basal cell carcinoma samples with matched whole-exome sequencing (WES)

data[39]. Using the matched WES data, the genome of each sample was first segmented

into regions of homogeneous bulk copy number (Fig. 4a, middle panel shows the

segmentation for SU008). Alleloscope was then applied to the scATAC-seq data to derive

allele-specific copy number estimates of each cell in each region. Scatterplots of $(\hat{\rho}, \hat{\theta})$

for five example CNA regions and one control region (chr12) from SU008 are shown in

Fig. 4a. For this sample, peak profiles characterizing chromatin accessibility separated

the cells confidently into three main clusters: 308 tumor cells, 259 fibroblasts and 218

endothelial cells with the cell type identity retrieved from the original study. Since the

normal fibroblast and epithelial cells are not expected to carry broad copy number events,

they provide a "null scenario" for comparison to the tumor cells. Density contours for each

cell type are shown in the $(\hat{\rho}, \hat{\theta})$- scatterplots (Fig. 4a). The $(\hat{\rho}, \hat{\theta})$ values clearly separate

the tumor cells from the normal cells for each CNA region, with the tumor cell cluster

positioned at canonical points in each case, indicating that these statistics used by

Alleloscope can accurately distinguish amplifications and loss-of-heterozygosity events

in scATAC-seq data. In particular, Alleloscope differentiated the cells that carry the

putative copy neutral LOH events in regions 4a, 6b, and 15b through shifts in major

haplotype proportion. Note that normal cells, which are not expected to carry broad

chromosome-scale CNAs, exhibit chromosome-level deviations in total coverage due to

broad chromatin remodeling but with no significant difference in their $\hat{\theta}_{ir}$ distribution, as

exemplified by the chr6b region. The coverage in large genomic bins across the genome

was further investigated for the two normal cell types in scATAC-seq data (Supplementary

Fig. 14; Supplementary Methods). Comparing one normal cell type against the other,

many regions exhibit broad shifts in relative coverage. Thus, relying solely on shifts in

coverage, without complementary shifts in major haplotype proportion, would lead to false

positive copy number detections for scATAC-seq data.

By assigning allele-specific CNA profiles to single cells in scATAC-seq data, Alleloscope

allows the integrative analysis of chromosomal instability and chromatin remodeling as

follows (Fig. 4b): The scATAC-seq data, paired with bulk or single-cell DNA sequencing

data, allows us to detect subclones. In parallel, a peak-by-cell matrix can be computed

following standard pipelines. Then, the subclone memberships or CNA profiles can be

visualized on the low-dimensional embedding of the peak matrix, and the subclones can

be further compared in terms of peak or transcription factor motif enrichment. Precise

haplotype profiles for each subclone then allow us to identify significantly

enriched/depleted peaks after accounting for copy number differences, thus delineating

events that are uniquely attributable to chromatin remodeling.

Hierarchical clustering using major haplotype proportion $\hat{\theta}$ identifies the tumor cells from

the normal cells for both SU006 (Supplementary Fig. 15) and SU008, and clearly

delineates a subclone in SU008 marked by a copy-neutral LOH event on chr4a (Fig. 4c).

Focusing on SU008, we call the cell lineage that carries the chr4a LOH event clone-2,

and the other lineage clone-1. In parallel, clustering by peaks cleanly separates the tumor

cells from the epithelial cells and fibroblasts (Fig. 4d: left), and further, demarcates two

distinct clusters in the tumor cells (peaks-1 and peaks-2) (Fig. 4d: middle). What is the

relationship between the peaks-1 and peaks-2 clusters obtained from peak signals to the

two clones delineated by chr4a LOH? Coloring by chr4a major haplotype proportion ($\hat{\theta}$)

on the peaks-derived UMAP shows that the LOH in this region is carried by almost all of

the cells in peaks-2 but only a subset of the cells in peaks-1 (Fig. 4d: middle). This can

also be clearly seen in the density of $\hat{\theta}$ (Fig. 4d: right): While $\hat{\theta}$ is heavily concentrated

near 1 for peaks-2, it is bimodal for peaks-1. Since clone-1 and clone-2 are differentiated

by a copy-neutral event, this separation by peaks into two clusters is not driven by broad

differences in total copy number. Since clone-2 is split into two groups of distinct peak

signals, we infer that the chromatin remodeling underlying the divergence of the peaks-2

cells is likely to have occurred in the clone-2 lineage, after the chr4a LOH event (Fig. 4e).

In this way, Alleloscope analysis of this scATAC-seq data set allowed us to overlay two

subpopulations defined by peak signals with two subpopulations defined by a subclonal

copy-neutral LOH, and infer their potential temporal order.

Integrative analysis of clonal evolution and altered chromatin accessibility for a

complex polyclonal gastric cancer cell line

The gastric cancer cell line SNU601 exhibits complex subclonal structure, as evidenced

by multiple multiallelic CNA regions (chr3e and chr18a are shown in Fig. 3c). In addition

to scDNA-seq, we also performed scATAC-seq on this sample to profile the chromatin

accessibility of 3,515 cells at mean coverage of 73,845 fragments per cell. This allows us

to compare the allele-specific copy number profiles obtained by scATAC-seq with those

given by scDNA-seq and integrate the two data types in a multi-omic characterization of

this complex tumor.

First, we segmented the genome and estimated the allele-specific copy number profiles

of single cells at each segment for both the scATAC-seq and scDNA-seq data, following

the procedure in Fig. 1 with some modifications due to the lack of normal cells to use as

control for this sample (see methods). Fig. 5a shows the relative total coverage, pooled

across cells from scDNA-seq. Fig. 5b shows $(\hat{\rho}, \hat{\theta})$-scatterplots for five example CNA

regions in scDNA-seq and scATAC-seq. Compared to the scATAC-seq data, the scDNA-

seq data has about 8-fold higher total read coverage and 7-fold higher heterozygous site

coverage per cell. Thus, while subclones corresponding to distinct haplotype profiles are

cleanly separated in the scDNA-seq data, they are much more diffuse in the scATAC-seq

data. Yet, cluster positions in scATAC-seq roughly match those in scDNA-seq. As

expected, the $(\hat{\rho}, \hat{\theta})$-scatterplots reveal the high level of chromosomal instability in this

sample, with each region exhibiting multiple clusters of different haplotype structures that

indicate the existence of subclones carrying mirrored events and, for some regions, the

variation of haplotype dosage over a gradient across cells.

Fig. 5c shows the hierarchical clustering of cells from scDNA-seq based on their allele-

specific copy number profiles, revealing the subclonal structure and the co-segregating

CNA events that mark each subclone. For each cell in each region, Alleloscope also

produces a confidence score for its assignment to different haplotype profiles

(Supplementary Fig. 16). Based on visual examination of the confidence scores at the

marker regions in both the ATAC and DNA sequencing data sets, we identified 6

subclones for further investigation (Clones 1-6 labeled at the right of the heatmap). The

allele-specific copy number profiles allow us to manually reconstruct the probable

evolutionary tree relating these 6 clones under the following three rules:

(1) Parsimony: The tree with the least number of copy number events is preferred.

(2) Monotonicity: For a multi-allelic region with escalating amplifications (e.g. Mm, MMm,

MMMm), the haplotype structures were produced in a monotonic order (e.g. Mm→

MMm→ MMMm) unless a genome doubling event occurred.

(3) Irreversibility of LOH: Once a cell completely loses an allele (i.e. copy number of that

allele becomes 0), it can no longer gain it back.

The evolutionary tree, thus derived, is shown in Fig. 6b. The mirrored-subclonal

amplifications on chr3q, the deletion on chr4p, and the multiallelic amplification on chr20q

allowed us to infer the early separation of clones 3-6 from clones 1-2. Subclones 3-6 are

confidently delineated by further amplifications on chr3q, chr20q, chr11, chr13, and chr17.

Note that high chromosomal instability led to concurrent gains of 1q and 7p in both the

Clone 1-2 and Clone 3-6 lineages. We also observed a large number of low-frequency

but high-confidence CNA events indicating that ongoing chromosomal instability in this

population is spawning new sporadic subclones that have not had the chance to expand.

We now turn to scATAC-seq data, focusing on the 10 marker regions which, together,

distinguish Clones 1-6: chr1b, 3b-d, 4b, 7a, 11b, 13b, and 20b-c. The $(\hat{\rho}, \hat{\theta})$ values

computed by Alleloscope allows us to directly assign allele-specific copy number profiles

to each cell for each region, as well as subclone labels to each cell, with posterior

confidence score. The subclone assignment utilizes a Bayesian mixture model that pools

information across the 10 marker regions. Despite the low accuracy in per-region

genotyping, when information is pooled across the 10 marker regions, 81.6% of the 2,753

cells after filtering can be assigned to a subclone with >95% posterior confidence

(Supplementary Fig. 17, the number of ATAC cells confidently assigned to each clone

are shown in Fig. 6a.). These subclone assignments for each cell, and cell-level haplotype

profiles for each region, can now be integrated with peak-level signals.

Following the scheme in Fig. 4b, we computed the Uniform Manifold Approximation and

Projection (UMAP) coordinates for the scATAC-seq cells based on their peak profiles,

which enables us to visualize the difference in chromatin accessibility profiles among

individual cells (Fig. 6a). UMAP scatterplots colored by clone assignment show that the

6 clones exhibit marked differences in their chromatin accessibility profiles (Fig. 6a): While

Clone 1 and Clone 2 are concentrated at the top half of the UMAP, Clones 3-5 are

positioned almost exclusively at the bottom half. Clone 6, which exhibits more variance,

is also significantly enriched at the bottom half of the UMAP. Among Clones 3-5, Clone 3

has a distinct chromatin accessibility profile that is mostly concentrated at the bottom tip,

Clone 4 is positioned higher, while Clone 5 contains cells that are similar to both clones

3 and 4. We expect some of these peak-level differences to be driven by CNAs.

To delineate the peaks that differ between clones, and to distinguish peak differences

that are not accountable by CNAs, we developed a statistical test to adjust for underlying

copy numbers to identify differentially accessibility peaks (DAPs) between the cell

populations on the two branches (Fig. 6b). The test is based on a generalized log-

likelihood ratio test with specific adjustment terms for the copy number profile of each of

the two subclones (Methods). The same test, but neglecting coverage adjustment, was

also performed for comparison. Along each branch, three numbers are shown for DAPs—

1. DAPs identified only before adjustment; 2. DAPs identified both before and after

adjustment; 3. DAPs identified only after adjustment (Fig. 6b). For the smaller subclones

(Clone 3,4,5), low coverage limits the detection power and thus limits the DAP counts in

both categories. Yet, juxtaposing DAP and CNA events along the tumor phylogeny yields

insights: Along most lineages, adjustment for CNA-induced broad shifts in coverage

removes a large number of peaks whose differential-accessibility signal is explainable by

the underlying change in copy numbers. Interestingly, CNA adjustment also identifies new

DAPs whose signals were obscured by a change in copy number in the opposite direction.

This argues for the importance of CNAs as a mechanism underlying subclonal differences

in chromatin accessibility in this tumor.

Nevertheless, along some branches we find a large number of DAPs remaining after

adjustment (i.e. DAPs identified both before and after adjustment), and thus must be due

to other mechanisms. Two example DAPs of this latter category are shown as insets in

Fig. 6b, with full list given in Supplementary Table 3. The first example is a peak at the

transcription start site (TSS) of the REC8 gene, which is located on chr14 where no

apparent CNAs were observed across the six major subclones. The TSS of REC8 is open

in clones 3-6 but closed in clones 1-2 (p-value<0.0001). REC8 is a gene encoding a

meiosis-specific cohesion component that is normally suppressed in mitotic proliferation,

and its role in cancer has recently gained increasing attention and controversy: While Yu

et al.[40] found the expression of this gene to suppress tumorigenicity in a gastric cancer

cell line, McFarlane et al.[41] postulated that it may be broadly activated in some cancers

where it generates LOH by reductional segregation. The opening of the TSS of REC8,

stably maintained in Clones 3-6, suggests that meiotic processes may underlie the

increased chromosomal instability of this multiclonal lineage. The second example is a

peak at the TSS of the WWOX gene, located on chr16, which is significantly depleted in

Clone 3 (p-value<0.0001). Although chr16 has LOH across all tumor cells, there are no

detectable subclonal differences, and thus we don't expect the decrease in accessibility

at WWOX for subclone 3 to be due to a large copy number event. Since WWOX is a well-

known tumor suppressor whose down-regulation is associated with more advanced

tumors[42, 43], its decrease in accessibility suggests a more aggressive phenotype for Clone

3. Overall, these two examples show how Alleloscope can be used to dissect the roles of

CNA and chromatin-level changes in the identification of gene targets for follow-up study.

Discussion

Despite recent advances in the application of single cell sequencing to cancer, we are

still far from understanding the diversity of genomes that are undergoing selection at the

single cell level. Notably, little is yet known about the intratumor diversity of allelic

configurations within CNA regions, and to what extent the diversity of cells in chromatin

accessibility can be attributed to diversity in allele-specific copy number. Here, we

presented Alleloscope, a new method for allele-specific copy number estimation that can

be applied to single cell DNA and ATAC sequencing data (separately or in combination).

Through a combination of matched linked-reads whole genome sequencing,

downsampling-based benchmark experiments, and simulations, we comprehensively

assessed the accuracy of Alleloscope and benchmarked against CHISEL, which is

currently the only other method for scDNA-seq allele-specific copy number profiling.

Detailed discussion about the two methods can be found in Supplementary Results.

We applied Alleloscope to a panel of breast, colorectal, and gastric tumors and cancer

cell lines, where it revealed an unprecedented level of allelic heterogeneity within

hypermutable CNA regions. In these regions, subclones reside on a gradient of allelic

ratios that is unobservable in total copy number analysis. In simple cases, these

hypermutable regions contain mirrored subclones, as previously identified[9, 10], but are

often much more complex. We observed multiple instances of convergent evolution

involving recurrent CNA events affecting the same region, some verified by linked read

sequencing. In accordance with the findings in Watkins et al.[44], we found using

Alleloscope that chromosomal instability drives the formation of subclones not only in

primary tumors but also after metastasis. Alleloscope was also applied to a scDNA-seq

dataset[15] generated from a different sequencing protocol (Supplementary Fig. 18;

Supplementary Methods), further demonstrating the general applicability of the approach.

Having established the allelic complexity of CNAs at single cell resolution, we next applied

Alleloscope to scATAC-seq data, thus enabling the combined study of clonal evolution

and chromatin accessibility. First, we considered the analysis of a public data set

consisting of two basal cell carcinoma samples, for which matched bulk whole-exome

sequencing data was used for initial genome segmentation upon which single cell CNA

genotyping was then conducted in the scATAC-seq data. Here we showed that

Alleloscope can detect amplifications, deletions, and copy-neutral LOH events accurately

in scATAC-seq data, and was able to find a subclone delineated by a copy-neutral LOH

event. Juxtaposing this subclone assignment with peak signals allowed us to detect a

wave of genome-wide chromatin remodeling in the lineage carrying the LOH. Next, we

applied Alleloscope to a complex polyclonal gastric cancer cell line with matched scDNA-

seq data. By overlaying peak signals with subclones delineated by allele-specific copy

number estimates, we developed a statistical test to adjust for copy numbers in identifying

DAPs. Comparison between the testing results before and after copy number adjustment

suggests that much of the intratumor heterogeneity in chromatin accessibility can be

attributed to CNAs. Focusing on subclone-enriched peaks with copy number adjustment

allowed the prioritization of genes for downstream follow-up.

Alleloscope can potentially be applied to the integration of single cell data of other

modalities, for example scATAC-seq and scRNA-seq data, to investigate the relationships

between clonal evolution, chromatin remodeling, and transcriptome. To facilitate

experimental design for single cell omics sequencing protocols, we investigated the

performance of Alleloscope under different experimental parameters (number of cells,

total per cell coverage, and total coverage at heterozygous SNP sites). Coverage at

heterozygous SNP sites is an especially important consideration for the design of scRNA-

seq and scATAC-seq studies, for which shifts in total coverage is an unreliable proxy for

underlying DNA copy number. For scATAC-seq, the lower heterozygosity within peak

regions led to lower number of reads mapping to heterozygous loci as compared to

scDNA-seq, and this resulted in noisier subclone detection. Most of the current scRNA-

seq technologies only sequence either the 3' or 5' end of the mRNA transcripts, which

limits the number of heterozygous SNP sites covered by reads. The latest developments

in single cell long read sequencing[45-47] and single cell multimodal sequencing[48] herald

new analysis opportunities with this method.

<u>Methods</u>

ScDNA-seq Data Sets and Pre-processing

The ten 10x scDNA-seq datasets analyzed in this study are summarized in

Supplementary Table 1. P5931, P5847 and P6461 scDNA-seq data were generated

using the method described in the previous study[35]. We applied the Cell Ranger DNA

pipeline (https://support.10xgenomics.com/single-cell-dna/software/overview/welcome;

beta version: 6002.16.0) for sample demultiplexing, read alignment, CNA calling and

visualization. Most data were aligned to the GRCh38 reference genome except for the

two processed BC10x datasets (GRCh37). For the tumor samples with a matched

normal sample, the GATK HaplotypeCaller was used to call heterozygous SNPs on the

matched normal samples. Otherwise, SNPs were retrieved on the tumor sample

themselves. Next, we applied VarTrix to efficiently generate SNP-by-cell matrices for

both reference and alternative alleles of the identified SNPs.

To select high-quality SNPs, we filtered out the SNPs with <5 reads for P5846 and

P5847, <10 reads for P5915 and P5931, <15 reads for P6335 and P6461, <20 reads for

P6198, SNU601, and ther BC10X samples based on the number of SNP detected for

each sample. Additionally, SNPs located in the centromeres and telomeres were

excluded. To exclude cells that might undergo apoptosis or cell cycles, noisy cells

labeled by the Cell Ranger tool were excluded.

Single-cell ATAC Data sets, Sequencing and Preprocessing

The scATAC-seq datasets analyzed in this study are summarized in Supplementary

Table 4. We used the following procedure to generate the SNU601 scATAC-seq

dataset.About 400,000 cells were washed with RPMI media and centrifuged (400g for 5

min at 4°C) twice. The supernatant was removed and resuspended with chilled PBS +

0.04% BSA solution. The resuspended pellet was centrifuged (400g for 5min at 4°C),

and the supernatant was removed again. Then 100 µL of chilled Lysis Buffer (10 mM

Tris-HCl (pH 7.4), 10 mM NaCl, 3 mM $MgCl_2$, 1% BSA, 0.1% Nonidet P40 Substitute,

0.1% Tween-20 and 0.01% digitonin) was added and carefully mixed 10 times. The tube

was incubated on ice for 7 min. After incubation, 1 mL of chilled Wash Buffer (10 mM

Tris-HCl (pH 7.4), 10 mM NaCl, 3 mM $MgCl_2$, 1% BSA and 0.1% Tween-20) was added

and mixed 5 times followed by centrifugation of nuclei (500g for 5 min at 4°C). After

removing the supernatant, nuclei were resuspended in chilled Nuclei Buffer (10X

Genomics), filtered by Flowmi Cell Strainer (40uM) and counted using a Countess II FL

Automated Cell Counter. Then the nuclei were immediately used to generate scATAC-

seq library.

ScATAC-seq library was generated using the Chromium Single Cell ATAC Library & Gel

Bead Kit (10X Genomics) following the manufacturer's protocol. We targeted 3000

nuclei with 12 PCR cycles for sample index PCR. Library was checked by 2% E-gel

(Thermofisher Scientific) and quantified using Qubit (Thermofisher Scientific).

Sequencing was performed on Illumina NextSeq500 using NextSeq 500/550 High

Output Kit v2.5 (Illumina).

Raw sequencing reads of the SNU601 scATAC-seq sample was de-multiplexed with the

10x Genomics Cell Ranger ATAC Software (v.1.2.0; https://support.10xgenomics.com/single-

cell-atac/software/pipelines/latest/algorithms/overview) and aligned to the GRCh38 reference

genome. The aligned scATAC-seq data of the two pre-treatment basal cell carcinoma

samples (BCC;SU006 and SU008) were retrieved from GSE129785[26]. To obtain all

potential SNPs for the SU006 and SU008 samples, GATK Mutect2 was used to call

single-nucleotide variants (SNVs) on the bam files deduplicated by the Picard toolkits of

both the t-cell dataset and the tumor dataset from the same tumor. All SNVs from the

paired tumor-normal datasets were combined and the read counts of these SNPs were

quantified for each cell in the tumor scATAC-seq dataset. For the SNU601 scATAC-seq

data, we instead quantified the read counts of the two alleles of the SNPs more reliably

called from the paired normal scDNA-seq data. Then the SNP-by-cell matrices for both

reference and alternative alleles were generated using Vartrix for all the scATAC-seq

datasets. To select potential germline SNPs, we further filtered out the SNVs <5 reads

for the SU008 sample and <10 reads for the SU006 sample. SNPs with extreme VAF

values <0.1 or >0.9 were also excluded for both samples. Because we used the phasing

information from the paired scDNA-seq data to assist the estimation of the haplotype

structures for the SNU601 scATAC-seq data, we did not filter out SNPs with extreme

VAF values for the scATAC-seq data. To improve the quality of the downstream

analysis for SNU601 scATAC-seq data, we filtered out cells <5 reads and the SNPs <5

reads.

Linked-read sequencing and data processing

The five samples with the linked-read sequencing data were acquired as surgical

resections following informed consent under an approved institutional review board

protocol from Stanford University. Samples were subjected to mechanical and

enzymatic dissociation as previously described, followed by cryopreservation of

dissociated cells[33].

Cryofrozen cells were rapidly thawed in a bead bath at 37 ºC. Cell counts were obtained

on a BioRad TC20 cell counter (Biorad, Hercules, CA) using 1:1 trypan blue dilution.

Between 1.5-2.5 million total cells were washed twice in PBS. Centrifugation was

carried out at 400g for 5 minutes. PBS was removed and cell pellets were frozen at -

80ºC. DNA extraction was carried out on cell pellets following thawing using either

MagAttract HMW DNA Kit (P5931) or AllPrep DNA/RNA Mini Kit (Qiagen Inc.,

Germantown, MD, USA) as per manufacturer's protocol. Quantification was carried out

using Qubit (Thermofisher Scientific).

Sequencing libraries were prepared from DNA using Chromium Genome Reagent Kit

(v2 Chemistry) (10X Genomics, Pleasanton, CA, USA) following manufacturer's

instructions. Sequencing was performed using Illumina HiSeq or NovaSeq sequencers

using 150x150 bp paired end sequencing and i7 index read of 8 bp. Long Ranger (10X

Genomics; v2.2.0) was used to align reads, call and phaseSNPs, indels and structural

variants.

Segmentation

The first step of Alleloscope is to segment the genome into regions with different CNA

profiles. The appropriate segmentation algorithm depends on what samples are

available. First, matched bulk DNA sequencing data (WGS/WES) or pseudo-bulk data

from scDNA-seq data can be segmented using FACLON[5], a segmentation method that

jointly models the bulk coverage and bulk VAF profiles, if a matched normal sample is

available. To accommodate segments from rare subclones, methods that integrate

shared cellular breakpoints in CNA detection for scDNA-seq[49] can improve sensitivity. If

a normal sample is unavailable, Alleloscope instead uses an HMM-based segmentation

method. The HMM method, which operates on the binned counts of pooled cells,

assumes a Markov transition matrix on four hidden states representing deletion, copy-

neutral state, single-copy amplification and double-copy amplification:

$$617 \quad \begin{pmatrix} 1-3t & t & t & t \\ t & 1-3t & t & t \\ t & t & 1-3t & t \\ t & t & t & 1-3t \end{pmatrix}, \text{ where } t = 1 \times 10^{-6} \text{ by default. The segmentation}$$

618 plots in the study were generated by using the HMM algorithm. With the paired sample,

619 the P6198 tumor sample was segmented using FALCON on the SNPs >30 reads with

620 default parameters. Besides FALCON and HMM methods, users can choose the

621 methods that work best for specific datasets. For example, we also tried ASCAT[11] on

622 the P6198 sample for comparison (Supplementary Fig. 10).

Whole-exome sequencing (WES) data processing

624 The WES data of the two paired tumor-normal samples (SU006 and SU008) were

625 obtained from PRJNA533341. Raw fastq files were aligned to the GRCh37 reference

626 genome using bwa-mem[50] with duplicate reads removed using the Picard toolkits[51]. The

627 copy number calls of paired normal-tumor samples were obtained using Varscan2[52].

628 Then the HMM algorithm was applied to segment the genome.

SNP Phasing and Single-cell Allele Profile Estimation per region

For each region r after segmentation, an expectation-maximization (EM)-based method

is used to iteratively phase each SNP and estimate cell-specific allele-specific copy

number states for scDNA-seq and scATAC-seq datasets. "Major haplotype" is defined

as the haplotype with higher aggregate copy number in the sample. Let I_j indicate

whether the reference allele of SNP j is located on the major haplotype and θ_{ir} denote

major haplotype proportion of cell i for region r. The EM model iterates the E-step and

the M-step. The complete log likelihood of the model is

637
$$l(\theta_{ir}) = \sum_{j=1}^{n} log P(A_{ij}, B_{ij}|\theta_{ir})$$

638
$$= \sum_{j=1}^{n} \{[A_{ij}log\theta_{ir} + B_{ij}\log(1-\theta_{ir})]I_j + [B_{ij}log\theta_{ir} + A_{ij}\log(1-\theta_{ir})](1-I_j)\}$$

639 where A_{ij} and B_{ij} are the observed read counts for the reference and alternative alleles

640 of cell i on SNP j. In the E-step, we calculate the expected value of the posterior

641 probability of the hidden variable I_j.

642
$$E_{\hat{\theta}_{ir}(t)}[I_j|A_{ij}, B_{ij}] = \hat{I}_j^{(t)} = \frac{\prod_i \hat{\theta}_{ir}^{(t)A_{ij}}\left(1-\hat{\theta}_{ir}^{(t)}\right)^{B_{ij}}}{\prod_i \hat{\theta}_{ir}^{(t)A_{ij}}\left(1-\hat{\theta}_{ir}^{(t)}\right)^{B_{ij}} + \prod_i\left(1-\hat{\theta}_{ir}^{(t)}\right)^{A_{ij}}\hat{\theta}_{ir}^{(t)B_{ij}}}$$

643 where $\hat{\theta}_{ir}^{(t)}$ is the parameter from the tth iteration. In the M-step, $\hat{\theta}_{ir}$ is updated by solving

644
$$\hat{\theta}_{ir}^{(t+1)} = argmax_{\theta_{ir}} E\left[l(\theta_{ir})\middle|A_{ij}, B_{ij}, \hat{\theta}_{ir}^{(t)}\right]$$

645
$$= \frac{\sum_j[A_{ij}\hat{I}_j^{(t)} + B_{ij}(1-\hat{I}_j^{(t)})]}{\sum_j[A_{ij}\hat{I}_j^{(t)} + B_{ij}(1-\hat{I}_j^{(t)})] + \sum_j[A_{ij}(1-\hat{I}_j^{(t)}) + B_{ij}\hat{I}_j^{(t)}]}$$

where $\hat{\theta}_{ir}^{(t)}$ and $\hat{\theta}_{ir}^{(t+1)}$ are from two successive iterations of EM. An initial estimate $\hat{I}_j^{(0)}$ is

first derived from the bulk VAF profile, and the two steps are iterated until converge. The

maximum number of SNPs was set as 30,000 by default. For the SNU601 scATAC-seq

dataset, the phases estimated in the paired scDNA-seq dataset ($\hat{I}_j$'s) were directly

applied to estimate the $\hat{\theta}_{ir}$'s of the cells in the scATAC-seq data. To have enough

power, we included the cells with ≥ 20 reads covering the identified SNPs for each

region.

Selecting normal cells and normal regions for single-cell Coverage Normalization

To compute the relative coverage change for each cell in region r ($\hat{\rho}_{ir}$), normal cells and

diploid regions are required for normalization. The $\hat{\theta}_{ir}$'s inferred from the EM-based

algorithm are used to help identify normal cells and diploid regions. To identify normal

cells, the cells were divided into k largest groups ($k = 5$ by default) by hierarchical

clustering. The c[th] cluster is considered normal if it has the minimum distance calculated

by

660
$$\sum_{r=1}^{R} \left| \frac{\sum_{\hat{\theta}_{ir} \in S_c} \hat{\theta}_{ir}}{n_c} - 0.5 \right|^2$$

661   where S_c represents $\hat{\theta}_{ir}$ values of the cells in the c[th] cluster, and n_c is total cell number

662   in the c[th] cluster. All cells in the c[th] cluster are considered as candidate normal cells.

663   Putative diploid regions are next identified in each cluster by considering both $\hat{\theta}_{ir}$ and

664   coverage. Alleloscope computes the first measurement (d_{cr}) as the sum $\hat{\theta}_{ir}$ distance of

665   the cells in the c[th] cluster for each region r.

666
$$d_{cr} = \sum_{\hat{\theta}_{ir} \in S_c} \left| \hat{\theta}_{ir} - 0.5 \right|^2$$

667   Because regions with both haplotypes equally amplified can also have small d_{cr},

668   adjusted raw coverages ($\tilde{\rho}_{ir}$; defined as below) of cell i and region r are also

669   considered in diploid region selection.

670
$$\tilde{\rho}_{ir} = \frac{N_{ir}}{N_i} \times \frac{L_r}{LL},$$

671   where for cell i and region r, N_{ir} is the total read counts and N_i is the total read counts

672   across regions. L_r is length of the region r and LL is total length of the genome. For

673   region r, cells with $\tilde{\rho}_{ir}$ values larger than the 99th percentile are assigned the $\tilde{\rho}_{ir}$ values

674   equal to the 99th percentile across the cells. The second measurement (m_{cr}) used to

675   select diploid regions in the cth cluster is the mean $\tilde{\rho}_{ir}$ for each region r

676
$$m_{cr} = \frac{\sum_{\tilde{\rho}_{ir} \in S_c} \tilde{\rho}_{ir}}{n_c}$$

where S_c here represents $\tilde{\rho}_{ir}$ values of the cells in the cth cluster. Then Alleloscope

identifies potential diploid regions for each cluster by raking the sums of d_{cr} ranks and

m_{cr} ranks. Excluding the cth cluster identified as the normal group, Alleloscope

proposed a list of candidate diploid regions across the clusters by selecting the majority

region.

Because coverage on scATAC-seq data is confounded by epigenetic signals,

chromosome 22 for SU008 and chromosome 18 for SU006 were directly selected as

normal regions based on the matched WES data. Cells were considered as normal or

tumor cells based on the epigenetic signals. For the SNU601 scATAC-seq dataset,

chromosome 10 was selected as the normal region based on the paired scDNA-seq

data.

Cell-level CNA state estimation

689   The cell-level allele-specific copy number profiles are defined by both relative coverage

690   change ($\hat{\rho}_{ir}$) and major haplotype proportion ($\hat{\theta}_{ir}$) for region r. After the normal cells and

691   a control region are identified, cell-specific relative coverage change in region r is

692   calculated as

$$\hat{\rho}_{ir} = \frac{N_{ir}}{N_{i0}} / median(\frac{N_{0r}}{N_{00}})$$

where N_{ir} is total read counts in region r and N_{i0} is total read counts in a reference

region for cell i. N_{0r} is a vector denoting total read counts in region r of all identified

normal cells and N_{00} is a vector denoting total read counts in the same reference region

of all identified normal cells. For P6461 with no diploid region, $\hat{\rho}_{ir}$ was adjusted by

timing 2 for the abnormal cells. Because SNU601 is a tumor cell line with no normal

cells in the dataset, N_{0r} and N_{00} were calculated from the cells in the normal P6198

sample instead for the scDNA-seq data. For SNU601 scATAC-seq data, we aligned the

distribution of the $\hat{\rho}_{ir}$ values in paired scDNA-seq data to the distribution of the $\frac{N_{ir}}{N_i}$

values for each region to get the normalized $\hat{\rho}_{ir}$ in the scATAC-seq data. The

normalized $\hat{\rho}_{ir}$ values for the scATAC-seq data were computed by

$$\hat{\rho}_{ir}^{atac} = \frac{N_{ir}}{N_{i0}} / median(\frac{N_{ir}}{N_{i0}}) \times median(\hat{\rho}_{ir}^{dna})$$

Cells with extreme $\hat{\rho}_{ir}$ values larger than the 99th percentile and smaller than the first

percentile across the cells were excluded for each region. With the $(\hat{\rho}_{ir}, \hat{\theta}_{ir})$ pairs, cells

in the scDNA-seq data can be classified into haplotype profile g with the expected

(ρ_g, θ_g) values based on minimum distance. With noisier signals in the scATAC-seq

data, the haplotype structures identified in the paired scDNA-seq data can help guide

the genotyping for each region. For region r, the posterior probability that cell i carries

haplotype profile g was

712
$$P\big(GT_{ir} = g_r \big| \hat{\rho}_{ir}, \hat{\theta}_{ir}\big) = \frac{P\big(\hat{\rho}_{ir}, \hat{\theta}_{ir} \big| GT_{ir} = g_r\big)\pi_{g_r}}{\sum_{g_{r'}} P\big(\hat{\rho}_{ir}, \hat{\theta}_{ir} \big| GT_{ir} = g'\big)\pi_{g'_r}},$$

713   where g_r denotes the haplotype profile observed in region r in the paired scDNA-seq

714   data and π_{g_r} denotes the prior probability that a randomly sampled cell carrying the g_r

715   haplotype profile. A uniform prior was used for π_{g_r}. In the formula,

716
$$P\big(\hat{\rho}_{ir}, \hat{\theta}_{ir} \big| GT_{ir} = g_r\big) = P\big(\hat{\rho}_{ir} \big| \mu = \rho_g; \sigma_\rho = 0.25\big) \times P\left(\hat{\theta}_{ir} \bigg| \mu = \theta_g; \sigma = \sqrt{\frac{\hat{\theta}_{ir}(1 - \hat{\theta}_{ir})}{n_{ir}}}\right),$$

where n_{ir} is total read counts in region r for cell i. For scATAC-seq data, the haplotype

profile of cell i in region r was estimated by maximizing the above posterior probability.

The haplotype profiles of each region are visualized using different colors in scatter

plots for both scDNA-seq and scATAC-seq data with the confidence scores calculated

using the distance of the points to the canonical centers and the standard deviations.

Validations using matched linked-read sequencing data for each region

We performed validations on five samples using matched linked-read sequencing data.

Two strategies were used for the validations. First, to assess the phasing accuracy of

region r, the largest haplotype block within region r in the matched linked-read

sequencing data was selected for comparison, ensuring that the included SNPs are

phased with respect to one another. Then the $\hat{I}_j$ estimates for SNPs within region r in

the scDNA-seq data were converted to binary indicator $\tilde{I}_j$

$$\tilde{I}_j = 1(\hat{I}_j > 0.5) \in \{0,1\}$$

Using the major/minor haplotype setting, all SNPs in the largest haplotype block in the

matched linked-read sequencing data were placed accordingly to retried I_j as the gold

standard. Comparing the binary estimates $\tilde{I}_j$ and I_j, the phasing accuracy for region r

was computed as follows:

$$ACC_{r,phasing} = \frac{\sum 1\,(\tilde{I}_j = I_j; \tilde{I}_j \in I_j)}{\sum 1(\tilde{I}_j \in I_j)},$$

Secondly, we evaluated the accuracy for cell-level CNA state estimation for region r. To

do this, for each cell in region r we compared the estimated haplotype profiles ($\hat{\theta}_{ir}$) to

the haplotype profiles (θ_{ir}) obtained by plugging in known phases provided by the

matched linked-read sequencing data as the gold standard. Similarly, the haplotype

block with the largest number of SNPs was used here. The θ_{ir}'s, used as the gold

standard, were retrieved by directly plugging in known phases I_j, provided by the

matched linked-read sequencing data (explained in the first strategy). Then the

accuracy for cell-level CNA state estimation of region r is computed as

743
$$ACC_{r,CNAstate} = \frac{\sum_i 1(\hat{\theta}_{ir} = \theta_{ir})}{\sum_i i}$$

Simulations and Power Analysis

745 For a simulated region, let n be the number of cells, m be the number of heterozygous

746 SNPs, θ be the major haplotype proportion, and μ_i be the total coverage of cell i sampled

747 from the cells on chr7 in the P5931 tumor sample. For cell i, we simulated total coverage

748 of SNP j (μ_{ij}) using a Poisson distribution

749
$$\mu_{ij} \sim Poisson(\mu_i),$$

750 where $i = 1 \sim n$. Parallelly, phases of SNP j (I_j) were simulated under a Bernoulli

751 distribution

752
$$I_j \sim Bernoulli(0.5),$$

753 where I_j indicates whether a reference allele is on the major haplotype for SNP j, and

754 j=1~m. Using μ_{ij} and I_j, simulated read counts of reference alleles of SNP j in cell i (A_{ij})

755 were simulated under a Binomial distribution

756
$$A_{ij} \sim Binomial(\mu_{ij}, p_{ij}),$$

757 where p_{ij} is the proportion of the reference allele at loci j in cell i with the values shown in

758 the following table

p_{ij}	cell i with CNA	cell i without CNA
$I_j = 1$	θ	0.5
$I_j = 0$	$1 - \theta$	0.5

Then simulated read counts of alternative alleles of SNP j in cell i (B_{ij}) were retrieved by

$$B_{ij} = \mu_{ij} - A_{ij}$$

In the first simulation used to illustrate distribution of the estimates from Alleloscope, we

fixed the cell number n to be 1,000, the SNP number m to be 10,000 which are typical in

real datasets. θ was set to be 1 and 0.66 for cells carrying deletion and one-copy

amplification respectively with the purity equal to 0.5. On the simulated A_{ij} and B_{ij}

matrices Alleloscope estimated phases for each SNP and CNA states for each cell.

Distribution of the estimated values versus the true values are visualized using boxplots.

To explore the effects of SNP numbers, cell coverage, cell numbers, and purity, power

analysis was performed for one-copy deletion and one-copy amplification scenarios. We

assessed the accuracy for phasing and cell-level CNA state estimation under the

following scenarios: SNP numbers from 1,000 to 50,000, mean coverage from 0.01 to

0,05 for each cell, cell number from 500 to 2500. For different scenarios, we assessed

the effect of three purity: 0.5, 0.1, and 0.01, reflecting from larger subclones to rare

subclones. All parameters remained the same as those in the previous paragraph except

for the parameters that were assessed. Phasing accuracy was calculated by comparing

true I_j's and estimated $\hat{I}_j$'s in the region. If $\hat{I}_j \geq 0.5$, the values were considered as 1.

Otherwise, the values were considered 0. On the other hand, the accuracy of cell CNA

state estimation was the clustering accuracy using the estimated $\hat{\theta}_i$ values. Cells with $\hat{\theta}_i$

values smaller than the midpoints between true θ of normal cells ($\theta_o = 0.5$) and true θ of

carriers ($\theta_{del} = 1$; $\theta_{amp} = 0.66$) were considered as normal cells; otherwise, cells were

considered as carriers. The clustering accuracy was calculated by comparing the clusters

to the true cell states.

Cell Lineage Reconstruction

For scDNA-seq data, cell-specific haplotype profiles from Alleloscope across the

genome are used to reconstruct cell lineage trees. The "Gower's distance" is calculated

using "cluster" R package on the nominal haplotype profiles between cells. The

'pheatmap' R package was used to perform hierarchical clustering on the distance using

the "ward.D2" method. Because fewer SNPs lead to higher variance in $\hat{\theta}_{ir}$'s, we by

default included segments with more than 2,000 SNPs identified. For visualization, each

segment was plotted with its length proportional to 5,000,000 bins, and the heights of

the clustering tree were log-transformed.

To explore tumor subclonal structure in the two BCC scATAC-seq datasets, we instead

clustered the cells using $\hat{\theta}_{ir}$ values, which are orthogonal to the CNA signals based on

total coverage, across the segments with more than 500 SNPs. Then hierarchical

clustering is performed on the Euclidean distance using the "ward.D2" method.

Because the subclones for the SNU601 sample were identified from the scDNA-seq

data, we adopted a supervised strategy to assign each cell in the SNU601 scATAC-seq

dataset into different subclones. First, we identified 10 marker regions-- chr1b, 3b-d, 4b,

7a, 11b, 13b, and 20b-c that help to differentiate the cells into the six major subclones

based on the subclone specific copy number profiles from the scDNA-seq data.

Combining the haplotype profiles across the ten regions for each cell enables

assignment of the cells into one of the six subclones with high confidence. The posterior

probability of cell i coming from clone k was

804
$$P(Clone_i = k | \hat{\rho}_i, \hat{\theta}_i) = \frac{P(\hat{\rho}_i, \hat{\theta}_i | Clone_i = k)\pi_k}{\sum_{k'} P(\hat{\rho}_i, \hat{\theta}_i | Clone_i = k')\pi_{k'}},$$

805   where $k \in \{1\sim6\}$ for the six clones and π_k is the prior probability that a randomly

806   sampled cell coming from the k^{th} clone, which was set to uniform in the analysis.. In the

807   formula,

808
$$P(\hat{\rho}_i, \hat{\theta}_i | Clone_i = k) = \prod_x P(\hat{\theta}_{ix}, \hat{\rho}_{ix} | Clone_i = k)$$

809
$$= \prod_x \phi\left(\frac{\hat{\theta}_{ix} - \theta_{kx}}{\sqrt{n_{ix}\theta_{kx}(1 - \theta_{kx})}}\right) \phi\left(\frac{\hat{\rho}_{ix} - \rho_{kx}}{\sigma_\rho}\right),$$

where x indexes the ten marker regions; $\hat{\theta}_{ix}$ and $\hat{\rho}_{ix}$ are the estimated major haplotype

proportion and relative coverage for cell i in the scATAC-seq data; θ_{kx} and ρ_{kx} are the

"known values" for specific haplotype profiles for clone k derived from the paired

scDNA-seq data; and n_{ix} is the number of total read counts in the x^{th} marker region for

cell i. Each cell was assigned into one of the six subclones by maximizing the posterior

probability with the confidence score being the posterior probability of the assigned

clone.

ScATAC-seq data analysis

To investigate the relationships between allele-specific CNAs and chromatin

accessibility, we processed the peak signals in addition to the allele-specific CNAs for

each cell in scATAC-seq data. For the two public BCC samples, the peak-by-cell

matrices were obtained from GSE129785. We log-transformed the count matrices,

selected the peaks in >10% cells, regressed out cell total coverage for each peak by

linear regression, and projected the cells onto the UMAP plot using genome-wide peak

signals[53]. The cell type identify for each cluster was retrieved from the previous study[26].

To further explore intratumor heterogeneity, we selected the cells labeled as tumor

cells, repeated the pre-processing steps described above, and projected the tumor cells

onto the UMAP plot. Then DNA level information and epigenetic signals for each cell

were integratively visualized.

For the SNU601 scATAC-seq dataset, scATAC-pro[54] was used to call peaks and

generate the peak-by-cell matrix. We first filtered out the cells that have proportions of

fragments on the detected peaks <0.4 and or total peaks outside of the range

15,000~100,000, and filtered out the peaks observed in less than 10% of cells. After the

matrix was log-transformed, we regressed out cell total coverage for each peak by

linear regression. Using Louvain clustering, the cells in two clusters with extreme total

fragments were removed. After cell filtering, we repeated the process and projected the

cells onto the UMAP plot. Then the clonal assignment based on the DNA information

and the peak signals were integrated at the single-cell level

Differentially accessible peaks (DAPs) identification with copy number

adjustment for scATAC-seq data

To identify differentially accessible peaks (DAPs) after accounting for copy number

differences between two clones, we developed a statistical test based on a generalized

log-likelihood ratio (LLR) statistic. We first define some necessary terms: Given a

segmentation of the genome, for a given clone c, let $\theta_c = \{(L_R, Z_R)\}$ be the copy number

profile, which can be expressed as a vector of region lengths (L_R) and average copy

numbers (Z_R) across all regions R. For peak k, we define function $f_k(\theta_c)$ to be the

proportion of genomic DNA in the peak region based on the copy number profile of

clone c, computed as follows:

848
$$f_k(\theta_c) = \frac{L_{r(k)} Z_{r(k)}}{\sum_r L_r Z_r} \times \frac{L_k}{L_{r(k)}},$$

849   where $r(k)$ is the region r to which peak k belongs and L_r is the length of peak k.

850   For peak k and clone c, let Y_{ck} be the total read count across the cells in the clone, and

851   let N_c be the total read count across all peaks summed across the cells in the clone. Y_{cg}

852   can be modeled by Binomial sampling,

853
$$Y_{ck} \sim Binomial(N_c, f_k(\theta_c) p_{ck}),$$

where, p_{ck} is the relative probability of detecting reads in the peak k region after

adjusting for the copy numbers. Between clones with different copy number profiles,

differences in p_{ck} is evidence for chromatin remodeling. Thus, pairwise comparisons

were performed for each branch (clone c_1 versus clone c_2) and for each peak, using the

generalized likelihood ratio test with the hypothesis $H_0: p_{c_1 k} = p_{c_2 k} = p_{0k}$ and $H_1: p_{c_1 k} \neq$

$p_{c_2 k}$.

Let $l(p)$ denote the log-likelihood function of Y_{ck} under this model, and $\hat{p}$ denote the

MLE. The LLR for GLRT follows the Chi-squared distribution as below with derivation of

MLE estimates in the Supplementary Methods.

863
$$LLR_k = -2 \log \frac{\max\limits_{p_{0k} \in p} l(p_{0k})}{\max\limits_{p_{c_1 k}; p_{c_2 k} \in p} l(p_{c_1 k}, p_{c_2 k})} = 2\ell(\hat{p}_{c_1 k}, \hat{p}_{c_2 k}) - 2\ell(\hat{p}_{0k}) \sim X_1^2,$$

864   Using the GLRT, a peak is considered as a DAP if its FDR adjusted p-value<0.01 for

865   the pairwise clonal comparison.

866   Simultaneously, the GLRT under the binomial model without copy number adjustment

867   was also performed for comparison:

868
$$Y_{ck} \sim Binomial(N_c, p_{ck}),$$

The same criterion was used to identify DAPs for each pairwise clonal comparison

(FDR adjusted p-values<0.01). Each DAP was further mapped to the genes that are

potentially regulated based on the $\pm$ 2,000 bp distance on the genome.

Data availability

The patient scDNA-seq and linked-read sequencing data generated for this study are

available under dbGAP identifier phs001711. The scATAC-seq dataset is available in the

National Institute of Health's SRA repository under accession PRJNA674903. There are

no restrictions on data availability or use. The other patient scDNA-seq data were

obtained from dbGAP under accession phs001818.v3.p1[35] and phs001711[12]. The cell

line scDNA-seq dataset was from the Sequence Read Archive (SRA) under accession

PRJNA498809. The public scATAC-seq data and whole exome sequencing data were

obtained from the SRA under accession PRJNA532774[26] and PRJNA533341[39].

<u>Code availability</u>

Alleloscope is available on GitHub at <u>https://github.com/seasoncloud/Alleloscope</u>.

<u>Reference</u>

1.   Baylin, S.B. & Jones, P.A. A decade of exploring the cancer epigenome - biological and
885   translational implications. *Nat Rev Cancer* **11**, 726-734 (2011).
886   2.   Sandoval, J. & Esteller, M. Cancer epigenomics: beyond genomics. *Curr Opin Genet Dev*
887   **22**, 50-55 (2012).
888   3.   Greaves, M. & Maley, C.C. Clonal evolution in cancer. *Nature* **481**, 306-313 (2012).
889   4.   Burrell, R.A., McGranahan, N., Bartek, J. & Swanton, C. The causes and consequences
890   of genetic heterogeneity in cancer evolution. *Nature* **501**, 338-345 (2013).
891   5.   Chen, H., Bell, J.M., Zavala, N.A., Ji, H.P. & Zhang, N.R. Allele-specific copy number
892   profiling by next-generation DNA sequencing. *Nucleic Acids Res* **43**, e23 (2015).
893   6.   Favero, F. et al. Sequenza: allele-specific copy number and mutation profiles from tumor
894   sequencing data. *Ann Oncol* **26**, 64-70 (2015).
895   7.   Ha, G. et al. TITAN: inference of copy number architectures in clonal cell populations
896   from tumor whole-genome sequence data. *Genome Res* **24**, 1881-1893 (2014).
897   8.   Shen, R. & Seshan, V.E. FACETS: allele-specific copy number and clonal heterogeneity
898   analysis tool for high-throughput DNA sequencing. *Nucleic Acids Res* **44**, e131 (2016).
899   9.   Jamal-Hanjani, M. et al. Tracking the Evolution of Non-Small-Cell Lung Cancer. *N Engl*
900   *J Med* **376**, 2109-2121 (2017).
901   10.   Zaccaria, S. & Raphael, B.J. Characterizing allele- and haplotype-specific copy numbers
902   in single cells with CHISEL. *Nat Biotechnol* (2020).
903   11.   Van Loo, P. et al. Allele-specific copy number analysis of tumors. *Proc Natl Acad Sci U*
904   *S A* **107**, 16910-16915 (2010).
905   12.   Andor, N. et al. Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines
906   reveals rules of in vitro evolution. *NAR Genom Bioinform* **2**, lqaa016 (2020).
907   13.   Bakker, B. et al. Single-cell sequencing reveals karyotype heterogeneity in murine and
908   human malignancies. *Genome Biol* **17**, 115 (2016).
909   14.   Garvin, T. et al. Interactive analysis and assessment of single-cell copy-number
910   variations. *Nat Methods* **12**, 1058-1060 (2015).
911   15.   Kim, C. et al. Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated
912   by Single-Cell Sequencing. *Cell* **173**, 879-893 e813 (2018).
913   16.   Laks, E. et al. Clonal Decomposition and DNA Replication States Defined by Scaled
914   Single-Cell Genome Sequencing. *Cell* **179**, 1207-1221 e1222 (2019).
915   17.   Navin, N. et al. Tumour evolution inferred by single-cell sequencing. *Nature* **472**, 90-94
916   (2011).

18. Velazquez-Villarreal, E.I. et al. Single-cell sequencing of genomic DNA resolves sub-
clonal heterogeneity in a melanoma cell line. *Commun Biol* **3**, 318 (2020).
19. Wang, Y. et al. Clonal evolution in breast cancer revealed by single nucleus genome
sequencing. *Nature* **512**, 155-160 (2014).
20. Klemm, S.L., Shipony, Z. & Greenleaf, W.J. Chromatin accessibility and the regulatory
epigenome. *Nat Rev Genet* **20**, 207-220 (2019).
21. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J. Transposition
of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).
22. Buenrostro, J.D. et al. Single-cell chromatin accessibility reveals principles of regulatory
variation. *Nature* **523**, 486-490 (2015).
23. Corces, M.R. et al. The chromatin accessibility landscape of primary human cancers.
Science **362** (2018).
24. Granja, J.M. et al. Single-cell multiomic analysis identifies regulatory programs in
mixed-phenotype acute leukemia. *Nat Biotechnol* **37**, 1458-1465 (2019).
25. Litzenburger, U.M. et al. Single-cell epigenomic variability reveals functional cancer
heterogeneity. *Genome Biol* **18**, 15 (2017).
26. Satpathy, A.T. et al. Massively parallel single-cell chromatin landscapes of human
immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* **37**, 925-
936 (2019).
27. Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
Methods* **14**, 975-978 (2017).
28. Cabal-Hierro, L. et al. Chromatin accessibility promotes hematopoietic and leukemia
stem cell activity. *Nat Commun* **11**, 1406 (2020).
29. Dravis, C. et al. Epigenetic and Transcriptomic Profiling of Mammary Gland
Development and Tumor Models Disclose Regulators of Cell State Plasticity. *Cancer
Cell* **34**, 466-482 e466 (2018).
30. Guilhamon, P. et al. Single-cell chromatin accessibility in glioblastoma delineates cancer
stem cell heterogeneity predictive of survival. *bioRxiv*, 370726 (2020).
31. Pan, D. et al. A major chromatin regulator determines resistance of tumor cells to T cell-
mediated killing. *Science* **359**, 770-775 (2018).
32. Qu, K. et al. Chromatin Accessibility Landscape of Cutaneous T Cell Lymphoma and
Dynamic Response to HDAC Inhibitors. *Cancer Cell* **32**, 27-41 e24 (2017).
33. Shaffer, S.M. et al. Rare cell variability and drug-induced reprogramming as a mode of
cancer drug resistance. *Nature* **546**, 431-435 (2017).
34. Shu, S. et al. Synthetic Lethal and Resistance Interactions with BET Bromodomain
Inhibitors in Triple-Negative Breast Cancer. *Mol Cell* **78**, 1096-1113 e1098 (2020).
35. Sathe, A. et al. The cellular genomic diversity, regulatory states and networking of the
metastatic colorectal cancer microenvironment. *bioRxiv* (2020).
36. Bell, J.M. et al. Chromosome-scale mega-haplotypes enable digital karyotyping of cancer
aneuploidy. *Nucleic Acids Res* **45**, e162 (2017).
37. Greer, S.U. et al. Linked read sequencing resolves complex genomic rearrangements in
gastric cancer metastases. *Genome Med* **9**, 57 (2017).
38. Zheng, G.X. et al. Haplotyping germline and cancer genomes with high-throughput
linked-read sequencing. *Nat Biotechnol* **34**, 303-311 (2016).

39. Yost, K.E. et al. Clonal replacement of tumor-specific T cells following PD-1 blockade.
Nat Med **25**, 1251-1259 (2019).
40. Yu, J. et al. REC8 functions as a tumor suppressor and is epigenetically downregulated in
gastric cancer, especially in EBV-positive subtype. *Oncogene* **36**, 182-193 (2017).
41. McFarlane, R.J. & Wakeman, J.A. Meiosis-like Functions in Oncogenesis: A New View
of Cancer. *Cancer Res* **77**, 5712-5716 (2017).
42. Aqeilan, R.I. et al. Loss of WWOX expression in gastric carcinoma. *Clin Cancer Res* **10**,
3053-3058 (2004).
43. Baryla, I., Styczen-Binkowska, E. & Bednarek, A.K. Alteration of WWOX in human
cancer: a clinical view. *Exp Biol Med (Maywood)* **240**, 305-314 (2015).
44. Watkins, T.B.K. et al. Pervasive chromosomal instability and karyotype order in tumour
evolution. *Nature* (2020).
45. Gupta, I. et al. Single-cell isoform RNA sequencing characterizes isoforms in thousands
of cerebellar cells. *Nat Biotechnol* (2018).
46. Lebrigand, K., Magnone, V., Barbry, P. & Waldmann, R. High throughput error
corrected Nanopore single cell transcriptome sequencing. *Nat Commun* **11**, 4025 (2020).
47. Singh, M. et al. High-throughput targeted long-read single cell sequencing reveals the
clonal and transcriptional landscape of lymphocytes. *Nat Commun* **10**, 3120 (2019).
48. Zhu, C., Preissl, S. & Ren, B. Single-cell multimodal omics: the power of many. *Nat*
Methods **17**, 11-14 (2020).
49. Wang, R., Lin, D.Y. & Jiang, Y. SCOPE: A Normalization and Copy-Number Estimation
Method for Single-Cell DNA Sequencing. *Cell Syst* **10**, 445-452 e446 (2020).
50. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).
51. McKenna, A. et al. The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297-1303 (2010).
52. Koboldt, D.C. et al. VarScan 2: somatic mutation and copy number alteration discovery
in cancer by exome sequencing. *Genome Res* **22**, 568-576 (2012).
53. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and
projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).
54. Yu, W., Uzun, Y., Zhu, Q., Chen, C. & Tan, K. scATAC-pro: a comprehensive
workbench for single-cell chromatin accessibility sequencing data. *Genome Biol* **21**, 94
(2020).
996

Acknowledgements

The work is supported by the National Institutes of Health [P01HG00205ESH to B.T.L.,

S.M.G. and H.P.J., 5R01-HG006137-07 and 1U2CCA233285-01 to C-Y.W. and to N.R.Z.,

1R35HG011292-01 to B.T.L.]. Additional support to HPJ came from the Research Scholar

Grant, RSG-13-297-01-TBG from the American Cancer Society, Clayville Foundation and

the Gastric Cancer Foundation.

Author contributions

C.-Y.W. and N.R.Z. conceived the computational methods and designed the study with

help from H.P.J. C.-Y.W. developed and implemented the computational methods and

conducted all data analyses. B.T.L. helped in data interpretation. B.T.L., H.K. and A.S.

performed all related sample preparation and sequencing. S.M.G. performed data pre-

processing and coordinated data transfer. H.P.J. advised all experiments and data

collection. C.-Y.W., N.R.Z, and H.P.J. wrote the manuscript. All authors read and

approved the final manuscript.

Competing interests

The authors declare no competing interests.

1013

Figure 1.

Fig. 1: Overview of allele-specific copy number estimation of single cells with Alleloscope. 1. The algorithm operates on raw read count matrices for reference allele (Ref) and alternative allele (Alt) computed from single cell DNA or ATAC sequencing. 2. First, we obtain a segmentation of the genome based on sample-matched whole genome or whole exome sequencing data using FALCON[5]. If scDNA-seq is available, cells can be pooled to derive a pseudo-bulk. 3. We first define $\hat{\theta}_{ir}$ as the major haplotype proportion for cell i for region r derived from the segmentation. Then for each region r, Alleloscope simultaneously phase SNPs ($\hat{I}_j$) and estimate cell major haplotype proportion ($\hat{\theta}_{ir}$) by expectation maximization (EM) algorithm. In the E-step, information is pooled across cells to estimate the phasing of each SNP. In the M-step, information is pooled across all SNPs in the region are pooled to estimate the major haplotype proportion $\hat{\theta}_{ir}$ for each cell. The toy example shows a scenario with two cells for a region containing 5 SNPs, with cell 2 carrying an amplification of the major haplotype (in pink). For each cell and each SNP, alleles that are observed in a sequenced read are bolded in black (we assume that only one read is observed, reflecting the sparsity of the data). The true phase (I_j) of the SNPs and the true major haplotype proportion (θ_{ir}) are shown. 4. After $\hat{\theta}_{ir}$ are estimated for each region r, $\hat{\theta}_{ir}$'s are pooled across all regions to identify candidate normal cells and candidate normal regions for computing a normalized coverage $\hat{\rho}_{ir}$ for region r in cell i. Matched normal can be specified if available. 5. Alleloscope assigns integer allele-specific copy numbers to each cell for each region based on the ($\hat{\rho}_{ir}, \hat{\theta}_{ir}$) pairs.

Figure 2.

Fig. 2: **Validation of the Alleloscope results on the P5931 gastric cancer patient sample and linked-read sequencing data.** (a) Illustration of the validation scheme using linked-read sequencing data. Phasing accuracy and accuracy in allele-specific copy number state estimation are used to access performance of the method. (b) Hierarchical clustering of cells in the P5931t sample based on allele-specific copy numbers given by Alleloscope, showing normal cells and 4 main clones, as well as a number of small clones marked by highly confident low-frequency mutations. M: Major haplotype, m: minor haplotype. (c) $(\hat{\rho}_{ir}, \hat{\theta}_{ir})$ estimated by Alleloscope for four regions, colored by the inferred haplotype profile. Note that clusters fall on canonical points corresponding to discrete allele-specific copy number configurations. Phasing accuracy for each region is shown in the plot title. In the color legend, M and m represent the "Major haplotype" and "minor haplotype" respectively. (d) Similar to (c), with $\tilde{\theta}_{ir}$ estimated using known SNP phases from matched linked-read sequencing data, colored by the haplotype profiles assigned in (c) using Alleloscope without the given phasing information. Accuracy in allele-specific copy number state estimation (Genotyping accuracy) is labeled in the plots.

Figure 3.

Fig. 3: Across multiple cancer types, Alleloscope detects loss-of-heterozygosity
events and multi-allelic copy number aberrations, delineating complex subclonal
structure which are invisible to total copy number analysis. (a) The Cell Ranger
hierarchical clustering result for P6198t with copy-neutral regions labeled (total 512 cells).
(b) Top: FALCON segmentation of P6198 chr5 into two regions with different allele-
specific copy number profiles. Bottom: Detailed haplotype profiles of the two regions from
Alleloscope, showing that the first region is diploid across cells and the second region has
a loss-of-heterozygosity for a subpopulation of cells. The a and b following the
chromosome number denote two ordered segments. (c) Single cell allele-specific
estimates $(\hat{\rho}, \hat{\theta})$, colored by assigned haplotype profiles, for select regions in the samples
P6198t (metastasized colorectal cancer sample), SNU601 (gastric cancer cell line),
P6335 (colorectal cancer sample), and BC10X (breast cancer cell line). In the color
legend, M and m represent the "Major haplotype" and "minor haplotype" respectively. The
lower-case letters following the chromosome number in the titles denote the ordered
genomic segments.

1069

Figure 4.

Fig. 4: Alleloscope multiomic analysis of scATAC-seq data of a basal cell carcinoma sample (SU008[26]). (a) Genotype profiles for six example regions for cells in scATAC-seq data. The regions are taken from segmentation of matched whole exome sequencing (WES) data. Each dot represents a cell-specific $(\hat{\rho}_{ir}, \hat{\theta}_{ir})$ pair. Cells are colored by annotation derived from peak signals[26], Tumor: tumor cells, Fibro: fibroblasts, Endo: Endothelial cells]. Density contours are computed for each cell type (tumor, fibroblasts, endothelial) separately and shown by color on the plot. The lower-case letters following the chromosome number in the titles denote the ordered genomic segments. (b) Pipeline for multi-omics analysis integrating allele-specific copy number estimates and chromatin accessibility peak signals on ATAC-seq data. (c) Hierarchical clustering of cells by major haplotype proportion ($\hat{\theta}$) allows the separation of tumor cells from normal cells, as well as the differentiation of a subclone within the tumor cells. The marker region on chr4a separating the two tumor subclones is highlighted. (d) Integrated visualization of

chr4a major haplotype proportion ($\widehat{\theta}_{ir}$) and genome-wide peak profile. Left: UMAP
projection of the 788 cells in the dataset by their genome-wide peak profile, colored by
$\widehat{\theta}_{ir}$. The cell type annotation (endothelial, fibroblasts, and tumor cells) is labeled in the
plot. Middle: UMAP projection of only the 308 tumor cells by their genome-wide peak
profile shows two well-separated clusters: peaks1 and peaks2. Right: Density of $\widehat{\theta}_{ir}$
values for the peaks1 and peaks2 subpopulations. (e) Intratumor heterogeneity of SU008
is shaped by a subclonal LOH of chr4a followed by subsequent genome-wide chromatin
remodeling leading to three subpopulations: Clone 1 which does not carry the chr4a LOH
(peaks cluster 1), Clone 2 carrying the chr4a LOH (peaks cluster 1), and remodeled clone
2 (peaks cluster 2).

1094

Figure 5.

Fig. 5: Alleloscope analysis of scDNA-seq and scATAC-seq data reveals complex subclonal heterogeneity in the SNU601 gastric cancer cell line. (a) Genome segmentation using HMM on the pooled total coverage profile computed from scDNA-seq data. (b) Single cell allele-specific copy number profiles $(\hat{\rho}, \hat{\theta})$ for five regions in scDNA-seq and scATAC-seq data. Cells are colored by haplotype profiles according to legend in Fig. 5c. (c) Tumor subclones revealed by hierarchical clustering of allele-specific copy number profiles from the scDNA-seq data. Genotypes of the five regions shown in Fig. 5b, for three example cells, are shown in the left. The haplotype structures for the 5 regions in Fig. 5b of three cells randomly chosen from Clone 1, 2, and 3, are shown to the left of the heatmap. In the color legend, M and m represent the "Major haplotype" and "minor haplotype" respectively. The six clones selected for downstream analysis in scATAC-seq data are labeled in the plot.

Fig. 6: Integrative analysis of allele-specific copy number and chromatin accessibility for SNU601 ATAC sequencing data. (a) UMAP projection of genome-wide scATAC-seq peak profile on 2,753 cells. The same group of cells were clustered into one of the six subclones based on their allele-specific copy number profiles across the 10 selected regions. Cells in different subclones are labeled with different colors, using the same color scheme as that for the subclone labels in Fig. 4c. The number of cells colored in each UMAP is shown at the bottom-right corners. (b) A highly probable lineage history of SNU601, with copy number alterations (CNAs) and differentially accessibility peaks (DAPs) marked along each branch. P-values of the tests for association between DAPs and CNAs are shown along each branch. For two example DAP genes, pooled peak signals for each subclone are shown as inset plots.

Table 1. Comparison between Alleloscope and CHISEL using two benchmark strategies.

Dataset	Method	Sensitivity	Specificity
Benchmark with matched linked-read sequencing data			
P5915	Alleloscope	0.9373	0.9915
	CHISEL (before correction)	0.7284	0.9757
	CHISEL (after correction)	0.7786	0.9722
P5931	Alleloscope	0.9402	0.9986
	CHISEL (before correction)	0.7520	0.9434
	CHISEL (after correction)	0.0112	0.9508
P6198	Alleloscope	0.9433	0.9666
	CHISEL (before correction)	0.9397	0.9311
	CHISEL (after correction)	0.9700	0.9359
P6335	Alleloscope	0.9671	0.9906
	CHISEL (before correction)	0.7858	0.9873
	CHISEL (after correction)	0.8404	0.9943
P6461	Alleloscope	0.8638	0.9904

	CHISEL (before correction)	0.7932	0.9624
	CHISEL (after correction)	0.8143	0.9411
Benchmark with subsampling			
50% subsample	Alleloscope	0.9261	0.9905
	CHISEL (before correction)	0.9267	0.9383
	CHISEL (after correction)	0.9496	0.9446
25% subsample	Alleloscope	0.8249	0.9525
	CHISEL (before correction)	0.6515	0.6412
	CHISEL (after correction)	0.6533	0.6475

1123