A computational approach to evaluate the combined effect of SARS-CoV-2 RBD mutations and ACE2 receptor genetic variants on infectivity: The COVID-19 host-pathogen nexus

Dana Ashoor [1,†], Noureddine Ben Khalaf [1,†], Maryam Marzouq [1], Hamdi Al Jarjanazi [1,3], Sadok Chelif [2] and M.Dahmani Fathallah [1,*]

[1] Department of Life Sciences, Health Biotechnology Program # King Fahad Chair for Health Biotechnology, College of Graduate Studies, Arabian Gulf University, PO Box 26671, Manama, Kingdom of Bahrain
[2] Department of Family and Community Medicine, College of Medicine and Medical Sciences, Arabian Gulf University, PO Box 26671, Manama, Kingdom of Bahrain
[3] Environmental Monitoring and Reporting Branch, Ontario Ministry of the Environment, Conservation and Parks, Toronto, Ontario, Canada.
† These authors contributed equally to this work.
* Correspondence: d.fathallah@agu.edu.bh; Tel.: Tel: +973 17 239602; Fax: +973 17 239484

Abstract: SARS-CoV-2 infectivity is largely determined by the virus Spike protein binding to the ACE2 receptor. Meanwhile, marked infection rate differences were reported between populations and individuals. To understand the disease dynamic, we developed a computational approach to study the implications of both SARS-CoV-2 RBD mutations and ACE2 polymorphism on the stability of the virus-receptor complex. We used the 6LZG PDB RBD/ACE2 3D model, the mCSM platform, the LigPlot+ and PyMol software to analyze the data on SARS-CoV-2 mutations and ACE variants retrieved from GISAID and Ensembl/GnomAD repository. We observed that out of 351 RBD point mutations, 83% destabilizes the complex according to free energy ($\Delta\Delta G$) differences. We also spotted variations in the patterns of polar and hydrophobic interactions between the mutations occurring in 15 out of 18 contact residues. Similarly, comparison of the effect on the complex stability of different ACE2 variants showed that the pattern of molecular interactions and the complex stability varies also according to ACE2 polymorphism. We infer that it is important to consider both ACE2 variants and circulating SARS-CoV-2 RBD mutations to assess the stability of the virus-receptor association and evaluate infectivity. This approach might offers a good molecular ground to mitigate the virus spreading.

Keywords: COVID-19; SARS-CoV-2; angiotensin converting enzyme 2 receptor; spike; receptor binding domain; virus-host interaction; SARS-CoV-2 mutations

1. Introduction

The emergent coronavirus, called Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has been spreading globally and efficiently since late 2019 (Elflein, 2020). SARS-CoV-2 is a new member of the coronaviruses family that includes SARS coronavirus (SARS-CoV) and Middle East Respiratory Syndrome coronavirus (MERS-CoV) (Chan et al., 2020). SARS-CoV-2 is the etiologic agent of the ongoing coronavirus disease (COVID-19) pandemic that claimed over two and a half millions life to date (EDCDP, 2020). The number of active cases per 100 thousand population estimates the prevalence of COVID-19 while the incidence is the number of new cases per time unit. The prevalence and incidence rate are good indicators for the tracking of infectious outbreaks (Noordzij et al., 2010). Indeed, they are very useful to evaluate the effectiveness of the mitigation efforts in the control of the disease spreading (Roser, 2020). Even though the spreading of the SARS-CoV-2 virus is global, the prevalence/incidence of the disease varies throughout the world, for some regions being more severely hit then others (EDCDP, 2020;Elflein, 2020;Roser,

2020). Even the overall case fatality ratio of COVID-19 varies between location and intensity of
transmission (Lewnard and Lo, 2020;Ruan, 2020;Verity et al., 2020). Such discrepancies may be due
to several factors that affect the virus infectious potency (infectivity) such as the health care
prevention system organization and the stringency of the confinement measures (Koo et al., 2020),
as well as the economic differences. Nevertheless, other important factors intrinsic to the virus
structure and the host genetic background are crucial determinants of the SARS-CoV-2 infectivity.
SARS-CoV-2 Invasion of human host cell is essentially but not exclusively mediated by the
interaction of the virus capsid spike protruding S protein with the Angiotensin Convertase Enzyme
2 (ACE2) (Belouzard et al., 2012;Alifano et al., 2020). ACE2 is expressed on the surface of various
human body cell-types (Walls et al., 2020;Zhao et al., 2020) where it displays a zinc
metalloproteinase activity (Tipnis et al., 2000). The interaction ACE2/SARS-CoV-2 occurs through
the receptor-binding domain (RBD) sequence of the S1 chain of the spike protein (Lan et al.,
2020;Yan et al., 2020). This functional domain is highly prone to mutations (Nelson-Sathi et al.,
2020). Meanwhile, the human ACE2 receptor gene that is located on chromosome X is highly
polymorphic with hundreds of genetic variants identified to date in various populations (Cao et al.,
2020). Noteworthy are the 1700 ACE2 variants reported in the Chinese populations though few
variants are highly frequent and the majority of them are very rare (Lippi et al., 2020). Interestingly
some ACE2 variants was shown to be significantly associated with the onset of diseases such
arterial hypertension, diabetes mellitus, cerebral stroke, coronary artery disease, heart septal wall
thickness and ventricular hypertrophy all considered as comorbidity factors of COVID-19
(Kowalczuk et al., 2008;Lu et al., 2012;Pinheiro et al., 2019;Calcagnile et al., 2020). The complexity of
the SARS-CoV-2 S protein association with the ACE2 receptor is highlighted in a number of X-ray
crystallography models. These are namely the 6LZG (SARS-CoV-2 Spike RBD /ACE2) complex, the
2AJF (SARS-CoV Spike RBD /ACE2) complex (Wang et al., 2020), the 6M0J (SARS-CoV-2 Spike RBD
/ACE2) complex, and the 6VW1 (chimeric SARS-CoV/SARS-CoV-2 Spike RBD /ACE2) complex
(Goodsell et al., 2020). These models have different resolution and are available at RCBS Protein
Data Bank (https://www.rcsb.org/). Studies using these models have shown that the affinity of the
RBD SARS-CoV-2 to the human ACE2 varies between ACE2 genetic variants suggesting that some
people might be genetically protected where others are susceptible to COVID-19 (Lippi et al.,
2020;Othman et al., 2020). Indeed, by measuring the ACE2 affinity for SARS-CoV-2 Spike using
docking simulations, Calcagnile et al. (Calcagnile et al., 2020) showed that ACE2 variant S19P,
which is more frequent in Africans and K26R more frequent in Europeans, are respectively
protective and predisposing genetic factors to COVID-19. Meanwhile, other studies denied the
association between ACE2 genetic polymorphism and the susceptibility to SARS infection (Chiu et
al., 2004) and to COVID-19 (Lopera Maya et al., 2020). Nevertheless, the virus-host-cell interaction
involves specific contacts between two structurally defined molecular entities. Therefore, it is very
likely that the interaction between the SARS-CoV-2 and the human cell surface receptor ACE2 and
thus the virus infectivity is determined by the genetic variations of both the pathogen and the
receptor interacting sequences. This can reflect a higher magnitude of complexity in the interplay
between this new viral pathogen and its human host.
In this paper, we report the development of a computational approach designed to examine
this postulate. We first compiled the mutations observed in different isolates of the SARS-CoV-2 S
protein RBD region as well as ACE2 receptor variants. Secondly, we assessed the stability of the
complex between the virus RBD mutants and ACE2 isoform 1. Then we studied the structural basis
of these molecular interactions. Comparison of the binding energy variations ($\Delta\Delta G$) between the
combinations of over 200 ACE2 genetic variants with 350 RBD mutated forms showed a gradient of
stability among these combinations. We also observed significant variations in the patterns of polar
and hydrophobic bonds between 18 RBD mutated contact residues and their partners at the
interface of the complex with ACE2 Isoform 1.
We inferred from these data that both ACE2 receptor genetic polymorphism and the circulating
SARS-CoV-2 RBD genetic type determine the stability of the virus-receptor association and might
affect the infectivity of SARS-CoV-2. Furthermore, these molecular interaction patterns highlight

the level of complexity of the interplay between SARS-CoV-2 and its major ACE2 receptor. The data
also point out to the benefit this computational approach may bring to the rapid evaluation of the
infectivity of a given viral genetic subtype in consideration of an individual or a group dominant
ACE 2 variant. This might help fine-tuning the prevention strategies against the spreading of the
virus.
104

2. Materials and Methods

2.1. SARS-CoV-2 S protein RBD mutations and ACE2 genetic variants data retrieval

We first retrieved the mutations located at the SARS-CoV-2 Spike protein RBD region (351
missense mutations), Wuhan strain (NCBI Reference: YP_009724390.1). The data was obtained from
the SARS-CoV-2 Spike protein mutation table compiled by the Singapore Bioinformatics Institute
(SBI)   at   (https://mendel.bii.a-star.edu.sg/METHODS/corona/beta/MUTATIONS/hCoV-
19_Human_2019_WuhanWIV04/hCoV-19_Spike_new_mutations_table.html) based on sequence
information provided by GISAID initiative (https://www.gisaid.org/). Data for genetic variants of
ACE2 was from the Ensembl dbSNP, GnomAd and UNIPROT databases.
To identify the amino acids differences in the spike RBD between the new Wuhan strain SARS-
CoV-2 and previously known SARS-CoV (NCBI Reference: YP_009825051.1), the sequence of the
RBD domain of both viruses was determined using Pfam sequence analysis (https://pfam.xfam.org)
of the S full -protein. SARS-CoV-2 RBD protein sequence was aligned with SARS-CoV RBD protein
using Clustal Omega alignment tool (https://www.ebi.ac.uk/Tools/msa/clustalo/). The locations of
the variant amino acids were identified and the reported mutations of SARS-CoV-2 at these
locations were examined for possible reversed mutations to the SARS-CoV sequence. The effect of
these mutations on the complex stability with the human ACE2 receptor was examined as
described in the next section.

2.2. RBD mutation effect analysis

The PDB structure (PDB ID 6LZG) of the complex formed between the SARS-CoV-2 Spike
protein RBD and the human ACE2 receptor isoform 1 was downloaded from the RCSB database
(https://www.rcsb.org). ACE2 isoform 1 sequence correspond to the ACE2 sequence that was used
to generate the 6LZG 3D model. This structural complex model was taken as reference model for
this study. The structure was cleaned from water and heteroatoms using PyMol software (PyMol
Molecular). The mCSM server, a web server (http://biosig.unimelb.edu.au/mcsm/) that predicts the
effects of mutations in proteins using graph-based signatures platform (Pires et al., 2020) was used
to evaluate the effect of the mutations reported in the RBD domain on the stability of the virus-
receptor complex. The prediction relies on graph-based signatures that encode distance patterns
between atoms and are used to represent the protein residue environment and to train predictive
models. The models predict the stability changes through free energy ($\Delta\Delta G$) difference due to a
single nonsynonymous substitution in the amino acid chain of the protein-protein complex. Finally,
mutations were ranked by their $\Delta\Delta G$ values. Site Directed Mutator server (SDM)
(http://marid.bioc.cam.ac.uk/sdm2/) (Worth et al., 2011) was used to create the 23 contact residues
mutants and chain replacement was used to build the complexes using PyMol (Schrödinger,
2010;Verity et al., 2020). Interacting residues in the generated structures, involving polar and
hydrophobic interactions from both receptor and ligand, were determined using LigPlot+ software
(Laskowski and Swindells, 2011) and PyMol software was used to illustrate the interacting interface
in presence of the most destabilizing mutations in the RBD.

2.3. hACE2 variants analysis

The protein sequence for human ACE2 Isoform 1 (hACE2) was downloaded from the Uniprot
data bank (https://www.uniprot.org/) (Uniprot ID: Q9BYF1-1) and used as a reference sequence in
this study. We analyzed 231 human ACE2 SNPs causing protein sequence variations collected from

the Ensembl dbSNP,GnomAD and UNIPROT databases. We used the PDB structure (PDB ID
6LZG), LigPlot+ software (Laskowski and Swindells, 2011), and mCSM server
(http://biosig.unimelb.edu.au/mcsm/) to identify possible amino acid substitutions that affect the
virus-receptor complex stability, and their interaction pattern. We then built protein complexes
(hACE2/RBD) using SDM server and Pymol software to analyze S protein mutations effect on
complex stability for selected hACE2 variants with predicted destabilizing effect on the original
complex. We used LigPlot+ and PyMol softwares to analyze the molecular interactions at the
complex interface.

3. Results

3.1. SARS-CoV-2 S protein RBD mutations retrieval

This study focused on the investigation of the structural variations of the Spike's RBD of the
SARS-CoV-2 virus from the first reported Wuhan strain (NCBI Reference: YP_009724390.1).
Alignment of the SARS-CoV surface glycoprotein and SARS-CoV-2 spike RBD sequence showed
74% identity and one insertion (V483) (data not shown). Random search of the GISAID data bank
yielded 351 non-synonymous mutations with some positions changing from one up to eight
different residues as shown in Figure 1. Out of these mutations, 24 are reverse mutations to the
SARS-CoV RBD sequence.

Figure 1. A Schematic diagram of the of SARS-CoV-2 S spike protein sequence (purple letters) and the mutations in the Receptor Binding Domain (RBD) aligned with
SARS-CoV sequence (green letters). The S protein is cleaved into S1 and S2 subunit at the polybasic furin cleavage site. Stabilizing and destabilizing mutations of the RBD
domains are represented with blue and red letters respectively. Asterisks display amino acid variations between SARS-CoV-2 and SARS-CoV RBDs. Magenta boxes
highlighted the positions of the 15 (out of 18) contact residues where mutations were reported. Colons indicate reverse mutations to SARS- CoV sequence. Empty box
with dashes (-) represent an insertion of a valine (V) at position 483 of SARS-CoV-2. Signal peptide (SP), N-terminal domain (NTD), Receptor-Binding Domain (RBD),
Receptor-Binding Motif (RBM), Fusion Peptide (FP), Heptad Repeat 1 (HR1), Heptad Repeat 2 (HR2), Transmembrane Domain (TM) and Cytoplasmic Tail (CT) a.a
(amino acid). The RBD sequence analyzed in this diagram corresponds to residues 333-527 of the S protein RBD in the structure PDB ID 6LZG.

3.2. Analysis of the RBD mutations effect on the S protein/ACE2 complex stability

The PDB ID 6LZG model of the complex RBD S protein (Wuhan strain sequence)/Human
ACE2 isoform 1 was selected for this study since it displayed the best resolution (2.5 Å) of the
complexes available in the PDB database. The LigPlot+ analysis of the interactions showed that
there are 18 residues of SARS-CoV-2 interacting with the hACE2 receptor, (Supplementary Table 1),
with ten of them forming polar bonds with the receptor. In addition, these 18 contact residues have
one or more hydrophobic interactions with the hACE2 receptor except for G446 as it forms only one
polar interaction with Q42 of the ACE2 receptor. A total of 12 polar bonds and 65 hydrophobic
interactions were identified at the complex interface as shown in Figure 2. These interactions
involve the 18 contact residues (Supplementary Table 1). Out of these residues, 15 showed multiple
mutations as highlighted in Figure 1.

Figure 2. Representation of the polar and hydrophobic interactions between SARS-CoV-2 spike RBD
of the Wuhan strain (chain B) and hACE2 isoform 1 receptor (chain A) of the complex structure
(PDB ID 6LZG). The dotted green lines show the polar interactions, and the dotted red lines the
hydrophobic interactions.

The complete set of data on the effect of the SARS-CoV-2 RBD 351 mutations on the complex
stability are displayed in Supplementary Material 1. Figure 3 shows the distribution of the Free
energy difference of all analyzed mutations. We observe that, out of the 351 mutations reported in
the GISAID server, 83% of the mutations were classified as "Destabilizing", and 17% were classified
as "Stabilizing" according to their $\Delta\Delta G$ value. Energy difference values ranged from 0.788 Kcal/mol
(T333Q, most stabilizing) to -2.482 Kcal/mol (G502R, most destabilizing). Among the destabilizing
mutations, three mutations were classified as "Highly destabilizing" with $\Delta\Delta G$ values less than -2.0
Kcal/mol (G502R, N501Y and Y505E) (Supplementary Material 1). Furthermore, this analysis

showed that among the mutations that reversed to SARS-CoV RBD sequence, mutation A372T is
predicted to have a relatively good stabilizing effect on the complex with hACE2 isoform 1 as
shown by the value of the ΔΔG (+0.508) Figure 2

Figure 3. Distribution of the free energy ΔΔG differences of the reported 351 nonsynonymous
substitution in the protein sequence of the hACE2-SARS-COV-2 S RBD complex.

SDM generated complexes for the S protein mutants were used to establish the molecular
interaction patterns using LigPlot+. Out of the mutations affecting the 18 interacting residues only
six mutations (K417N, G446A, Y449N, Y453R, Q493R, and T500I), predicted as destabilizing
mutations, cause polar bond differences (Supplementary Table 1). Three out of these residues were
shown to be conserved residues between SARS-CoV-2 and SARS- CoV (Y449, Y453, and T500).
Figure 4 shows the differences in polar interactions between theses mutants and the human ACE2
receptor as compared to the reference complex. Inter-chain polar bonds are shown to be missing in
all the mutated structures at the complex interface level except for Q493R, as a new polar
interaction with E35 of ACE2 receptor was introduced. For the hydrophobic interactions, the most
stabilizing and destabilizing mutations were analyzed (40 mutants) and the results showed that the
interaction patterns did not change for the mutations that occurred out of the contact residues.
Figure 5 shows the hydrophobic interaction changes in comparison to Wuhan SARS-CoV-2 strain
for the Spike/ACE2 complex for mutations that occurred in contact residues. Mutations in contact
residue N501, G502 and Y505 were shown to be the most destabilizing mutations according to ΔΔG
predictions. Ranked from the highest destabilizing to the lowest these mutations are G502R, N501Y,
Y505E, N501T and N501S with the following respective ΔΔGs (-2.482, -2.317, -2.014, -1.689, -1.589
Kcal/mol). The least destabilizing mutant among the contact residues is L455F with a ΔΔG of -0.124
Kcal/mol. Interestingly, all of these mutants have no missing polar interactions. However, they
contain extra hydrophobic interactions compared to the original complex. Polar bonds seem to be
important for maintaining the complex stability between RBD and hACE2 receptor, which explain
the high-energy difference, predicted by mCSM, however gaining extra hydrophobic interactions
highly affect the complex stability (Supplementary Table 1).

Figure 4. Mutations in the SARS-COV-2 RBD sequence causing changes in the pattern of polar interactions with the human ACE2 receptor. Original and mutated residues (K417N, G446A, Y449N, Y453R, Q493R, and T500I) are represented in sticks. Protein chains are shown in cartoon representation (RBD; Blue / hACE2; Green). Polar interactions are shown with yellow dashes.

Figure 5. Representation of the hydrophobic interaction of SARS-CoV-2 spike RBD (chain B) with hACE2 receptor (chain A) of the 6LZG complex. Hydrophobic bonds are represented in Red lines and Salt bridges in dotted green lines.

3.3. Analysis of the effect of ACE2 genetic polymorphism on the S protein/ACE2 complex stability

The analysis of the effect of hACE2 variants on the SARS-CoV2/hACE2 complex stability
showed that out of 231 variants, 203 (88%) are destabilizing (Supplementary Material 2). Out of
these, six variants (D355A, W163R, D355N, G405E, G377E and Y252N) have highly destabilizing
effect with a ΔΔG less than -2.0 Kcal/mol (Supplementary Material 2).
The analysis of the hACE2 residues involved in the interaction of the receptor with the SARS-
CoV-2 Spike's protein, showed that variants D355A, D355N, E35K, F40L, E35D, M82I, S19P and
T27A are destabilizing variants (Supplementary Material 2), with the D355A and D355N, predicted
as being highly destabilizing (-2.775 Kcal/mol and-2.342 Kcal/mol, respectively). Interaction analysis
showed that D355 establishes a network of intramolecular polar interactions with hACE2′ Y41,
G326, N331, L351 and R357 in addition to the polar interaction with the Spike′ T500 Figure 6 A.
Meanwhile, variant D355A Figure 6 B causes the loss of this interaction network, suggesting that
this mutation strongly destabilizes the complex, which is in line with the ΔΔG prediction. The
D355N mutation interaction analysis Figure 6 C shows that N355 maintains only three polar bonds
with hACE2 R357, Y41 and Spike T500.

Figure 6. Genetic variations in the human ACE2 receptor sequence causing changes in the pattern of
polar interactions with the SARS-COV-2 Wuhan strain RBD sequence. Original D355 (A) and
variant D355A (B) and D355N (C) are shown in sticks representation. Protein chains are shown in
cartoon representation (RBD; Blue / hACE2; Green). Polar interactions are represented by yellow
dashes.

The eight hACE2 variants in residues involved in the interaction interface of the receptor with
the SARS-CoV-2 Spike's protein were screened for their effect against all the reported S protein
mutations (Supplementary Material 2). Results showed a significant change in energy prediction
(threshold set at 20% change in ΔΔG) for some variants against Spike mutants compared to the
hACE2 isoform 1, as shown in Figure 7. In fact, D355N was predicted to have more destabilizing
effect with mutants N501T, N501S and G502D with ΔΔG values reduced by 34, 39 and 22%
respectively, compared to hACE2 isoform 1. In addition, the variant is predicted to be less
destabilizing with the mutant G502C with a ΔΔG value increased by 37%. Similarly, variants T27A,
M82I, and S19P show a less destabilizing effect with Spike mutants F456L, F486L and G476S/A,
respectively.

266
Figure 7. **Gradient of binding stability between SARS-CoV-2 mutants and ACE2 variants at the contact**
residues (a) 3D ribbon chart showing the ΔΔG distribution depending on mutations and variants. Each ribbon
represents the values of ΔΔG for a specific variant (bold labels) with the different mutants. For a given line,
most values are equal to Isoform1 variant, which drives the trend (yellow ribbon). The chart shows the spikes
having a distance ≥ 0.2 to Isoform1. The value of 0.2 is selected empirically for visual convenience and as a
threshold to label a spike (combination mutation/ACE2 variant) as important. **(b)** Matrix showing the ΔΔG
difference between Variants and Isoform1 for various contact residues mutations. The difference represents an
algebraic measure ranging from -0.62 to 0.46. The matrix cells are colored using color gradient (d) to illustrate
the intensity and direction of the gap. The matrix lines represent the mutations with at least one non-null
difference. The non-null values are shown inside the cells and can be used to assess the difference with
Isoform1. **(c)** Table showing the difference between variants ΔΔG and Isoform1 ΔΔG for specific mutations in
term of a unidimensional scale. It is obtained by discarding all null cells and by sorting remaining cells in
ascending order. A bold line is used to distinguish important (having a distance ≥ 0.2 to Isoform1). **(d)** A
gradient palette to visually weight a continuous change from -1 (Green) to +1(Red) through 0 (White). The

choice of numerical limits (-1 and +1) is made to facilitate the visual reading of the intensity and sense of the
gap to Isoform1.

Table 1 summarizes the variations of molecular interactions numbers and patterns between the
SARS-CoV-2 RBD mutations occurring in the 18 residues the virus engages to interact with the
ACE2 isoform1 and the eight variants found to be significantly destabilizing when tested against
the Wuhan RBD sequence in comparison to hACE2 isoform1. The number and position of polar
and hydrophobic bonds in addition to the salt bridges characterize each interaction pattern. The
most striking interactions variations concern SARS-CoV-2 RBD mutants N501T and N501S and
ACE2 variant D355N. Interactions with corresponding residues are detailed in Supplementary
Table 2.

291
292
293

294 **Table 1**. Variation of the number of polar and hydrophobic interactions between 24 different SARS-CoV-2
295 spike RBD mutations that occurred in the 18 contact residues with nine hACE2 genetic variants. P: polar, H:
296 hydrophobic. The red and the green show respectively the most and the least divergent interaction from the
297 Wuhan virus RBD sequence and ACE2 isoform1 interaction considering both the ΔΔG and the pattern of
298 molecular interactions.

RBD sequence residues		ACE2 variants																
		Isoform 1		D355A		D355N		E35D		E35K		F40L		M82I		T27A		S19P	
		P	H	P	H	P	H	P	H	P	H	P	H	P	H	P	H	P	H
Ref. sequence	SARS-CoV-2 Wuhan strain	12	65	12	64	13	68	13	65	12	65	12	65	12	65	12	64	11	64
Mutations	K417N	11	64	11	63	12	67	12	64	11	64	11	64	11	64	11	63	10	63
	K417R	12	68	12	67	13	71	13	68	12	68	12	68	12	68	12	67	11	67
	G446A	11	65	11	64	12	68	12	65	11	65	11	65	11	65	11	64	10	64
	Y449N	10	62	10	61	11	65	11	62	10	62	10	62	10	62	10	61	9	61
	Y453F	11	59	11	58	12	62	12	59	11	59	11	59	11	59	11	58	9	58
	L455F	12	70	12	69	13	73	13	70	12	70	12	70	12	70	12	69	11	69
	F456L	12	64	12	63	13	67	13	64	12	64	12	64	12	64	12	62	11	63
	A475V	12	68	12	67	13	71	13	68	12	68	12	68	12	68	12	66	11	66
	G476S	12	64	12	63	13	67	13	64	12	64	12	64	12	64	12	63	11	63
	G476A	12	64	12	63	13	67	13	64	12	64	12	64	12	64	12	63	11	63
	F486L	12	62	12	61	13	65	13	62	12	62	12	62	12	62	12	61	11	61
	Q493R	13	66	13	65	14	69	12+ 1salt	62	12	62	13	66	13	66	13	65	12	65
	Q493L	12	64	12	63	13	67	12	64	12	64	12	64	12	64	12	63	11	63
	G496C	12	71	12	70	13	74	13	71	12	71	12	71	12	71	12	70	11	70
	T500I	11	61	11	58	11	60	12	59	11	59	11	59	11	59	11	58	10	58
	N501T	12	66	12	65	13	69	13	66	12	66	12	66	12	66	12	65	11	65
	N501S	12	65	12	64	13	68	13	65	12	65	12	65	12	65	12	64	11	64
	N501Y	12	72	12	71	13	75	13	72	12	72	12	72	12	72	12	71	11	71
	G502D	12	69	12	68	13	72	13	69	12	69	12	69	12	69	12	68	11	68
	G502C	12	68	12	67	13	71	13	68	12	68	12	68	12	68	12	67	11	67
	G502R	12	70	12	69	13	73	13	70	12	70	12	70	12	70	12	69	11	69
	Y505H	12	61	12	60	13	64	13	61	12	61	12	61	12	61	12	60	11	60
	Y505E	12+ 1salt	66	12+ 1salt	65	13+ 1salt	69	13+ 1salt	66	12+ 1salt	66	12+ 1salt	66	12+ 1salt	66	12+ 1salt	65	11+ 1salt	65

4. Discussion

As the COVID-19 pandemic progresses, it is becoming more and more important to
understand the molecular basis of viral infectivity of the SARS-CoV-2, the etiologic agent of the
disease. This is crucial to evaluate correctly the virus capacity to invade the host cell and exploit its
molecular machinery to replicate and produce infectious viral progeny that may invade other cells
and lead to disease progression (Yan et al., 2020). It is also important to investigate the difference in
susceptibility to the infection considering the differences observed in the disease prevalence
between various regions of the world as well as inter individual variability (Hoffmann et al.,
2020;Hou et al., 2020;Hussain et al., 2020). This necessitates a multidisciplinary approach that would
be the most appropriate to get the best insights into SARS-CoV-2 infectivity. These insights will
likely guide the appropriate response of public health authorities worldwide to combat the
pandemic (Frieden and Lee, 2020;Hellewell et al., 2020). The genetic variability of a specific virus
receptor that affect its infective power was demonstrated for the human immunodeficiency virus
(HIV). Indeed, in humans the CD4 receptor variant (C868T) and the CCR5 receptor CCR5-Δ32
variant confer respectively susceptibility and resistance against certain HIV strains (Marmor et al.,
2006). Other studies on group 2 coronavirus that causes hepatitis in mouse showed that the viral
receptor allelic variants were associated with altered virus binding and hence with host
susceptibility (Ohtsuka and Taguchi, 1997). Similarly, it was reported that rs73635825 (S19P) and
rs143936283 (E329G) alleles of the hACE2 receptor may offer some level of resistance/protection
against the infection because of a low affinity to the virus RBD region (Calcagnile et al.,
2020;Hussain et al., 2020). Nevertheless, most of the accumulated evidences suggesting that genetic
polymorphism of the ACE2 gene may be associated to susceptibility or protection from SARS-CoV-
2 (Chiu et al., 2004;Calcagnile et al., 2020;Lippi et al., 2020) are based on comparative studies of the
molecular interactions between ACE2 variants and a single genetic type of the virus. Hence, not
considering the high number of mutations in the RBD region of the virus. In this work, we have
considered both the SARS-CoV-2 variability introduced by spontaneous mutagenesis and the
natural genetic variation of the ACE2 receptor (Li et al., 2020). Indeed, we have compiled 351 non-
synonymous natural mutations that occurred in the SARS-CoV-2 Wuhan strain S protein RBD
region. This high number of mutations in a relatively short time suggest a genetic instability of the
virus receptor-binding domain knowing that RNA-viruses have generally low mutagenesis rate.
We showed that over eighty percent of these mutations destabilizes the complex with ACE2
isoform 1. This data suggests that the crushing number of SARS-CoV-2 naturally occurring genetic
changes are most likely making the virus less infective. We also found that 24 mutations are reverse
mutations to the SARS-CoV RBD sequence. Fourteen of these reversed mutations showed a
destabilizing effect on the SARS-CoV-2 RBD/hACE2 complex while four of them were stabilizing
mutations. This would partially explain the difference in infectivity of SARS-CoV and SARS-CoV-2
observed from epidemiological and clinical data (EDCDP, 2020).
On another hand, studies of SARS-CoV-2 RBD/ACE2-isoform 1 complexes focusing on the
mutations in the RBD contact residues showed that only six mutants (K417N, G446, Y449N, Y453R,
Q493 and T500I), predicted as destabilizing mutations, cause polar bond changes. Previous studies
reported 17 residues in the SARS-CoV-2 RBD to be involved in the interaction with the ACE2
receptor. Fourteen of these residues are conserved between SARS-CoV and SARS-CoV-2 with eight
residues being contact residues (Lan et al., 2020;Walls et al., 2020). In this study, we found that 18
residues in the RBD region are involved in the interaction with the ACE2 receptor. The interaction
pattern did not change for the mutations that occurred in non-contact residues, which suggest that
these residues act as a structural scaffold that is important for the virus binding to its receptor. Polar
bonds seem to be important for maintaining the complex stability between RBD and hACE2
receptor, which explain the high-energy difference, predicted by mCSM. In addition, out of the 18

described contact residues, ten different mutations occurring in conserved residues between SARS-
CoV-2 and SARS-CoV (Y449N, Y453R, G496C, T500I, G502D, G502C, G502R, Y505H, Y505E, and
Y505W) are predicted to be destabilizing mutations. This is consistent with the fact that conserved
residues are important to maintain the overall correct RBD domain three-dimensional fold needed
for efficient binding to the ACE2 receptor. The loss of polar bonds constitutes an important factor in
protein-protein interaction.

Besides, the analysis of the effect of hACE2 variants on the SARS-CoV-2 Wuhan Strain/ACE2
complex stability showed that out of 231 variants, 88% are predicted to be destabilizing. Eight
ACE2 variants of ACE2 residues involved in the interaction interface of the receptor with the SARS-
CoV-2 Spike's protein are destabilizing. Variants D355A and D355N predicted as highly
destabilizing are potentially protective which is consistent with previously reported experimental
data. This *in silico* study of the molecular bonds showed that D355 establishes a network of polar
interactions with ACE2 residues Y41, G326, N331, L351 and R357 and Spike' T500. The substitution
of the 355 aspartic acid by an alanine (D355A) causes the loss of this interaction network. 292,
mutation D355N shows that Asparagine 355 maintains only three polar bonds with hACE2 R357,
Y41 and Spike T500, which likely makes this variant the most, destabilizing one of the RBD/ACE2
complex of the RBD mutations studied. This result suggest that this variant is potentially more
resistant to the virus attachment. In addition, in our study ACE2 variant S19P more frequent in
Africans and reported to be a protective genetic variant (Calcagnile et al., 2020), showed the lowest
number of interactions with the different spike RBD mutants Table 1. Therefore, if we consider our
findings together with the literature report on the genetic susceptibility to SARS-CoV-2 infection,
we can argue that there is no absolute susceptible or protective ACE2 genetic variant. Indeed, the
same given ACE2 variant might protect from a virus genetic type and be susceptible to another
depending on the molecular interactions engaged between the genetic type of the circulating virus
and an individual's receptor genetic variant. This may explain the striking individual difference
observed globally in clinical presentations with some patients remaining asymptomatic, whilst
others developing the disease (COVID-19) considering that all have the same comparable baseline
risk. Hence, using a computational approach to assess SARS-CoV-2 infectivity through the
evaluation of the binding of the circulating genotype with individual or population-dominant
ACE2 variant could be a rapid and reasonable way to estimate the spreading of the disease. Indeed,
experimental measurement of the S protein-ACE2 affinity is time consuming and cannot be
deployed for a large number of RBD mutants-ACE2 variants combinations. Nevertheless, it should
be noted that accumulation of several mutations in the same virus type have usually cumulative
effects that cannot be measured by the approach we used in this work unless a more refined
RBD/ACE structure with high resolution [less than 2 Angstrom] becomes available. Therefore, the
approach developed in this work can be applied using a platform software such as MutaBind2
(https://lilab.jysw.suda.edu.cn/research/mutabind2/). Furthermore, combining *in silico* information
with clinical and other conventional epidemiological data could be very helpful in stratifying the
risk of infection and for fine-tuning of the mitigation efforts to circumvent disease progress.

Author Contributions: Dana Ashoor and Noureddine Ben Khalaf: *In Silico* analysis, methodology, data
curation, writing and editing (both authors contributed equally to this work). Maryam Marzouq: Mutations'
review, Illustrations figures and tables, revision. Hamdi Al Jarjanazi: primary and secondary data retrieval
formatting and management. Sadok Chlif: Illustrations and mathematical data analysis. M. Dahmani
Fathallah: Project conception, work design, data analysis, writing, editing, and supervision. All authors have
read and agreed to the submitted version of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

Alifano, M., Alifano, P., Forgez, P., and Iannelli, A. (2020). Renin-angiotensin system at the heart of COVID-19 pandemic. *Biochimie* 174, 30-33.

Belouzard, S., Millet, J.K., Licitra, B.N., and Whittaker, G.R. (2012). Mechanisms of coronavirus cell entry mediated by the viral spike protein. *Viruses* 4, 1011-1033.

Calcagnile, M., Forgez, P., Iannelli, A., Bucci, C., Alifano, M., and Alifano, P. (2020). ACE2 polymorphisms and individual susceptibility to SARS-CoV-2 infection: insights from an in silico study. *bioRxiv*, 2020.2004.2023.057042.

Cao, Y., Li, L., Feng, Z., Wan, S., Huang, P., Sun, X., Wen, F., Huang, X., Ning, G., and Wang, W. (2020). Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* 6, 11.

Chan, J.F., Kok, K.H., Zhu, Z., Chu, H., To, K.K., Yuan, S., and Yuen, K.Y. (2020). Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect* 9, 221-236.

Chiu, R.W., Tang, N.L., Hui, D.S., Chung, G.T., Chim, S.S., Chan, K.A., Sung, Y.-M., Chan, L.Y., Tong, Y.-K., and Lee, W.-S.J.C.C. (2004). ACE2 gene polymorphisms do not affect outcome of severe acute respiratory syndrome. 50, 1683-1686.

Edcdp (2020). *COVID-19 situation update worldwide, as of 13 October 2020* [Online]. European Centre for Disease Prevention and Control Available: https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases [Accessed].

Elflein, J. (2020). *Coronavirus (COVID-19) disease pandemic- Statistics & Facts* [Online]. statista. Available: https://www.statista.com/topics/5994/the-coronavirus-disease-covid-19-outbreak/ [Accessed].

Frieden, T.R., and Lee, C.T. (2020). Identifying and Interrupting Superspreading Events-Implications for Control of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* 26, 1059-1066.

Goodsell, D.S., Zardecki, C., Di Costanzo, L., Duarte, J.M., Hudson, B.P., Persikova, I., Segura, J., Shao, C., Voigt, M., Westbrook, J.D., Young, J.Y., and Burley, S.K. (2020). RCSB Protein Data Bank: Enabling biomedical research and drug discovery. *Protein Sci* 29, 52-65.

Hellewell, J., Abbott, S., Gimma, A., Bosse, N.I., Jarvis, C.I., Russell, T.W., Munday, J.D., Kucharski, A.J., Edmunds, W.J., Funk, S., and Eggo, R.M. (2020). Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Glob Health* 8, e488-e496.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A., Muller, M.A., Drosten, C., and Pohlmann, S. (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 181, 271-280 e278.

Hou, Y., Zhao, J., Martin, W., Kallianpur, A., Chung, M.K., Jehi, L., Sharifi, N., Erzurum, S., Eng, C., and Cheng, F. (2020). New insights into genetic susceptibility of COVID-19: an ACE2 and TMPRSS2 polymorphism analysis. *BMC Med* 18, 216.

Hussain, M., Jabeen, N., Raza, F., Shabbir, S., Baig, A.A., Amanullah, A., and Aziz, B.J.J.O.M.V.
(2020). Structural variations in human ACE2 may influence its binding with SARS-CoV-2
spike protein.

Koo, J.R., Cook, A.R., Park, M., Sun, Y., Sun, H., Lim, J.T., Tam, C., and Dickens, B.L. (2020).
Interventions to mitigate early spread of SARS-CoV-2 in Singapore: a modelling study.
Lancet Infect Dis 20, 678-688.

Kowalczuk, S., Broer, A., Tietze, N., Vanslambrouck, J.M., Rasko, J.E., and Broer, S. (2008). A
protein complex in the brush-border membrane explains a Hartnup disorder allele. *FASEB J*
22, 2880-2887.

Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., Zhang, Q., Shi, X., Wang, Q., Zhang, L., and Wang,
X. (2020). Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2
receptor. *Nature* 581, 215-220.

Laskowski, R.A., and Swindells, M.B. (2011). LigPlot+: multiple ligand-protein interaction diagrams
for drug discovery. *J Chem Inf Model* 51, 2778-2786.

Lewnard, J.A., and Lo, N.C. (2020). Scientific and ethical basis for social-distancing interventions
against COVID-19. *Lancet Infect Dis* 20, 631-633.

Li, Q., Cao, Z., and Rahman, P. (2020). Genetic variability of human angiotensin-converting enzyme
2 (hACE2) among various ethnic populations. *Mol Genet Genomic Med* 8, e1344.

Lippi, G., Lavie, C.J., Henry, B.M., and Sanchis-Gomar, F. (2020). Do genetic polymorphisms in
angiotensin converting enzyme 2 (ACE2) gene play a role in coronavirus disease 2019
(COVID-19)? *Clin Chem Lab Med* 58, 1415-1422.

Lopera Maya, E.A., Van Der Graaf, A., Lanting, P., Van Der Geest, M., Fu, J., Swertz, M., Franke, L.,
Wijmenga, C., Deelen, P., Zhernakova, A., and Sanna, S. (2020). Lack of Association
Between Genetic Variants at ACE2 and TMPRSS2 Genes Involved in SARS-CoV-2 Infection
and Human Quantitative Phenotypes. *Front Genet* 11, 613.

Lu, N., Yang, Y., Wang, Y., Liu, Y., Fu, G., Chen, D., Dai, H., Fan, X., Hui, R., and Zheng, Y. (2012).
ACE2 gene polymorphism and essential hypertension: an updated meta-analysis involving
11,051 subjects. *Mol Biol Rep* 39, 6581-6589.

Marmor, M., Hertzmark, K., Thomas, S.M., Halkitis, P.N., and Vogler, M. (2006). Resistance to HIV
infection. *J Urban Health* 83, 5-17.

Nelson-Sathi, S., Umasankar, P.K., Sreekumar, E., Radhakrishnan Nair, R., Joseph, I., Nori, S.R.C.,
Philip, J.S., Prasad, R., Navyasree, K.V., Ramesh, S., Pillai, H., Ghosh, S., Santosh Kumar,
T.R., and Radhakrishna Pillai, M. (2020). Mutational landscape and in silico structure
models of SARS-CoV-2 Spike Receptor Binding Domain reveal key molecular determinants
for virus-host interaction. *bioRxiv*, 2020.2005.2002.071811.

Noordzij, M., Dekker, F.W., Zoccali, C., and Jager, K.J. (2010). Measures of disease frequency:
prevalence and incidence. *Nephron Clin Pract* 115, c17-20.

Ohtsuka, N., and Taguchi, F.J.J.O.V. (1997). Mouse susceptibility to mouse hepatitis virus infection
is linked to viral receptor genotype. 71, 8860-8863.

Othman, H., Bouslama, Z., Brandenburg, J.T., Da Rocha, J., Hamdi, Y., Ghedira, K., Srairi-Abid, N.,
and Hazelhurst, S. (2020). Interaction of the spike protein RBD from SARS-CoV-2 with
ACE2: Similarity with SARS-CoV, hot-spot analysis and effect of the receptor
polymorphism. *Biochem Biophys Res Commun* 527, 702-708.

Pinheiro, D.S., Santos, R.S., Jardim, P., Silva, E.G., Reis, A.a.S., Pedrino, G.R., and Ulhoa, C.J. (2019).
The combination of ACE I/D and ACE2 G8790A polymorphisms revels susceptibility to
hypertension: A genetic association study in Brazilian patients. *PLoS One* 14, e0221248.
Pires, D.E.V., Rodrigues, C.H.M., and Ascher, D.B. (2020). mCSM-membrane: predicting the effects
of mutations on transmembrane proteins. *Nucleic Acids Res* 48, W147-W153.
Roser, M., Ritchie, H., , Ortiz-Ospina, E., and Hasell, J. (2020). *Coronavirus Pandemic (COVID-19)*
[Online]. Published online at OurWorldInData.org. Available:
https://ourworldindata.org/coronavirus [Accessed].
Ruan, S. (2020). Likelihood of survival of coronavirus disease 2019. *Lancet Infect Dis* 20, 630-631.
Schrödinger, LLC. (2010). The PyMOL Molecular Graphics System, Version 2.0
Tipnis, S.R., Hooper, N.M., Hyde, R., Karran, E., Christie, G., and Turner, A.J. (2000). A human
homolog of angiotensin-converting enzyme. Cloning and functional expression as a
captopril-insensitive carboxypeptidase. *J Biol Chem* 275, 33238-33243.
Verity, R., Okell, L.C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., Cuomo-Dannenburg, G.,
Thompson, H., Walker, P.G.T., Fu, H., Dighe, A., Griffin, J.T., Baguelin, M., Bhatia, S.,
Boonyasiri, A., Cori, A., Cucunuba, Z., Fitzjohn, R., Gaythorpe, K., Green, W., Hamlet, A.,
Hinsley, W., Laydon, D., Nedjati-Gilani, G., Riley, S., Van Elsland, S., Volz, E., Wang, H.,
Wang, Y., Xi, X., Donnelly, C.A., Ghani, A.C., and Ferguson, N.M. (2020). Estimates of the
severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* 20, 669-677.
Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., Mcguire, A.T., and Veesler, D. (2020). Structure,
Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* 181, 281-292 e286.
Wang, Q., Zhang, Y., Wu, L., Niu, S., Song, C., Zhang, Z., Lu, G., Qiao, C., Hu, Y., Yuen, K.Y., Zhou,
H., Yan, J., and Qi, J. (2020). Structural and Functional Basis of SARS-CoV-2 Entry by Using
Human ACE2. *Cell* 181, 894-904 e899.
Worth, C.L., Preissner, R., and Blundell, T.L. (2011). SDM--a server for predicting effects of
mutations on protein stability and malfunction. *Nucleic Acids Res* 39, W215-222.
Yan, R., Zhang, Y., Li, Y., Xia, L., Guo, Y., and Zhou, Q. (2020). Structural basis for the recognition of
SARS-CoV-2 by full-length human ACE2. *Science* 367, 1444-1448.
Zhao, Y., Zhao, Z., Wang, Y., Zhou, Y., Ma, Y., and Zuo, W. (2020). Single-Cell RNA Expression
Profiling of ACE2, the Receptor of SARS-CoV-2. *Am J Respir Crit Care Med* 202, 756-759.
520

A Schematic diagram of the of SARS-CoV-2 S spike protein sequence (purple letters) and the mutations in the Receptor
RBD) aligned with SARS-CoV sequence (green letters). The S protein is cleaved into S1 and S2 subunit at the polyba
site. Stabilizing and destabilizing mutations of the RBD domains are represented with blue and red letters respectively. A
mino acid variations between SARS-CoV-2 and SARS-CoV RBDs. Magenta boxes highlighted the positions of the 15 (ou
sidues where mutations were reported. Colons indicate reverse mutations to SARS- CoV sequence. Empty box with da
an insertion of a valine (V) at position 483 of SARS-CoV-2. Signal peptide (SP), N-terminal domain (NTD), Receptor-Binding
ceptor-Binding Motif (RBM), Fusion Peptide (FP), Heptad Repeat 1 (HR1), Heptad Repeat 2 (HR2), Transmembrane Dom
lasmic Tail (CT) a.a (amino acid). The RBD sequence analyzed in this diagram corresponds to residues 333-527 of the S prot
cture PDB ID 6LZG.

Figure 2. Representation of the polar and hydrophobic interactions between SARS-CoV-2 spike RBD of the Wuhan strain (chain B) and hACE2 isoform 1 receptor (chain A) of the complex structure (PDB ID 6LZG). The dotted green lines show the polar interactions, and the dotted red lines the hydrophobic interactions.

Figure 3. Distribution of the free energy DDG differences of the reported 351 nonsynonymous substitution in the protein sequence of the hACE2-SARS-COV-2 S RBD complex.

utations in the SARS-COV-2 RBD sequence
nges in the pattern of polar interactions with
ACE2 receptor. Original and mutated
417N, G446A, Y449N, Y453R, Q493R, and
represented in sticks. Protein chains are
rtoon representation (RBD; Blue / hACE2;
lar interactions are shown with yellow

Figure 5. Representation of the hydrophobic interaction of SARS-CoV-2 spike RBD (chain B) with hACE2 receptor (chain A) of the 6LZG complex. Hydrophobic bonds are represented in Red lines and Salt bridges in dotted green lines.

Figure 6. Genetic variations in the human ACE2 receptor sequence causing changes in the pattern of polar interactions with the SARS-COV-2 Wuhan strain RBD sequence. Original D355 (A) and variant D355A (B) and D355N (C) are shown in sticks representation. Protein chains are shown in cartoon representation (RBD; Blue / hACE2; Green). Polar interactions are represented by yellow dashes.

...radient of binding stability between SARS-CoV-2 mutants and ACE2 ...the contact residues (a) 3D ribbon chart showing the DDG distribution ...n mutations and variants. Each ribbon represents the values of DDG for a ...nt (bold labels) with the different mutants. For a given line, most values are ...orm1 variant, which drives the trend (yellow ribbon). The chart shows the ...g a distance ≥ 0.2 to Isoform1. The value of 0.2 is selected empirically for ...nience and as a threshold to label a spike (combination mutation/ACE2 ...mportant. **(b)** Matrix showing the DDG difference between Variants and ...various contact residues mutations. The difference represents an algebraic ...nging from -0.62 to 0.46. The matrix cells are colored using color gradient ...te the intensity and direction of the gap. The matrix lines represent the ...th at least one non-null difference. The non-null values are shown inside ...can be used to assess the difference with Isoform1. **(c)** Table showing the ...tween variants DDG and Isoform1 DDG for specific mutations in term of a ...al scale. It is obtained by discarding all null cells and by sorting remaining ...nding order. A bold line is used to distinguish important (having a distance ≥ ...m1). **(d)** A gradient palette to visually weight a continuous change from -1 ...·1(Red) through 0 (White). The choice of numerical limits (-1 and +1) is ...itate the visual reading of the intensity and sense of the gap to Isoform1.