Silver nanoparticles induce a *triclosan-like* antibacterial action
mechanism in multi-drug resistant *Klebsiella pneumoniae*

Vikram Pareek[1, 2], Stéphanie Devineau[3], Sathesh K. Sivasankaran[4], Arpit Bhargava[2], Jitendra
Panwar[2]*, Shabarinath Srikumar[5] and Séamus Fanning[1,6]*

[1] UCD-Centre for Food Safety, UCD School of Public Health, Physiotherapy and Sports Science,
University College Dublin, Dublin D04 V1W8, Ireland

[2] Department of Biological Sciences, Birla Institute of Technology & Science, Pilani 333031, India

[3] Université de Paris, BFA, UMR 8251, CNRS, F-75013 Paris, France

[4] Genome Informatics Facility, Iowa State University, 448 Bessy Hall, Ames, IA, 50011, USA

[5] Department of Food, Nutrition and Health, College of Food and Agriculture, UAE University, Al
Ain, UAE

[6] Institute for Global Food Security, Queen's University Belfast, 19 Chlorine Gardens, Belfast BT9
5DL, United Kingdom

* jpanwar@pilani.bits-pilani.ac.in

* sfanning@ucd.ie

Abstract

Infections associated with antimicrobial-resistant bacteria now represent a significant threat to
human health using conventional therapy, necessitating the development of alternate and more
effective antibacterial compounds. Silver nanoparticles (Ag NPs) have been proposed as potential
antimicrobial agents to combat infections. A complete understanding of their antimicrobial activity
is required before these molecules can be used in therapy. Lysozyme coated Ag NPs were
synthesized and characterized by TEM-EDS, XRD, UV-vis, FTIR spectroscopy, zeta potential and
oxidative potential assay. Biochemical assays and deep level transcriptional analysis using RNA
sequencing were used to decipher how Ag NPs exert their antibacterial action against multi-drug
resistant *Klebsiella pneumoniae* MGH78578. RNAseq data revealed that Ag NPs induced a
triclosan-like bactericidal mechanism responsible for the inhibition of the type II fatty acid
biosynthesis. Additionally, released Ag^+ generated oxidative stress both extra- and intracellularly in
K. pneumoniae. The data showed that *triclosan-like* activity and oxidative stress cumulatively
underpinned the antibacterial activity of Ag NPs. This result was confirmed by the analysis of the
bactericidal effect of Ag NPs against the isogenic *K. pneumoniae* MGH78578 *ΔsoxS* mutant, which
exhibits a compromised oxidative stress response compared to the wild type. Silver nanoparticles
induce a *triclosan-like* antibacterial action mechanism in multi-drug resistant *K. pneumoniae*. This
study extends our understanding of anti-*Klebsiella* mechanisms associated with exposure to Ag
NPs. This allowed us to model how bacteria might develop resistance against silver nanoparticles,
should the latter be used in therapy.

Keywords: *Klebsiella pneumoniae*; silver nanoparticles; RNA sequencing; soxS; triclosan.
37

Introduction

Antimicrobial resistance (AMR) is responsible for approximately 700,000 deaths annually across the globe and this number is expected to increase further if new measures are not adopted and antibacterial compounds discovered.(1) The emergence of multi-drug resistance (MDR) in various pathogenic bacterial species represents a serious public health challenge,(2, 3) leading to hospital- and community-acquired infections, which are difficult to treat and control.(4, 5) Since AMR is estimated to overtake cancer as the main cause of death in 50 years, innovative approaches including the development of novel antimicrobial strategies using silver nanoparticles (Ag NPs) are required.(6, 7)

Klebsiella pneumoniae is one of the members of the ESKAPE pathogens (representing *Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumannii, Pseudomonas aeruginosa* and *Enterobacter* spp.).(4) It is a member of Enterobacteriaceae family - Gram-negative, non-motile, and rod-shaped. This bacterium is considered an opportunistic pathogen commonly found in the intestine, mouth, and skin of humans. It is mainly associated with hospital-acquired infections (nosocomial infections) and responsible for respiratory/urinary tract infections, pneumonia, and sepsis.(1, 8, 9) This pathogen can also form biofilms on indwelling medical devices leading to persistent nosocomial infection.(10, 11) About 25 % of nosocomial *K. pneumoniae* were found to be resistant to carbapenem-based compounds.(12, 13) *K. pneumoniae* were also found to be resistant to colistin, a last-line antibiotic.(14, 15) The emergence of AMR in *K. pneumoniae* against critically important classes of antibiotics represents a major threat to conventional clinical therapy.(11, 16) Novel antibacterial strategies are required to overcome this challenge.(17, 18)

Silver and other metals such as copper and zinc have historically been used as potential antibacterial agents.(19, 20) Antibacterial activity of silver can vary depending on its chemical form.(21–23) Metallic forms continuously release small numbers of ions, which makes it a slow-acting agent, whilst the ionic form is more efficient. Although Ag^+ is reported to exhibit better antibacterial activity,(24) direct exposure to mammalian cells has toxic side effects that limit its application in therapy.(25, 26) In contrast, Ag NPs provide a greater surface area, leading to a more controlled release of Ag^+.(27) Hence, this form has potential as an antibacterial compound.(28, 29) Despite this advantage, little is known about the antibacterial action mechanisms of Ag NP based formulations.

We synthesized lysozyme coated Ag NPs (L-Ag NPs) and characterized their antibacterial mechanism against MDR *K. pneumonia* MGH78578 using chemical analysis, biochemical assays and deep-level RNA sequencing. Our data revealed that L-Ag NPs induced a *triclosan-like* antibacterial effect against MDR *K. pneumoniae.* The inhibition of the type II fatty acid biosynthesis along with Ag^+ induced oxidative stress were responsible for the anti-*K. pneumoniae* effect. Our

74 results provide molecular insights into how bacteria might deploy antibacterial strategies to
75 counteract the toxic effects of Ag NPs. This allowed us to model how *K. pneumoniae* might develop
76 resistance against Ag NPs.(30)

Results and discussion

Synthesis and characterization of L-Ag NPs

L-Ag NPs were synthesized by a co-reduction method wherein lysozyme functions both as a
reducing and stabilizing agent using heat reflux action at 120 °C.(31, 32) Lysozyme acts as a
capping material devoid of enzymatic activity. We selected L-Ag NPs based on their bactericidal
activity against MDR *K. pneumoniae* MGH78578. The synthesis of L-Ag NPs was associated with
the appearance of the plasmon peak at 414 nm (Figure 1A).(33) The average diameter of L-Ag
NPs measured by TEM was 5.2 ± 1.2 nm (Figures 1B-C). The X-ray diffraction pattern obtained
corresponds to the face-centered cubic lattice structure of crystalline silver (JCPDS file 04-0783)
(Figure 1D, S1).(34, 35) The surface charge of L-Ag NPs was estimated by measuring their zeta
potential in water and modified LB (mLB) medium (devoid of NaCl) at 37° C. L-Ag NPs were found
to be negatively charged in both conditions with ζ = -38.2 ± 1.6 mV in water and ζ = -24.2 ± 0.9 mV
in mLB medium. The 1,632 cm^{-1} peak in the FTIR spectrum corresponds to the amide I vibration
characteristic of the protein backbone, confirming the presence of lysozyme associated with Ag
NPs (Figure 1E).(36)

L-Ag NPs inhibit the proliferation of *K. pneumoniae* MGH78578

In media with high salt content like chloride and phosphate, Ag NPs may aggregate and free Ag^+
can precipitate, potentially reducing their bactericidal activity.(37–40) Hence, the bactericidal effect
of L-Ag NPs against *K. pneumoniae* MGH78578 was assessed in mLB medium (LB medium devoid
of NaCl).(41, 42) No significant difference between the growth of *K. pneumoniae* in mLB and LB
media was observed (Figure 2A) confirming mLB had no phenotypic effect. Approximately 3×10^7
CFU mL^{-1} log_{10} phase bacterial cells were exposed to L-Ag NPs. The MIC of L-Ag NPs was 21 µg
(Ag) mL^{-1} (Figure 2B). To determine the bactericidal efficiency of L-Ag NPs, *K. pneumoniae*
MGH78578 exposed to L-Ag NPs were spread plated and incubated for 12 h at 37 °C. The MBC
of L-Ag NPs was 45 µg (Ag) mL^{-1} (Figure S2). These results show that L-Ag NPs inhibit the
proliferation of *K. pneumoniae* MGH78578 at low concentrations.

L-Ag NPs generate ROS and limited membrane damage in *K. pneumoniae* MGH78578

Reactive oxygen species (ROS) could be generated outside the bacterial cells by L-Ag NPs and
released Ag^+.(43) Limited dissolution of L-Ag NPs was observed in mLB media for 8 h at 37°C
(Figure 2C). Extracellular ROS production was evaluated by measuring the oxidative potential of
L-Ag NPs in a simplified synthetic respiratory tract lining fluid. The depletion of antioxidants (uric

acid, acetic acid, and reduced glutathione GSH) was also noted when measured by HPLC after 4

h incubation at 37 °C (Figure 3A).(44) We observed a dose-dependent depletion in ascorbic acid

and GSH (up to 100%), showing that, at low NP concentration, L-Ag NPs can generate ROS

extracellularly.

Then, we measured the concentration of silver that enter $K. pneumoniae$ MGH78578 by ICP-OES

following treatment with L-Ag NPs for 5, 30, and 60 min. The uptake of silver, either in the form of

L-Ag NPs or Ag^+, was proportional to the time duration of treatment (Figure S3). To determine

whether ROS were also produced inside the bacterial cells following exposure to L-Ag NPs, we

used the DCFH-DA assay.(45) A dose-dependent increase in the intracellular ROS was observed,

a reflection of the oxidative stress external to the bacterial cell (Figure 3B).

The severe oxidative stress induced by L-Ag NPs could lead to many downstream phenotypes in

bacteria including compromised membrane integrity.(23, 27, 28, 46) TEM analysis of $K.$

$pneumoniae$ did not show any change in the membrane integrity of the exposed bacterial cells

(Figures 3C-D). To verify this observation, we explored the degree of bacterial membrane damage

in the whole population using an MDA assay. A limited though significant concentration-dependent

increase in the level of lipid peroxidation was observed (Figure 3E). Additionally, a concentration-

dependent increase of carbohydrate release was observed following L-Ag NP exposure (Figure

3F). Overall, our results show that L-Ag NPs induced limited bacterial membrane damage, a

phenotype that was not observed by TEM.

Transcriptomic profiling of L-Ag NPs exposed $K. pneumoniae$ MGH78578 using RNA-seq

We used RNA-seq to understand how $K. pneumoniae$ MGH78578 responded to L-Ag NP exposure

following a previously standardized protocol investigating the transcriptome of $K. pneumoniae$

MGH78578 exposed to sub-inhibitory concentrations of a chemosensitizer.(47) Here, we exposed

$K. pneumoniae$ MGH78578 to a sub-inhibitory concentration of L-Ag NPs at MIC_{75} (15.8 µg (Ag)

mL^{-1}) for a period of 5- and 30-min. The 5 min time point was used to identify the transcriptional

signals associated with early exposure, whilst the 30 min time point demonstrated the adaptive

responses of L-Ag NP exposed $K. pneumoniae$ (replication time is 20 min). Approximately 336

million reads were obtained across all 8 libraries with an average of 42 million reads per library

(Table S1) sufficient for downstream transcriptomic analysis.(47, 48) The $VOOM$ function in the

$limma$ package(49) was used to identify differentially regulated genes. We obtained statistically

significant data (p ≤ 0.05) for a total of 3,760 and 3,945 genes following 5 min and 30 min exposure

respectively (Figure 4A) (Dataset WS1). Statistically significant genes with a log_2 fold change

expression of ≥ 1.0 and ≤ -1.0 (exposed $versus$ unexposed cells) were considered to be up- and

down-regulated respectively.(47, 50) For the post-5 min exposure, 1,008 genes were found to be

upregulated and 1,063 genes were downregulated. At 30 min post exposure, 1,205 genes were

found to be upregulated and 1,246 genes were downregulated (Figure 4B). Among all the

upregulated genes at the 5- and 30-min, 705 genes were found to be commonly upregulated,

whereas 303 and 500 genes were uniquely upregulated at 5 min and 30 min respectively. For

downregulated genes, 810 genes were found to be commonly downregulated at both time points,

whereas 253 and 436 genes were specifically downregulated at 5 min and 30 min (Figure 4C).

We selected representative genes related to the bacterial defence system (*soxS*), transcriptional

regulator (*ramA*), outer membrane porin proteins (*ompC*) and virulence (*rfaH*) to validate our

transcriptomic data using qRT-PCR. Gene expression analysis reflected a similar pattern of results

when compared to RNA-seq data (Figure S4).

Bacterial oxidative stress response following exposure to L-Ag NP

Our RNA-seq data shows that the oxidative stress response induced by ROS generation is

choreographed by both *soxS* and *oxyR,* two major transcriptional factors that respond to oxidative

stress in Enterobacteriaceae.(51) The OxyR system functions as a global regulator of peroxide

stress whilst the SoxRS system is involved in the control of superoxide stress.(51, 52) Increased

intracellular ROS levels induce the oxidation of OxyR protein, which in turn activates detoxifying

processes including heme biosynthesis, thiol-disulfide isomerization, among others.(53) In the case

of the SoxRS regulon, oxidative stress causes the oxidization of the SoxR protein, which activates

the *soxS* transcriptional regulator, thereby triggering a bacterial defense mechanism involving

various efflux pumps and redox counter measures.(51) In L-Ag NP exposed *K. pneumoniae*

MGH78578, *soxS* was highly upregulated (300- and 88-fold in 5 and 30 min) and in contrast *oxyR*

was moderately upregulated (approximately 5-fold in both time points). This suggested that the

oxidative stress regulon was active throughout the 30 min of L-Ag NP exposure time, particularly

for the *soxS* regulon.

To confirm this, we compared our RNA-seq dataset to the *K. pneumoniae* MGH78578 oxidative

soxS regulon generated earlier.(54) We found that of the 254 genes belonging to the oxidative *soxS*

regulon, 129 (51%) had similar expression patterns in both datasets, confirming that the *soxS*

regulon is activated in L-Ag NP exposed *K. pneumoniae* (Dataset WS2). A typical example of *soxS*

induction can be visualized in the expression of *fpr* [ferredoxin-NADP$^+$ reductase], which was

induced at 22- and 9-fold at 5 and 30 min. In response to *soxS* activation, a redox neutralization

process is triggered related to the overexpression of *fpr* by 4.5-fold at 5 min, that then reduced to

3.2-fold at 30 min post-treatment.(55)

Activation of efflux pumps following exposure to L-Ag NP

Bacteria activate various efflux pumps to expel the toxic Ag$^+$.(56, 57) Classic efflux pump-encoding

genes such as *acrAB-tolC* (58, 59) and *marRAB* were found to be highly upregulated following

treatment with L-Ag NPs, a feature that was sustained throughout the testing period of 30 min,
possibly driven by the positive regulation *via* SoxS.(51, 54) Another efflux pump encoded by the
silCFBA operon conferring a silver resistance phenotype was also upregulated.(57, 60)
Transcription of the *silCFBA* operon and *silP* are controlled by the *silRS*, which encodes a two-
component system wherein SilR acts as a response regulator and SilS acts as a histidine
kinase.(61, 62) SilP is a P-type ATPase efflux pump, which facilitates the passage of Ag$^+$ from
cytoplasm to the periplasm. SilF acts as a chaperone and transfers Ag$^+$ from periplasm to the
SilCBA complex, a three-protein dependent cation/proton antiporter system. Another protein from
silver resistance system is SilE, present downstream to the *silRS* (Figure 5).(61, 62) Upregulation
of *silC/B/E/R/S/P* and KPN_pKPN3p05946 genes were noted in our RNA-seq data showing the
activation of efflux pumps, encoding a possible silver resistance in response to L-Ag NP exposure.
193

The type II fatty acid biosynthesis genes were down-regulated in L-Ag NP exposed *K.*
***pneumoniae* MGH78578**

Since *K. pneumoniae* MGH78578 expressed an MDR phenotype,(13) L-Ag NP exposure
dependent activation of *acrAB-tolC, marRAB,* and *sil* genes should, in principle, confer silver
resistance in *K. pneumoniae*. Additionally, the strong oxidative stress response should enable the
bacterium to counter the oxidative stress induced by L-Ag NPs. Both of these mechanisms should
render *K. pneumoniae* MGH78578 resistant against silver. However, *K. pneumoniae* MGH78578
was found to be susceptible to L-Ag NPs. We hypothesized that the bactericidal activity of L-Ag
NPs could be due to a hitherto unknown mechanism, rather than the most commonly reported
oxidative stress based bactericidal activity.(27) To investigate, we compared our RNA-seq data to
the transcriptional response of *E. coli* exposed to 37 antimicrobial compounds.(63) This global
transcriptional profile of *E. coli* exposed to 37 antibiotics generated a list of 447 genes whose
signature expression pattern characterized the mechanism of action associated with each
antibiotic, leading the authors to hypothesize that the antibacterial action mechanism of any
unknown/uncharacterized compounds can be deduced by comparing the transcriptional profile of
a compound of interest with that of these 447 *E. coli* MG1665 K12 genes. We compared our
exposed *K. pneumoniae* MGH78578 transcriptional data set with each individual dataset obtained
from *E. coli* global transcriptional profile. Our comparative analysis showed that the highest number
of similarly expressed gene pairs were obtained from the <u>triclosan exposure</u> dataset, suggesting
that exposure to L-Ag NPs induced a *triclosan-like* exposure response in *K. pneumoniae*
MGH78578. Some 76% of the gene pairs (73/96) at 5 min L-Ag NP exposure (Dataset WS3 and
WS4) and 74% (87/117) at 30 min exposure (Dataset WS5 and WS6) had similar expression
patterns compared with the *E. coli* triclosan dataset.

Triclosan is a broad-spectrum antimicrobial compound that acts by inhibiting FabI (a NADH-
dependent enoylacyl carrier), a protein belonging to the type II fatty acid biosynthesis, part of a
well-conserved pathway that is essential for bacterial survival. Inhibition of fatty acid biosynthesis
compromises the bacterial cell membrane. During exposure to L-Ag NPs, some *K. pneumoniae*
MGH78578 genes associated with type II fatty acid biosynthesis including *fabA/H/D/G/F/B* were
significantly downregulated particularly during the adaptive response at 30 min. The
downregulation of the different *fab* genes signals a compromised type II fatty acid biosynthesis
underpinning the *triclosan-like* L-Ag NP bactericidal activity. Importantly, *fadL* and *fadD* were
downregulated. FadL is a porin that transports extracellular fatty acids across the outer membrane
to the inner membrane where they are activated by acyl CoA synthetase FadD. Downregulation of
both *fadL* and *fadD* shows that extracellular fatty acids are not transported leading to subsequent
suppression of type II fatty acid biosynthesis pathway. We, however, did not observe any differential
expression in the *fabI* gene, possibly because the triclosan affects FabI post-translationally.
Recently, the TraDIS-Xpress approach involving a transposon mutant library and massively parallel
sequencing of transposon chromosome junctions was used to identify *E. coli* genes that respond
to triclosan exposure.(64) In comparison with these data, we identified genes that were both
enriched in the *E. coli* TraDIS-Xpress dataset and that were found to be differentially expressed in
our *K. pneumoniae* MGH78578 dataset. Common genes like *purL, purH, waeL, wzx* were
downregulated at either one or both time points while *metB* was upregulated. Similarly, *K.*
pneumoniae MGH78578 genes like *trkA, pcnB, infB,* and *ubiB/F* were downregulated at least in
one or both time points, while *lonH* was upregulated (Dataset WS1). TraDIS-Xpress selected *rbs*
in their triclosan exposure screen. In *E. coli,* RbsABC forms the ABC-type high-affinity D-ribose
transporter, while RbsD/K phosphorylates D-ribose to D-ribose 5-phosphate.(65) Though we did
not observe a statistically significant differential expression for *rbsB,* down-regulation was observed
for *rbsD* and *rbsC,* showing similarly compromised D-ribose uptake. These common observations
across different datasets gave confidence to our earlier observation that L-Ag NP exposure elicits
a *triclosan-like* bactericidal effect in *K. pneumoniae* inhibiting type II fatty acid biosynthesis.
244

The L-Ag NP based antibacterial action is cumulative of oxidative stress and fatty acid
biosynthesis inhibition
We investigated whether the L-Ag NP based antibacterial effect was primarily due to type II fatty
acid biosynthesis inhibition or as a cumulative effect of both oxidative stress and fatty acid
biosynthesis inhibition. We exposed the isogenic *K. pneumoniae MGH78578 ΔsoxS* mutant, which
has a compromised oxidative stress response mechanism,(54) to L-Ag NPs. The MIC of L-Ag NPs
reduced to 15 µg mL^{-1} from 21 µg mL^{-1} showing that oxidative stress did contribute to a significant

antibacterial effect. The antibacterial effect of L-Ag NPs is, therefore, a cumulative effect of both
type II fatty acid biosynthesis inhibition and oxidative stress responses.
254

Concluding observations

L-Ag NPs were very efficient in killing MDR *K. pneumoniae* MGH78578, triggering oxidative stress
and a *triclosan-like* mechanism to exert their anti-*Klebsiella* effect. With toxicity studies in
appropriate cell models, L-Ag NPs may be a future candidate as a novel antimicrobial agent. We
hypothesized that our RNA-seq data could provide clues as to how *K. pneumoniae* might develop
resistance against L-Ag NPs. Genes including *rbsD/C, trkA, pcnB* and *infB* were downregulated
during the adaptive response at 30 min. Transposon insertional mutants in *E. coli* genes *trkA, pcnB*
and *infB* exhibit better survival in the presence of triclosan.[64] Further, transposon insertion in
rbsB, inhibiting the periplasmic ribose-binding domain of the RbsABC ribose importer, gave a
selective survival advantage in the presence of triclosan. We hypothesize that by downregulating
these genes during the adaptive response, *K. pneumoniae* MGH78578 could elicit strategies to
combat the effects of exposure whilst resistance against L-Ag NPs. Exposure to sub-inhibitory
concentrations of antimicrobial compounds is one of the main drivers in the evolution of AMR. By
continuous exposure to sub-inhibitory concentrations of silver and the subsequent downregulation
of these genes, *K. pneumoniae* might offer a transient resistance to L-Ag NPs. This hysteresis
effect might be the prelude to developing a fully expressed antimicrobial resistance mechanism
against silver.
272

Materials and Methods

Bacteria and media

Klebsiella pneumoniae MGH78578 (ATCC®700721) is a clinical isolate chosen for its MDR
phenotype[47, 66] and the availability of its whole genome sequence (NC_009648.1). Bacteria
were grown in Luria Bertani (LB) and modified LB (mLB) media constituted without NaCl.[39, 67]
Bacteria were recovered from long-term storage (at -80°C in glycerol stocks) in mLB medium
(casein enzyme hydrolysate 10 g L^{-1} and yeast extract 5 g L^{-1}, pH 7.2 ± 0.2) for 12 h at 37 °C with
shaking (150 rpm). For the preparation of inoculum, an overnight grown bacterial culture was
inoculated into freshly prepared mLB medium and grown until the mid-log phase ($OD_{600 nm}$ 0.5 to
0.6). An overnight grown bacterial culture (approximately 10^7 CFU mL 1) was inoculated separately
in freshly prepared LB and mLB broth medium. Bacterial growth was determined by measuring the
optical density (OD) at 600 nm at 2 h time intervals. Biological experiments were carried out in both
technical and biological duplicates.

Synthesis of L-Ag NPs

L-Ag NPs were synthesized following a protocol adapted from Ashraf et al (see SI).(31) To
determine the size, shape, crystallinity and surface capping, L-Ag NPs were characterized by
Transmission Electron Microscopy (TEM), Energy Dispersive Spectroscopy (EDS), X-ray
Diffraction (XRD), UV-vis spectroscopy, Fourier Transform Infrared spectroscopy (FTIR). The zeta
potential of L-Ag NPs was measured with a Zetasizer Nano ZS (Malvern Instruments). The acellular
oxidative potential of L-Ag NPs was assessed by measuring the depletion in antioxidants (uric acid,
ascorbic acid, reduced glutathione) by HPLC following incubation of L-Ag NPs in a simplified
synthetic respiratory tract lining fluid for 4 h at 37 °C.(68) The dissolution kinetics of L-Ag NPs in
mLB was measured by inductively coupled plasma optical emission spectrometry (ICP-OES) (Avio
200, PerkinElmer). All measurements were performed in duplicate. The detailed protocols are
described in the supplementary information (SI) file. Freshly prepared L-Ag NPs were used for all
the experiments.

Antibacterial susceptibility determinations

The minimum inhibitory concentration (MIC) was determined using the broth microdilution assay
following CLSI guidelines. Bacterial cells were exposed to L-Ag NPs (ranging from 0.5 to 64 µg
(Ag) mL^{-1}) and incubated at 37 °C in the dark for 24 h. Bacterial growth was determined by
measuring the $OD_{600 nm}$. In addition, the free lysozyme was tested in parallel for any antibacterial
activity. Bacterial cells exposed to L-Ag NPs were spread plated and incubated for 12 h at 37 °C to
determine the minimum bactericidal concentration (MBC). Media without L-Ag NPs and bacterial
cells were used as positive- and negative-controls respectively.(67) All experiments were done in
biological duplicates and the results are represented as mean ± standard deviation.

Mode of action studies-measurement of reactive oxygen species (ROS)

The generation of ROS inside the bacterial cell following exposure to L-Ag NPs was measured
using a 2,7-dichlorodihydrofluorescein diacetate (DCFH-DA) assay.(69) DCFH-DA was added to
the bacterial cell suspension at a final concentration of 10 µM and incubated for 1 h at 37 °C in the
dark. Free dye was then separated from the DCFH-DA loaded bacterial cells by centrifugation at
8,000 rpm for 15 min followed by washing with PBS. Bacterial cells were exposed to different (sub)-
MIC concentrations of L-Ag NPs (i.e. MIC_{25}, MIC_{50}, MIC_{75}, and MIC_{100}) for 30 min at 37 °C in fresh
mLB medium with shaking (150 rpm). The fluorescence intensity of dichlorodihydrofluorescein
(DCF) was detected using a Perkin Elmer VICTOR X Multilabel Plate Reader (USA) at an excitation
and emission wavelength of 485 and 535 nm respectively. The experiments were performed in
duplicates and the results expressed as mean ± standard deviation.

The effect of L-Ag NPs on the bacterial cell envelope was examined by TEM. To prepare sample
for TEM analysis, bacterial cells exposed to MIC_{75} L-Ag NPs were centrifuged at 8,000 rpm for 10
min. Subsequently, the supernatant was discarded, and the bacterial pellet was washed twice with
PBS followed by fixation in 2.5% v/v electron microscopy grade glutaraldehyde in 0.05 M sodium

cacodylate buffer pH 7.2 for 2 h at 4 °C. TEM samples were prepared by drop-casting the bacterial cell suspension on to a carbon-coated copper grid that was later imaged on a Hitachi H-7650 TEM instrument (Hitachi High-Technologies Corporation) at an acceleration voltage of 100 kV. Bacterial cells that were not exposed to L-Ag NPs were used as the control. A minimum number of 50 bacterial cells were analyzed on different TEM images in each condition.

The membrane damage was analyzed using the MDA and anthrone assays. Bacterial cells were treated with different (sub)-MIC concentrations of L-Ag NPs, i.e. MIC_{25}, MIC_{50}, MIC_{75}, and MIC_{100}. The determination of MDA concentration was done as described previously.[70] The anthrone assay was done as described previously.[67] The experiments were performed in biological duplicates and the results expressed as mean ± standard deviation.

The intracellular concentration of Ag was measured by ICP-OES following exposure to MIC_{75} L-Ag NPs for 5, 30 and 60 min at 37 °C. A freshly grown bacterial culture (approximately 10^7 CFU mL^{-1}) was exposed to MIC_{75} L-Ag NPs followed by incubation for 5, 30 and 60 min at 37 °C in mLB. After incubation, bacterial cells were pelleted by centrifugation at 8,000 rpm for 10 min at 4 °C then dried and digested in a 1 mL mixture of H_2O_2:HNO_3 (50:50) for 2 h. After acid digestion, the final volume was made up to 10 mL and filtered using 0.22 µm syringe filter. The silver concentration of the sample was measured by ICP-OES (Avio 200, PerkinElmer, USA).[38] Untreated bacterial cells were taken as a control for the respective time points. The experiments were performed in biological duplicates and the results expressed as mean ± standard deviation. Detailed protocols are described in SI.

RNA sequencing

A freshly grown culture of *K. pneumoniae* MGH78578 from mid-log_{10} phase (approximately 10^7 CFU mL^{-1}) was exposed to MIC_{75} L-Ag NPs for 5 and 30 min at 37 °C in mLB broth. Bacterial cells not exposed to L-Ag NPs were selected as a control for each time point. RNA was isolated using RNAeasy extraction kit (Qiagen) and treated with Turbo DNase kit (Ambion's). RNA integrity was assessed using a Bioanalyzer 2100 RNA 6000 nanochip (Agilent Technologies). RNA library sequencing was performed at the Centre for Genomic Research, University of Liverpool, UK. Ribosomal RNA was removed with a Ribo-Zero rRNA removal kit (Illumina). Libraries were prepared with NEBNext Directional RNA Library Prep Kit (BioLabs). Pooled libraries (Table S1) were loaded on cBot (Illumina) and cluster generations was performed. Single-sequencing using 150 bp read length was performed on lane of the HiSeq 4000 sequencer (Illumina). Raw sequencing data was processed as described in SI. The RNAseq data produced from the present work were deposited to the NCBI-GEO database and are available under the accession number GSE151953.

Comparative gene expression analysis with E. coli MGH1655 exposed to individual antibiotics

A comparison of our transcriptional data with the global transcriptome profiling of *E. coli* MGH1665

K12 strain exposed to 37 antibiotics was carried out. Only those genes that gave statistically

significant differential expression patterns were selected from both datasets. Our gene expression

dataset was compared to each of the 37 *E. coli* transcriptional profiles and genes that had similar

expression patterns (both up-and down-regulation) were counted.

Validation of RNA-seq data by quantitative RT-PCR analysis. RNA isolated from the samples

were converted to cDNA by using high-capacity RNA to cDNA kit (Thermo fisher, Ireland). Primers

based on the selected genes of interest were designed with 6-FAM/ZEN/IBFQ double-quenched

probes and synthesized commercially by Integrated DNA Technologies (IDT, Belgium) (Table S2).

The cDNA was used as a template and analysis was done by the addition of PrimeTime Gene

Expression Master Mix. qRT-PCR was performed in an Eppendorf Mastercycler realplex ep

gradient S (Eppendorf, UK). This analysis was carried out in two biological replicates each along

with three technical replicates. The fold-change in the expression of the genes of interest was

determined by the method of Livek and Schmittgen(71), i.e. $2^{-\Delta\Delta Ct}$ method using *rho* as a

housekeeping gene.

Statistical analysis. The results for the biochemical assay were analysed using unpaired Student

t-test as appropriate for the dataset. The qRT-PCR measurements data were statistically analysed

using Prism software (v. 8.0 GraphPad Software) following the two-way analysis of variance.

Boneferroni method was used to analyse the multiple comparisons. The symbol 'ns' used in the

graphs corresponds to statistically non-significant with $p > 0.05$. The asterisk symbols in the graphs

correspond to * $p \leq 0.05$, ** $p \leq 0.01$, and *** $p \leq 0.001$. All the data points represent the mean of

two independent measurements. The uncertainties were represented as standard deviations. In

RNAseq results, NSE represents non-significant expression.

Acknowledgments

Authors are thankful to BITS Pilani, University College Dublin, and Université de Paris for providing lab facilities. VP acknowledges CSIR-SRF [09/719(0090)/2018-EMR-I] and EMBO Short Term Fellowship (ESTF 7654) for the financial support. This project was supported, in part, by the Ulysses PHC Scheme (CNRS). We acknowledge Justine Renault, Linh-Chi Bui, and the Bioprofiler Facility (BFA) for the HPLC analysis.

Author Contributions

SD, SS, SF, JP designed the study. VP, AB, SD characterized the silver NPs. VP and SS carried out the RNA-seq experiments. SKS generated the bioinformatics based gene expression dataset.

SS, SD, VP, SF carried out the detailed RNA-seq dataset analysis that led to the triclosan observation. All authors read and approved the manuscript.

ORCID

Vikram Pareek 0000-0003-3980-1238
Stéphanie Devineau 0000-0002-1133-5223
Sathesh K. Sivasankaran 0000-0003-3037-6001
Arpit Bhargava 0000-0001-6450-6551
Jitendra Panwar 0000-0002-0750-9745
Shabarinath Srikumar 0000-0003-3775-2831
Séamus Fanning 0000-0002-1922-8836

Competing interests

The authors declare no competing interests. None to declare.

References

1. Organization WH. 2019. No time to wait: securing the future from drug-resistant infections. Report to the Secretary-General of the United Nations. World Heal Organ Geneva.
2. Logan LK, Weinstein RA. 2017. The epidemiology of Carbapenem-resistant enterobacteriaceae: The impact and evolution of a global menace. J Infect Dis 215:S28–S36.
3. Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, Colomb-Cotinat M, Kretzschmar ME, Devleesschauwer B, Cecchini M, Ouakrim DA, Oliveira TC, Struelens MJ, Suetens C, Monnet DL, Strauss R, Mertens K, Struyf T, Catry B, Latour K, Ivanov IN, Dobreva EG, Tambic Andraševic A, Soprek S, Budimir A, Paphitou N, Žemlicková H, Schytte Olsen S, Wolff Sönksen U, Märtin P, Ivanova M, Lyytikäinen O, Jalava J, Coignard B, Eckmanns T, Abu Sin M, Haller S, Daikos GL, Gikas A, Tsiodras S, Kontopidou F, Tóth Á, Hajdu Á, Guólaugsson Ó, Kristinsson KG, Murchan S, Burns K, Pezzotti P, Gagliotti C, Dumpis U, Liuimiene A, Perrin M, Borg MA, de Greeff SC, Monen JC, Koek MB, Elstrøm P, Zabicka D, Deptula A, Hryniewicz W, Caniça M, Nogueira PJ, Fernandes PA, Manageiro V, Popescu GA, Serban RI, Schréterová E, Litvová S, Štefkovicová M, Kolman J, Klavs I, Korošec A, Aracil B, Asensio A, Pérez-Vázquez M, Billström H, Larsson S, Reilly JS, Johnson A, Hopkins S. 2019. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. Lancet Infect Dis 19:56–66.
4. Pendleton JN, Gorman SP, Gilmore BF. 2013. Clinical relevance of the ESKAPE pathogens. Expert Rev Anti Infect Ther 11:297–308.
5. Klemm EJ, Wong VK, Dougan G. 2018. Emergence of dominant multidrug-resistant bacterial clades: Lessons from history and whole-genome sequencing. Proc Natl Acad Sci 115:12872–12877.
6. Gupta A, Mumtaz S, Li CH, Hussain I, Rotello VM. 2019. Combatting antibiotic-resistant bacteria using nanomaterials. Chem Soc Rev 48:415–427.
7. Huh AJ, Kwon YJ. 2011. "Nanoantibiotics": A new paradigm for treating infectious diseases using nanomaterials in the antibiotics resistant era. J Control Release 156:128–145.
8. Podschun R, Ullmann U. 1998. Klebsiella spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev 11:589–603.
9. Lee GC, Burgess DS. 2012. Treatment of Klebsiella pneumoniae carbapenemase (KPC)

infections: a review of published case series and case reports. Ann Clin Microbiol Antimicrob 11:32.

10. Jagnow J, Clegg S. 2003. Klebsiella pneumoniae MrkD-mediated biofilm formation on extracellular matrix-and collagen-coated surfaces. Microbiology 149:2397–2405.

11. Li B, Zhao Y, Liu C, Chen Z, Zhou D. 2014. Molecular pathogenesis of Klebsiella pneumoniae. Future Microbiol 9:1071–1081.

12. Han JH, Goldstein EJC, Wise J, Bilker WB, Tolomeo P, Lautenbach E. 2016. Epidemiology of carbapenem-resistant Klebsiella pneumoniae in a network of long-term acute care hospitals. Clin Infect Dis 64:839–844.

13. Anes J, Hurley D, Martins M, Fanning S. 2017. Exploring the genome and phenotype of multi-drug resistant Klebsiella Pneumoniae Of clinical origin. Front Microbiol 8.

14. Antoniadou A, Kontopidou F, Poulakou G, Koratzanis E, Galani I, Papadomichelakis E, Kopterides P, Souli M, Armaganidis A, Giamarellou H. 2007. Colistin-resistant isolates of Klebsiella pneumoniae emerging in intensive care unit patients: first report of a multiclonal cluster. J Antimicrob Chemother 59:786–790.

15. Neuner EA, Yeh J-Y, Hall GS, Sekeres J, Endimiani A, Bonomo RA, Shrestha NK, Fraser TG, van Duin D. 2011. Treatment and outcomes in carbapenem-resistant Klebsiella pneumoniae bloodstream infections. Diagn Microbiol Infect Dis 69:357–362.

16. Hu Y, Anes J, Devineau S, Fanning S. 2020. Klebsiella pneumoniae: Prevalence, Reservoirs, Antimicrobial Resistance, Pathogenicity, and Infection: A Hitherto Unrecognized Zoonotic Bacterium. Foodborne Pathog Dis https://doi.org/10.1089/fpd.2020.2847.

17. Adamo R, Margarit I. 2018. Fighting Antibiotic-Resistant Klebsiella pneumoniae with "Sweet" Immune Targets. MBio 9:e00874-18.

18. Mulani MS, Kamble EE, Kumkar SN, Tawre MS, Pardesi KR. 2019. Emerging strategies to combat ESKAPE pathogens in the era of antimicrobial resistance: A review. Front Microbiol 10.

19. Alexander JW. 2009. History of the medical use of silver. Surg Infect (Larchmt) 10:289–292.

20. Rai MK, Deshmukh SD, Ingle AP, Gade AK. 2012. Silver nanoparticles: The powerful nanoweapon against multidrug-resistant bacteria. J Appl Microbiol 112:841–852.

21. Mijnendonckx K, Leys N, Mahillon J, Silver S, Van Houdt R. 2013. Antimicrobial silver: uses, toxicity and potential for resistance. Biometals 26:609–621.

22. Li R, Chen J, Cesario TC, Wang X, Yuan JS, Rentzepis PM. 2016. Synergistic reaction of silver nitrate, silver nanoparticles, and methylene blue against bacteria. Proc Natl Acad Sci U S A 113:13612–13617.

23. Zheng K, Setyawati MI, Leong DT, Xie J. 2018. Antimicrobial silver nanomaterials. Coord Chem Rev 357:1–17.

24. Randall CP, Oyama LB, Bostock JM, Chopra I, O'Neill AJ. 2012. The silver cation (Ag+): antistaphylococcal activity, mode of action and resistance studies. J Antimicrob Chemother 68:131–138.

25. Chernousova S, Epple M. 2013. Silver as antibacterial agent: ion, nanoparticle, and metal. Angew Chemie Int Ed 52:1636–1653.

26. Vimbela G V., Ngo SM, Fraze C, Yang L, Stout DA. 2017. Antibacterial properties and toxicity from metallic nanomaterials. Int J Nanomedicine 12:3941–3965.

27. Durán N, Durán M, de Jesus MB, Seabra AB, Fávaro WJ, Nakazato G. 2016. Silver nanoparticles: A new view on mechanistic aspects on antimicrobial activity. Nanomedicine Nanotechnology, Biol Med 12:789–799.

28. Kang J, Dietz MJ, Hughes K, Xing M, Li B. 2019. Silver nanoparticles present high intracellular and extracellular killing against Staphylococcus aureus. J Antimicrob Chemother 74:1578–1585.

29. Roe D, Karandikar B, Bonn-Savage N, Gibbins B, Roullet J baptiste. 2008. Antimicrobial surface functionalization of plastic catheters by silver nanoparticles. J Antimicrob Chemother 61:869–876.

30. Chopra I. 2007. The increasing use of silver-based products as antimicrobial agents: A useful development or a cause for concern? J Antimicrob Chemother 59:587–590.
31. Ashraf S, Chatha MA, Ejaz W, Janjua HA, Hussain I. 2014. Lysozyme-coated silver nanoparticles for differentiating bacterial strains on the basis of antibacterial activity. Nanoscale Res Lett 9:1–10.
32. Eby DM, Schaeublin NM, Farrington KE, Hussain SM, Johnson GR. 2009. Lysozyme catalyzes the formation of antimicrobial silver nanoparticles. ACS Nano 3:984–994.
33. Singh DK, Kumar J, Sharma VK, Verma SK, Singh A, Kumari P, Kharwar RN. 2018. Mycosynthesis of bactericidal silver and polymorphic gold nanoparticles: physicochemical variation effects and mechanism. Nanomedicine 13:191–207.
34. Jain N, Bhargava A, Majumdar S, Tarafdar JC, Panwar J. 2011. Extracellular biosynthesis and characterization of silver nanoparticles using Aspergillus flavus NJP08: a mechanism perspective. Nanoscale 3:635–641.
35. Pareek V, Bhargava A, Panwar J. 2020. Biomimetic approach for multifarious synthesis of nanoparticles using metal tolerant fungi: A mechanistic perspective. Mater Sci Eng B 262:114771.
36. Baker MJ, Trevisan J, Bassan P, Bhargava R, Butler HJ, Dorling KM, Fielden PR, Fogarty SW, Fullwood NJ, Heys KA. 2014. Using Fourier transform IR spectroscopy to analyze biological materials. Nat Protoc 9:1771.
37. Lok C-N, Ho C-M, Chen R, He Q-Y, Yu W-Y, Sun H, Tam PK-H, Chiu J-F, Che C-M. 2007. Silver nanoparticles: partial oxidation and antibacterial activities. JBIC J Biol Inorg Chem 12:527–534.
38. McQuillan JS, Groenaga Infante H, Stokes E, Shaw AM. 2012. Silver nanoparticle enhanced silver ion stress response in Escherichia coli K12. Nanotoxicology 6:857–866.
39. Chambers BA, Afrooz ARMN, Bae S, Aich N, Katz L, Saleh NB, Kirisits MJ. 2013. Effects of chloride and ionic strength on physical morphology, dissolution, and bacterial toxicity of silver nanoparticles. Environ Sci Technol 48:761–769.
40. Pareek V, Gupta R, Panwar J. 2018. Do physico-chemical properties of silver nanoparticles decide their interaction with biological media and bactericidal action? A review. Mater Sci Eng C 90:739–749.
41. Pelletier DA, Suresh AK, Holton GA, McKeown CK, Wang W, Gu B, Mortensen NP, Allison DP, Joy DC, Allison MR. 2010. Effects of engineered cerium oxide nanoparticles on bacterial growth and viability. Appl Environ Microbiol 76:7981–7989.
42. Chatterjee S, Bandyopadhyay A, Sarkar K. 2011. Effect of iron oxide and gold nanoparticles on bacterial growth leading towards biological application. J Nanobiotechnology 9:34.
43. Le Ouay B, Stellacci F. 2015. Antibacterial activity of silver nanoparticles: A surface science insight. Nano Today 10:339–354.
44. Crobeddu B, Aragao-Santiago L, Bui LC, Boland S, Baeza Squiban A. 2017. Oxidative potential of particulate matter 2.5 as predictive indicator of cellular stress. Environ Pollut 230:125–133.
45. Aranda A, Sequedo L, Tolosa L, Quintas G, Burello E, Castell J V, Gombau L. 2013. Dichloro-dihydro-fluorescein diacetate (DCFH-DA) assay: a quantitative method for oxidative stress assessment of nanoparticle-treated cells. Toxicol Vitr 27:954–963.
46. Zou L, Wang J, Gao Y, Ren X, Rottenberg ME, Lu J, Holmgren A. 2018. Synergistic antibacterial activity of silver with antibiotics correlating with the upregulation of the ROS production. Sci Rep 8:1–11.
47. Anes J, Sivasankaran SK, Muthappa DM, Fanning S, Srikumar S. 2019. Exposure to sub-inhibitory concentrations of the chemosensitizer 1-(1-naphthylmethyl)-piperazine creates membrane destabilization in multi-drug resistant Klebsiella pneumoniae. Front Microbiol 10:1–14.
48. Haas BJ, Chin M, Nusbaum C, Birren BW, Livny J. 2012. How deep is deep enough for RNA-Seq profiling of bacterial transcriptomes? BMC Genomics 13:734.
49. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res 43:e47–e47.

50. Dash S, Sarashetti PM, Rajashekar B, Chowdhury R, Mukherjee S. 2018. TGF-β2-induced EMT is dampened by inhibition of autophagy and TNF-α treatment. Oncotarget 9:6433.

51. Seo SW, Kim D, Szubin R, Palsson BO. 2015. Genome-wide Reconstruction of OxyR and SoxRS Transcriptional Regulatory Networks under Oxidative Stress in Escherichia coli K-12 MG1655. Cell Rep 12:1289–1299.

52. Zheng M, Åslund F, Storz G. 1998. Activation of the OxyR transcription factor by reversible disulfide bond formation. Science (80-) 279:1718–1722.

53. Dubbs JM, Mongkolsuk S. 2012. Peroxide-sensing transcriptional regulators in bacteria. J Bacteriol 194:5495–5503.

54. Anes J, Dever K, Eshwar A, Nguyen S, Cao Y, Sivasankaran SK, Sakalauskaite S, Lehner A, Devineau S, Daugelavicius R. 2020. Analysis of the oxidative stress regulon identifies soxS as a genetic target for resistance reversal in multi-drug resistant Klebsiella pneumoniae. bioRxiv.

55. Krapp AR, Rodriguez RE, Poli HO, Paladini DH, Palatnik JF, Carrillo N. 2002. The flavoenzyme ferredoxin (flavodoxin)-NADP (H) reductase modulates NADP (H) homeostasis during the soxRS response of Escherichia coli. J Bacteriol 184:1474–1480.

56. Gurbanov R, S. Ozek N, Tunçer S, Severcan F, Gozen AG. 2018. Aspects of silver tolerance in bacteria: infrared spectral changes and epigenetic clues. J Biophotonics 11:e201700252.

57. Randall CP, Gupta A, Jackson N, Busse D, O'Neill AJ. 2014. Silver resistance in Gram-negative bacteria: A dissection of endogenous and exogenous mechanisms. J Antimicrob Chemother 70:1037–1046.

58. Hirakata Y, Kondo A, Hoshino K, Yano H, Arai K, Hirotani A, Kunishima H, Yamamoto N, Hatta M, Kitagawa M. 2009. Efflux pump inhibitors reduce the invasiveness of Pseudomonas aeruginosa. Int J Antimicrob Agents 34:343–346.

59. Baugh S, Phillips CR, Ekanayaka AS, Piddock LJ V, Webber MA. 2013. Inhibition of multidrug efflux as a strategy to prevent biofilm formation. J Antimicrob Chemother 69:673–681.

60. Elkrewi E, Randall CP, Ooi N, Cottell JL, O'Neill AJ. 2017. Cryptic silver resistance is prevalent and readily activated in certain Gram-negative pathogens. J Antimicrob Chemother 72:3043–3046.

61. Starodub ME, Trevors JT. 1989. Silver resistance in Escherichia coli R1. J Med Microbiol 29:101–110.

62. Massani MB, Klumpp J, Widmer M, Speck C, Nisple M, Lehmann R, Schuppler M. 2018. Chromosomal Sil system contributes to silver resistance in E. coli ATCC 8739. BioMetals 31:1101–1114.

63. O'Rourke A, Beyhan S, Choi Y, Morales P, Chan AP, Espinoza JL, Dupont CL, Meyer KJ, Spoering A, Lewis K. 2020. Mechanism-of-action classification of antibiotics by global transcriptome profiling. Antimicrob Agents Chemother 64.

64. Yasir M, Turner AK, Bastkowski S, Baker D, Page AJ, Telatin A, Phan M-D, Monahan L, Savva GM, Darling A. 2020. TraDIS-Xpress: a high-resolution whole-genome assay identifies novel mechanisms of triclosan action and resistance. Genome Res 30:239–249.

65. Shimada T, Kori A, Ishihama A. 2013. Involvement of the ribose operon repressor RbsR in regulation of purine nucleotide synthesis in Escherichia coli. FEMS Microbiol Lett 344:159–165.

66. Ogawa W, Li D-W, Yu P, Begum A, Mizushima T, Kuroda T, Tsuchiya T. 2005. Multidrug resistance in Klebsiella pneumoniae MGH78578 and cloning of genes responsible for the resistance. Biol Pharm Bull 28:1505–1508.

67. Bhargava A, Pareek V, Roy Choudhury S, Panwar J, Karmakar S. 2018. Superior Bactericidal Efficacy of Fucose-Functionalized Silver Nanoparticles against *Pseudomonas aeruginosa* PAO1 and Prevention of Its Colonization on Urinary Catheters. ACS Appl

Mater Interfaces 10:29325–29337.

68.   Boland S, Baeza-squiban A. 2015. Acellular antioxidant depletion assay. SOP WP3 nanOxiMet.

69.   Wang H, Joseph JA. 1999. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. Free Radic Biol Med 27:612–616.

70.   Buege JA, Aust SD. 1978. [30] Microsomal lipid peroxidation, p. 302–310. *In* Methods in enzymology. Elsevier.

71.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2− ΔΔCT method Methods 25: 402–408. Find this Artic online.

Figures and Tables

Figure 1. Characterization of L-Ag NPs. (A) UV-visible spectrum (inset shows a picture of the L-Ag NP suspension after synthesis). (B) TEM image. (C) Distribution of L-Ag NP diameter determined from TEM analysis. (D) XRD diffraction spectrum. (E) FTIR spectrum.

Figure 2. (A) Bacterial growth curve analysis of *K. pneumoniae* MGH78578 in LB and mLB media. (B) Determination of *K. pneumoniae* L-Ag NP MIC in mLB medium. (C) Dissolution kinetics of L-Ag NPs in mLB medium at 37°C.

Figure 3. (A) Acellular oxidative potential of L-Ag NPs measured by the depletion of antioxidants, uric acid (UA), ascorbic acid (AA), and reduced glutathione (GSH), in simplified synthetic lung lining fluid at 37°C. (B) Analysis of the intracellular ROS generation in *K. pneumoniae* MGH78578 following exposure to L-Ag NPs by the DCFH-DA assay. (C,D) Representative TEM images of untreated *K. pneumoniae* (control) (C) and *K. pneumoniae* exposed to MIC75 of L-Ag NPs (D). (E,F) Membrane damage analysis of *K. pneumoniae* following exposure to L-Ag NPs. Analysis by MDA assay (E) and anthrone assay (F).

Figure 4. Overview of the RNAseq analysis of *K. pneumoniae* MGH78578 following treatment with MIC$_{75}$ L-Ag NPs. (A) Statistically significant and insignificant genes recorded at 5 and 30 min post-treatment. (B) Statistically significant upregulated, downregulated and neutral genes identified at 5 and 30 min post-treatment. (C) Comparison of the upregulated, downregulated, and neutral genes identified at 5 and 30 min post-treatment.

Figure 5. A model depicting the bactericidal action of L-Ag NPs on *K. pneumoniae* MGH78578. L-Ag NPs and silver ions (Ag⁺) enter the bacterial cell *via* outer membrane porins. Increasing cytoplasmic Ag⁺ concentration elicits the production of reactive oxygen species (ROS) including superoxide radicals, which then activate the *soxS*-mediated oxidative stress response. The SilE efflux system functions to eliminate Ag⁺ from the cytoplasm. RNA-seq data identified 7 genes (denoted by the green arrowheads) associated with type II fatty acid biosynthesis that are down-regulated leading to reduced phospholipid biosynthesis. This reduction in phospholipid turnover will affect the membrane stability leading to cell lysis underpinning the bactericidal activity. Grey arrows represent genes that are activated. White arrows represent genes that are either not identified in the *K. pneumoniae* MGH78578 genome or not differentially regulated when the bacteria are exposed to L-Ag NPs.