Parasitic success and venom composition evolve upon specialization

of parasitoid wasps to different host species

Fanny Cavigliasso[‡], Hugo Mathé-Hubert[‡], Jean-Luc Gatti, Dominique Colinet*, Marylène

Poirié*

Université Côte d'Azur, INRA, CNRS, ISA, France

[‡]Present addresses:
F.C.: Department of Ecology and Evolution, University of Lausanne, Switzerland. H. M-H.: TIMC-
IMAG - Techniques de l'Ingénierie Médicale et de la Complexité - Informatique, Mathématiques et
Applications, Grenoble - UMR 5525, France.
11

12 * these authors should be considered joint senior author

14 Corresponding author: Marylène Poirié, Evolution and Specificity of Multitrophic Interactions (ESIM),
15 Institut Sophia Agrobiotech, 400 route des Chappes, BP167, 06903 Sophia Antipolis, France. Tel:
16 +33492386409; E-mail: marylene.poirie@univ-cotedazur.fr
17

Abstract

Female endoparasitoid wasps usually inject venom into hosts to suppress their immune response

and ensure offspring development. However, the parasitoid's ability to evolve towards increased

success on a given host simultaneously with the evolution of the composition of its venom has

never been demonstrated. Here, we designed an experimental evolution to address this question.

We crossed two parasitoid lines of *Leptopilina boulardi* differing both in parasitic success on

different *Drosophila* hosts and venom composition. F_2 descendants were reared on three

different *Drosophila* species for nine generations. We tested for evolution of parasitic success

over the generations and for the capacity of parasitoids selected on a given host to succeed on

another host. We also tested whether the venom composition - based on a statistic analysis of

the variation in intensity of the venom protein bands on SDS-PAGE 1D - evolved in response

to different host species. Results showed a specialization of the parasitoids on their selection

host and a rapid and differential evolution of the venom composition according to the host.

Overall, data suggest a high potential for parasitoids to adapt to a new host, which may have

important consequences in the field as well in the context of biological control.

Keywords: parasitoid wasps, *Drosophila*, parasitic success, experimental evolution, venom

evolution, specialization

1. Introduction

Endoparasitoid wasps are insects whose larvae develop inside the host, generally leading to its death [1]. High selection pressures are therefore exerted to maximize the parasitic success that has been shown to evolve rapidly according to host resistance [2–4] and host species [5,6]. The ability to succeed in multiple host species and adapt to a new host is important for the abundance and survival of parasitoids in the event of environmental changes, such as the local extinction of a host. The host range of parasitoids is generally not limited to a single species, even for those considered specialists. For example, *Leptopilina boulardi,* considered a specialist of *D. melanogaster* and *D. simulans,* can develop in other species of the *melanogaster* subgroup of Drosophilidae, including *D. yakuba* [7,8].

Parasitoids have evolved different strategies to circumvent the host's immune defense, which usually consists of the formation of a multicellular melanized capsule around the parasitoid egg, together with the activation of the phenol-oxidase cascade which leads to the production of melanin and the release of toxic radicals [9–11]. The most common strategy is the injection of venom with the egg, which suppresses the encapsulation process [12–14]. To our knowledge, only two studies have analyzed the ability of parasitoid venom to evolve according to the host. Both *Lysiphlebus fabarum* parasitic success and expression of venom genes rapidly and differentially evolved depending on the strains of a defensive symbiont hosted by the host *Aphis fabae* [4]. More recently, a rapid and differential evolution of the venom protein composition of *L. boulardi* was described according to the susceptible/resistant phenotype of two *D. melanogaster* host strains [15]. However, the relationship to parasitic success was not analyzed. Since these studies involved a single host species, our goal here was to investigate whether the parasitic success and the venom protein composition could evolve according to different host species. We also sought to determine whether the venom of a

parasitoid wasp successful in several host species contains broad-spectrum factors or a

combination of specialized factors, each specific to a given host.

We used *L. boulardi* as a model because of its intraspecific variability in both venom

composition and parasitic success on different hosts. The ISm line always succeeds on *D.*

melanogaster but is consistently encapsulated by *D. yakuba* whereas the ISy line can succeed

on both *Drosophila* species but only on certain genotypes [16]. The venom of these lines differs

widely, mainly due to quantitative differences in the venomous proteins. As an example, a

RhoGAP domain-containing protein named LbGAP is in a significantly higher amount in the

venom of ISm than in that of ISy [17,18]. LbGAP would be necessary for ISm parasitic success

on an ISy-resistant *D. melanogaster* strain through the induction of morphological changes in

the lamellocytes, host immune cells devoted to the encapsulation [17,19–22]. LbSPN, a serine

protease inhibitor of the serpin superfamily [18] illustrates the qualitative variation of venom

proteins between ISm and ISy. LbSPNy, which inhibits the activation of the phenoloxidase

cascade in *D. yakuba* [23] and LbSPNm are one of the most abundant proteins in the ISy and

ISm venom, respectively. Although they are encoded by allelic forms of the same gene, they

differ in molecular weight and the sequence of the active site, suggesting they have different

targets and/or function [18].

Here, we report data from an experimental evolution designed to evaluate the

evolvability of *L. boulardi* (i) parasitic success and (ii) the venom protein composition

according to different *Drosophila* host strains and species. The experiment was designed to

characterize the venom allowing the parasitoid to develop on different hosts and to inform on

whether such venom contains rather broad-spectrum factors or a combination of factors specific to

each host. For this, we crossed *L. boulardi* ISm and ISy individuals and reared their descendants

for nine generations independently on the hosts species and strains differing for their resistance

to these wasps. We then analyzed the parasitic success on the selection host and the venom

composition in three generations: after the first (F_3) and the last (F_{11}) generation under selection and after an intermediate generation (F_7). We also tested for a specialization of parasitoids by measuring the parasitic success of individuals from the latest generation of selection on all the different hosts. Overall, parasitoids have shown specialization to their selection host as well as a rapid and differential evolution of the venom composition. Changes in the intensity of the venom protein bands were mainly observed in response to selection on a single host, although some changes occurred under selection by several hosts. This suggests that most of the venom factors of this wasp are host-specific while a few may have a wider spectrum.

2. Material and methods

2.1. Biological material

L. boulardi ISm (Gif stock number 431) and ISy (Gif stock number 486) isofemale lines originate from populations collected in Nasrallah (Tunisia) and Brazzaville (Congo), respectively [24]. Both lines were reared on their susceptible *Drosophila melanogaster* maintenance strain (Nasrallah from Tunisia, Gif stock number 1333, here called S_{Nasr}) at 25°C. Emerged adults were kept at 20°C on agar medium and fed with honey.

Five *Drosophila* host strains from three different species differing in their resistant/susceptible phenotype to ISm and ISy were used (Figure 1A). The *D. melanogaster* R strain (Gif stock number 1088), originating from isofemale lines obtained from a population of Brazzaville, Congo [25] through subsequent genetic approaches [26,27], is resistant to ISy and susceptible to ISm [28,29]. The *D. simulans* Japan strain, from a population collected in Japan, is susceptible to ISm and ISy. *D. yakuba* 1907 (Gif stock number 1907) and 307 (Gif stock number 307.14), originate from Tanzania and from the São Tomé island, respectively. *D. yakuba* 1907 is resistant to both parasitoid lines while 307 is resistant to ISm and susceptible to ISy.

     2.2. Experimental evolution protocol

Three wasp mass rearings were created from ($♀$ISm x $♂$ISy and $♀$ISy x $♂$ISm) crosses

of 12 virgin females and six males on the *D. melanogaster* S_{Nasr} host (Figure 1B). A total of 15

replicates were then created from these mass rearings using six F_1 hybrid females and three F_1

males from each direction of crossing, still on *D. melanogaster* S_{Nasr}. Finally, F_2 individuals

from each of the 15 replicates were used to produce five groups that were then reared separately

on the five different hosts until the F_{11} generation (F_6 only for *D. yakuba* 1907, therefore

excluded for further analyses; Figure 1B). For each of the 75 populations (15 replicates x five

hosts), up to 10 females and five males (mean: 9.51 ♀, 4.31 ♂) were randomly chosen to

produce the F_3 generation, and up to 20 females and 10 males (mean: 18.48 ♀, 8.99 ♂) to

produce F_4 to F_{11} generations.

The parasitic success and venom composition were analyzed for females of three

generations: the first generation of selection on different hosts (F_3), an intermediate generation

(F_7), and the last generation (F_{11}). Since the venom composition and the parasitoid behavior

could vary between females depending of the number of eggs previously laid, only females

never allowed to oviposit were used for the analyses.

2.3. Analysis of the outcome of the Drosophila – L. boulardi *interaction*

Parasitic tests and dissection:

Twenty second-instar larvae of the investigated host species were deposited in small

medium-containing dishes with one parasitoid female. The parasitoid was removed after 4h at

25°C and the dishes kept at 25°C for 48h until dissection of the *Drosophila* larvae . They were

then categorized as (i) non-parasitized, (ii) mono-parasitized and (iii) multi-parasitized (see

Figure S1 for proportions of each). Only mono-parasitized host larvae were considered for the

analysis to avoid unpredictable effects of the presence of several parasitoid larvae in a single

host.

Three possible outcomes were then recorded: i) free parasitoid larva alone, ii) free

parasitoid larva together with an open capsule and iii) complete closed capsule (Figure 2A).

Among these, the percentage of outcomes for i) and iii) were generally used to assess the

parasitoid's immune suppression ability and the host encapsulation capacity [5,25,30]. In this

work, we added the outcome ii) to evaluate the parasitoid's ability to escape the encapsulation

process initiated by the host after recognition of the parasitoid egg. Since the escaped parasitoid

larva was alive, we considered this outcome to be a parasitic success, similar to a free parasitoid.

Statistical analysis:

We used a generalized linear mixed model (GLMM) to analyze each parameter with the

continuous generation or the selection host as a fixed effect, the replicates as a random effect

and a binomial error distribution. The GLMMs were fitted with the *glmer* function in "lme4" R

package [31], except for unbalanced data for which they were fitted with the *bglmer* function

implemented in the "blme" R package [32]. As for all the binomial GLMMs performed,

overdispersion was tested with the *overdisp_fun* function [33] and accounted for when

necessary ($p < 0.05$) by adding a random factor corresponding to the observations number [34].

GLMMs were followed by post-hoc Tukey tests with the "multcomp" R package [35] to

compare hosts two by two.

2.4. Venom separation, western blot and data acquisition

L. boulardi venom reservoirs were dissected individually in 15 μl of insect Ringer

supplemented with protease inhibitors cocktail (PI; Roche), mixed with an equivalent volume

of Laemmli reducing buffer and heated (95°C, 10 min). The individual protein samples were

then split in two, half being used for the global analysis, half for the specific analysis (see the

two next paragraphs). Protein separation was done by 1D SDS-PAGE electrophoresis using

commercial gels to ensure reproducibility (8–16% Criterion™ TGX™ Precast Midi Protein

Gel, Bio-Rad). A sample of venom of ISm and ISy lines, equivalent to half a female reservoir

(from a pool of venom from 60 individuals collected in 1.8 ml) was also loaded on each gel and

served as a reference.

The global analysis was performed as described in [36] and used in [15]. Briefly, the

gels were silver-stained and their digital pictures analyzed with the Phoretix 1D software (now

CLIQS, TotalLab, UK) to extract the intensity profile of each lane (individual sample). We then

used R functions to obtain for each lane the intensities of a set of "reference bands" of known

molecular weight. These intensities were normalized and estimated with the following

combination of parameters: without background – peak height – quantile normalization (more

details in [36]). The normalized intensities of the reference bands are the variables

characterising the venom composition.

The specific analysis was performed using Western blots and antibodies targeting

previously characterized *L. boulardi* venom proteins as described in [15]. The LbSPN, LbGAP

and LbGAP2 polyclonal antibodies are described in [19,20,37,38]. The secondary was a goat

anti-rabbit IgG horseradish peroxidase conjugate (1:10,000; Sigma). Western blots were

revealed with a luminescent substrate (Luminata Crescendo; Millipore) and digitalized. Data

were then recorded as follows for each individual:

i) LbSPN: the genotype at the *lbspn* locus was determined by the presence/absence of

the LbSPNy (54 kDa) and LbSPNm (45 kDa) bands.

(ii) LbGAP: a strong signal was observed in the venom of ISm or (ISm × ISy) F_1 only.

The presence/absence of a signal at the expected size was then used to distinguish *lbgap*

homozygotes and *lbgap*/*lbgap$_y$* heterozygotes from *lbgap$_y$* homozygotes.

(iii) LbGAP2: the variation in quantity was more continuous. The normalized quantity

of LbGAP2 was estimated as the ratio between the signal intensity on the Western blot without

background and the sum of the intensities of the reference bands (used as a proxy for the amount

of protein in the venom samples) in the corresponding lane on silver stained gels.

2.5. Statistical analysis for the global analysis of the venom

<u>Evolution of the venom composition.</u> We analyzed the evolution of the venom

composition using PERMANOVAs (permutational MANOVAs; "vegan" R package; function

adonis2; [39,40]). PERMANOVAs mesure the association between the multidimensional

variation of some explained variables (venom) and some explanatory variables, but at the

difference of MANOVAs, they don't compare the correlation structure among different groups.

For this analysis, we measured the multidimensional variation through the Euclidean distance,

and we tested the significance of explanatory variables' marginal effects with 5,000

permutations. To determine whether the venom composition has evolved on each host

separately, PERMANOVAs were fitted with the generation (F_3, F_7 or F_{11}; continuous variable)

and the population (same as replicates in this situation) as explanatory variables. For *D. yakuba*

307, we fitted an additional PERMANOVA without F_3 individuals due to their small number.

We then determined whether the venom composition has evolved differently depending on the

host by fitting a PERMANOVA with the following explanatory variables: the generation, the

host, the interaction of both and the 44 experimental populations (sum of replicates maintained

on each host until the end of the experiment) to account for the effect of genetic drift. Then,

PERMANOVAs were performed to compare the venom composition two by two between hosts

to determine whether a venom composition evolved on one host differed from that evolved on

another.

To characterize the evolutions detected by the PERMANOVAs, we used linear

discriminant analyses (LDA). Specifically, we performed LDA to characterize (i) the evolution

of the venom composition on each host separately (LDAs performed with the three groups of

individuals i.e. from the three analyzed generations) and (ii) the differential evolution of venom

composition depending on the host (LDAs performed with the 12 groups of individuals (three

generations × four hosts) and each two by two comparison of venom composition between

hosts. We used the "ade4" R package [41] with the individual venom compositions as

continuous variables and the generation or the "host × generation" interaction as a factor. Since

LDA does not account for the variation between replicates, they were centered before the

analysis using the *wca* function (within class analysis) in the "ade4" R package. With this

additional step, all replicates had the same mean for each variable and thus the variation among

generations and hosts cannot result from the variation from replicates andunbalanced data. The

LDA axes were labeled with the biological meaning determined from the position of the 12

groups of individuals (generations × hosts) on these axes. To describe the venom evolution

trend, we plotted for each host in each LDA an arrow representing the linear regression fitted

to coordinates of the three centroids corresponding to the F_3, F_7 and F_{11} generations and

weighted by the number of individuals per generation and host. The only exception was the

LDA on *D. yakuba* 307 for which the linear regression was calculated from centroid points of

the F_7 and F_{11} generations only.

<u>Evolution of venom protein bands</u>. Non-parametric Spearman rank correlation tests

were performed for each LDA done for each host separately to identify the protein bands that

correlated with the regression describing the trend of venom evolution. P-values were

Bonferroni corrected using the *p.adjust* R function.

As some protein bands were probably indirectly selected due to their correlation with

other bands, a combination of clustering and partial correlation analyses was used to identify

bands undergoing direct selection. We first performed an UPGMA clustering analysis ("hclust"

R package) with $1 - |\rho|$ as the metric distance, where ρ is the spearman correlation. Then, we

used a threshold correlation of 0.4 to construct band clusters for which false detection of certain

bands as "correlated" could have occurred. For each cluster with at least two protein bands

correlated to the regression (represented by an arrow), we performed partial correlations with

the *pcor* function in the "ppcor" R package to determine if the residual variation in the intensity

of each band, independent from the other bands of the cluster, was still correlated to the

regression describing the evolution. P-values were also Bonferroni corrected using the *p.adjust*

R function.

<u>Comparison of the venom composition evolved on each host to that of the parental</u>

<u>lines.</u> To make this comparison, we computed the averages of venom composition evolved on

each host separately for each replicate. Each of the 44 populations from the first and last

generation of selection (F_3 and F_{11}) was scaled between 0 and 1 depending on its distance to the

ISm or ISy venom using the formula $D_{ISy}/(D_{ISm} + D_{ISy})$, D_{ISy} and D_{ISm} being the Euclidean

distances between the average venom composition of the population and the venom

composition of ISm and ISy. Then, we tested whether the venom composition of parasitoids

reared on each host evolved toward a parental line by comparing the scaled distances between

F_3 and F_{11} generations with paired Wilcoxon rank tests. Finally, to determine whether a protein

band selected on a given host corresponded to an ISm or ISy band, we assigned it a value from

0 to 1 in relation to a higher intensity in ISy or ISm, respectively. This value is the intensity of

the band in the venom of ISm divided by the sum of its intensities in the venom of ISm and ISy.

2.6. Statistical analysis for the specific analysis of venom

The three variables describing LbSPN, LbGAP and LbGAP2 are of different type

(categorical for LbSPN with two different alleles, presence/absence for LbGAP, continuous for

LbGAP2 with the relative intensity). We therefore used different approaches to analyse them.

LbSPN is a codominant marker with two alleles (*lbspnm* and *lbspny*) encoding proteins

of distinct molecular weight. To determine if the frequency of these alleles had evolved on each

host, we fitted one generalized linear mixed model (GLMM) per host with generation as a fixed

continuous effect and replicates as a random effect with a binomial distribution using the

"lme4" R package [31].

LbGAP is a dominant marker with two alleles (*lbgap* and *lbgap_y*) determining the

presence in the venom of the LbGAP protein in detectable quantity (*lbgap* homozygotes and

lbgap/*lbgapy* heterozygotes) or not (*lbgapy* homozygotes). The evolution of the

presence/absence of the LbGAP/LbGAPy proteins independently on each host was tested using

the same procedure as for LbSPN.

For LbGAP2 (continuous variation), we used linear mixed models (LMM) with

generation as a fixed continuous effect and replicates as a random effect to determine if the

quantity of LbGAP2 had evolved, separately for each host. The models were fitted with the

"nlme" R package [42] on the box cox-transformed (lambda = 0.25) standardized intensity of

LbGAP2 to normalize the residues.

3. Results

3.1. Experimental evolution protocol

The variability on which the selection took place was generated by reciprocal crosses

between the ISm and ISy parasitoid lines. F_1 hybrids were then used to produce the 15 F_2

replicates from which five groups were formed and reared independently on *D. melanogaster*

S_{Nasr}, *D. melanogaster* R, *D. yakuba* 307, *D. yakuba* 1907 and *D. simulans* (Figure 1B). Of

these 75 starting F_2 experimental wasp populations, 44 were successfully maintained until the

F_{11} generation. Among these, none of the *D. yakuba* 1907 replicates survived after the F_6,

preventing further analyses for this host. 12 and 15 replicates survived on *D. melanogaster* S_{Nasr}

and *D. simulans*, respectively (Table S1), and 13 on *D. melanogaster* R, whereas only four

could be maintained on *D. yakuba* 307 until the F_{11}, the other replicates mainly becoming

extinct at the first generation of selection (F_3) (Table S1).

3.2. Evolution of the interaction outcome according to the selection host

We tested the evolution of the parasitoids capacity to bypass the encapsulation response

of their selection host after five and nine generations of selection (i.e. F_7 and F_{11}), using two

females per experimental population, except for *D. yakuba* 307 for which four females were

tested since fewer replicates were available. Three parameters were analyzed: (i) the parasitic

success, i.e. among mono-parasitized host the proportion of hosts containing a free parasitoid

larva alone or together with an open capsule, (ii) the capacity of the parasitoid to inhibit the

capsule formation, i.e. among mono-parasitized host the proportion of *Drosophila* larvae

containing a free parasitoid larva alone and (iii) the capacity of the parasitoid to escape from

the capsule, i.e. among mono-parasitized hosts showing an encapsulation response, the

proportion of hosts containing a free parasitoid larva together with an open capsule (Figure 2A).

The parasitic success of parasitoids reared on *D. melanogaster* S_{Nasr} or *D. simulans* on

this same host remained close to 100% throughout the experimental evolution (Figure 2B, Table

S2, GLMM, $p > 0.30$ for both hosts). In contrast, it increased with the generation on *D.*

melanogaster R (from about 80% at F_3 to 90% at F_{11}) and *D. yakuba* 307 (from about 20% at

F_3 to 65% at F_{11}) (Figure 2B, Table S2, GLMM, $p = 0.002$ for *D. melanogaster* R and $p <$

0.001for *D. yakuba* 307). For *D. melanogaster* R this increase seemed to result solely from the

increased capacity to escape from the capsule (Figure 2B, Table S2, GLMM, $p = 0.001$) since

no significant change was observed for its ability to inhibit encapsulation (Figure 2B, Table S2,

GLMM, p = 0.33). For *D. yakuba* 307, the success increase seemed to result mainly from a

higher capacity to escape from the capsule (Figure 2B, Table S2, GLMM, p <0.001) but also,

to a lesser extent, to inhibit encapsulation at F_{11} (Figure 2B, Table S2, GLMM, p = 0.003).

Finally, a much lower parasitic success was observed at F_3 for *D. yakuba* 307 (about 20%) than

for the other hosts (about 80% to 100% depending on the host) (Figure 2B).

3.3. Specialization of parasitoids to their selection host

To determine whether the change in parasitoid ability to bypass host encapsulation was

specific to the selection host, we compared the success of F_{11} parasitoids on their own selection

host *vs.* the three other hosts. The parasitic success on *D. melanogaster* S_{Nasr} and *D. simulans*

was close to 100% regardless of the selection host (Figure 2C, Table S3, GLMM, Tukey post

hoc-test, p > 0.70 for both hosts). In contrast, parasitoids reared on *D. melanogaster* R were

more successful on this same host than those reared on *D. yakuba* 307 (Figure 2C, Table S3,

GLMM, Tukey post hoc-test, p <0.001). This can be explained by a higher capacity of

parasitoids selected on *D. melanogaster* R to both inhibit encapsulation and escape from a

capsule compared to those selected on *D. yakuba* 307 (Figure 2C, Table S3, GLMM, Tukey

post hoc-test, p = 0.015 for encapsulation inhibition and p < 0.001 for escape capacity). Finally,

parasitoids selected on *D. melanogaster* R had a higher capacity to escape from a capsule of

this same host compared to parasitoids selected on *D. melanogaster* S_{Nasr} or *D. simulans*, despite

a similar parasitic success (Figure 2C, Table S3, GLMM, Tukey post hoc-test, p < 0.001for the

escape capacity between *D. melanogaster* R and the two other hosts; p = 0.09 and p = 0.30 and

for the parasitic success between *D. melanogaster* R vs *D. melanogaster* S_{Nasr} or vs *D. simulans*,

respectively).

Parasitoids reared on *D. yakuba* 307 had a higher parasitic success on this same host

than parasitoids reared on all the other hosts (Figure 2C, Table S3, GLMM, Tukey post hoc-

test, $p < 0.005$ for all comparisons involving *D. yakuba* 307). The parasitoids capacity to inhibit

encapsulation by *D. yakuba* 307 was very low, with no significant difference between the

selection hosts (Figure 2C, Table S3, GLMM, Tukey post hoc-test, $p > 0.60$). Therefore, the

difference in parasitic success between selection hosts was most probably explained by an

increase capacity to escape from the capsule (Figure 2C, Table S3, GLMM, Tukey post hoc-

test, $p < 0.05$ for each pairwise comparison except for *D. melanogaster* R and *D. simulans* for

which $p = 0.76$).

3.4. Differential evolution of venom composition according to the host

The venom analysis was performed on 50 individuals per host and generation distributed

over all replicates, except for *D. melanogaster* S_{Nasr} at F_3 (47 ♀), R at F_{11} (49 ♀) and *D. yakuba*

307 (10, 39 and 40 ♀ at F_3, F_7 and F_{11}, respectively). In total, the venom protein content was

analysed for 535 females and 36 reference protein bands were identified whose intensities

represent the variables describing the venom composition (Figure S2).

The PERMANOVAs performed for each host separately revealed a strong generation

effect for *D. melanogaster* S_{Nasr} and R, and *D. simulans* (Table S4, $p < 0.01$), suggesting that

the venom composition evolved in response to each of these hosts. This was confirmed by a

linear discriminant analysis (LDA) discriminating the groups of individuals based on generation

for the three hosts (Figure 3). Of note, the generation effect revealed by the PERMANOVA

was only significant for *D. yakuba* 307 when removing F_3 individuals from the analysis (Table

S4, $p = 0.042$), likely because only few females were available at F_3. LDA may overfit groups

composed of few individuals [43,44] and was therefore performed for *D. yakuba* 307 only

between F_7 and F_{11}. LDA discriminated these two groups, therefore confirming the generation

effect revealed by the PERMANOVA (Figure 3).

A PERMANOVA fitted to all hosts together evidenced a significant effect of the "host"

($p < 0.001$), the "generation" ($p < 0.001$) and their interaction ($p < 0.001$) on the variation of

bands intensity, demonstrating that the venom composition evolved differently on the different

hosts (Table S5). Yet, although significant, the combined effects of the "host", "the generation",

and the interaction of the two only accounted for a small part of the variance of the venom

composition ($R^2 = 8.7\%$; Table S5) compared to that explained by the experimental

"population" ($R^2 = 26.6\%$; Table S5). The LDA confirmed the differential venom evolution

according to the host (Figures 4A and 4B, $p < 0.001$) but the 12 groups (four hosts x three

generations) largely overlapped, which confirmed the small part of the variance explained by

the host and the generation. The first four out of the 11 discriminant axes were the only ones to

be biologically meaningful (Figures 4A and 4B). The first two axes discriminated the F_7 and

F_{11} generation for parasitoids reared on the different hosts, except for the two *D. melanogaster*

strains (Figure 4A), evidencing a rapid and differential evolution of the venom composition

according to the host species. This was confirmed by the arrows describing the venom evolution

that were orthogonal or in opposite direction for the different host species except for the two *D.*

melanogaster strains (Figure 4A). Axis 3 discriminated the three generations for parasitoids

reared on *D. melanogaster* strains and *D. simulans* while individuals reared on *D. yakuba* 307

were discriminated by axis 4 (Figure 4B). This suggests that the venom of parasitoids reared on

D. yakuba 307 evolved differently than that of parasitoids reared on the other hosts as confirmed

by the direction of the arrow describing the venom evolution for individuals raised on this

species (Figure 4B).

To better identify the hosts on which the venom composition has evolved differentially,

PERMANOVAs and LDAs were performed to compare them two by two. There was a strong

effect of the "generation × host" interaction for all comparisons involving *D. yakuba* 307 (Table

S5, p < 0.01) and the comparison between *D. melanogaster* S_{Nasr} and *D. simulans* (Table S5, p

= 0.004). This was confirmed by the LDA since groups of individuals were separated at F_7 and

F_{11} and arrows representing the direction of evolution were almost orthogonal (Figures 4C to

4F). No effect of the "generation × host" interaction was observed in the two PERMANOVAs

comparing evolution on *D. melanogaster* R to either *D. melanogaster* S_{Nasr} or *D. simulans*

(Table S5, p = 0.06 for the comparison with *D. melanogaster* S_{Nasr}; p = 0.35 for the comparison

with *D. simulans*). The LDAs nevertheless suggested a trend of differential evolution although

the separation of groups at generations F_7 and F_{11} was lesser between the two strains of *D.*

melanogaster than between *D. melanogaster* R and *D. simulans* (Figures 4G and 4H).

3.5. Identification of protein bands that evolved on each host

Selected protein bands were identified based on their correlation to the linear regressions

(calculated from centroid points of generation groups and represented by arrows) describing

the venom evolution in the LDAs performed for each host separately (Figure 3). However, some

of these protein bands could have been selected only indirectly because of their correlation with

other directly selected bands, either due to their migration proximity on the gel or to a linkage

disequilibrium. To disentangle them, we used a combination of clustering (Figure S3) and

partial correlation analyses (Table S6) as described in Cavigliasso et al. (2019). The analysis

revealed that (i) seven protein bands had evolved on *D. melanogaster* S_{Nasr} (four selected, three

counter-selected), (ii) six on *D. melanogaster* R (three selected, three counter-selected), (iii)

seven on *D. simulans* (five selected, two counter-selected) and (iv) two on *D. yakuba* 307 (one

selected, one counter-selected) (Table 1, Table S6 and Figure S5).

Of the 17 protein bands that evolved in response to at least one host, only three evolved

in the same direction on several hosts. Indeed, bands #10 and #19 were respectively selected

and counter-selected on *D. melanogaster* S_{Nasr}, *D. melanogaster* R and *D. simulans* (Table 1)

and band #27 was selected on *D. simulans* and *D. yakuba* 307. Overall, this confirmed that the

venom composition evolved rapidly and differentially between hosts.

3.6. Putative identification of proteins that evolved on each host

The tentative identification of the proteins contained in the bands under selection was

performed by matching these bands with those in 1D electrophoresis gels used for *L. boulardi*

ISm and ISy venom proteomics [18]. The level of intensity of a band can result from that of

several proteins having migrated at the same position. However, since the initial composition

of the venom resulted from crosses between the ISm and ISy lines, the proteins responsible for

a high intensity of a band were likely to be the most abundant proteins in the corresponding

protein band of ISm or ISy venom. We therefore used the number of peptides matches from a

previous mass spectrometry [18] to classify the proteins in the bands as abundant or not (Table

2). We could identify at least one abundant protein for 12 out of the 17 bands under selection,

which are most likely responsible for the observed changes in the overall band intensity.

Although their coding sequence has been previously determined[18], the most abundant

proteins in five out of these 12 bands had no similarity to known proteins (Table 2). Their

biochemical function thus remains to be determined as for a majority of the proteins contained

in the ISm and ISy venoms[18]. Among the bands that evolved on one host only, a Glucose-

Methanol-Choline (GMC) oxidoreductase was the most abundant protein in band #12 selected

on *D. melanogaster* R (Table 2), and a RhoGAP, LbGAP, was the most abundant protein in

band #23 counter-selected on *D. melanogaster* S_{Nasr}. We performed a specific analysis for

LbGAP that confirmed the counter-selection on *D. melanogaster* S_{Nasr} but also on *D.*

melanogaster R (Figure 5A, GLMM, p = 0.009 for *D. melanogaster* S_{Nasr}, p = 0.021 for *D.*

melanogaster R). Without a good specific antibody effective against the GMC oxidoreductase,

no specific analysis could be performed for this protein. The most abundant protein in band

#25, counter-selected on *D. simulans*, was another RhoGAP, LbGAPy4. A third RhoGAP,

LbGAP2, was found as the most abundant protein in bands #31, selected on *D. melanogaster*

R and #32, selected on *D. simulans*, as well as in band #27, selected on *D. simulans* and *D.*

yakuba 307 (Table 2). However, the selection/counter-selection of LbGAP2 was not detected

in the specific analysis (Figure 5B, LMM, p = 0.14 for *D. melanogaster* R and *D. simulans*, p

= 0.20 for *D. melanogaster* S_{Nasr} and p = 0.44 for *D. yakuba* 307).

Regarding the two other bands that evolved on more than one host, a Sushi/SCR/CCP

domain-containing protein (SCR: Short Consensus Repeat; CCP: Complement Control Protein)

was the most abundant protein in band #10, selected on *D. melanogaster* S_{Nasr}, *D. melanogaster*

R and *D. simulans* (Table 2). Finally, the most abundant protein in band #19, counter-selected

on all hosts except *D. yakuba* 307, was the serpin LbSPNy. The counter-selection of the *lbspny*

allele encoding LbSPNy was confirmed by the specific analysis on *D. melanogaster* S_{Nasr}, R

and *D. simulans* (Figure 5C, GLMM, p = 0.022 for *D. melanogaster* S_{Nasr}, p = 0.018 for *D.*

melanogaster R, p = 0.015 for *D. simulans*). It also revealed a selection of *lbspny* on *D. yakuba*

307 (Figure 5C, GLMM, p = 0.005) although not detected by the global approach.

3.7.Trends of venom evolution

The average venom composition of each of the 44 experimental populations was scaled

between 0 and 1 according to their distance to the venom composition of ISm and ISy (Figure

6). When comparing the venom composition in F_3 and F_{11} generations, we observed a trend for

evolution towards the venom composition of ISm for all hosts, except for *D. yakuba* 307 for

which the evolution of the venom did not change the relative distance to parental strains (Figure

6, paired Wilcoxon test, p = 0.011 for *D. melanogaster* S_{Nasr}, p = 0.010 for *D. melanogaster* R,

p = 0.045 for *D. simulans*, p = 0.60 for *D. yakuba* 307).).

We then assigned a value between 0 and 1 to each protein band: 0 when a band was present

in ISy but absent in ISm, 1 for the opposite (Table 1). In agreement with the above results, most

selected protein bands were of the ISm type, whereas most of the counter-selected bands were

of the ISy type (Table 2).

4. Discussion

This study aimed at testing simultaneously the evolution of (i) the parasitic success and

(ii) the venom composition of *L. boulardi* on *Drosophila* hosts from three different species. It

was also expected to be informative on whether a venom allowing to develop on different hosts

would rather contain broad-spectrum factors or a combination of factors specific to each

host.We used an experimental evolution in which selection acted on the standing genetic

variation, not on new mutations appeared during the experiment, as is usually done when using

fast-growing organisms [45]. The experimental evolution was initiated by crossing *L. boulardi*

ISm and ISy parasitoid lines, which differ in their parasitic success on *D. melanogaster* and *D.*

yakuba species [8] and their venom composition [18]. Since both lines have spent many

generations in the laboratory, the initial variation on which natural selection could act during

the experimental evolution resulted mainly from the variation between them. The F_2 female

offsprings, whose venom contains new combinations of proteins, were separated in independent

populations and reared up to the F_{11} generation on the four different host strains / species on

which ISm and/or ISy are successful.

Among these four hosts, *D. melanogaster* S_{Nasr} and *D. simulans* are both susceptible to

ISm and ISy. We previously evidenced an evolution of the venom composition of *L. boulardi*

not only on a resistant host strain but also on a susceptible one [15]. *D. melanogaster* S_{Nasr} and

D. simulans were therefore used to determine whether two hosts could exert a different selection

pressure on the venom composition of *L. boulardi* despite their susceptibility. Our objective

was also i) to confirm that the creation of new combinations of venom factors still allowed

parasitic success on these susceptible host strains and ii) to assess whether the venom

composition could nevertheless evolve, possibly by selecting still effective but less expensive

factors.

A fifth host was used in the experimental evolution, *D. yakuba* 1907 on which neither

ISm nor ISy are successful. Interestingly, parasitoids managed to develop on this host until the

F_6 generation suggesting that the creation of new combinations of venom factors at F_2 has

allowed them to develop in a clearly refractory strain for a certain time. However, these new

combinations were not successfully selected in any of the replicates since all populations raised

on this host ended up being extinct. This may suggest that the virulence factors responsible for

this temporary parasitic success are encoded by the same loci leading to a higher fitness of

heterozygotes (overdominance), such alleles having been lost due to the small population size.

4.1. Parasitic success and venom evolution according to the host

Although the parasitic success remained close to 100% on *D. melanogaster* S_{Nasr} and *D.*

simulans, the composition of the parasitoid venom has nevertheless evolved on these

susceptible hosts. Although genetic drift due to the small number of individuals per replicate

necessarily impacted the venom composition, the observed changes most likely occurred under

selection. Genetic drift is a random phenomenon that differently affects each population

whereas changes in venom composition were common to most of the replicates suggesting a

selection strong enough not to be masked by the drift. The evolution of venom in response to

susceptible hosts suggests a selection of some venom components over others, potentially less

costly to produce. The venom is not only important to overcome host immune defenses but also

to ensure the quality of offspring developement [46]. The differential venom evolution on the

two susceptible hosts could therefore also result from a selection to increase the fitness of the

developing offspring that are facing different susceptible host strains. This scenario would

suggest a fine tuning of the host physiology to ensure the best match with the parasitoid larvae

requirements.

In contrast, the success of parasitoids selected on *D. melanogaster* R and *D. yakuba* 307

significantly increased with the generation. This rapid increase was not surprising since the

selection pressures on parasitoids are very high and it has previously been documented for other

host – parasitoid interactions. Accordingly, the parasitic success of *Asobara tabida* increased

after seven generations of selection on *D. melanogaster* [47] and that of two aphid parasitoids

increased in response to the symbiont-associated resistance of the host [2,3]. A suprising result

was the much higher increase in parasitic success for the parasitoids reared on *D. yakuba* 307

than on *D. melanogaster* R, probably because the former was much less successful at F_3 (about

20% versus 80%). This difference in parasitic success at F_3 is quite surprising since the ISm

line always succeeds on *D. melanogaster* R and ISy on *D. yakuba* 307. It was however

consistent with the high extinction rate of F_3 replicates on *D. yakuba* 307, suggesting a lower

virulence of F_2 parasitoids on this host than on the others. Accordingly, [5,24] identified the

virulence of F_1 (ISm x ISy) hybrids as semi-dominant on *D. melanogaster* and recessive on *D.*

yakuba, although they used different strains of *D. melanogaster* and *D. yakuba* than ours.

4.2. Specialization of parasitoids to their selection host and differential venom evolution

The greater parasitic success on resistant hosts (*D. yakuba* and *D. melanogaster* R)

proved to be specific to the selection host (Figure 2C). Parasitoids maintained on a given

resistant host were more successful on this host than those reared on another host. The

differential evolution of both parasitic success and venom evolution suggests a specialization

of parasitoids on their selection host, as previously reported experimentaly in relation to host resistance [3,4] and predicted by simulation results [48]. Such a differential evolution of venom has also been observed in a vertebrate model – the snake of genus *Echis* – in response to diet [49]. The selection of certain protein bands for each host, suggests a role in parasitic success, while the counter-selection suggests a cost associated with their production or function on a given host. These differential selection and counter-selection of the venom content could explain the lesser success of a parasitoid on a resistant host (such as *D. melanogaster* R or *D. yakuba* 307) after a selection on another host strain. For example, the counter-selection of the protein band #19 in the venom of parasitoids selected on *D. melanogaster* S_{Nasr}, R, and *D. simulans* may have reduced their success on *D. yakuba* 307. Consistently, another study with other host strains showed that F_1 (ISm x ISy) hybrids experienced a decrease in virulence against *D. yakuba* after being reared on *D. melanogaster*, suggesting that virulence factors selected on *D. melanogaster* were costly for parasitism on *D. yakuba* [5,50].

A possible hypothesis to explain the differential evolution of venom between parasitoids selected on the two susceptible hosts, *D. melanogaster* S_{Nasr} and *D. simulans* is the occurence of a host-specific cost to produce certain venom components, resulting in the counter-selection of different proteins. Interestingly, the success on *D. yakuba* of parasitoids selected on *D. melanogaster* S_{Nasr} is lower than those selected on *D. simulans*, confirming the selection of specific venom components to bypass host defenses. It also suggests a difference in immune defenses between each host. As an example, the prophenol-oxidase (PPO) sequences, essential proteins for the melanization process, differ between host species and might be targeted by different venom proteins [51–53].

4.3. A combination of mostly host-specific proteins and a few broad-spectrum proteins

in venom to succeed on several hosts

The change in intensity observed for most of the 17 venom protein bands which evolved

occurred in response to selection by a single host only. This is in accordance with the study of

[4] in which most of the differentially expressed genes of selected parasitoids were lineage-

specific. As a counter-example, two protein bands evolved in the same direction for *D.*

melanogaster S_{Nasr} and R, and *D. simulans* and one for *D. simulans* and *D. yakuba* 307.

Therefore, although the number of evolving protein bands may have been underestimated due

to a lack of power of the global approach or because they contain several proteins of opposite

evolution, our data suggest that the venom of the selected parasitoids contain a combination of

mostly host-specific proteins and a few broad-spectrum proteins to succeed on these hosts.

4.4. The venom of parasitoids selected on D. yakuba 307 evolved more specifically

All PERMANOVAs involving *D. yakuba* 307 showed a significant effect of the

"generation × host" interaction suggesting that the venom composition of parasitoids reared on

this host evolved more differently than that of those reared on the other hosts. Moreover,

parasitoids selected on *D. melanogaster* strains and *D. simulans* showed a trend for evolution

towards the venom composition of ISm between F_3 and F_{11} unlike those selected on *D. yakuba*

307. Likewise, the protein bands #10 and 19, selected and counter-selected on all hosts except

D. yakuba 307 are of ISm and ISy origin, respectively. In addition, the specific analysis revealed

a selection of the *lbspny* allele on *D. yakuba* 307 and a counter-selection on all other hosts.

Since *lbspnm* and *lbspny* are two alleles of a co-dominant marker, LbSPN, this could also be

interpreted as a selection of *lbspnm* on *D. melanogaster* and *D. simulans* although its role in

parasitic success is yet to be demonstrated. These differential trends in the venom composition

towards ISm and ISy types may reflect the geographic distribution of parasitoids and hosts - the

ISm type of *L. boulardi* being Mediterranean and therefore not encountering *D. yakuba*, mainly

present in tropical regions of Africa [5,16,50], as well as the host preferences of the *L. boulardi*

lines. ISy was previously shown to oviposit preferentially in *D. yakuba* while ISm preferred *D.*

melanogaster [54].

4.5. Proteins whose quantity evolved

For many protein bands identified as evolving, the most abundant protein – which is

likely responsible for changes in the band intensity – had no predicted function although its

coding sequence has been previously determined [18]. These proteins have thus been little or

not studied so far but could nevertheless play an essential role in parasitic success and would

deserve more attention. Our method without *a priori* seems therefore relevant to identify new

candidate proteins, possibly involved in the parasitic success.

Among the proteins with a predicted function and identified as the most abundant one

in their evolving band, we found the RhoGAPs LbGAP and LbGAP2 and the serpin LbSPN for

which a Western blot analysis with specific antibodies was also performed. The global approach

and the specific analysis agreed on the counter-selection of LbGAP on *D. melanogaster* S_{Nasr}.

This was also evidenced on the resistant strain of *D. melanogaster* R but with the specific

analysis only, probably because of the lower power of the global approach. This was surprising

since a selection of LbGAP on *D. melanogaster* R was observed in a previous work [15].

However, in this current study, the proportion of LbGAP individuals at F_3 on *D. melanogaster*

hosts was much higher than expected under Hardy-Weinberg equilibrium (90% vs 75%). We

therefore cannot exclude a decrease of the frequency of LbGAP individuals to reach equilibrium

at F_{11} instead of a counter-selection. Moreover, the initial crosses were done in both directions,

instead of only one in the previous study, leading to more balanced frequencies for the overall

ISm / ISy alleles. The selection may therefore have acted differently in the two experiments and

other alleles than *lbgap* been selected on *D. melanogaster* R. LbGAP is involved in changes of

the morphological shape of *D. melanogaster* R lamellocytes, possibly preventing the

encapsulation by the host [17,19–22]. We could hypothesize that LbGAP is involved in a

strategy to suppress encapsulation. However, we did not observe the increase in parasitoid

ability to suppress encapsulation that would therefore be expected under this assumption, but

rather an increase of the parasitoid ability to escape from a capsule (Figure 2B). The venom

contains several other RhoGAPs including LbGAP2, all mutated on their catalytic site, which

might "somehow" compensate for the reduced LbGAP quantity by acting in parasitism success.

The selection of LbGAP2 on *D. melanogaster* R, *D. simulans* and *D. yakuba* 307 based on the

global approach was however not supported by the specific analysis. This suggests that other

proteins in the LbGAP2-containing bands are responsible of the changes in intensity detected

by the global approach.

Interestingly, we evidenced the selection of LbSPNy on *D. yakuba* 307, although with

the specific analysis only, probably because of the small number of individuals at F_3 on that

host. This is in line with the demonstrated involvement of LbSPNy in the inhibition of the

phenoloxidase cascade activation of this same species of *Drosophila* [23]. The phenoloxidase

cascade leads to the melanization of the capsule and the release of cytotoxic radicals to kill the

parasitoid [23,51–53]. The melanisation process being involved in the strengthening of the

capsule, this selection would be rather consistent with the increase of the escape ability of the

formed capsule in parasitoids selected on *D. yakuba* associated with the increase in parasitic

success (Figure 2B). The global approach and the specific analysis also showed the counter-

selection of LbSPNy on *D. melanogaster* S_{Nasr} and R and *D. simulans* in agreement with

previous results for *D. melanogaster* hosts [15]. This suggests a possible cost of either the

production or the presence of LbSPNy in their venom.

In conclusion, the parasitoid model is very relevant for an experimental evolutionary

approach compared to other models of venomous animals such as scorpions or snakes although

promising advances were obtained from studies of venomics and virulence on such models

[49,55]. Our results have highlighted a specialization of parasitoids on their selection host, in

link with a rapid differential evolution of the composition of the venom according to the host.

Most of the evolving venom proteins evolved in response to selection by a single host,

suggesting that parasitoids use at least partially different mechanisms to bypass the defenses of

different hosts, and therefore that these host species may also implement partly different defense

mechanisms. *D. melanogaster* and *D. simulans* might share some of them so that part of the

venom proteins to succeed on these host species would be common. *D. yakuba* is more apart,

the maintenance of parasitoids on this species resulting in the selection of more specific venom

factors. From these data, we end up with no universal answer to the question of the venom

content of a "generalist" parasitoid in terms of broad-spectrum or host-specific factors. The

venom of ISm may contain a mixture of both, allowing success on *D. melanogaster* and *D.*

simulans, species usually found in sympatry, while the question remains more open for ISy

venom. Finally, in a more general context, the rapid evolution of the venom highlights the

strong capacity of parasitoids to possibly adapt to their environment [56]. It is notably striking

that crosses between parasitoids creating new combinations of venomous proteins could help

them succeed on initially refractory hosts.

References

1.   Godfray HCJ. 1994 *Parasitoids: behavioral and evolutionary ecology*. Princeton Univ.

Press, Princeton, NJ.

2.   Dion E, Zélé F, Simon J-C, Outreman Y. 2011 Rapid evolution of parasitoids when

faced with the symbiont-mediated resistance of their hosts. *J. Evol. Biol.* **24**, 741–750.

(doi:10.1111/j.1420-9101.2010.02207.x)

3.   Rouchet R, Vorburger C. 2014 Experimental evolution of parasitoid infectivity on

symbiont-protected hosts leads to the emergence of genotype specificity. *Evolution (N.*

Y). **68**, 1607–1616. (doi:10.1111/evo.12377)

4.   Dennis AB, Patel V, Oliver KM, Vorburger C. 2017 Parasitoid gene expression

changes after adaptation to symbiont-protected hosts. *Evolution (N. Y).* **71**, 2599–2617.

(doi:10.1111/evo.13333)

5.   Dupas S, Boscaro M. 1999 Geographic variation and evolution of immunosuppressive

genes in a *Drosophila* parasitoid. *Ecography (Cop.).* **22**, 284–291. (doi:10.1111/j.1600-

0587.1999.tb00504.x)

6.   Fors L, Markus R, Theopold U, Ericson L, Hambäck PA. 2016 Geographic variation

and trade-offs in parasitoid virulence. *J. Anim. Ecol.* **85**, 1595–1604.

(doi:10.1111/1365-2656.12579)

7.   Carton Y, Bouletreau M, Alphen J, Van Lenteren J. 1986 The *Drosophila* parasitic

wasps. In *The Genetics and Biology of Drosophila*, pp. 347–394. Academic Press Inc.

(London) Ltd.

8.   Dubuffet A, Dupas S, Frey F, Drezen JM, Poirié M, Carton Y. 2007 Genetic

interactions between the parasitoid wasp *Leptopilina boulardi* and its *Drosophila* hosts.

Heredity (Edinb). **98**, 21–27. (doi:10.1038/sj.hdy.6800893)

9.   Carton Y, Poirié M, Nappi AJ. 2008 Insect immune resistance to parasitoids. *Insect Sci.*

15, 67–87. (doi:10.1111/j.1744-7917.2008.00188.x)

10.   Nappi AJ. 2010 Cellular immunity and pathogen strategies in combative interactions

involving *Drosophila* hosts and their endoparasitic wasps. *Invertebr. Surviv. J.* **7**, 198–

210.

11.   Vlisidou I, Wood W. 2015 *Drosophila* blood cells and their role in immune responses.

FEBS J. **282**, 1368–1382. (doi:10.1111/febs.13235)

12.   Asgari S, Rivers DB. 2011 Venom proteins from polydnavirus-producing

endoparasitoids: Their role in host-parasite interactions. *Annu. Rev. Entomol.* **56**, 313–

35. (doi:10.1146/annurev-ento-120709-144849)

13. Poirié M, Colinet D, Gatti JL. 2014 Insights into function and evolution of parasitoid

wasp venoms. *Curr. Opin. Insect Sci.* **6**, 52–60. (doi:10.1016/j.cois.2014.10.004)

14. Moreau SJM, Asgari S. 2015 Venom proteins from parasitoid wasps and their

biological functions. *Toxins (Basel).* **7**, 2385–2412. (doi:10.3390/toxins7072385)

15. Cavigliasso F, Mathé-Hubert H, Kremmer L, Rebuf C, Gatti J, Malausa T, Colinet D,

Poirié M. 2019 Rapid and differential evolution of the venom composition of a

parasitoid wasp depending on the host strain. *Toxins (Basel).* **11**, 629.

16. Dubuffet A, Colinet D, Anselme C, Dupas S, Carton Y, Poirié M. 2009 Variation of

Leptopilina boulardi success in *Drosophila* hosts. What is inside the black box? *Adv.*

Parasitol. **70**, 147–188. (doi:10.1016/S0065-308X(09)70006-5)

17. Colinet D, Schmitz A, Cazes D, Gatti JL, Poirié M. 2010 The origin of intraspecific

variation of virulence in an eukaryotic immune suppressive parasite. *PLoS Pathog.* **6**,

1–11. (doi:10.1371/journal.ppat.1001206)

18. Colinet D, Deleury E, Anselme C, Cazes D, Poulain J, Azema-Dossat C, Belghazi M,

Gatti JL, Poirié M. 2013 Extensive inter- and intraspecific venom variation in closely

related parasites targeting the same host: The case of *Leptopilina* parasitoids of

Drosophila. *Insect Biochem. Mol. Biol.* **43**, 601–611. (doi:10.1016/j.ibmb.2013.03.010)

19. Labrosse C, Stasiak K, Lesobre J, Grangeia A, Huguet E, Drezen JM, Poirié M. 2005 A

RhoGAP protein as a main immune suppressive factor in the *Leptopilina boulardi*

(Hymenoptera, Figitidae)-*Drosophila melanogaster* interaction. *Insect Biochem. Mol.*

Biol. **35**, 93–103. (doi:10.1016/j.ibmb.2004.10.004)

20. Labrosse C, Eslin P, Doury G, Drezen JM, Poirié M. 2005 Haemocyte changes in *D.*

melanogaster in response to long gland components of the parasitoid wasp *Leptopilina*

boulardi: a Rho-GAP protein as an important factor. *J. Insect Physiol.* **51**, 161–170.

(doi:10.1016/j.jinsphys.2004.10.004)

21.   Colinet D, Schmitz A, Depoix D, Crochard D, Poirié M. 2007 Convergent use of

RhoGAP toxins by eukaryotic parasites and bacterial pathogens. *PLoS Pathog.* **3**, e203.

(doi:10.1371/journal.ppat.0030203)

22.   Wan B, Goguet E, Ravallec M, Pierre O, Lemauf S, Volkoff A-N, Gatti J-L, Poirié M.

2019 Venom atypical extracellular vesicles as interspecies vehicles of virulence factors

involved in host specificity: the case of a *Drosophila* parasitoid wasp. *Front. Immunol.*

10, 1–14. (doi:10.3389/fimmu.2019.01688)

23.   Colinet D, Dubuffet A, Cazes D, Moreau S, Drezen JM, Poirié M. 2009 A serpin from

the parasitoid wasp *Leptopilina boulardi* targets the *Drosophila* phenoloxidase cascade.

Dev. Comp. Immunol. **33**, 681–689. (doi:10.1016/j.dci.2008.11.013)

24.   Dupas S, Frey F, Carton Y. 1998 A single parasitoid segregating factor controls

immune suppression in *Drosophila*. *J. Hered.* **89**, 306–311.

(doi:10.1093/jhered/89.4.306)

25.   Carton Y, Frey F, Nappi A. 1992 Genetic determinism of the cellular immune reaction

in *Drosophila melanogaster*. *Heredity (Edinb).* **69**, 393–399.

(doi:10.1038/hdy.1992.141)

26.   Carton Y, Nappi AJ. 1997 *Drosophila* cellular immunity against parasitoids. *Parasitol.*

Today **13**, 218–227.

27.   Poirié M, Frey F, Hita M, Huguet E, Lemeunier F, Periquet G, Carton Y. 2000

Drosophila resistance genes to parasitoids: Chromosomal location and linkage

analysis. *Proc. R. Soc. B Biol. Sci.* **267**, 1417–1421. (doi:10.1098/rspb.2000.1158)

28.   Kim-Jo C, Gatti JL, Poirié M. 2019 *Drosophila* cellular immunity against parasitoid

wasps: A complex and time-dependent process. *Front. Physiol.* **10**, 603.

(doi:10.3389/fphys.2019.00603)

29.   Hita M, Espagne E, Lemeunier F, Pascual L, Carton Y, Periquet G, Poirié M. 2006

Mapping candidate genes for *Drosophila melanogaster* resistance to the parasitoid

wasp *Leptopilina boulardi*. *Genet. Res.* **88**, 81–91. (doi:10.1017/S001667230600841X)

30.   Fellowes MDE, Kraaijeveld AR, Godfray HCJ. 1998 Trade-off associated with

selection for increased ability to resist parasitoid attack in *Drosophila melanogaster*.

Proc. R. Soc. B Biol. Sci. **265**, 1553–1558. (doi:10.1098/rspb.1998.0471)

31.   Singmann H, Bolker B, Westfall J, Aust F. 2019 afex: analysis of factorial

experiments.

32.   Chung Y, Rabe-Hesketh S, Dorie V, Gelman A, Liu J. 2013 A nondegenerate

penalized likelihood estimator for variance parameters in multilevel models.

Psychometrika **78**, 685–709.

33.   Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JSS.

2009 Generalized linear mixed models: a practical guide for ecology and evolution.

Trends Ecol. Evol. **24**, 127–135. (doi:10.1016/j.tree.2008.10.008)

34.   Harrison XA. 2015 A comparison of observation-level random effect and Beta-

Binomial models for modelling overdispersion in Binomial data in ecology &

evolution. *PeerJ* **2015**. (doi:10.7717/peerj.1114)

35.   Hothorn T, Bretz F, Westfall P. 2008 Simultaneous inference in general parametric

models. *Biometrical J.* **50**, 346–363. (doi:10.1002/bimj.200810425)

36.   Mathé-Hubert H, Gatti JL, Colinet D, Poirié M, Malausa T. 2015 Statistical analysis of

the individual variability of 1D protein profiles as a tool in ecology: An application to

parasitoid venom. *Mol. Ecol. Resour.* **15**, 1120–1132. (doi:10.1111/1755-0998.12389)

37.   Colinet D, Mathé-Hubert H, Allemand R, Gatti JL, Poirié M. 2013 Variability of

venom components in immune suppressive parasitoid wasps: From a phylogenetic to a

population approach. *J. Insect Physiol.* **59**, 205–212.

(doi:10.1016/j.jinsphys.2012.10.013)

38.   Colinet D, Kremmer L, Lemauf S, Rebuf C, Gatti JL, Poirié M. 2014 Development of

RNAi in a *Drosophila* endoparasitoid wasp and demonstration of its efficiency in

impairing venom protein production. *J. Insect Physiol.* **63**, 56–61.

(doi:10.1016/j.jinsphys.2014.02.011)

39.   Dixon P. 2003 VEGAN , a package of R functions for community ecology. *J. Veg. Sci.*

14, 927–930.

40.   Anderson MJ. 2017 Permutational Multivariate Analysis of Variance

(PERMANOVA). *Wiley StatsRef Stat. Ref. Online* , 1–15.

(doi:10.1002/9781118445112.stat07841)

41.   Dray S, Dufour A-B. 2007 The ade4 package: implementing the duality diagram for

ecologists. *J. Stat. Softw.* **22**, 1–20.

42.   Pinheiro J, Bates D, DebRoy S, D S, Team RDC. 2011 nlme: linear and nonlinear

mixed effects models. *R Packag. version* **3**, 1–97.

43.   Qiao Z, Zhou L, Huang J. 2008 Effective linear discriminant analysis for high

dimensional, low sample size data. *Proceeding World Congr. Eng.* **2**, 2–4.

44.   Luo D, Ding C, Huang H. 2011 Linear discriminant analysis: New formulations and

overfit analysis. *Proc. Natl. Conf. Artif. Intell.* **1**, 417–422.

45.   Lenski RE. 2017 Experimental evolution and the dynamics of adaptation and genome

evolution in microbial populations. *ISME J.* **11**, 2181–2194.

(doi:10.1038/ismej.2017.69)

46.   Mrinalini, Siebert AL, Wright J, Martinson E, Wheeler D, Werren JH. 2015 Parasitoid

venom induces metabolic cascades in fly hosts. *Metabolomics* **11**, 350–366.

(doi:10.1007/s11306-014-0697-z)

47.   Kraaijeveld AR, Hutcheson KA, Limentani EC, Godfray HCJ. 2001 Costs of

counterdefenses to host resistance in a parasitoid of *Drosophila*. *Evolution (N. Y).* **55**,

1815–1821. (doi:10.1111/j.0014-3820.2001.tb00830.x)

48.   Lapchin L. 2002 Host-parasitoid association and diffuse coevolution: When to be a

generalist? *Am. Nat.* **160**, 245–254. (doi:10.1086/341020)

49.   Barlow A, Pook CE, Harrison RA, Wüster W. 2009 Coevolution of diet and prey-

specific venom activity supports the role of selection in snake venom evolution. *Proc.*

R. Soc. B Biol. Sci. **276**, 2443–2449. (doi:10.1098/rspb.2009.0048)

50.   Dupas S, Dubuffet A, Carton Y, Poirié M. 2009 Local, geographic and phylogenetic

scales of coevolution in *Drosophila* -parasitoid interactions. *Adv. Parasitol.* **70**, 281–

295. (doi:10.1016/S0065-308X(09)70011-9)

51.   Dudzic JP, Kondo S, Ueda R, Bergman CM, Lemaitre B. 2015 *Drosophila* innate

immunity: Regional and functional specialization of prophenoloxidases. *BMC Biol.* **13**,

1–16. (doi:10.1186/s12915-015-0193-6)

52.   Nappi AJ, Christensen BM. 2005 Melanogenesis and associated cytotoxic reactions:

Applications to insect innate immunity. *Insect Biochem. Mol. Biol.* **35**, 443–459.

(doi:10.1016/j.ibmb.2005.01.014)

53.   Cerenius L, Lee BL, Söderhäll K. 2008 The proPO-system: pros and cons for its role in

invertebrate immunity. *Trends Immunol.* **29**, 263–271. (doi:10.1016/j.it.2008.02.009)

54.   Dubuffet A, Álvarez CIR, Drezen JM, Van Alphen JJM, Poirié M. 2006 Do parasitoid

preferences for different host species match virulence? *Physiol. Entomol.* **31**, 170–177.

(doi:10.1111/j.1365-3032.2006.00505.x)

55.   Evans ERJ, Northfield TD, Daly NL, Wilson DT. 2019 Venom costs and optimization

in scorpions. *Front. Ecol. Evol.* **7**, 1–7. (doi:10.3389/fevo.2019.00196)

56.   Mathé-Hubert H, Kremmer L, Colinet D, Gatti JL, Van Baaren J, Delava É, Poirié M.

2019 Variation in the venom of parasitic wasps, drift, or selection? Insights from a

multivariate QST analysis. *Front. Ecol. Evol.* **7**, 256. (doi:10.3389/fevo.2019.156)

Author contributions: F.C., H.M-H, J-L.G., D.C. and M.P. designed and planned the

experiments. F.C. carried out the experiments. F.C. and H.M-H. analyzed the data. F.C. and

D.C. wrote the first draft. All authors were involved in editing of the final version. All authors

approve the final version of the manuscript. Authors declare no conflict of interest.

Acknowledgments: We are grateful to Christian Rebuf for help in insect rearing and to Dr.

Dominique Joly (Gif/Yvette) for kindly providing the *D. yakuba* strain 307.14. We also thank

the "Plateau de microscopie ISA" and Maya Belghazi (CNRS, INP, Marseille, France) for

updating the mass spectrometry analyzes. Fanny Cavigliasso PhD was funded by the French

Ministry of Higher Education and Research. The work received support from the University of

Nice Sophia Antipolis" (Incentive Scientific Credits) and was supported by the French

Government (National Research Agency, ANR) through the "Investments for the Future"

programs LABEX SIGNALIFE ANR-11-LABX-0028-01 and IDEX UCAJedi ANR-15-

IDEX-01. We thank the three reviewers for their comments on an earlier version of the

manuscript.

Data Accessibility: The data sets supporting the results are available in a figShare repository

(https://doi.org/10.6084/m9.figshare.13139090.v1). Drosophila strains/species and parasitoid

lines are maintained in the laboratory and available upon request.

826
827

Band Number	Band Origin	Protein bands evolution in response to the host			
		D. melanogaster S$_{Nasr}$	*D. melanogaster* R	*D. yakuba* 307 (F$_7$ *vs* F$_{11}$)	*D. simulans*
2	0.60	↗			
4	0.26			↘	
7	0.44	↗			
10	0.51	↗	↗		↗
12	0.62		↗		
17	0.36		↘		
18	0.23		↘		
19	0.18	↘	↘		↘
22	0.28	↘			
23	0.58	↘			
25	0.32				↘
27	0.64			↗	↗
31	0.66		↗		
32	0.63				↗
34	0.54				↗
35	0.54				↗
36	0.63	↗			

Table 2. Correspondence between evolving bands and their putative protein content determined from the comparison with data from Colinet et al. (2013a). Only proteins for which at least 10 peptide matches were found in Mascot searches to unisequences identified in transcriptomics of the venom apparatus [18] were considered as abundant and therefore listed. The number of proteins found in the band, their predicted function and the number of peptide matches for each unisequence are provided. Data on the band origin and direction of evolution are from Table 1. Up ("↗") and down ("↘") arrows indicate a selection or a counter-selection, respectively. *D. mel.* S_{Nasr}: *D. melanogaster* S_{Nasr}; *D. mel.* R: *D. melanogaster* R. NA: data not available. On *D. yakuba* 307, the evolution reflects changes between F_7 and F_{11} only.

Reference band	Number of proteins in the band	Putative function	Number of peptide matches	Band origin	Band evolution on			
					D. mel. S_{Nasr}	*D. mel.* R	*D. yakuba* 307 (F_7 *vs* F_{11})	*D. simulans*
2	1	Unknown	12	ISm	↗			
4	3	Unknown	55	ISy			↘	
		Sushi/SCR/CCP domain containing protein	11					
		Unknown	10					
7 (no abundant protein found)	NA	NA	NA	ISy	↗			
10	2	Sushi/SCR/CCP domain containing protein	10	ISm	↗	↗		↗
		Unknown	10					
12	1	GMC oxidoreductase	40	ISm		↗		
17	1	Unknown	20	ISy			↘	
18 (not sequenced in [18])	NA	NA	NA	ISy			↘	

Reference band	Number of proteins in the band	Putative function	Number of peptide matches	Band origin	Band evolution on			
					D. mel. S_{Nasr}	*D. mel.*R	*D. yakuba* 307 (F_7 *vs* F_{11})	*D. simulans*
19	1	Serpin (LbSPNy)	81	ISy	↘	↘		↘
22	2	Unknown Serpin (LbSPNy)	19 10	ISy	↘			
23	5	RhoGAP (LbGAP) Unknown Serpin (LbSPNm) Unknown Unknown	52 21 17 12 11	ISm	↘			
25	1	RhoGAP (LbGAPy4)	24	ISy				↘
27	3	RhoGAP (LbGAP2) RhoGAP (LbGAP1) Serpin (LbSPNm)	43 23 15	ISm			↗	↗
31 or 32 (not separated in [18])	2	RhoGAP (LbGAP2) Unknown	20 18	ISm		↗ (#31)		↗ (#32)
34 (no abundant protein found)	NA	NA	NA	ISm				↗
35	1	Unknown	18	ISm				↗
36 (no abundant protein found)	NA	NA	NA	ISm	↗			

858
859
Figure 1. Biological model and experimental evolution protocol. A. Outcome of interaction
between the five host strains (*D. melanogaster* S_{Nasr}, *D. melanogaster* R, *D. yakuba* 307, *D.*
yakuba 1907 and *D. simulans*) and the two *L. boulardi* lines (ISm and ISy). The black arrow
shows an encapsulated parasitoid egg inside a *Drosophila* larva. B. Design of the experimental
evolution. ISm and ISy: ISm and ISy lines of *L. boulardi*; S_{Nasr}: *D. melanogaster* S_{Nasr}; R: *D.*
melanogaster R; y307: *D. yakuba* 307; simu: *D. simulans*. The red cross indicates the extinction
of parasitoids reared on *D. yakuba* 1907, thus no analysis was performed on individuals from
this host. F_3, F_7 and F_{11}: the three generations of *L. boulardi* that were analyzed for venom
composition and parasitic success. The F_3 is the first generation under selection.

Figure 2. Evolution of the parasitoid's ability to bypass the encapsulation response of the host. A. Possible outcomes observed in *Drosophila* larvae 48 hours after parasitism and details of the analyzed parameters: parasitic success, parasitoid capacity to inhibit encapsulation and parasitoid capacity to escape from a capsule. B. Evolution of (i) parasitic success, (ii) parasitoid capacity to inhibit the egg encapsulation by the host or (iii) parasitoid capacity to escape from a capsule, depending of the selection host. . C. Capacity of parasitoids from the F_{11} generation to bypass encapsulation by four different hosts. The hosts listed on the

877   topare the "selection hosts", those listed on the left are the "assayed hosts", used for

878   parasitism assays. *D. mel.*: *D. melanogaster*. Letters above bars indicates the significance of

879   the difference. Different colors of letters indicate different statistical tests. Error bars indicate

880   standard errors.

Figure 3. Evolution of the venom composition according to the selection host. Position of the individuals (shown as dots) on the two discriminant axes for each selection host. Individuals are grouped and coloured according to the generation (F_3, F_7 and F_{11}). The arrows represent the trend of the venom evolution. The dotted arrows represent the linear regressions calculated from coordinates of the three centroid points corresponding to the F_3, F_7 and F_{11} generations and weighted by the number of individuals per generation. For *D. yakuba* 307, the linear regression was calculated from centroid points of F_7 and F_{11} only due to the low number of females available in F_3. P-values obtained from the PERMANOVA for the "generation" effect are provided at the bottom right of the LDA (see Table S4).

Figure 4. Differential evolution of venom composition. Discriminante analysis showing elipses

and centroids (intersection of dashed lines) of each group formed by the host and the generation

are represented (see Figure S4 for more details on the position of each individual). The name

of each group is written on the centroid. S_{Nasr}: *D. melanogaster* S_{Nasr}; simu: *D. simulans*; R: *D.*

melanogaster R; yak: *D. yakuba* 307. A-B. Venom evolution in response to the four hosts at F_3,

F_7 and F_{11} generations on discriminant axes 1 and 2 (A) and discriminant axes 3 and 4 (B). C-

H. Discriminant analysis performed on the two first discriminant axes for each two-by-two

comparison of venom composition between parasitoids selected on different hosts. For each of

these LDAs, the pair of host considered is indicated at the left. Individuals are grouped and

coloured according to their selection host and generation. The dotted arrows representing the

direction of the venom evolution are the linear regressions fitted to the coordinates of the three

centroid points (F_3, F_7 and F_{11}). P-values obtained from the PERMANOVA for effects of the

"generation", "host" and "generation $\times$ host" interaction are provided at the bottom right of the

LDA (see Table S5 for more details).

Figure 5. Specific analysis of the evolution of LbGAP, LbGAP2 and LbSPN proteins of parasitoids selected on *D. melanogaster* S_{Nasr} (in yellow), *D. melanogaster* R (in red), *D. simulans* (in purple) and *D. yakuba* 307 (in grey-green). A. Evolution of the frequency of individuals harbouring a high quantity of the LbGAP protein. B. Evolution of the LbGAP2 protein amount, relative to the total amount of proteins in the venom samples. C. Evolution of the frequency of the *lbspny* allele.

Figure 6. Relative distance between the venom composition of parasitoids selected on *D. melanogaster* S_{Nasr} (in yellow), *D. melanogaster* R (in red), *D. simulans* (in purple) and *D. yakuba* 307 (in grey-green) in F_3 and F_{11} and that of *L. boulardi* ISm (1) and ISy (0) lines. The horizontal dashed black line indicates the proportion of ISm alleles after the crossing between ISm and ISy lines (0.5 at the beginning of the experiment). In *D. yakuba* 307, one replicate is missing in F_3 because not enough individuals were available to produce the next generation, perform parasitic tests and do the venom analysis.