An Algorithmic framework for genome-wide identification of Sugarcane (*Saccharum officinarum*)-encoded microRNA targets against SCBV

Muhammad Aleem Ashraf[1, 2*, †], Xiaoyan Feng[1, †], Xiaowen Hu[3, †], Fakiha Ashraf[1], Linbo Shen[1], and Shuzhen Zhang[1, *]

[1] Institute of Tropical Bioscience and Biotechnology, Sugarcane Research Centre of Chinese Academy of Tropical Agricultural Sciences, Haikou, China

[2] Department of Bioscience and Biotechnology, Khwaja Fareed University of Engineering and Information Technology, Rahim Yar Khan, Pakistan

[3] Zhanjiang Experimental Station, Chinese Academy of Tropical Agricultural Sciences, Zhanjiang, Guandong, China

* Correspondence: aleem.ashraf@kfueit.edu.pk   Tel.:   +92-68-588-2431   (M.A.A.); zhangshuzhen@itbb.org.cn; Tel.: +86-898-6621-5109 (S.Z).

[†] These authors contributed equally to this work.

Abstract:

Sugarcane Bacilliform Virus (SCBV) is considered an economically the most damaging pathogen for sugarcane production worldwide. Three ORFs are characterized in a single molecule of circular, ds-DNA genome of the SCBV, encoding for hypothetical protein (ORF1), DNA binding protein (ORF2) and Polyprotein (ORF3). The study was aimed to predict and comprehensively evaluate sugarcane miRNAs for the silencing of SCBV genome using *in-silico* algorithms. Computational methods were used for prediction of candidate miRNAs from sugarcane (*S. officinarum* L.) to silence the expression of SCBV genes through translational inhibition by mRNA cleavage. Mature sugarcane miRNAs were retrieved and were assessed to hybridization with the SCBV genome. A total of fourteen potential candidate miRNAs from

sugarcane were computed by all the algorithms used for the silencing of SCBV. A consensus of three algorithms predicts hybridization sites of sof-miR159e at common locus 5534. The miRNA-mRNA interaction was estimated by computing free-energy of miRNA-mRNA duplex using RNAcofold algorithm. Regulatory network of predicted candidate miRNAs of sugarcane with SCBV ORFs, generated using Circos, identify novel targets. Consequently, detecting and discarding inefficient amiRNAs prior to cloning would help suppressed mutants faster. The efficacy of predicted candidate miRNAs was evaluated to test the survival rate of the *in vitro* amiRNA-mediated effective badnaviral silencing and resistance in sugarcane cultivars.

Keywords: miRNAs; *Saccharum officinarum*; *Sugarcane bacilliform virus;* Genome silencing; RNAhybrid; free-energy; psRNATarget; RNAi; computational approach; badnavirus.

1. Introduction

Sugarcane bacilliform viruses (SCBVs) are classified into Badnavirus genus of Caulimoviridae family. These viruses are composed of monopartite, circular, non-enveloped bacilliform with dimensions (30nm ×120-150 nm), double-stranded DNA (ds-DNA)-genome of approximately 7.2-9.2 Kbp in size [1]. The genome of SCBV constitutes three major open reading frames (ORFs) that are located on the 'plus DNA strand' with a single discontinuity [2]. ORF1 encodes a small hypothetical protein which has unknown function. ORF2 encodes a virion-associated DNA-binding protein. ORF3 encodes a largest polyprotein which is represented as (P3.) that was composed of multiple functional sub-units. The polyprotein (P3) is cleaved by a viral aspartic protease to give rise multiple functional small proteins: intracellular movement, capsid, aspartic protease, RT, and ribonuclease [1-6]. H. RT-RNaseH-coding region was considered the most common taxonomic marker for identification of badnaviral genomic

components. This coding region is a standard source to compare the sequence diversity of the

badnaviral genomes [7]. The *Badnavirus* genome is replicated using reverse transcription of a

more-than-genome length RNA. It works as a standard template to complete the translation

process of the badnavirus proteins. It is also used as reverse transcription for replication of the

badnavirus genome [8, 9].

First report of SCBV infection was observed in the Cuban sugarcane cultivar B34104 in

1985 [10].These viruses are disseminated worldwide and decrease crop production significantly

because of accessibility and exchange of biological materials globally. SCBV is a source of

infection for several bioenergy crop sugarcane cultivars, varieties and species. Broad host range

of SCB includes diverse and economically important members of *Poaceae* (sugarcane, rice) and

Musaceae (banana) families. Natural transmission of SCBV is disseminated by sap-feeding

mealy bug species by vegetative cutting [11]. SCBV disease symptoms include chlorosis and leaf

freckle. Infected sugarcane plants were also monitored to have no symptoms. SCBV-infection in

sugarcane plants causes much reduced sucrose content, juice, stalk weigh, purity and gravity in

recent years in China [6]. Plant employs multiple defense strategies to control virus invasion,

replication and transmission in the cell. RNAi-mediated response of plants against invading

viruses is highly significant during infection period [12]. RNAi mechanism inhibits the protein

translation at mRNA level using a highly sequence specific strategy [13]. Sugarcane plant

species were imparted natural inherited active immunity consisting of small, non-coding

microRNAs (miRNAs) to control viral diseases. The miRNA-mediated genome silencing was

considered to certify the activity of positive or negative immune-based regulation. It is also

considered as key activator of immune defense in plants [14]. These miRNAs belong to a class

of 21-23 nucleotides long, endogenous, non-coding RNA molecules that plays a major role to

control the gene expression at post-transactional level. RNA silencing in the form of miRNAs

within the host plant, is a source of natural immunity. It provides resistance to the host plant after

infection of foreign genetic elements including plant viruses [15-17].

Artificial microRNA (amiRNA)-mediated silencing of invading viruses in plants was first

time reported by Niu [18]. This amiRNA-based silencing strategy was applied effectively in

many plants to combat plant viruses: cotton leaf curl Kokhran virus (CLCuKoV-Bu) [19],

Cucumber mosaic virus (CMV)[20], CymMV and ORSV [21]. Serological method, ELISA,

immunosorbent electron microscopy (ISEM) [22], IC-PCR [23], RT-PCR [24], rolling circle

amplification (RCA) [25] are the most widely used reliable detection techniques for SCBV. New

strategies are under development for improved diagnoses and cure of badnaviruses [26].

Prediction and validation of host-virus interactions is the basic step to understand the

critical functional efficacy of miRNAs in the broader range of miRNA-mRNA regulatory

networks controlling biological processes. Each target miRNA possesses potential hybridization

binding sites. In order to validate potential miRNA target experimentally is highly costly and

laborious. An *In Silico* approach to identify miRNA targets concludes potential target binding

sites for laboratory research. Computational strategies determine how miRNAs target to a

desirable mRNA. A large number of computational algorithms are publically available for

miRNA target prediction. It is highly advantageous to get biological access to several

computational tools with different features. The researcher is challenged with an important

choice for selecting suitable tools for prediction [27].

The current study was designed to implement miRNA prediction algorithms and identify

potential targets of sugarcane-derived miRNAs against SCBV as precedent for creating

resistance to sugarcane cultivars using RNAi technology. We have also screened the potential

sugarcane miRNAs for understanding sugarcane-badnavirus interactions. The novel amiRNA-
mediated approach is credible to generate SCBV-resistance sugarcane plants through genetic
engineering.

2. Materials and Methods

2.1. Retrieval of Sugarcane MicroRNAs

The sugarcane mature microRNAs (miRNAs) and stem-loop hairpin precursor sequences
were retrieved from the miRNA biological sequence database, miRBase (v22)
(http://mirbase.org/). miRBase serve as primary public repository and standard online reference
resource for all published miRNA sequences, textual annotation and gene nomenclature [28-30].
In this study, 16 *S.officinarum* (MI0001756-MI0001769) and 19 Saccharum spp. (MI0018180-
MI18197) miRNA sequences were downloaded (Supplementary Table 1).

2.2. SCBV Genome Retrieval and Annotation

The full-length transcript of the SCBV-BRU genome (Genbank accession number
JN377537.1, size 7884 bp) was isolated from the S.officinarum cultivar, cloned, sequenced and
submitted to the NCBI database [31]. Expected size and abundance of ORFs along nucleotide
distribution of the above mentioned NCBI retrieved SCBV-BRU genome was estimated using
pDRAW32 DNA analysis software (v 1.1.129). The SCBV-BRU genome annotation is designed
which represents ORFs of varying lengths.

2.3. Target Prediction in SCBV Genome

Target prediction is considered a key feature towards identification of credible miRNA-
mRNA interactions hybridization. A present, many target prediction algorithms was designed to

predict and identify best candidate miRNA targets. Each tool uses a specific criteria and

stringency for miRNA target prediction. We used four target prediction algorithms cited in

literature — miRanda, RNA22, RNAhybrid and psRNATarget to find the most relevant

sugarcane miRNAs for silencing of SCBV genome (Table1). We designed the most effective

computational approach to analyze miRNA targets at three different prediction levels: individual,

union and intersection. A detailed workflow pipeline was present in (Figure 1).

2.4. miRanda

miRanda is considered more frequently used standard miRNA-target predictor scanning

algorithm. It was implemented first time in 2003 [32] and updated into a web-based tool for

miRNA analysis [33]. It considers following algorithmic features; seed-based interaction,

minimum free energy (MFE), sequence complementarity and interspecies conservation in the

target genome. The latest version of miRanda software was accessed using online source

website (http://www.microrna.org/). SCBV genome and sugarcane mature miRNA sequences

were processed after defining default setting. These include energy threshold = -15kcal/mol,

score threshold = 130, scaling parameter = 4.00, gap-open penalty= -9.00 and gap-extend

penalty= -4.00.

2.5. RNA22

RNA22 is a friendly user, highly sensitive, web-based (http://cm.jefferson.edu/rna22v1.0/)

novel pattern-recognition algorithm that serves as interactive exploration for predicting target

sites with corresponding hetero-duplexes. Non-seed based interaction, pattern recognition, site

complementarity and folding energy are key parameters of RNA22 algorithm [34]. Final scoring

obviates to use cross-species conservation sequence filter [35]. The minimum number of paired-

up bases in heteroduplex, maximum folding energy for heteroduplex and maximum number of

G: U wobbles allowed in the seed region are user-defined parameters. RNA22 was processed

under default parameters: specificity (61%), sensitivity (63%), and seed size (7) with unpaired

base (1), no limit for G: U wobbles, paired-up bases (12) and folding energy (-12.00 kcal/mol).

2.6. RNAhybrid

RNAhybrid is easy, fast, flexible web-based (http://bibiserv.techfak.uni-

bielefeld.de/rnahybrid) intermolecular hybridization algorithm used to estimate miRNA-mRNA

interaction as well target prediction based on MFE hybridization. The major parameters include

MFE (energy threshold), target-site abundance and site complementarity. A p-value is assigned

to assess RNA-RNA interaction based hybridization sites in the 3′ UTR sequence [36].

RNAhybrid is widely used to estimate the MFE of the consensus miRNA-target pair and the

mode of target inhibition as suggested [37]. MFE threshold (-20Kcal/mol), hit per target (1), no

G: U binding in seed region is fixed default parameters.

2.7. psRNATarget

psRNATarget is a newly developed web server (http://plantgrn.noble.org/psRNATarget/)

to identify target genes of the plant miRNAs based on complementary matching scoring schema.

It was used to discover validated miRNA-mRNA interactions [38]. The plant psRNATarget was

designed to integrate key function for miRNA target prediction using complementarity scoring

and secondary-structure prediction [39]. Target-site accessibility was evaluated by estimating

unpaired energy (UPE) to unfold secondary structure [37]. We consider following unique

parameter for analysis: (Expectation= 8.5, penalty for extending gap = 0.5, penalty opening gap

= 2.0, penalty for G.U pair = 0.5) The HSP size = 19 and seed region = 2–13 nucleotides were

set. Translation inhibition range was set 10-11 nucleotides.

2.8. Mapping of miRNA-Target Interaction

Interaction map was created between sugarcane miRNAs and SCBV ORFs using the Circos algorithm [40].

2.9. RNAfold

RNAfold is a new web-based algorithm applied for the prediction of stable secondary structure of pre-miRNAs based on the MFE [41].

2.10. Free energy (ΔG) Estimation of Duplex Binding

RNAcofold is a novel web-based server applied for the estimation of free energy (ΔG) associated with miRNA-mRNA interactions [42]. Free energy of miRNA: miRNA duplex is considered a key predictor for miRNA targets during hybridization. FASTA sequences of sugarcane consensus miRNAs and their corresponding targets were processed to RNAcofold using default parameters.

2.10.1. Graphical Representation

All the computational data was processed into graphical representations using R language (v3.1.1) [43].

3. Results

3.1. Genome Assembly of SCBV

SCBV is a plant pararetrovirus, was classified in the genus *Badnavirus* of the family *Caulimoviridae*. The genomic ds-DNA molecule of SCBV is comprised of three ORFs, separated by an intergenic region (IR). ORF1 is composed of 557 nucleotides (618-1175 nt) encoding a hypothetical protein (P1) with 185 amino acids (aa). While ORF2 composed of 370

nucleotides (1176-1546 nt) codes for a virion-associated DNA binding protein (P2) with 123 aa.

The precise functional capabilities of these proteins (encoded by ORF1 and ORF2) have not been

explored. A large polyprotein (1977 amino acids) is encoded by ORF3 (1547-7479 nt) to cleave

by a viral aspartic protease. The resulting proteins obtained are named as movement, capsid

protein, aspartyl proteinase, reverse transcriptase and ribonuclease H. The IR is composed of

1022 nucleotides (7479-618) and is located between 3'-ORF3 to 5'-ORF1. The IR works as a

promoter and controls the transcription and regulation of the SCBV genome. Genome

organization of the SCBV with three ORFs is shown in (Figure 2).

188 3.2. miRNA Target Prediction

All the computational algorithms predicted potential hybridization sites, of *Saccharum*

officinarum miRNAs in the SCBV genome. We compared the efficiency of target prediction

algorithms mentioned above----miRanda, RNA22, RNAhybrid and psRNATarget. We designed

top effective approach for evaluating predicted sugarcane miRNA targets at individual, union

and intersection levels. A detailed workflow pipeline of this novel approach is represented

(Figure 1). miRanda (Figure 3A), RNA22 (Figure 3B), RNAhybrid (Figure 3C) and

psRNATarget (Figure 3D) identified putative target sites in the SCBV genome. Union plot

indicates entire genome-binding sites identified by candidate miRNAs using target prediction

tools (Figure 4, Table S2). The Venn diagram represents union and intersection of genomic loci

identified from all the four algorithms used in this study (Figure 5, Table S3 and File S1).

199 3.3. ORF1 encoding Hypothetical Protein

The hypothetical protein of the SCBV genome was encoded by the ORF1 which has

unknown function [2]. miRanda predicted only two target sites of sugarcane miRNAs (sof-

miR156 and sof-miR168) at nucleotide positions (818-837 and 617-638) to target ORF1 (Figure

3A). RNA22 predicted binding sites of miRNAs (sof-miR156 and sof-miR168a) at two different

locus positions 817 and 834 respectively (Figure 3B). RNAhybrid algorithm predicted multiple

potential binding sites of sugarcane miRNAs (sof-miR168 (a, b), ssp-mi827, ssp-miR1128) at

multiple nucleotide positions (612-632, 1170-1192, 1137-1157) respectively (Figure 3C). In

addition, psRNATarget identified potential hybridization sites of sof-miR159(c, e) at locus

positions 1003 and 820 respectively (Figure 3D).

3.4. ORF2 encoding DNA Binding Protein

Nucleic acid (DNA)-binding protein of the SCBV genome was encoded by ORF2 [6, 44].

RNAhybrid and miRanda predicted potential target binding site of ssp-miR166 at locus position

(1449-1470) (Figure 3A, 3C). Suitable candidate miRNAs from sugarcane (ssp-miR444 (a, b,

3p) was observed to target the ORF2 at single loci nucleotide position (1301-1326) determined

by miRanda algorithm (Figure 3A). No sugarcane miRNAs were predicted to target ORF2 gene

by RNA22 tool (Figure 3B). Similarly, RNAhybrid predicted bindings of ssp-miR166 at locus

1450 as shown in (Figure 3C). miRNA prediction results revealed that no candidate miRNA was

identified to have potential genome-binding sites in the ORF2 region as predicted by

psRNATarget as shown in (Figure 3D).

3.5. ORF3 encoding Polyprotein (CP, AP, RT and RNase H)

The poly proteins constitute the largest portion of the SCBV genome was encoded by the

ORF3 [2, 6]. Potential candidate miRNAs from sugarcane were identified by miRanda algorithm

to target the ORF3: (sof-miR159 (a, b, c, d, and e) at common locus 5534, sof-miR167 (a, b) at

locus 2273, sof-miR168b at locus 4588, sof-miR408 (a, b, c, d, and e) at two common locus

4595 and 6695, ssp-miR166 at locus 1986, ssp-miR827 at locus 2816 and ssp-miR444 (a, b and

c-3p) at common locus 6184. Multiple loci interactions were predicted for sof-miR159, sof-

miR408 and ssp-miR444 families at nucleotide positions (5534–5552, 5576–5596), (4595–4615,

6695–6715) and (1679-1701, 3293-3313) of ORF3, respectively (Figure 3A). Potential target

binding sites were determined in the ORF3 of the SCBV genome by RNA22 algorithm: (sof-

miR168a at locus 3263, sof-miR168b at nucleotide positions (1693, 3263), sof-miR396 at locus

2050 and ssp-827 at locus 2796 (Figure 3B). Multiple loci interactions were also identified for

sof-miR159, sof-miR408 and ssp-miR444 families at nucleotide positions (5532, 6536), (5645,

6695) and (5246, 6793) respectively (Figure 3B).

Suitable miRNAs that potentially targeted the ORF3 were hybridized to understand

miRNA-mRNA interaction, by RNAhybrid. These are sof-miR159 (a, b, d and e) at common

locus 5535, sof-miR159c at locus 6518, sof-miR167 (a, b) at locus 2826, sof-miR169 at locus

7362, ssp-miR473 (a, b, c) at common locus 6438, ssp-miR444 (a, b) at locus 6796, ssp-miR444

c-3p at locus 2899 and ssp-miR1432 at locus 7314 (Figure 3C). ORF3 was targeted by candidate

miRNAs: (sof-miR159e at locus 2647, sof-miR396 at locus 5363, ssp-miR166 at locus 1986,

ssp-miR437 (a, c) at 2647 and ssp-miR827 at locus 7337 and ssp-miR444 (a, b and c-3p) at locus

6797) identified by psRNATarget. Multiple loci interactions were observed for sof-miR408 and

ssp-miR444 families at nucleotide positions (1766-1786, 3669-3689, 5683-5702) and (4466–

4486, 6797–6816, 6865-6885, 7079-7099) respectively (Figure 3D).

3.6. Visualization and Analysis of miRNA-Target Interaction Network

Though initially, Circos plotting tool was designed to analyze mutations in comparative

metagenomics and transcriptomic biological data [45]. To study a comprehensive visualization

of host-virus interaction, we created Circos plot to integrate biological data from sugarcane

miRNAs and their predicted SCBV genomic target genes (ORFs) (Figure 5). In order to reduce

visual graphical complexity and permit improved readability, we only used selected sugarcane

miRNAs and their SCBV targets obtained from miRanda analysis. The results suggest that

biological data visualization of candidate miRNAs from sugarcane, with SCBV-encoded ORFs

determines credible information of desirable preferred targets of SCBV ORFs using consensus

miRNAs. We have combined sugarcane miRNA data and their predicted SCBV targets

simultaneously in this manner.

3.7. Prediction of Consensus Secondary Structures

The validation of consensus sugarcane-miRNAs was confirmed by the prediction of their

stable secondary structures using RNAfold algorithm. Precursors of mature sugarcane miRNAs

were manually curated. MFE is the key factor to determine the stable secondary structures of

precursors. All the predicted consensus sugarcane miRNA precursors were observed to possess a

lower MFE value (ranging from −57.70 to −114.70 kcal/mol) (Table 2). The predicted secondary

structures of six precursors of pre-miRNAs are shown (Figure 6) as predicted by intersection of

three consensus algorithms at the same locus. Top stable secondary structure of sof-MIR159e

precursor was predicted with standard features (MFE: 107.50 Kcal/mol, MFEI: 1.06 Kcal/mol).

The predicted secondary structures of 14 consensus sugarcane miRNAs passed the standard

criteria as mentioned. We have determined the salient characteristics of six consensus precursor

miRNAs in this study such as MFE, AMFE, MFEI, length precursor and GC contents. In our

studies, length precursor range from 105-266 nucleotides, MFE (-57.70 to −110.70 kcal/mol),

AMFE (-39.92 to 60.09) and GC content (38–47%) and MFEI ranges from −0.83 to −1.26.

3.8. Predicting Consensus Sugarcane miRNAs for Silencing SCBV Genome

269 Based on predicted targeting miRNAs from sugarcane to silence the SCBV genome,

270 fourteen miRNAs (sof-miR156, sof-miR159c, sof-miR159e, sof-miR168a, sof-miR396, sof-

miR408a, sof-miR408b, sof-miR408c, sof-miR408d, sof-miR408e, ssp-miR827, ssp-miR444a, ssp-miR444b and sof-miR444c-3p) was detected by union of consensus multiple algorithms (miRanda, RNA22, RNAhybrid and psRNATarget) used in this study (Figure 7). Moreover, SCBV genomic components (ORF1, ORF2, ORF3 and LIR) were observed to be targeted by a total of eleven sugarcane miRNAs which were hybridized at unique positions within the ORF1: (sof-miR156 (locus 818), sof-miR168 (a, b) (locus 617) ORF2: ssp-miR166 (locus 1450) and ORF3: sof-miR159c (locus 5534), sof-miR408 (a, b, c, d and e) (locus 6695) and LIR: sof-miR396 (locus 79) according to intersection of two consensus algorithms. Out of 28 sugarcane miRNAs, only six sugarcane miRNA (sof-miR159 (a, b, d and e) at common locus position 5535 and ssp-miR444 (a, b) at locus 6797) were predicted at the common locus by at least three algorithms used (Figure 8 and Table 2). Out of 14 consensus miRNAs, only one miRNA of *S.officinarum* (sof-miR159e at locus 5535) with MFE -26.7 Kcal/mol, was considered as the top effective candidate; deliver more efficient silencing of the SCBV genome. The efficacy of sof-miR159e target against SCBV was validated by suppression of RNAi-mediated viral combat through cleavage of viral mRNA or translational inhibition. Secondary structure of sof-MIR159e precursor has been displayed in (Figure 6). Multiple loci interactions were observed for sof-miR159e at nucleotide positions 5534-5552 (consensus of three algorithms; miRanda, RNA22 and RNAhybrid) and 2647 (psRNATarget) of ORF3.

3.9. Assessment of Free Energy (ΔG) of miRNA-mRNA Interaction

The predicted consensus sugarcane miRNAs were validated by estimating free-energies of miRNA/target duplexes (table 2). The free-energies (ΔG) of six consensus sugarcane miRNAs are estimated: sof-miR159 (a, b, d) (ΔG: -20.10 kcal/mol), sof-miR15e (ΔG: -20.40 kcal/mol), and ssp-miR444 (a, b) (ΔG: -14.50 kcal/mol).

4. Discussion

For Filtering of false-positive results, we study the performance of computational algorithms to validate the miRNA target prediction data. We designed the most effective approach for validation of miRNA target prediction results at individual, union and intersection levels. Computational prediction algorithms offer a rapid method to predict potential host-derived miRNA targets in the virus genome. We evaluated the efficiency of four frequently used computational tools cited in literature—miRanda, RNA22, RNAhybrid and psRNATarget to identify miRNA-target prediction and to analyze miRNA-mRNA interactions as summarized in (Table 1). miRanda is widely used algorithms, includes main aspects of miRNA-target prediction; conservation level and miRNA 3' site [46]. RNA22 adopts a sequence-based pattern recognition mechanism to identify potential targets independently of conservation status and also include 3' UTR, 5'UTR and CDS regions for target prediction [34]. RNA22 algorithm is a novel alternative option for exploring new miRNA: mRNA interactions, because of its unique capabilities but likelihood of high rate of potential false-positive [47]. RNAhybrid is widely used to find the energetically most favorable hybridization binding sites [36, 48]. We calculate the MFE and determine the target inhibition as recommended from Broderson conclusion using RNAhybrid [37]. psRNATarget is a novel plant candidate miRNAs prediction algorithm used for assessment of miRNA-mRNA interaction [38].

Several potential sugarcane miRNA targets and miRNA-mRNA interaction could be consensually predicted by all the algorithms (Figure 7). miRanda and psRNATarget are two powerful plant miRNA prediction algorithms for identifying hybridization sites in the viral genome. Plant miRNAs are responsible to induce the degradation of the target gens using perfect or imperfect complementarity base pairing [49]. The current study demonstrates that

SCBV genome components (ORF1, ORF2 and ORF3) are susceptible to be targeted by a set of

consensus sugarcane miRNAs. In addition, sof-miR159 (a, b, d and e) was found to target the

ORF3 at consensus hybridization site by at least three algorithms (Figure 8). Free energy

assessment is a dynamic feature of miRNA and target binding. Previous study revealed a

significant correlation of free energy between the translational repression and the hybridization

binding of the seed region [50]. The thermodynamic stability of the miRNA-mRNA duplex was

estimated by assessment of free energy to monitor site accessibility for determination of the

secondary structure duplex [27]. In order to validate miRNA-mRNA interaction, free energy of

duplex was assessed as shown in (Table 2). Our prediction results are outstanding to show high

stability of sugarcane-encoded miRNA-SCBV-mRNA duplex at low level of free energy at low

free energy level as shown in the (Table 2 and Figure 6). RNA duplex is considered more stable

due to stronger binding of miRNA to mRNA [51, 52].

We used union and intersection approaches to reduce false-positive prediction. Union

approach relies on combining more than one target prediction tool: true and false targets.

Sensitivity level of predicted targets increases due to decrease of specificity. Intersection

approach is entirely different and depends upon the combining of two or more computational

algorithmic tools which enhances the specificity level of predicted targets due to decrease of

sensitivity [53]. Our target prediction results revealed that both the computational approaches

achieved the best outcome with maximum of performance for predicting and estimating the best

targets as shown in (Figure 7 and 8). Previous studies have also reported the silencing of plant

viruses using host-derived miRNAs applying a set of computational algorithms. Identification

and evaluation of best-fit candidate miRNA targets of different crops plants was concluded

successfully in PVY [54], maize chlorotic mottle virus (MCMV) [55], CLCuKoV-Bu [56],

RYMV [57] and SCBGAV to find miRNA-target interaction [58]. We have designed the same

novel bioinformatics approach for target prediction in the SCBV genome to control the emerging

badnavirus in sugarcane cultivars.

In our previous study, we identified the most potential consensus sugarcane miRNA

(sof-miR396) to target ORF3 of SCBGAV genome using multiple computational algorithms

[58]. The quantity of false-positive miRNA-target interaction estimated by multiple algorithms

depends upon the mode of miRNA-target recognition. MFE is also another one of the important

factors to affect the miRNA-target interaction for result validation [59]. To set a lower MFE

value, give rise high probability of miRNA-target complex formation [60] . In the current study,

for miRanda analysis, a stringent cut-off −15 kcal/mol was set for narrowing down the miRNA

candidates. Similarly to validate host-virus interaction, a MFE cut-off -20 kcal/mol applied for

RNAhybrid analysis [32]. In the previous study, consensus prediction results were sorted based

on a stringent cut-off −20 kcal/mol by intersection of two potential algorithms to explore

possible interaction between *Aedes aegypti* cellular miRNA and arboviruses [61]. Although MFE

has considerable role for development miRNA-mRNA complex but it does not certify that the

interaction will lead to functional changes.

In the current study, we identify six potential miRNA hybridization binding sites that

have exhibited low MFE and free-energy of duplex formation. These predicted miRNAs not only

have potential target SCBV genome at transgenic level but also have stronger probability to

develop miRNA-viral mRNA complex formation.   These miRNAs also have chance to

participate in SCBV replication mechanism: a consensus sugarcane miRNA (sof-miR396) has

binding site within the SCBV large intergenic region (LIR) at locus 79 predicted by miRanda

and RNAhybrid algorithms. In the previous study, we predicted that the sof-miRNA396 is an

effective candidate to target SCBGAV genome [58]. The sof-miR159e was predicted by all the

algorithms. The miR159 was explored to have strong role for silencing of *GAMYB* to enable

normal growth [62]. Phe-MIR159 was involved to regulate gene responsible to secondary

thickening in *Phyllostachys edulis* plant [63]. It is important to assess the function of predicted

potential consensus miRNAs for identification of badnavirus replication to demonstrate SCBV

replication experimentally. A hypothetical model was designed to show sugarcane-derived

miRNAs can inhibit SCBV mRNA and sugarcane genes against SCBV virus as shown in (Figure

9). It facilitates to plant-encoded miRNAs to cleavage of SCBV miRNA.

RNAi screens is novel technology to discover various cellular functions and to identify

host-derived factors of viruses [64]. We selected 28 experimental validated sugarcane miRNAs

which had annotated targets that are part of the SCBV. We have designed amiRNA-based gene

construct of SCBV which contains modified miRNA/miRNA* sequence in duplex of precursor

(sof-MIR159) with artificial sequence to create mature amiRNA as shown in (Figure 10). The

amiRNA-based silencing technology is successfully validated in many crop plants to control

emerging plant viruses [18, 19, 21]. In summary, our computational work for the silencing of the

SCBV genome could offer new approach for the remodel of existing strategy as antiviral agents.

We demonstrated to minimize novel antiviral effects of host-derived miRNAs against SCBV.

5. Conclusions

The SCBV has appeared as a major problem in China. SCBV diminishes the quantitative

yield in all sugarcane cultivars. In the current study, we applied computational tools to predict

and comprehensively analyse the candidate miRNA from sugarcane against the SCBV, prior to

cloning. Among them, sof-miR159e was predicted as top effective candidate that could target the

vital gene (ORF3) of the SCBV genome. Our results conclude an alternative strategy to existing

molecular approaches could be re-purposed to control badnaviral infections. The current findings

provide a proof of novel scheme to construct amiRNA-based genome silencing therapeutics to

combat SCBV.

Supplementary Materials: The following are available (Table S1, S2, S3 and File S1).

Author Contributions: Conceptualization, M.A.A. and S.Z.; Methodology, M.A.A., X.H and

F.A; Software, M.A.A. F.A and X.H; Validation, F.A, L.S and X.F.; Formal Analysis, M.A.A,

X.H.; Investigation, M.A.A., L.S and X.F.; Resources, S.Z. and X.F.; Data Curation, S.Z, L.S.

and X.F; Writing—Original Draft Preparation, M.A.A and F.A; Writing—Review and Editing,

M.A.A, F.A and S.Z.; Visualization, L.S; Supervision, S.Z.; Project Administration, S.Z and

X.F.; Funding Acquisition, S.Z, X.F and X.H. All authors have read and agree to the published

version of the manuscript.

Funding: This work was supported by the Central Public-interest Scientific Institution Basal

Research Fund for Chinese Academy of Tropical Agricultural Sciences (Grant number:

19CXTD-33), National Natural Science Foundation of China (Grant number: 31771865), the

Sugar Crop Research System (Grant ID: CARS-170301) and the Talented Young Scientist

Program of China (Grant ID: Pakistan-18-004).

Acknowledgments: We are highly thankful to our lab colleagues for their assistance in data

analysis. We thank to Dr. Zhiqiang Xia (ITBB) for assistance in constructing Circos plot.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the

design of the study; in the collection, analyses, or interpretation of data, in the writing of the

manuscript, or in the decision to publish the results.

References

1. Bouhida, M., B. Lockhart, and N.E. Olszewski, *An analysis of the complete sequence of a sugarcane bacilliform virus genome infectious to banana and rice.* Journal of General Virology, 1993. **74**(1): p. 15-22.

2. Sun, S.-R., et al., *Molecular characterization of two divergent variants of sugarcane bacilliform viruses infecting sugarcane in China.* European Journal of Plant Pathology, 2016. **145**(2): p. 375-384.

3. King, A.M., et al., *Virus taxonomy: ninth report of the International Committee on Taxonomy of Viruses.* Vol. 9. 2011: Elsevier.

4. Geijskes, R.J., et al., *Sequence analysis of an Australian isolate of sugarcane bacilliform badnavirus.* Archives of virology, 2002. **147**(12): p. 2393-2404.

5. Muller, E., et al., *High molecular variability of sugarcane bacilliform viruses in Guadeloupe implying the existence of at least three new species.* Virus research, 2011. **160**(1-2): p. 414-419.

6. Ahmad, K., et al., *Presence of diverse sugarcane bacilliform viruses infecting sugarcane in China revealed by pairwise sequence comparisons and phylogenetic analysis.* The plant pathology journal, 2019. **35**(1): p. 41.

7. Bousalem, M., E.J. Douzery, and S. Seal, *Taxonomy, molecular phylogeny and evolution of plant reverse transcribing viruses (family Caulimoviridae) inferred from full-length genome and reverse transcriptase sequences.* Archives of virology, 2008. **153**(6): p. 1085-1102.

8. Borah, B.K., et al., *Bacilliform DNA-containing plant viruses in the tropics: commonalities within a genetically diverse group.* Molecular plant pathology, 2013. **14**(8): p. 759-771.

9. Iskra-Caruana, M.-I., et al., *The common evolutionary history of badnaviruses and banana.* Infection, Genetics and Evolution, 2014. **21**: p. 83-89.

10. Autrey, L., et al. *The distribution of sugarcane bacilliform virus in various geographical regions.* in *Proc Int Soc Sugar Cane Technol.* 1995.

11. Lockhart, B., M. Irey, and J. Comstock. *Sugarcane bacilliform virus, sugarcane mild mosaic virus and sugarcane yellow leaf syndrome.* in *Sugarcane germplasm conservation and exchange: report of an international workshop held in Brisbane, Queensland, Australia, 28-30 June 1995.* 1996. Canberra, Australia: Australian Centre for International Agricultural ....

12. Muhammad, T., et al., *RNA interference: a natural immune system of plants to counteract biotic stressors.* Cells, 2019. **8**(1): p. 38.

13. Khvorova, A., A. Reynolds, and S.D. Jayasena, *Functional siRNAs and miRNAs exhibit strand bias.* Cell, 2003. **115**(2): p. 209-216.

14. Wang, W. and G. Galili, *Tuning the Orchestra: miRNAs in plant immunity.* Trends in plant science, 2019. **24**(3): p. 189-191.

15. Gaffar, F.Y. and A. Koch, *Catch me if you can! RNA silencing-based improvement of antiviral plant immunity.* Viruses, 2019. **11**(7): p. 673.

16. Qu, J., J. Ye, and R. Fang, *Artificial microRNA-mediated virus resistance in plants.* Journal of virology, 2007. **81**(12): p. 6690-6699.

17. Rosa, C., et al., *RNA interference mechanisms and applications in plant pathology.* Annual review of phytopathology, 2018. **56**: p. 581-610.

18. Niu, Q.-W., et al., *Expression of artificial microRNAs in transgenic Arabidopsis thaliana confers virus resistance.* Nature biotechnology, 2006. **24**(11): p. 1420-1428.

19. Ali, I., et al., *Artificial microRNA-mediated resistance against the monopartite begomovirus Cotton leaf curl Burewala virus.* Virology journal, 2013. **10**(1): p. 231.

452	20.	Zhang, X., et al., *Expression of artificial microRNAs in tomato confers efficient and stable virus*
453		*resistance in a cell-autonomous manner.* Transgenic research, 2011. **20**(3): p. 569-581.
454	21.	Petchthai, U., C.S. Le Yee, and S.-M. Wong, *Resistance to CymMV and ORSV in artificial*
455		*microRNA transgenic Nicotiana benthamiana plants.* Scientific reports, 2018. **8**(1): p. 1-8.
456	22.	Seal, S., et al., *The prevalence of badnaviruses in West African yams (Dioscorea cayenensis-*
457		*rotundata) and evidence of endogenous pararetrovirus sequences in their genomes.* Virus
458		Research, 2014. **186**: p. 144-154.
459	23.	Harper, G., et al., *Detection of episomal banana streak badnavirus by IC-PCR.* Journal of
460		Virological Methods, 1999. **79**(1): p. 1-8.
461	24.	Sun, S.-R., et al., *Development of quantitative real-time PCR assays for rapid and sensitive*
462		*detection of two badnavirus species in sugarcane.* BioMed research international, 2018. **2018**.
463	25.	Sukal, A.C., et al., *Assessment and optimization of rolling circle amplification protocols for the*
464		*detection and characterization of badnaviruses.* Virology, 2019. **529**: p. 73-80.
465	26.	Bömer, M., et al., *A sequence-independent strategy for amplification and characterisation of*
466		*episomal badnavirus sequences reveals three previously uncharacterised yam badnaviruses.*
467		Viruses, 2016. **8**(7): p. 188.
468	27.	Peterson, S.M., et al., *Common features of microRNA target prediction tools.* Frontiers in
469		genetics, 2014. **5**: p. 23.
470	28.	Kozomara, A., M. Birgaoanu, and S. Griffiths-Jones, *miRBase: from microRNA sequences to*
471		*function.* Nucleic acids research, 2019. **47**(D1): p. D155-D162.
472	29.	Griffiths-Jones, S., et al., *miRBase: microRNA sequences, targets and gene nomenclature.* Nucleic
473		acids research, 2006. **34**(suppl_1): p. D140-D144.
474	30.	Kozomara, A. and S. Griffiths-Jones, *miRBase: annotating high confidence microRNAs using deep*
475		*sequencing data.* Nucleic acids research, 2014. **42**(D1): p. D68-D73.
476	31.	Karuppaiah, R., R. Viswanathan, and V.G. Kumar, *Genetic diversity of Sugarcane bacilliform virus*
477		*isolates infecting Saccharum spp. in India.* Virus genes, 2013. **46**(3): p. 505-516.
478	32.	Enright, A.J., et al., *MicroRNA targets in Drosophila.* Genome biology, 2003. **5**(1): p. R1.
479	33.	John, B., et al., *Human microRNA targets.* PLoS biol, 2004. **2**(11): p. e363.
480	34.	Miranda, K.C., et al., *A pattern-based method for the identification of MicroRNA binding sites*
481		*and their corresponding heteroduplexes.* Cell, 2006. **126**(6): p. 1203-1217.
482	35.	Loher, P. and I. Rigoutsos, *Interactive exploration of RNA22 microRNA target predictions.*
483		Bioinformatics, 2012. **28**(24): p. 3322-3323.
484	36.	Krüger, J. and M. Rehmsmeier, *RNAhybrid: microRNA target prediction easy, fast and flexible.*
485		Nucleic acids research, 2006. **34**(suppl_2): p. W451-W454.
486	37.	Brodersen, P., et al., *Widespread translational inhibition by plant miRNAs and siRNAs.* Science,
487		2008. **320**(5880): p. 1185-1190.
488	38.	Dai, X., Z. Zhuang, and P.X. Zhao, *psRNATarget: a plant small RNA target analysis server (2017*
489		*release).* Nucleic acids research, 2018. **46**(W1): p. W49-W54.
490	39.	Dai, X. and P.X. Zhao, *psRNATarget: a plant small RNA target analysis server.* Nucleic acids
491		research, 2011. **39**(suppl_2): p. W155-W159.
492	40.	Krzywinski, M., et al., *Circos: an information aesthetic for comparative genomics.* Genome
493		research, 2009. **19**(9): p. 1639-1645.
494	41.	Lorenz, R., et al., *ViennaRNA Package 2.0.* Algorithms for molecular biology, 2011. **6**(1): p. 26.
495	42.	Bernhart, S.H., et al., *Partition function and base pairing probabilities of RNA heterodimers.*
496		Algorithms for Molecular Biology, 2006. **1**(1): p. 3.
497	43.	Gandrud, C., *Reproducible research with R and R studio.* 2013: CRC Press.
498	44.	Jacquot, E., et al., *The open reading frame 2 product of cacao swollen shoot badnavirus is a*
499		*nucleic acid-binding protein.* Virology, 1996. **225**(1): p. 191-195.

| 500 | 45. | Naquin, D., et al., *CIRCUS: a package for Circos display of structural genome variations from paired-end and mate-pair sequencing data.* BMC bioinformatics, 2014. **15**(1): p. 1-6. |

45. Naquin, D., et al., *CIRCUS: a package for Circos display of structural genome variations from*
paired-end and mate-pair sequencing data. BMC bioinformatics, 2014. **15**(1): p. 1-6.
46. Betel, D., et al., *Comprehensive modeling of microRNA targets predicts functional non-conserved*
and non-canonical sites. Genome biology, 2010. **11**(8): p. R90.
47. Riffo-Campos, Á.L., I. Riquelme, and P. Brebi-Mieville, *Tools for sequence-based miRNA target*
prediction: what to choose? International journal of molecular sciences, 2016. **17**(12): p. 1987.
48. Min, H. and S. Yoon, *Got target?: computational methods for microRNA target prediction and*
their extension. Experimental & molecular medicine, 2010. **42**(4): p. 233-244.
49. Jones-Rhoades, M.W., D.P. Bartel, and B. Bartel, *MicroRNAs and their regulatory roles in plants.*
Annu. Rev. Plant Biol., 2006. **57**: p. 19-53.
50. Doench, J.G. and P.A. Sharp, *Specificity of microRNA target selection in translational repression.*
Genes & development, 2004. **18**(5): p. 504-511.
51. Lewis, B.P., C.B. Burge, and D.P. Bartel, *Conserved seed pairing, often flanked by adenosines,*
indicates that thousands of human genes are microRNA targets. cell, 2005. **120**(1): p. 15-20.
52. Huang, Y., et al., *A study of miRNAs targets prediction and experimental validation.* Protein &
cell, 2010. **1**(11): p. 979-986.
53. M Witkos, T., E. Koscianska, and W. J Krzyzosiak, *Practical aspects of microRNA target prediction.*
Current molecular medicine, 2011. **11**(2): p. 93-109.
54. Iqbal, M.S., et al., *Prediction of host-derived miRNAs with the potential to target PVY in potato*
plants. Frontiers in Genetics, 2016. **7**: p. 159.
55. Iqbal, M.S., et al., *In silico MCMV silencing concludes potential host-derived miRNAs in maize.*
Frontiers in plant science, 2017. **8**: p. 372.
56. Akhter, Y. and J.A. Khan, *Genome wide identification of cotton (Gossypium hirsutum)-encoded*
microRNA targets against Cotton leaf curl Burewala virus. Gene, 2018. **638**: p. 60-65.
57. Jabbar, B., et al., *Target prediction of candidate miRNAs from Oryza sativa for silencing the*
RYMV genome. Computational biology and chemistry, 2019. **83**: p. 107127.
58. Ashraf, F., et al., *A novel computational approach to the silencing of Sugarcane Bacilliform*
Guadeloupe A Virus determines potential host-derived MicroRNAs in sugarcane (Saccharum
officinarum L.). PeerJ, 2020. **8**: p. e8359.
59. Pinzón, N., et al., *microRNA target prediction programs predict many false positives.* Genome
research, 2017. **27**(2): p. 234-245.
60. Kertesz, M., et al., *The role of site accessibility in microRNA target recognition.* Nature genetics,
2007. **39**(10): p. 1278-1284.
61. Yen, P.-S., et al., *Assessing the potential interactions between cellular miRNA and arboviral*
genomic RNA in the yellow fever mosquito, aedes aegypti. Viruses, 2019. **11**(6): p. 540.
62. Millar, A.A., A. Lohe, and G. Wong, *Biology and Function of miR159 in Plants.* Plants, 2019. **8**(8):
p. 255.
63. Cheng, Z., et al., *Integrated mRNA, MicroRNA Transcriptome and Degradome Analyses Provide*
Insights into Stamen Development in Moso Bamboo. Plant and Cell Physiology, 2020. **61**(1): p.
76-87.
64. Hirsch, A.J., *The use of RNAi-based screens to identify host proteins involved in viral replication.*
Future microbiology, 2010. **5**(2): p. 303-311.

547   Fig1. A flowchart designed for predicting candidate miRNAs from the SCBV genome.

549   Fig 2. Genomic organization of the *Sugarcane Bacilliform Virus* showing the predicted ORFs.

Fig 3. Computational prediction of candidate miRNA targets in the genome of the SCBV. A) miRNA-targets obtained from miRanda. B) RNA22 predicted potential hybridization sites C) Target sites of sugarcane miRNAs as identified by RNAhybrid. D) Prediction results of target sites of sugarcane miRNAs obtained by psRNATarget. The color code is associated with the corresponding ORFs in each graph.

Fig 4. Union plot summarizes miRNA Target candidate prediction represented as a union from

all the algorithms used in this study.

569 Fig 5. Circos plot of the genomic regulatory network interaction predicted to be targeted by the

570 sugarcane-miRNAs. The red, green and blue coloured lines represent SCBV genome

571 components (ORFs). The synergetic counterparts of sugarcane miRNAs and their target genes

572 (ORFs) of the SCBV genome are interconnected with coloured lines.

Fig 6. Six pre-miRNA secondary structures, precursor of sugarcane miRNAs were identified   in

this study by a consensus of three algorithms. Sugarcane mature miRNA name IDs, accession

IDs MFE and MFEI are given as follow: A) sof-MIR159a (MI0001756), - 110.30 kcal/mol, -0.87

B) sof-MIR159b (MI0001757), -110.30 kcal/mol, -0.87 C) sof-MIR159d (MI0001758), -105.80

kcal/mol, -0.83 D) sof-MIR159e (MI0001759), -107.50 kcal/mol, -1.06 E) ssp-MIR444a

(MI0018185), -57.70 kcal/mol, -1.15 F) ssp-MIR444b (MI0018186), -63.70kcal/mol, -1.26.

590   Fig 7. Venn diagram plot of SCBV genome targeted by sugarcane miRNAs. In total, 28 loci are

591   targeted by sugarcane-miRNAs, as predicted from four unique algorithms.

Fig 8.   Intersection plot was created miRNAs, predicted from at least three algorithms: (miRanda, RNA22 and RNAhybrid). The color code has been given with the figure.

606   Fig 9. Schematic model designed for amiRNA-based silencing mechanism. SCBV can activate

607   the production of sugarcane endogenous miRNAs after infection. The sugarcane miRNAs can

608   target SCBV mRNA for degradation.

616 Fig 10. Systematic mechanism of amiRNA-mediated gene silencing in transgenic sugarcane

617 plants. Candidate amiRNA is implanted into sof-MIR159 precursor after miRNA/miRNA duplex

618 replacement. This pre-amiRNA is processed to generate mature amiRNA/amiRNA* duplex.

623   Table 1. Comparison of distinctive parameters used in the common target prediction tools.

Tools	Algorithms	Seed pairing	Target site accessibility	Multiple sites	Translation Inhibition	Availability
miRanda	Local alignment	+	+	+	+	Web server and source code
RNA22	FASTA	_	+	+	_	Only web server
RNAhybrid	Interamolecular hybridization	+	+	+	+	Web server and source code
psRNATarget	Smith-Waterman	_	+	+	+	Only web server
Tapirhybrid	FASTA	+	+	+	_	Web server and source code
Targetfinder	FASTA	+	_	_	_	Only source code
Target-align	Smith-Waterman	-	_	+	_	Web server and source code
Targetscan	Custom made	+	_	+	+	Only source code

625   Table 2. Salient parameters of precursor miRNAs determined to develop secondary structures.

miRNA ID	Length miRNA	Length precursor	MFE (Kcal/mol)	AMFE[1]	MFEI[2]	(G+C) %	ΔG[3] (Kcal/mol)
sof-miR156	20	137	-66.20	-48.32	-0.96	50.00	-14.30
sof-miR159c	21	238	-110.60	-46.47	-0.88	52.38	-19.70
sof-miR159e	21	264	-107.50	-40.71	-1.06	38.09	-20.40
sof-miR168a	21	104	-66.20	-63.65	-1.02	61.90	-18.20
sof-miR396	21	134	-67.40	-50.29	-1.17	42.85	-19.60
sof-miR408a	21	283	-114.70	-40.53	-0.60	66.66	-16.00
sof-miR408b	21	286	-113.20	-39.58	-0.59	66.66	-16.00
sof-miR408c	21	286	-115.80	-40.48	-0.60	66.66	-16.00
sof-miR408d	21	215	-79.00	-36.76	-0.55	66.66	-16.00
sof-miR408e	21	283	-99.00	-34.98	-0.56	61.90	-16.00
ssp-miR827	21	130	-64.00	-49.23	-1.29	38.09	-17.90
ssp-miR444a	21	105	-57.70	-54.95	-1.15	47.62	-14.50
ssp-miR444b	21	106	-63.70	-60.09	-1.26	47.62	-14.50
ssp-miR444c	21	108	-61.80	-57.22	-1.33	42.85	-15.30