An Algorithmic framework for genome-wide identification of Sugarcane (*Saccharum officinarum* L.)-encoded microRNA targets against SCBV

Muhammad Aleem Ashraf[1, 2, *, †], Xiaoyan Feng[1, †], Xiaowen Hu[3, †], Fakiha Ashraf[1], Linbo Shen[1], and Shuzhen Zhang[1, *]

[1] Institute of Tropical Bioscience and Biotechnology, Sugarcane Research Centre of Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China

[2] Department of Biosciences and Technology, Khwaja Fareed University of Engineering and Information Technology, Rahim Yar Khan 64200, Pakistan.

[3] Zhanjiang Experimental Station, Chinese Academy of Tropical Agricultural Sciences, Zhanjiang 524000, China

* Correspondence: aleem.ashraf@kfueit.edu.pk (M.A.A), zhangshuzhen@itbb.org.cn (S.Z)

[†] These authors contributed equally to this work.

Abstract:

Sugarcane Bacilliform Virus (SCBV) is considered an economically the most damaging pathogen for sugarcane production worldwide. Three ORFs are characterized in a single molecule of circular, ds-DNA genome of the SCBV, encoding for hypothetical protein (ORF1), DNA binding protein (ORF2) and Polyprotein (ORF3). The study was aimed to predict and comprehensively evaluate sugarcane miRNAs for the silencing of SCBV genome using *in-silico* algorithms. Computational methods were used for prediction of candidate miRNAs from sugarcane (*S. officinarum* L.) to silence the expression of SCBV genes through translational inhibition by mRNA cleavage. Mature sugarcane miRNAs were retrieved and were assessed to hybridization with the SCBV genome. A total of fourteen potential candidate miRNAs from sugarcane were computed by all the algorithms used for the silencing of SCBV. A consensus of

three algorithms predicts hybridization sites of sof-miR159e at common locus 5534. The miRNA-mRNA interaction was estimated by computing free-energy of miRNA-mRNA duplex using RNAcofold algorithm. Regulatory network of predicted candidate miRNAs of sugarcane with SCBV ORFs, generated using Circos, identify novel targets. Consequently, detecting and discarding inefficient amiRNAs prior to cloning would help suppressed mutants faster. The efficacy of predicted candidate miRNAs was evaluated to test the survival rate of the *in vitro* amiRNA-mediated effective badnaviral silencing and resistance in sugarcane cultivars.

Keywords: miRNAs; *Saccharum officinarum*; *Sugarcane bacilliform virus;* Genome silencing; RNAhybrid; free-energy; psRNATarget; RNAi; computational approach; badnavirus.

1. Introduction

Sugarcane bacilliform viruses (SCBVs) are classified into Badnavirus genus of Caulimoviridae family. These viruses are composed of monopartite, circular, non-enveloped bacilliform with dimensions (30nm ×120-150 nm), double-stranded DNA (ds-DNA)-genome of approximately 7.2-9.2 Kbp in size [1]. The genome of SCBV constitutes three major open reading frames (ORFs) that are located on the 'plus DNA strand' with a single discontinuity [2]. ORF1 encodes a small hypothetical protein which has unknown function. ORF2 encodes a virion-associated DNA-binding protein. ORF3 encodes a largest polyprotein which is represented as (P3.) that was composed of multiple functional sub-units. The polyprotein (P3) is cleaved by a viral aspartic protease to give rise multiple functional small proteins: intracellular movement, capsid, aspartic protease, RT, and ribonuclease [1-6]. H. RT-RNaseH-coding region was considered the most common taxonomic marker for identification of badnaviral genomic components. This coding region is a standard source to compare the sequence diversity of the

badnaviral genomes [7]. The *Badnavirus* genome is replicated using reverse transcription of a more-than-genome length RNA. It works as a standard template to complete the translation process of the badnavirus proteins. It is also used as reverse transcription for replication of the badnavirus genome [8, 9].

First report of SCBV infection was observed in the Cuban sugarcane cultivar B34104 in 1985 [10].These viruses are disseminated worldwide and decrease crop production significantly because of accessibility and exchange of biological materials globally. SCBV is a source of infection for several bioenergy crop sugarcane cultivars, varieties and species. Broad host range of SCB includes diverse and economically important members of *Poaceae* (sugarcane, rice) and *Musaceae* (banana) families. Natural transmission of SCBV is disseminated by sap-feeding mealy bug species by vegetative cutting [11]. SCBV disease symptoms include chlorosis and leaf freckle. Infected sugarcane plants were also monitored to have no symptoms. SCBV-infection in sugarcane plants causes much reduced sucrose content, juice, stalk weigh, purity and gravity in recent years in China [6].

Plant employs multiple defense strategies to control virus invasion, replication and transmission in the cell. RNAi-mediated response of plants against invading viruses is highly significant during infection period [12]. RNAi mechanism inhibits the protein translation at mRNA level using a highly sequence specific strategy [13]. Sugarcane plant species were imparted natural inherited active immunity consisting of small, non-coding microRNAs (miRNAs) to control viral diseases. The miRNA-mediated genome silencing was considered to certify the activity of positive or negative immune-based regulation. It is also considered as key activator of immune defense in plants [14]. These miRNAs belong to a class of 21-23 nucleotides long, endogenous, non-coding RNA molecules that plays a major role to control the gene expression at post-

transactional level. RNA silencing in the form of miRNAs within the host plant, is a source of

natural immunity. It provides resistance to the host plant after infection of foreign genetic

elements including plant viruses [15-17]. Artificial microRNA (amiRNA)-mediated silencing of

invading viruses in plants was first time reported by Niu [18]. This amiRNA-based silencing

strategy was applied effectively in many plants to combat plant viruses: cotton leaf curl Kokhran

virus (CLCuKoV-Bu) [19], Cucumber mosaic virus (CMV)[20], CymMV and ORSV [21].

Serological method, ELISA, immunosorbent electron microscopy (ISEM) [22], IC-PCR [23],

RT-PCR [24], rolling circle amplification (RCA) [25] are the most widely used reliable detection

techniques for SCBV. New strategies are under development for improved diagnoses and cure of

badnaviruses [26].

Prediction and validation of host-virus interactions is the basic step to understand the

critical functional efficacy of miRNAs in the broader range of miRNA-mRNA regulatory

networks controlling biological processes. Each target miRNA possesses potential hybridization

binding sites. In order to validate potential miRNA target experimentally is highly costly and

laborious. An *In Silico* approach to identify miRNA targets concludes potential target binding

sites for laboratory research. Computational strategies determine how miRNAs target to a

desirable mRNA. A large number of computational algorithms are publically available for

miRNA target prediction. It is highly advantageous to get biological access to several

computational tools with different features. The researcher is challenged with an important

choice for selecting suitable tools for prediction [27]. The current study was designed to

implement miRNA prediction algorithms and identify potential targets of sugarcane-derived

miRNAs against SCBV as precedent for creating resistance to sugarcane cultivars using RNAi

technology. We have also screened the potential sugarcane miRNAs for understanding

sugarcane-badnavirus interactions. The novel amiRNA-mediated approach is credible to generate

SCBV-resistance sugarcane plants through genetic engineering.

2. Materials and Methods

2.1. Retrieval of Sugarcane MicroRNAs

The sugarcane mature microRNAs (miRNAs) and stem-loop hairpin precursor sequences

were retrieved from the miRNA biological sequence database, miRBase (v22)

(http://mirbase.org/). miRBase serve as primary public repository and standard online reference

resource for all published miRNA sequences, textual annotation and gene nomenclature [28-30].

In this study, 16 *S.officinarum* (MI0001756-MI0001769) and 19 Saccharum spp. (MI0018180-

MI18197) miRNA sequences were downloaded (Table S1).

2.2. SCBV Genome Retrieval and Annotation

The full-length transcript of the SCBV-BRU genome (Genbank accession number

JN377537.1, size 7884 bp) was isolated from the S.officinarum cultivar, cloned, sequenced and

submitted to the NCBI database [31]. Expected size and abundance of ORFs along nucleotide

distribution of the above mentioned NCBI retrieved SCBV-BRU genome was estimated using

pDRAW32 DNA analysis software (v 1.1.129). The SCBV-BRU genome annotation is designed

which represents ORFs of varying lengths.

2.3. Target Prediction in SCBV Genome

Target prediction is considered a key feature towards identification of credible miRNA-

mRNA interactions hybridization. A present, many target prediction algorithms was designed to

predict and identify best candidate miRNA targets. Each tool uses a specific criteria and

stringency for miRNA target prediction. We used four target prediction algorithms cited in

literature — miRanda, RNA22, RNAhybrid and psRNATarget to find the most relevant

sugarcane miRNAs for silencing of SCBV genome (Table 1). We designed the most effective

computational approach to analyze miRNA targets at three different prediction levels: individual,

union and intersection. A detailed workflow pipeline was present in (Fig 1).

Fig 1. A flowchart embodies the methodology of miRNA prediction from the SCBV

genome.   A flowchart designed for predicting candidate miRNAs from the SCBV genome

pipeline. Biological data composed of sugarcane miRNAs retrieved from the miRBase database

and SCBV genome from NCBI genbank database. An algorithmic framework consists of three

kinds of tools used for identification of sugarcane-encoded miRNA targets, prediction of

secondary structures and visualization of miRNA-target interaction. R language was used to

make plots and select data using in-house scripts/codes.

Table1. Comparison of distinctive parameters used in the common target prediction tools.

Tools	Algorithms	Seed pairing	Target site accessibility	Multiple sites	Translation Inhibition	Availability
miRanda	Local alignment	+	+	+	+	Web server and source code
RNA22	FASTA	_	+	+	_	Only web server
RNAhybrid	Interamolecular hybridization	+	+	+	+	Web server and source code
psRNATarget	Smith-Waterman	_	+	+	+	Only web server
Tapirhybrid	FASTA	+	+	+	_	Web server and source code
Targetfinder	FASTA	+	_	_	_	Only source code
Target-align	Smith-Waterman	_	_	+	_	Web server and source code
Targetscan	Custom made	+	_	+	+	Only source code

127   '+' Represent feature used, '-'indicates that these features were not used.

2.4. miRanda

miRanda is considered more frequently used standard miRNA-target predictor scanning algorithm. It was implemented first time in 2003 [32] and updated into a web-based tool for miRNA analysis [33]. It considers following algorithmic features; seed-based interaction, minimum free energy (MFE), sequence complementarity and interspecies conservation in the target genome. The latest version of miRanda software was accessed using online source website (http://www.microrna.org/). SCBV genome and sugarcane mature miRNA sequences were processed after defining default setting. These include energy threshold = -15kcal/mol, score threshold = 130, scaling parameter = 4.00, gap-open penalty= -9.00 and gap-extend penalty= -4.00.

2.5. RNA22

RNA22 is a friendly user, highly sensitive, web-based (http://cm.jefferson.edu/rna22v1.0/) novel pattern-recognition algorithm that serves as interactive exploration for predicting target sites with corresponding hetero-duplexes. Non-seed based interaction, pattern recognition, site complementarity and folding energy are key parameters of RNA22 algorithm [34]. Final scoring obviates to use cross-species conservation sequence filter [35]. The minimum number of paired-up bases in heteroduplex, maximum folding energy for heteroduplex and maximum number of G: U wobbles allowed in the seed region are user-defined parameters. RNA22 was processed under default parameters: specificity (61%), sensitivity (63%), and seed size (7) with unpaired base (1), no limit for G: U wobbles, paired-up bases (12) and folding energy (-12.00 kcal/mol).

2.6. RNAhybrid

RNAhybrid is easy, fast, flexible web-based (http://bibiserv.techfak.uni-bielefeld.de/rnahybrid) intermolecular hybridization algorithm used to estimate miRNA-mRNA interaction as well target prediction based on MFE hybridization. The major parameters include MFE (energy threshold), target-site abundance and site complementarity. A p-value is assigned to assess RNA-RNA interaction based hybridization sites in the 3′ UTR sequence [36]. RNAhybrid is widely used to estimate the MFE of the consensus miRNA-target pair and the mode of target inhibition as suggested [37]. MFE threshold (-20Kcal/mol), hit per target (1), no G: U binding in seed region is fixed default parameters.

2.7. psRNATarget

psRNATarget is a newly developed web server (http://plantgrn.noble.org/psRNATarget/) to identify target genes of the plant miRNAs based on complementary matching scoring schema. It was used to discover validated miRNA-mRNA interactions [38]. The plant psRNATarget was designed to integrate key function for miRNA target prediction using complementarity scoring and secondary-structure prediction [39]. Target-site accessibility was evaluated by estimating unpaired energy (UPE) to unfold secondary structure [37]. We consider following unique parameter for analysis: (Expectation= 8.5, penalty for extending gap = 0.5, penalty opening gap = 2.0, penalty for G.U pair = 0.5) The HSP size = 19 and seed region = 2–13 nucleotides were set. Translation inhibition range was set 10-11 nucleotides.

2.8. Mapping of miRNA-Target Interaction

Interaction map was created between sugarcane miRNAs and SCBV ORFs using the Circos algorithm [40].

2.9. RNAfold

RNAfold is a new web-based algorithm applied for the prediction of stable secondary structure of pre-miRNAs based on the MFE [41].

2.10. Free energy (ΔG) Estimation of Duplex Binding

RNAcofold is a novel web-based server applied for the estimation of free energy (ΔG) associated with miRNA-mRNA interactions [42]. Free energy of miRNA: miRNA duplex is considered a key predictor for miRNA targets during hybridization. FASTA sequences of sugarcane consensus miRNAs and their corresponding targets were processed to RNAcofold using default parameters.

2.10.1. Graphical Representation

All the computational data was processed into graphical representations using R language (v3.1.1) [43].

3. Results

3.1. Genome Assembly of SCBV

SCBV is a plant pararetrovirus, was classified in the genus *Badnavirus* of the family *Caulimoviridae*. The genomic ds-DNA molecule of SCBV is comprised of three ORFs, separated by an intergenic region (IR). ORF1 is composed of 557 nucleotides (618-1175 nt) encoding a hypothetical protein (P1) with 185 amino acids (aa). While ORF2 composed of 370 nucleotides (1176-1546 nt) codes for a virion-associated DNA binding protein (P2) with 123 aa. The precise functional capabilities of these proteins (encoded by ORF1 and ORF2) have not been explored. A large polyprotein (1977 amino acids) is encoded by ORF3 (1547-7479 nt) to cleave

by a viral aspartic protease. The resulting proteins obtained are named as movement, capsid

protein, aspartyl proteinase, reverse transcriptase and ribonuclease H. The IR is composed of

1022 nucleotides (7479-618) and is located between 3'-ORF3 to 5'-ORF1. The IR works as a

promoter and controls the transcription and regulation of the SCBV genome. Genome

organization of the SCBV with three ORFs is shown in (Fig 2).

Fig 2. Genomic organization of the Sugarcane Bacilliform Virus. Genome organization of the

Sugarcane Bacilliform Virus showing the predicted ORFs with arrows was composed dsDNA of

7884 bp in size.

3.2. ORF1 encoding Hypothetical Protein

The hypothetical protein of the SCBV genome was encoded by the ORF1 which has

unknown function [2]. miRanda predicted only two target sites of sugarcane miRNAs (sof-

miR156 and sof-miR168) at nucleotide positions (818-837 and 617-638) to target ORF1 (Fig

3A). RNA22 predicted binding sites of miRNAs (sof-miR156 and sof-miR168a) at two different

locus positions 817 and 834 respectively (Fig 3B). RNAhybrid algorithm predicted multiple

potential binding sites of sugarcane miRNAs (sof-miR168 (a, b), ssp-mi827, ssp-miR1128) at

multiple nucleotide positions (612-632, 1170-1192, 1137-1157) respectively (Fig 3C). In

addition, psRNATarget identified potential hybridization sites of sof-miR159(c, e) at locus

positions 1003 and 820 respectively (Fig 3D).

Fig 3. Target prediction of sugarcane miRNAs in the SCBV genome. Computational

prediction of candidate miRNA targets in the genome of the SCBV. A) miRNA-targets obtained

from miRanda. B) RNA22 predicted potential hybridization sites C) Target sites of sugarcane

miRNAs as identified by RNAhybrid. D) Prediction results of target sites of sugarcane miRNAs

obtained by psRNATarget.

3.3. ORF2 encoding DNA Binding Protein

Nucleic acid (DNA)-binding protein of the SCBV genome was encoded by ORF2 [6, 44].

RNAhybrid and miRanda predicted potential target binding site of ssp-miR166 at locus position

(1449-1470) (Fig 3A and 3C). Suitable candidate miRNAs from sugarcane (ssp-miR444 (a, b,

3p) was observed to target the ORF2 at single loci nucleotide position (1301-1326) determined

by miRanda algorithm (Fig 3A). No sugarcane miRNAs were predicted to target ORF2 gene by

RNA22 tool (Fig. 3B). Similarly, RNAhybrid predicted bindings of ssp-miR166 at locus 1450 as

shown in (Fig 3C). miRNA prediction results revealed that no candidate miRNA was identified

to have potential genome-binding sites in the ORF2 region as predicted by psRNATarget as

shown in (Fig 3D).

3.4. ORF3 encoding Polyprotein (CP, AP, RT and RNase H)

The poly proteins constitute the largest portion of the SCBV genome was encoded by the

ORF3 [2, 6]. Potential candidate miRNAs from sugarcane were identified by miRanda algorithm

to target the ORF3: (sof-miR159 (a, b, c, d, and e) at common locus 5534, sof-miR167 (a, b) at

locus 2273, sof-miR168b at locus 4588, sof-miR408 (a, b, c, d, and e) at two common locus

4595 and 6695, ssp-miR166 at locus 1986, ssp-miR827 at locus 2816 and ssp-miR444 (a, b and

c-3p) at common locus 6184. Multiple loci interactions were predicted for sof-miR159, sof-

miR408 and ssp-miR444 families at nucleotide positions (5534–5552, 5576–5596), (4595–4615,

6695–6715) and (1679-1701, 3293-3313) of ORF3, respectively (Fig 3A).

Potential target binding sites were determined in the ORF3 of the SCBV genome by RNA22

algorithm: (sof-miR168a at locus 3263, sof-miR168b at nucleotide positions (1693, 3263), sof-

miR396 at locus 2050 and ssp-827 at locus 2796 (Fig 3B). Multiple loci interactions were also

identified for sof-miR159, sof-miR408 and ssp-miR444 families at nucleotide positions (5532,

6536), (5645, 6695) and (5246, 6793) respectively (Fig 3B).

Suitable miRNAs that potentially targeted the ORF3 were hybridized to understand miRNA-

mRNA interaction, by RNAhybrid. These are sof-miR159 (a, b, d and e) at common locus 5535,

sof-miR159c at locus 6518, sof-miR167 (a, b) at locus 2826, sof-miR169 at locus 7362, ssp-

miR473 (a, b, c) at common locus 6438, ssp-miR444 (a, b) at locus 6796, ssp-miR444 c-3p at

locus 2899 and ssp-miR1432 at locus 7314 (Fig 3C). ORF3 was targeted by candidate miRNAs:

(sof-miR159e at locus 2647, sof-miR396 at locus 5363, ssp-miR166 at locus 1986, ssp-miR437

(a, c) at 2647 and ssp-miR827 at locus 7337 and ssp-miR444 (a, b and c-3p) at locus 6797)

identified by psRNATarget. Multiple loci interactions were observed for sof-miR408 and ssp-

miR444 families at nucleotide positions (1766-1786, 3669-3689, 5683-5702) and (4466–4486,

6797–6816, 6865-6885, 7079-7099) respectively (Fig 3D). Union plot indicates entire genome-

binding sites identified by candidate miRNAs using target prediction tools (Fig 4, Table S2).

Fig 4.   Union plot represents all the predicted sugarcane miRNA Targets in the SCBV

genome. Union plot summarizes miRNA Target candidate prediction represented as a union

from all the algorithms used in this study.

3.5. Visualization and Analysis of miRNA-Target Interaction Network

253   Though initially, Circos plotting tool was designed to analyse mutations in comparative

254   metagenomics and transcriptomic biological data [45]. To study a comprehensive visualization

of host-virus interaction, we created Circos plot to integrate biological data from sugarcane miRNAs and their predicted SCBV genomic target genes (ORFs) (Fig 5). In order to reduce visual graphical complexity and permit improved readability, we only used selected sugarcane miRNAs and their SCBV targets obtained from miRanda analysis. The results suggest that biological data visualization of candidate miRNAs from sugarcane, with SCBV-encoded ORFs determines credible information of desirable preferred targets of SCBV ORFs using consensus miRNAs. We have combined sugarcane miRNA data and their predicted SCBV targets simultaneously in this manner.

Fig 5. Circos plot representing miRNA-target interaction. Circos plot of the genomic regulatory network interaction predicted to be targeted by the sugarcane-miRNAs. The red, green and blue coloured lines represent SCBV genome components (ORFs). The synergetic counterparts of sugarcane miRNAs and their target genes (ORFs) of the SCBV genome are interconnected with coloured lines.

3.6. Predicting Common Sugarcane miRNAs

Based on predicted targeting miRNAs from sugarcane to silence the SCBV genome, fourteen miRNAs (sof-miR156, sof-miR159c, sof-miR159e, sof-miR168a, sof-miR396, sof-miR408a, sof-miR408b, sof-miR408c, sof-miR408d, sof-miR408e, ssp-miR827, ssp-miR444a, ssp-miR444b and sof-miR444c-3p) was detected by union of consensus multiple algorithms (miRanda, RNA22, RNAhybrid and psRNATarget) used in this study (Fig 6). Moreover, SCBV genomic components (ORF1, ORF2, ORF3 and LIR) were observed to be targeted by a total of eleven sugarcane miRNAs which were hybridized at unique positions within the ORF1: (sof-miR156 (locus 818), sof-miR168 (a, b) (locus 617) ORF2: ssp-miR166 (locus 1450) and ORF3:

sof-miR159c (locus 5534), sof-miR408 (a, b, c, d and e) (locus 6695) and LIR: sof-miR396

(locus 79) according to intersection of two consensus algorithms (Table 2, Table S3).

Fig 6. Venn diagram plot of SCBV genome targeted by sugarcane miRNAs. Venn diagram

plot of SCBV genome targeted by sugarcane miRNAs. In total, 28 loci are targeted by

sugarcane-miRNAs, as predicted from four unique algorithms.

Table2: Sugarcane miRNAs and their target positions in the SCBV identified by algorithms.

Sugarcane miRNAs	Position miRanda	Position RNA22	Position RNAhybrid	Position psRNATarget	MFE[*] miRanda	MFE[**] RNA22	MFE RNAhybrid	Expectation psRNATarget
sof-miR156	818	817	7608	7609	-17.23	-12.7	-23.9	8
sof-miR159a	**5534**	**5532**	**5535**		**-21.45**	**-19.9**	**-26.7**	
sof-miR159a(1)	5576	6536			-17.54	-12.5		
sof-miR159b	**5534**	**5532**	**5535**		**-21.45**	**-19.9**	**-26.7**	
sof-miR159b(1)	5576	6536			-17.54	-12.5		
sof-miR159c	5534	5532	6518	1003	-20.02		-28	6
sof-miR159c(1)		6533				-12.1		
sof-miR159d	**5534**	**5532**	**5535**		**-21.45**	**-19.9**	**-26.7**	
sof-miR159d(1)	5576	6536			-17.54	-12.5		
sof-miR159e	**5534**	**5532**	**5535**		**-21.45**	**-19.9**	**-26.7**	
sof-miR159e(1)	3633	6536			-16	-12.1		
sof-miR167a	2273		2826		-15.24		-27	
sof-miR167b	2273		2826		-15.24		-27	
sof-miR168a	617	834	612	4046	-19.53	-12.8	-29.9	8.5
sof-miR168a(1)		3263				-13.6		
sof-miR168b	617	1693	612		-19	-26.7	-28.7	
sof-miR168b(1)	4588	4907			-15.49	-12.8		
sof-miR396	79	2050	79	5563	-20.44	-13.3	-25.1	8.25
sof-miR408a	6695	6695	174	3669	-19.19	-16.1	-27.3	8
sof-miR408a(1)	4595	5645		1766	-21.35	-13.6		8
sof-miR408b	6695	6695	174	3669	-19.19	-16.1	-27.3	8
sof-miR408b(1)	4595	5645		1766	-21.35	-13.6		8
sof-miR408c	6695	6695	174	3669	-19.19	-16.1	-27.3	8
sof-miR408c(1)	4595	5645		1766	-21.35	-13.6		8
sof-miR408d	6695	6695	174	3669	-19.19	-16.1	-27.3	8
sof-miR408d(1)	4595	5645		1766	-21.35	-13.6		8
sof-miR408e	6695	6695	174	5683	-19.19	-16.1	-29.9	7.5
sof-miR408e (1)	4595	5645		1766	-19.53	-13.6		8
sof-miR408e (2)	242				-17.01			
ssp-miR166	1449		1450	7750	-27.85		-28.7	6
ssp-miR166(1)	1986			1986	-20.95			7
ssp-miR169	7748		7362		-17.38		-27.7	
ssp-miR437a			6438	2646			-20.1	7
ssp-miR437b			6438				-20.1	
ssp-miR437c			6437	2647			-21.8	7.5
ssp-miR437c(1)				2974				6
ssp-miR528			7619				-26.2	
ssp-miR827	2816	2796	1170	7337	-19.73	-13.6	-23.7	7.5
ssp-miR444a	6184	**6793**	**6796**	**6797**	-18.04	**-13.2**	**-26.9**	**6**
ssp-miR444a(1)	3293	5246		6865	-15.23	-15.6		6.5
ssp-miR444a(2)	1301				-17.44			

ssp-miR444b	6184	**6793**	**6796**	**6797**	-18.04	**-13.2**	**-26.9**	**6**
ssp-miR444b(1)	3293	5246		7079	-15.23	-15.6		6
ssp-miR444b(2)	1301			4466	-17.44			7.5
ssp-miR444b(3)	1676				-16.25			
ssp-miR444c-3p	6184	5246	2899	6797	-18.95	-17.1	-27.9	6
ssp-miR444c-3p(1)	328			6865		-16.8		7
ssp-miR444c-3p(2)	1301					-18.59		
ssp-miR444c-3p(3)	1680					-16.96		
ssp-miR1128			1137				-23	
ssp-miR1432			7314				-23.6	

*MFE: Minimum free energy measured in /Kcal/mol where [*]MFE represents minimum folding energy measured in Kcal/mol.

3.7. Predicting Consensus Sugarcane miRNAs for Silencing SCBV Genome

Out of 28 sugarcane miRNAs, only six sugarcane miRNA (sof-miR159 (a, b, d and e) at common locus position 5535 and ssp-miR444 (a, b) at locus 6797) were predicted at the common locus by at least three algorithms used (Fig 7, Table 2 and SF1). Out of 14 consensus miRNAs, only one miRNA of *S.officinarum* (sof-miR159e at locus 5535) with MFE -26.7 Kcal/mol, was considered as the top effective candidate; deliver more efficient silencing of the SCBV genome. The efficacy of sof-miR159e target against SCBV was validated by suppression of RNAi-mediated viral combat through cleavage of viral mRNA or translational inhibition. Multiple loci interactions were observed for sof-miR159e at nucleotide positions 5534-5552 (consensus of three algorithms; miRanda, RNA22 and RNAhybrid) and 2647 (psRNATarget) of ORF3.

Fig 7. Intersection plot of sugarcane miRNAs predicted from at least three algorithms. Intersection plot was created miRNAs, predicted from at least three algorithms: (miRanda, RNA22 and RNAhybrid). The color code has been given with the figure.

3.8. Prediction of Consensus Secondary Structures

The validation of consensus sugarcane-miRNAs was confirmed by the prediction of their stable secondary structures using RNAfold algorithm. Precursors of mature sugarcane miRNAs were manually curated. MFE is the key factor to determine the stable secondary structures of

precursors. All the predicted consensus sugarcane miRNA precursors were observed to possess a

lower MFE value (ranging from −57.70 to −114.70 kcal/mol) (Table 3).

The predicted secondary structures of six precursors of pre-miRNAs are shown (Fig 8) as

predicted by intersection of three consensus algorithms at the same locus. Top stable secondary

structure of sof-MIR159e precursor was predicted with standard features (MFE: 107.50

Kcal/mol, MFEI: 1.06 Kcal/mol). The predicted secondary structures of 14 consensus sugarcane

miRNAs passed the standard criteria as mentioned. We have determined the salient

characteristics of six consensus precursor miRNAs in this study such as MFE, AMFE, MFEI,

length precursor and GC contents.   In our studies, length precursor range from 105-266

nucleotides, MFE (-57.70 to −110.70 kcal/mol), AMFE (-39.92 to 60.09) and GC content (38–

47%) and MFEI ranges from −0.83 to −1.26.

Fig 8. Prediction of secondary structure of stem-loop sequences of sugarcane miRNAs. Six

pre-miRNA secondary structures, precursor of sugarcane miRNAs were identified   in this study

by a consensus of three algorithms. Sugarcane mature miRNA name IDs, accession IDs MFE

and MFEI are given as follow: A) sof-MIR159a (MI0001756), - 110.30 kcal/mol, -0.87 B) sof-

MIR159b (MI0001757), -110.30 kcal/mol, -0.87   C) sof-MIR159d (MI0001758), -105.80

kcal/mol, -0.83 D) sof-MIR159e (MI0001759), -107.50 kcal/mol, -1.06 E) ssp-MIR444a

(MI0018185), -57.70 kcal/mol, -1.15 F) ssp-MIR444b (MI0018186), -63.70kcal/mol, -1.26.

3.9. Assessment of Free Energy (ΔG) of miRNA-mRNA Interaction

The predicted consensus sugarcane miRNAs were validated by estimating free-energies

of miRNA/target duplexes (Table 3). The free-energies (ΔG) of six consensus sugarcane

miRNAs are estimated: sof-miR159 (a, b, d) (ΔG: -20.10 kcal/mol), sof-miR15e (ΔG: -20.40

kcal/mol), and ssp-miR444 (a, b) (ΔG: -14.50 kcal/mol).

Table 3. Salient parameters of precursor miRNAs was determined and estimation of free energy.

miRNA ID	Length miRNA	Length precursor	MFE[1] (Kcal/mol)	AMFE[2]	MFEI[3]	(G+C) %	ΔG[4] (Kcal/mol)
sof-miR156	20	137	-66.20	-48.32	-0.96	50.00	-14.30
sof-miR159c	21	238	-110.60	-46.47	-0.88	52.38	-19.70
sof-miR159e	21	264	-107.50	-40.71	-1.06	38.09	-20.40
sof-miR168a	21	104	-66.20	-63.65	-1.02	61.90	-18.20
sof-miR396	21	134	-67.40	-50.29	-1.17	42.85	-19.60
sof-miR408a	21	283	-114.70	-40.53	-0.60	66.66	-16.00
sof-miR408b	21	286	-113.20	-39.58	-0.59	66.66	-16.00
sof-miR408c	21	286	-115.80	-40.48	-0.60	66.66	-16.00
sof-miR408d	21	215	-79.00	-36.76	-0.55	66.66	-16.00
sof-miR408e	21	283	-99.00	-34.98	-0.56	61.90	-16.00
ssp-miR827	21	130	-64.00	-49.23	-1.29	38.09	-17.90
ssp-miR444a	21	105	-57.70	-54.95	-1.15	47.62	-14.50
ssp-miR444b	21	106	-63.70	-60.09	-1.26	47.62	-14.50
ssp-miR444c	21	108	-61.80	-57.22	-1.33	42.85	-15.30

326　[1]MFE is minimum free energy. [2]AMFE represents adjusted minimum free energy. [3]MFEI defines as minimum free
327　energy index. [4]ΔG represents minimum free energy of duplex formation.

4. Discussion

For Filtering of false-positive results, we study the performance of computational

algorithms to validate the miRNA target prediction data. We designed the most effective

approach for validation of miRNA target prediction results at individual, union and intersection

levels. Computational prediction algorithms offer a rapid method to predict potential host-

derived miRNA targets in the virus genome. We evaluated the efficiency of four frequently used

computational tools cited in literature—miRanda, RNA22, RNAhybrid and psRNATarget to

identify miRNA-target prediction and to analyze miRNA-mRNA interactions as summarized in

(Table1). miRanda is widely used algorithms, includes main aspects of miRNA-target prediction;

conservation level and miRNA 3' site [46]. RNA22 adopts a sequence-based pattern recognition

mechanism to identify potential targets independently of conservation status and also include 3'

UTR, 5'UTR and CDS regions for target prediction [34]. RNA22 algorithm is a novel alternative

option for exploring new miRNA: mRNA interactions, because of its unique capabilities but

likelihood of high rate of potential false-positive [47]. RNAhybrid is widely used to find the

energetically most favorable hybridization binding sites [36, 48]. We calculate the MFE and

determine the target inhibition as recommended from Broderson conclusion using RNAhybrid

[37]. psRNATarget is a novel plant candidate miRNAs prediction algorithm used for assessment

of miRNA-mRNA interaction [38].

Several potential sugarcane miRNA targets and miRNA-mRNA interaction could be

consensually predicted by all the algorithms (Fig 7). miRanda and psRNATarget are two

powerful plant miRNA prediction algorithms for identifying hybridization sites in the viral

genome. Plant miRNAs are responsible to induce the degradation of the target gens using

perfect or imperfect complementarity base pairing [49]. The current study demonstrates that

SCBV genome components (ORF1, ORF2 and ORF3) are susceptible to be targeted by a set of

consensus sugarcane miRNAs. In addition, sof-miR159 (a, b, d and e) was found to target the

ORF3 at consensus hybridization site by at least three algorithms (Fig 8). Free energy assessment

is a dynamic feature of miRNA and target binding. Previous study revealed a significant

correlation of free energy between the translational repression and the hybridization binding of

the seed region [50]. The thermodynamic stability of the miRNA-mRNA duplex was estimated

by assessment of free energy to monitor site accessibility for determination of the secondary

structure duplex [27]. In order to validate miRNA-mRNA interaction, free energy of duplex was

assessed as shown in (Table 2). Our prediction results are outstanding to show high stability of

sugarcane-encoded miRNA-SCBV-mRNA duplex at low level of free energy at low free energy

level as shown in the (Table 3 and Fig 8). RNA duplex is considered more stable due to stronger

binding of miRNA to mRNA [51, 52].

We used union and intersection approaches to reduce false-positive prediction. Union

approach relies on combining more than one target prediction tool: true and false targets.

Sensitivity level of predicted targets increases due to decrease of specificity. Intersection

approach is entirely different and depends upon the combining of two or more computational

algorithmic tools which enhances the specificity level of predicted targets due to decrease of

sensitivity [53]. Our target prediction results revealed that both the computational approaches

achieved the best outcome with maximum of performance for predicting and estimating the best

targets as shown in (Fig. 6 and 7). Previous studies have also reported the silencing of plant

viruses using host-derived miRNAs applying a set of computational algorithms. Identification

and evaluation of best-fit candidate miRNA targets of different crops plants was concluded

successfully in PVY [54], maize chlorotic mottle virus (MCMV) [55], CLCuKoV-Bu [56],

RYMV [57] and SCBGAV to find miRNA-target interaction [58]. We have designed the same

novel bioinformatics approach for target prediction in the SCBV genome to control the emerging

badnavirus in sugarcane cultivars.

In our previous study, we identified the most potential consensus sugarcane miRNA

(sof-miR396) to target ORF3 of SCBGAV genome using multiple computational algorithms

[58]. The quantity of false-positive miRNA-target interaction estimated by multiple algorithms

depends upon the mode of miRNA-target recognition. MFE is also another one of the important

factors to affect the miRNA-target interaction for result validation [59]. To set a lower MFE

value, give rise high probability of miRNA-target complex formation [60] . In the current study,

for miRanda analysis, a stringent cut-off -15 kcal/mol was set for narrowing down the miRNA

candidates. Similarly to validate host-virus interaction, a MFE cut-off -20 kcal/mol applied for

RNAhybrid analysis [32].

In the previous study, consensus prediction results were sorted based on a stringent cut-off −20

kcal/mol by intersection of two potential algorithms to explore possible interaction between

Aedes aegypti cellular miRNA and arboviruses [61]. Although MFE has considerable role for

development miRNA-mRNA complex but it does not certify that the interaction will lead to

functional changes. In the current study, we identify six potential miRNA hybridization binding

sites that have exhibited low MFE and free-energy of duplex formation. These predicted

miRNAs not only have potential target SCBV genome at transgenic level but also have stronger

probability to develop miRNA-viral mRNA complex formation. These miRNAs also have

chance to participate in SCBV replication mechanism: a consensus sugarcane miRNA (sof-

miR396) has binding site within the SCBV large intergenic region (LIR) at locus 79 predicted by

miRanda and RNAhybrid algorithms. In the previous study, we predicted that the sof-

miRNA396 is an effective candidate to target SCBGAV genome [58]. The sof-miR159e was

predicted by all the algorithms. The miR159 was explored to have strong role for silencing of

GAMYB to enable normal growth [62]. Phe-MIR159 was involved to regulate gene responsible

to secondary thickening in *Phyllostachys edulis* plant [63]. It is important to assess the function

of predicted potential consensus miRNAs for identification of badnavirus replication to

demonstrate SCBV replication experimentally. A hypothetical model was designed to show

sugarcane-derived miRNAs can inhibit SCBV mRNA and sugarcane genes against SCBV virus

as shown in (Fig 9). It facilitates to plant-encoded miRNAs to cleavage of SCBV miRNA.

RNAi screens is novel technology to discover various cellular functions and to identify

host-derived factors of viruses [64]. We selected 28 experimental validated sugarcane miRNAs

which had annotated targets that are part of the SCBV. We have designed amiRNA-based gene construct of SCBV which contains modified miRNA/miRNA* sequence in duplex of precursor (sof-MIR159) with artificial sequence to create mature amiRNA as shown in (Fig 10). The amiRNA-based silencing technology is successfully validated in many crop plants to control emerging plant viruses [18, 19, 21]. In summary, our computational work for the silencing of the SCBV genome could offer new approach for the remodel of existing strategy as antiviral agents. We demonstrated to minimize novel antiviral effects of host-derived miRNAs against SCBV.

Fig 9. Schematic model designed for amiRNA-based silencing mechanism. amiRNA based gene silencing mechanism was designed in schematic manner. SCBV can activate the production of sugarcane endogenous miRNAs after infection. The sugarcane miRNAs can target SCBV mRNA for degradation.

Fig 10. Systematic mechanism of amiRNA-mediated gene silencing in transgenic sugarcane. Systematic mechanism was designed to impart amiRNA-mediated gene silencing in transgenic sugarcane plants. Candidate amiRNA is implanted into sof-MIR159 precursor after miRNA/miRNA duplex replacement. This pre-amiRNA is processed to generate mature amiRNA/amiRNA* duplex.

5. Conclusions

The SCBV has appeared as a major problem in China. SCBV diminishes the quantitative yield in all sugarcane cultivars. In the current study, we applied computational tools to predict and comprehensively analyse the candidate miRNA from sugarcane against the SCBV, prior to cloning. Among them, sof-miR159e was predicted as top effective candidate that could target the vital gene (ORF3) of the SCBV genome. Our results conclude an alternative strategy to existing

molecular approaches could be re-purposed to control badnaviral infections. The current findings

provide a proof of novel scheme to construct amiRNA-based genome silencing therapeutics to

combat SCBV.

Supplementary Materials: The following are available (Table S1, S2, S3 and File S1).

Funding: This work was supported by the Central Public-interest Scientific Institution Basal

Research Fund for Chinese Academy of Tropical Agricultural Sciences (Grant number:

19CXTD-33), National Natural Science Foundation of China (Grant number: 31771865), the

Sugar Crop Research System (Grant ID: CARS-170301) and the Talented Young Scientist

Program of China (Grant ID: Pakistan-18-004). The funders had no role in the design of the

study; in the collection, analyses, or interpretation of data, in the writing of the manuscript, or in

the decision to publish the results.

Acknowledgments: We are highly thankful to our lab colleagues for their assistance in data

analysis. We wish to thank Dr. Zhiqiang Xia (ITBB) for providing facilities and assistance in the

construction of the Circos plot.

Conflicts of Interest: The authors declare no conflict of interest.

Author Contributions: Conceptualization, M.A.A. and S.Z.; Methodology, M.A.A., X.H and

F.A; Software, M.A.A. F.A and X.H; Validation, F.A, L.S and X.F.; Formal Analysis, M.A.A,

X.H.; Investigation, M.A.A., L.S and X.F.; Resources, S.Z. and X.F.; Data Curation, S.Z, L.S.

and X.F; Writing—Original Draft Preparation, M.A.A and F.A; Writing—Review and Editing,

M.A.A, F.A and S.Z.; Visualization, L.S; Supervision, S.Z.; Funding Acquisition, S.Z, X.F and

X.H. All authors have read and agree to the published version of the manuscript.

References

1. Bouhida, M., B. Lockhart, and N.E. Olszewski, *An analysis of the complete sequence of a sugarcane bacilliform virus genome infectious to banana and rice.* Journal of General Virology, 1993. **74**(1): p. 15-22.

2. Sun, S.-R., et al., *Molecular characterization of two divergent variants of sugarcane bacilliform viruses infecting sugarcane in China.* European Journal of Plant Pathology, 2016. **145**(2): p. 375-384.

3. King, A.M., et al., *Virus taxonomy: ninth report of the International Committee on Taxonomy of Viruses.* Vol. 9. 2011: Elsevier.

4. Geijskes, R.J., et al., *Sequence analysis of an Australian isolate of sugarcane bacilliform badnavirus.* Archives of virology, 2002. **147**(12): p. 2393-2404.

5. Muller, E., et al., *High molecular variability of sugarcane bacilliform viruses in Guadeloupe implying the existence of at least three new species.* Virus research, 2011. **160**(1-2): p. 414-419.

6. Ahmad, K., et al., *Presence of diverse sugarcane bacilliform viruses infecting sugarcane in China revealed by pairwise sequence comparisons and phylogenetic analysis.* The plant pathology journal, 2019. **35**(1): p. 41.

7. Bousalem, M., E.J. Douzery, and S. Seal, *Taxonomy, molecular phylogeny and evolution of plant reverse transcribing viruses (family Caulimoviridae) inferred from full-length genome and reverse transcriptase sequences.* Archives of virology, 2008. **153**(6): p. 1085-1102.

8. Borah, B.K., et al., *Bacilliform DNA-containing plant viruses in the tropics: commonalities within a genetically diverse group.* Molecular plant pathology, 2013. **14**(8): p. 759-771.

9. Iskra-Caruana, M.-I., et al., *The common evolutionary history of badnaviruses and banana.* Infection, Genetics and Evolution, 2014. **21**: p. 83-89.

10. Autrey, L., et al. *The distribution of sugarcane bacilliform virus in various geographical regions.* in *Proc Int Soc Sugar Cane Technol.* 1995.

11. Lockhart, B., M. Irey, and J. Comstock. *Sugarcane bacilliform virus, sugarcane mild mosaic virus and sugarcane yellow leaf syndrome.* in *Sugarcane germplasm conservation and exchange: report of an international workshop held in Brisbane, Queensland, Australia, 28-30 June 1995.* 1996. Canberra, Australia: Australian Centre for International Agricultural ….

12. Muhammad, T., et al., *RNA interference: a natural immune system of plants to counteract biotic stressors.* Cells, 2019. **8**(1): p. 38.

13. Khvorova, A., A. Reynolds, and S.D. Jayasena, *Functional siRNAs and miRNAs exhibit strand bias.* Cell, 2003. **115**(2): p. 209-216.

14. Wang, W. and G. Galili, *Tuning the Orchestra: miRNAs in plant immunity.* Trends in plant science, 2019. **24**(3): p. 189-191.

15. Gaffar, F.Y. and A. Koch, *Catch me if you can! RNA silencing-based improvement of antiviral plant immunity.* Viruses, 2019. **11**(7): p. 673.

16. Qu, J., J. Ye, and R. Fang, *Artificial microRNA-mediated virus resistance in plants.* Journal of virology, 2007. **81**(12): p. 6690-6699.

17. Rosa, C., et al., *RNA interference mechanisms and applications in plant pathology.* Annual review of phytopathology, 2018. **56**: p. 581-610.

18. Niu, Q.-W., et al., *Expression of artificial microRNAs in transgenic Arabidopsis thaliana confers virus resistance.* Nature biotechnology, 2006. **24**(11): p. 1420-1428.

19. Ali, I., et al., *Artificial microRNA-mediated resistance against the monopartite begomovirus Cotton leaf curl Burewala virus.* Virology journal, 2013. **10**(1): p. 231.

20. Zhang, X., et al., *Expression of artificial microRNAs in tomato confers efficient and stable virus*
resistance in a cell-autonomous manner. Transgenic research, 2011. **20**(3): p. 569-581.

21. Petchthai, U., C.S. Le Yee, and S.-M. Wong, *Resistance to CymMV and ORSV in artificial*
microRNA transgenic Nicotiana benthamiana plants. Scientific reports, 2018. **8**(1): p. 1-8.

22. Seal, S., et al., *The prevalence of badnaviruses in West African yams (Dioscorea cayenensis-*
rotundata) and evidence of endogenous pararetrovirus sequences in their genomes. Virus
Research, 2014. **186**: p. 144-154.

23. Harper, G., et al., *Detection of episomal banana streak badnavirus by IC-PCR.* Journal of
Virological Methods, 1999. **79**(1): p. 1-8.

24. Sun, S.-R., et al., *Development of quantitative real-time PCR assays for rapid and sensitive*
detection of two badnavirus species in sugarcane. BioMed research international, 2018. **2018**.

25. Sukal, A.C., et al., *Assessment and optimization of rolling circle amplification protocols for the*
detection and characterization of badnaviruses. Virology, 2019. **529**: p. 73-80.

26. Bömer, M., et al., *A sequence-independent strategy for amplification and characterisation of*
episomal badnavirus sequences reveals three previously uncharacterised yam badnaviruses.
Viruses, 2016. **8**(7): p. 188.

27. Peterson, S.M., et al., *Common features of microRNA target prediction tools.* Frontiers in
genetics, 2014. **5**: p. 23.

28. Kozomara, A., M. Birgaoanu, and S. Griffiths-Jones, *miRBase: from microRNA sequences to*
function. Nucleic acids research, 2019. **47**(D1): p. D155-D162.

29. Griffiths-Jones, S., et al., *miRBase: microRNA sequences, targets and gene nomenclature.* Nucleic
acids research, 2006. **34**(suppl_1): p. D140-D144.

30. Kozomara, A. and S. Griffiths-Jones, *miRBase: annotating high confidence microRNAs using deep*
sequencing data. Nucleic acids research, 2014. **42**(D1): p. D68-D73.

31. Karuppaiah, R., R. Viswanathan, and V.G. Kumar, *Genetic diversity of Sugarcane bacilliform virus*
isolates infecting Saccharum spp. in India. Virus genes, 2013. **46**(3): p. 505-516.

32. Enright, A.J., et al., *MicroRNA targets in Drosophila.* Genome biology, 2003. **5**(1): p. R1.

33. John, B., et al., *Human microRNA targets.* PLoS biol, 2004. **2**(11): p. e363.

34. Miranda, K.C., et al., *A pattern-based method for the identification of MicroRNA binding sites*
and their corresponding heteroduplexes. Cell, 2006. **126**(6): p. 1203-1217.

35. Loher, P. and I. Rigoutsos, *Interactive exploration of RNA22 microRNA target predictions.*
Bioinformatics, 2012. **28**(24): p. 3322-3323.

36. Krüger, J. and M. Rehmsmeier, *RNAhybrid: microRNA target prediction easy, fast and flexible.*
Nucleic acids research, 2006. **34**(suppl_2): p. W451-W454.

37. Brodersen, P., et al., *Widespread translational inhibition by plant miRNAs and siRNAs.* Science,
2008. **320**(5880): p. 1185-1190.

38. Dai, X., Z. Zhuang, and P.X. Zhao, *psRNATarget: a plant small RNA target analysis server (2017*
release). Nucleic acids research, 2018. **46**(W1): p. W49-W54.

39. Dai, X. and P.X. Zhao, *psRNATarget: a plant small RNA target analysis server.* Nucleic acids
research, 2011. **39**(suppl_2): p. W155-W159.

40. Krzywinski, M., et al., *Circos: an information aesthetic for comparative genomics.* Genome
research, 2009. **19**(9): p. 1639-1645.

41. Lorenz, R., et al., *ViennaRNA Package 2.0.* Algorithms for molecular biology, 2011. **6**(1): p. 26.

42. Bernhart, S.H., et al., *Partition function and base pairing probabilities of RNA heterodimers.*
Algorithms for Molecular Biology, 2006. **1**(1): p. 3.

43. Gandrud, C., *Reproducible research with R and R studio.* 2013: CRC Press.

44. Jacquot, E., et al., *The open reading frame 2 product of cacao swollen shoot badnavirus is a*
nucleic acid-binding protein. Virology, 1996. **225**(1): p. 191-195.

45.   Naquin, D., et al., *CIRCUS: a package for Circos display of structural genome variations from*
paired-end and mate-pair sequencing data. BMC bioinformatics, 2014. **15**(1): p. 1-6.

46.   Betel, D., et al., *Comprehensive modeling of microRNA targets predicts functional non-conserved*
and non-canonical sites. Genome biology, 2010. **11**(8): p. R90.

47.   Riffo-Campos, Á.L., I. Riquelme, and P. Brebi-Mieville, *Tools for sequence-based miRNA target*
prediction: what to choose? International journal of molecular sciences, 2016. **17**(12): p. 1987.

48.   Min, H. and S. Yoon, *Got target?: computational methods for microRNA target prediction and*
their extension. Experimental & molecular medicine, 2010. **42**(4): p. 233-244.

49.   Jones-Rhoades, M.W., D.P. Bartel, and B. Bartel, *MicroRNAs and their regulatory roles in plants.*
Annu. Rev. Plant Biol., 2006. **57**: p. 19-53.

50.   Doench, J.G. and P.A. Sharp, *Specificity of microRNA target selection in translational repression.*
Genes & development, 2004. **18**(5): p. 504-511.

51.   Lewis, B.P., C.B. Burge, and D.P. Bartel, *Conserved seed pairing, often flanked by adenosines,*
indicates that thousands of human genes are microRNA targets. cell, 2005. **120**(1): p. 15-20.

52.   Huang, Y., et al., *A study of miRNAs targets prediction and experimental validation.* Protein &
cell, 2010. **1**(11): p. 979-986.

53.   M Witkos, T., E. Koscianska, and W. J Krzyzosiak, *Practical aspects of microRNA target prediction.*
Current molecular medicine, 2011. **11**(2): p. 93-109.

54.   Iqbal, M.S., et al., *Prediction of host-derived miRNAs with the potential to target PVY in potato*
plants. Frontiers in Genetics, 2016. **7**: p. 159.

55.   Iqbal, M.S., et al., *In silico MCMV silencing concludes potential host-derived miRNAs in maize.*
Frontiers in plant science, 2017. **8**: p. 372.

56.   Akhter, Y. and J.A. Khan, *Genome wide identification of cotton (Gossypium hirsutum)-encoded*
microRNA targets against Cotton leaf curl Burewala virus. Gene, 2018. **638**: p. 60-65.

57.   Jabbar, B., et al., *Target prediction of candidate miRNAs from Oryza sativa for silencing the*
RYMV genome. Computational biology and chemistry, 2019. **83**: p. 107127.

58.   Ashraf, F., et al., *A novel computational approach to the silencing of Sugarcane Bacilliform*
Guadeloupe A Virus determines potential host-derived MicroRNAs in sugarcane (Saccharum
officinarum L.). PeerJ, 2020. **8**: p. e8359.

59.   Pinzón, N., et al., *microRNA target prediction programs predict many false positives.* Genome
research, 2017. **27**(2): p. 234-245.

60.   Kertesz, M., et al., *The role of site accessibility in microRNA target recognition.* Nature genetics,
2007. **39**(10): p. 1278-1284.

61.   Yen, P.-S., et al., *Assessing the potential interactions between cellular miRNA and arboviral*
genomic RNA in the yellow fever mosquito, aedes aegypti. Viruses, 2019. **11**(6): p. 540.

62.   Millar, A.A., A. Lohe, and G. Wong, *Biology and Function of miR159 in Plants.* Plants, 2019. **8**(8):
p. 255.

63.   Cheng, Z., et al., *Integrated mRNA, MicroRNA Transcriptome and Degradome Analyses Provide*
Insights into Stamen Development in Moso Bamboo. Plant and Cell Physiology, 2020. **61**(1): p.
76-87.

64.   Hirsch, A.J., *The use of RNAi-based screens to identify host proteins involved in viral replication.*
Future microbiology, 2010. **5**(2): p. 303-311.

sof-MIR159a   sof-MIR159b   sof-MIR159d   sof-MIR159e   ssp-MIR444a   ssp-MIR444b

A   B   C   D   E   F

A miRNA

miRNA★

sof-MIR159
(miRNA Precursor)

miRNA/miRNA* duplex replacement

B amiRNA

amiRNA★

Development of plant expression cassette

C Promoter   Pre-amiRNA   NOS Ter

Expression evaluation in transgenic plants

D amiRNA

amiRNA★

Pre-amiRNA

Cleavage of SCBV target mRNA in sugarcane cultivar

E amiRNA

m⁷G ——————————————— AAAAA

F

Artificial miRNAs