Biotransformation of lindane (γ-hexachlorocyclohexane) to non-toxic end products by sequential

treatment with three mixed anaerobic microbial cultures

Luz A. Puentes Jácome[a], Line Lomheim[a], Sarra Gaspard[b], and Elizabeth A. Edwards[a,c,*]

[a]Department of Chemical Engineering and Applied Chemistry, University of Toronto, Toronto,

Ontario, Canada

[b]Laboratory COVACHIMM2E, Université des Antilles, Pointe à Pitre, Guadeloupe, French

West-Indies, France

[c]Department of Cell and Systems Biology, University of Toronto, Toronto, Ontario, Canada

[*]Address correspondence to Elizabeth A. Edwards, elizabeth.edwards@utoronto.ca

Abstract

The γ isomer of hexachlorocyclohexane (HCH), also known as lindane, is a carcinogenic

persistent organic pollutant. Lindane was used worldwide as an agricultural insecticide. Legacy

soil and groundwater contamination with lindane and other HCH isomers is still a big concern.

The biotic reductive dechlorination of HCH to non-desirable and toxic lower chlorinated

compounds such as monochlorobenzene (MCB) and benzene, among others, has been broadly

documented. Here, we demonstrate for the first time that complete biotransformation of lindane

to non-toxic end products is attainable using a sequential treatment approach with three mixed

anaerobic microbial cultures referred to as culture I, II, and III. Biaugmentation with culture I

achieved dechlorination of lindane to MCB and benzene. Culture II was able to dechlorinate

MCB to benzene, and finally, culture III carried out methanogenic benzene degradation. Distinct

Dehalobacter populations, corresponding to different 16S rRNA amplicon sequence variants in

culture I and culture II, were responsible for lindane and MCB dechlorination, respectively. This

study continues to highlight key roles of *Dehalobacter* spp. as chlorobenzene- and HCH-

organohalide-respiring bacteria and demonstrates that sequential treatment with specialized

anaerobic cultures may be explored at field sites in order to address legacy soil and groundwater

contamination with HCH.

Introduction

Hexachlorocyclohexane (HCH) isomers are persistent organic pollutants. Lindane, the γ isomer

of hexachloroycyclohexane (γ-HCH), is the only HCH isomer exhibiting insecticidal properties,

and it is a known carcinogen. During the manufacture of "pure" lindane, large amounts of HCH

isomers were produced as waste by-products. Lindane (γ-HCH) and other HCH isomers (α-, β-,

δ-, and ε-HCH) were components of the agriproduct known as "technical HCH". Technical HCH

and lindane were extensively applied to a variety of crops for pest control. In some instances,

HCH manufacture by-products were also improperly disposed of in landfills or waste dumps.

Although most countries have banned the use of HCH, groundwater, soil, and sediment

contaminated with HCH is of worldwide concern. There is extensive evidence of HCH

contamination at a variety of sites in Spain,[1, 2] Germany,[3] China,[4] and the French West Indies,[5]

among others. Legacy HCH contamination, estimated at ~ 4 to 7 million tons,[6-8] has been

difficult to address and manage, likely because of the extent of the contamination and the

physicochemical characteristics of each HCH isomer.

Hexachlorocyclohexane isomers are poorly soluble in water. Estimates of water solubility for γ-

HCH (lindane) vary; some sources report it to be around 1 to 2 mg/L[9] and others estimate it to be

as high as 7 to 10 mg/L[10, 11] at temperatures near 20$\circ$C. HCH isomers partition into organic

compounds (log K_{ow}~3.3) and organic carbon (log K_{oc}~3.0).[12] Thus, HCHs sorb readily to

sediment and soil fractions. HCHs are persistent in the environment, yet the amount of detectable

HCH that remains in agricultural soils varies largely depending on the characteristics of the

contaminated site; for a review on the persistence of HCH isomers refer to Bhatt et al. (2009).[13]

Long-range global atmospheric transport and deposition of HCHs has also been demonstrated,[14,

15] resulting in pollution of pristine environments including the Canadian Artic. Natural

attenuation of HCH contamination over years or decades has been documented; a summary of

these studies is also included in Bhatt et al. (2009).[13] In light of the extent of HCH contamination

and the associated risk posed to human health and the environment, active remediation of

contaminated sites may be desirable. Bioremediation could be a suitable approach for HCH

transformation as demonstrated at a field study in the Netherlands.[16]

The aerobic biodegradation of HCH is well documented: pathways, microorganisms, and

enzymes are known.[17] The anaerobic biodegradation of HCH has been studied to a lesser extent,

yet anaerobic conditions often prevail in HCH-impacted sediments and groundwater. In such

environments, HCH isomers can undergo biologically-mediated dechlorination reactions.

Dechlorination of HCH isomers has been observed in sediment microcosms,[18] co-cultures,[19]

anaerobic sludge,[20] and enrichment cultures.[21] Known dechlorinating bacteria, such as

Dehalobacter, *Dehalococcoides*, and *Clostridium* have been linked to HCH dechlorination,[19, 20,]

[22, 23] *Dehalobacter* sp. E1 can metabolically dechlorinate β-HCH to MCB and benzene.[19] Partial

HCH dechlorination (in which only a fraction of the added HCH mass was dechlorinated) was

observed in pure cultures of *Dehalococcoides mccartyi* strains BTF08 and 195 after being pre-

grown with tetrachloroethene (PCE);[23, 24] it has been suggested that *Dehalococcoides mccartyi*

strain 195 may be able to respire γ-HCH.[23] Monochlorobenzene (MCB) and benzene are

frequently reported as stable end products of HCH dechlorination. However, since MCB is toxic

and benzene is a known carcinogen, these are undesirable biotransformation products. Although

MCB and benzene can be biodegraded aerobically, they are often found at contaminated sites

along with HCH.[1, 3, 16] HCH bioremediation could be feasible as long as any potentially toxic by-

products can be biodegraded as well. At some sites, anaerobic biodegradation of these

dechlorination by-products may be challenging. However, the sustained anaerobic conversion of

MCB and benzene was previously demonstrated in laboratory microcosm experiments[25] and

could be coupled with HCH-dechlorinating bioremediation technologies to achieve complete

detoxification.

MCB and benzene, the products of anaerobic HCH dechlorination, have been successfully

biotransformed anaerobically.[26-30] Independent long term investigations in our laboratory have

led to the development of three distinct sets of enrichment cultures of interest for HCH

bioremediation. The first set of cultures are HCH-dechlorinating enrichment cultures

(dechlorinating α-, β-, δ-, and γ-HCH), derived from contaminated sediments from Guadeloupe

(French West Indies), which completely transform HCH to MCB and benzene.[21] Here, the

culture that dechlorinates γ-HCH will be referred as culture I. The second culture, or culture II, is

a KB-1-derived enrichment culture that dechlorinates MCB to benzene.[26] The third culture, or

culture III, is a methanogenic enrichment culture in which benzene is converted to methane and

carbon dioxide[29] that has been enriched in the Edwards lab for over 20 years. We hypothesized

that a combined treatment using the three cultures would be able to completely transform HCH

to non-toxic products. Rather than simply mixing the 3 specific cultures together, we tested the

hypothesis via sequential addition of each culture. Sequential addition would allow us to

minimize any detrimental interactions between the different microbial populations, i.e. the

benzene-degrading culture (culture III) uses benzene as electron donor and its activity could be

impaired by the presence of alternative electron donors which are required for cultures I and II.

Using the sequential approach, we were able to demonstrate the anaerobic bioconversion of

lindane (γ-HCH) all the way to non-toxic end products. In addition, we demonstrate that distinct

native *Dehalobacter* populations, corresponding to different 16S rRNA amplicon sequence

variants in culture I and culture II, are responsible for the biotransformation of lindane to MCB

and benzene in culture I, and for the dechlorination of MCB to benzene in culture II. Here, we

also discuss the implications of our findings which together support the development of

anaerobic bioremediation technologies for HCH, MCB, and benzene.

Materials and Methods

Chemicals

Chemicals were purchased through Millipore Sigma (Canada) at the highest purity available.

Ethanol and methanol for the preparation of feedstocks were HPLC grade. Gases for anaerobic

culturing were purchased through Praxair (Canada).

Enrichment cultures used in this study

Culture I is an anaerobic mixed microbial culture, enriched from soil and river sediment

microcosms originating from lindane-contaminated sites in Guadeloupe (French West Indies). [21]

The microcosms were first set up in 2010, and details of the history of culture enrichment are

shown in the Supplementary Information (Figure S1). Culture I dechlorinates lindane (γ-HCH,

electron acceptor) stoichiometrically to a mixture of MCB and benzene (see **Error! Reference**

source not found., panel A, and Figure S2). Ethanol is currently supplied as the electron donor

at a ratio of 5 to 1 electron equivalents of donor to acceptor.

Culture II is a KB-1-derived anaerobic enrichment culture that dechlorinates MCB (electron

acceptor) to benzene (see **Error! Reference source not found.**, panel B). The culture was

derived from an enrichment culture that dechlorinates 1,2,4-trichlorobenzene to a mixture of

dichlorobenzene, MCB, and benzene.[26] Culture II is supplied with methanol (electron donor) at a

ratio of 5 to 1 electron equivalents of donor to acceptor, but additional re-amendments of donor

are generally needed for complete dechlorination of MCB to benzene.

Culture III is an anaerobic methanogenic enrichment culture that transforms benzene (electron

donor) through a series of syntrophic microbial interactions to methane and carbon dioxide (see

Figure 1, panel C).[29] This culture is currently being scaled-up at the SiREM Lab

(https://www.siremlab.com/) as DGG-B® and pilot tests are underway to evaluate its

effectiveness at field scale.

Sequential treatment with specialized anaerobic microbial enrichment cultures

The biotransformation experiment was performed in three phases, each phase initiated by the

addition of one of the three distinct cultures described above (see illustration of experiment setup

in Figure S3). A bicarbonate-buffered mineral medium[26], pre-reduced with an amorphous iron

(II) sulfide (FeS) slurry (~0.8 g/L), was used to set up and top up the experimental bottles. All

the steps involving feeding, inoculation, and sampling were performed in an anaerobic chamber

(Coy Lab Products, USA). All the experiments were set up using 115 mL clear glass bottles with

screw cap Mininert® valves to a final liquid volume of 90 mL. During phase I (~150 days), nine

active bottles (labelled as experimental bottles 1 to 9 in the supplementary Excel file) containing

lindane (~4.3 µmol or 1.25 mg per bottle) were inoculated (on Day 0) with a 10% transfer of the

lindane-dechlorinating culture (culture I), resulting in a final concentration of 1×10^7 bacterial

cells/mL in each experimental bottle. Three additional bottles were inoculated with 10%

autoclaved inoculum (negative controls); the inoculum was autoclaved in a liquid cycle held at

121°C and 15 psi for 45 min. Glass bottles were pre-loaded with lindane prior to bringing them

into the anaerobic chamber as follows: lindane was added to autoclaved empty bottles as a

concentrated lindane/acetone solution which was completely evaporated under a stream of

nitrogen. The bottles were then transferred to the anaerobic chamber for a minimum of 2 days

prior to inoculation. During inoculation, sterile mineral medium and culture I were aliquoted

directly into each bottle using sterile pipettes. On Day 93, after all lindane had been converted to

MCB and benzene, cultures were transferred into new bottles with additional lindane (~8.6 µmol

or 2.25 mg per bottle, prepared as described above). Electron donor, ethanol, was added every

four weeks to experimental and control bottles (on Day 0, 27, 59, 93, and 123) at a concentration

of 0.3 mM. The total amount of donor added during phase I is equivalent to 20 times the total

electron equivalents of acceptor added (assuming 6 electron equivalents per mole of lindane).

During phase II, from Day 148 to 337 (~190 days), six of the nine active bottles from phase I

(labelled as experimental bottles 1 to 6 in the supplementary Excel file) were bioaugmented on

two separate dates (Day 148 and Day 275) with the MCB-dechlorinating culture (culture II). For

each bioaugmentation event of phase II, the experimental bottles were inoculated with 1×10^6

bacterial cells of culture II per mL. From phase II onwards, cultures were added to each bottle as

concentrated inoculum using sterile plastic syringes via the septum on the screw cap to minimize

the potential loss of MCB and benzene and to replenish the experimental working volume (90

mL). For the first and second phase-II bioaugmentation events, 5 mL of 10X-concentrated

culture II and 10 mL of 4X-concentrated culture II were added to each bottle, respectively (refer

to the SI for concentration protocol). Culture II-bioaugmented bottles received methanol as

electron donor. Methanol was fed at a concentration of 0.1 mM (Day 148 and Day 198) and 0.2

mM (Day 218, 258, 275). The total amount of methanol added to these bottles throughout phase

II was equivalent to 50 times the electron equivalents of acceptor (total MCB produced during

the experiment; 2 electron equivalents per mole of MCB). The three remaining active bottles

from phase I were not inoculated with culture II, but were monitored as controls. Ethanol was

added to these control bottles on Day 198, 218, and 296.

During phase III (Day 148 to 337), three of the six active bottles that were bioaugmented in

phase II (labelled as experimental bottles 4, 5, and 6 in the supplementary Excel file) were

inoculated with 4×10^8 bacterial cells of the benzene-degrading culture (culture III) per mL,

while the remaining three bottles were monitored as controls. Additionally, two positive controls,

inoculated with culture III in the same fashion as bottles 4, 5, and 6, were set up in parallel to

monitor the activity of the benzene-degrading culture; neat benzene was added directly to fresh

mineral medium. Similarly to phase II, culture III was added to each bioaugmented bottle as

concentrated inoculum (6 mL at 6X) using sterile plastic syringes via the septum on the screw

cap (refer to the SI for concentration protocol).

Analytical methods

MCB, benzene, and methane were quantified using GC-FID (gas chromatography with flame

ionization detection). For the analysis, aqueous samples (1 mL) were added to 5 mL of acidified

(using 6N HCl) deionized water (pH < 2). Calibration standards for MCB and benzene were

prepared gravimetrically by adding a known mass of a combined MCB/benzene stock prepared

in methanol to bottles containing known volumes of water and headspace. Methane standards

were prepared by adding known volumes of pure methane gas, using gas-tight syringes, to

bottles containing known volumes of water and headspace. External calibrations were prepared

to convert area counts into concentrations (refer to the SI for details on GC method). Analysis of

γ-hexachlorocyclohexane (lindane) was not performed for the sequential biotransformation

experiment since that would have required removal of more than 1 mL of culture volume per

bottle per sampling point during phase I. However, lindane was measured in a separate

experiment shown in Figure S2, in which it was verified that culture I biotransforms lindane to

MCB and benzene stoichiometrically (moles of lindane degraded = moles of MCB + benzene

produced). This stoichiometric relationship between lindane fed and MCB and benzene produced

has consistently been observed in this culture throughout its enrichment process.

DNA sampling and quantitative PCR (qPCR)

DNA was extracted from 2 mL culture samples. Cells were harvested by centrifugation at 10,000x g for 15 min at 4°C. Cell pellets were resuspended in 100 μl of supernatant and DNA was extracted using the DNeasy PowerSoil kit following the manufacturer's recommendations (during steps 9 and 12 of the protocol, all supernatant was transferred). Real-time polymerase chain reaction (qPCR) assays were performed to track the gene copies of total Bacteria, *Dehalobacter* and Desulforomonadales using specific 16S rRNA gene primers (see Table S1 and Table S2 in the SI for qPCR primers and calibration information, respectively). For this study, a new primer set that targets representative sequences classified as either Desulforomonadales and/or *Geobacter* (a genus of the order Desulforomonadales) was designed; the target sequences were obtained from previous 16S rRNA gene amplicon sequencing studies of culture I[21] (refer to the SI for the qPCR prep and quantification protocol).

16S rRNA gene amplicon sequencing

DNA was extracted as described above and sent for Illumina amplicon sequencing at the Genome Quebec Innovation Centre (McGill University). DNA was sequenced using the Illumina MiSeq PE300 platform with modified versions of primers 926f and 1392r (926f-modified: AAACTYAAAKGAATWGRCGG; 1392r-modified: ACGGGCGGTGWGTRC) which target the V6−V8 variable region of the 16S rRNA gene from bacteria and archaea, as well as the 18S rRNA gene in eukarya. After sequencing, raw data was processed using the QIIME 2[31] Core 2020.2 distribution. Primers were removed using the cutadapt[32] plugin. Sequences were denoised and merged into amplicon sequence variants (ASVs) using DADA2[33] (forward sequence truncated at 260 bp, reverse sequence truncated at 240 bp). Taxonomy was assigned to ASVs using the feature-classifier plugin[34] with the classify-sklearn method using a pre-trained Naïve

Bayes Classifier with the SILVA 132 99% OTUs full-length sequences database (available at

https://docs.qiime2.org/2020.2/data-resources/). Data was taken to MS Excel and RStudio for

further processing. An Excel supplementary file is included with the resulting taxonomy and

summary of ASVs of interest. Raw 16S rRNA amplicon sequencing data has been deposited to

the NCBI Sequence Read Archive under bioproject PRJNA661282.

Estimation of absolute abundance of amplicon sequence variants (ASVs)

Absolute abundance of ASVs of interest was calculated as the product of the qPCR-determined

Bacteria 16S rRNA gene copy numbers (determined using the general Bacteria primer shown in

Table S1) and the relative abundance of each ASV (included in the Supplementary Excel file)

determined by 16S rRNA gene amplicon sequencing via the QIIME 2 pipeline (described

above). This approach was used to obtain the absolute abundance of bacterial groups of interest

during phase III and for the different *Dehalobacter* ASVs.

Results and Discussion

Inoculum samples from enrichment cultures used during the biotransformation

experiment

Samples from cultures I, II, and III were used as inoculum for each respective phase of the

biotransformation experiment. **Error! Reference source not found.** shows the observed

biotransformation step (top diagram), the bacterial community composition determined by 16S

rRNA amplicon sequencing at the time of inoculation (middle), and culture maintenance data

(bottom) for each culture; cultures I, II, and III, are shown in panels A, B, and C, respectively.

For culture I (**Error! Reference source not found.**A), lindane is dechlorinated to MCB and

benzene. The proposed series of reactions that leads to the formation of MCB and benzene in

culture I are discussed in Qiao et al.[21] Dechlorination rates for this culture range from ~ 6.0 to 13

µmol/L/day (1.8 – 3.8 mg/L/day). Detected abundant bacterial genera (> 2% relative abundance)

that are known organohalide-respirers include *Geobacter*, and *Dehalobacter.* Other abundant

bacterial genera include representatives of the fermentative and sulfate-reducing genus

Desulfovibrio, members of the recently revealed candidate phyla Berkelbacteria, as well as

members of the *Syntrophobacter*, and the Desulforomonadales family. Culture II, which

dechlorinates MCB to benzene, contains abundant bacteria classified as *Dehalobacter,*

Cloacimonas, *Synergistaceae, Pelolinea,* and *Spirochaetacea* (**Error! Reference source not**

found.B). MCB dechlorination to benzene is observed a rate of ~ 2.0 ± 0.3 µmol/L/day (0.16 ±

0.02 mg/L/day). Culture III, degrades benzene to methane and carbon dioxide (**Error!**

Reference source not found.C). Culture III is highly enriched in bacteria classified as

Deltaproteobacterial candidate Sva0485 which represents the key microorganism, historically

termed "ORM2" for Oil Refinery Methanogenic Clone 2, responsible for the initial attack on the

benzene ring.[30, 35] Other abundant bacteria in this culture belong to the phyla Omnitrophicaeota

and Yanofskybacteria. The ASVs classified as Yanofskybacteria correspond to another

previously described abundant bacterium in the benzene-degrading cultures, referred to as OD1.

[29] The benzene degradation rate for culture III used in this study was ~ 6.4 µmol/L/day (~ 0.5

mg/L/d).

Error! Reference source not found.

Phase I: dechlorination of lindane (γ-HCH) to MCB and benzene

Error! Reference source not found.A shows the dechlorination of lindane from Day 0 to Day 173, in the 9 experimental bottles inoculated with culture I. Within 14 days, about 96% of the theoretical mass of lindane added to the bottles was dechlorinated at an average rate of 3.3 ± 0.4 µmol/L/d (1.0 ± 0.1 mg/L/d). By Day 23, all lindane was transformed to MCB and benzene. Even though electron donor (ethanol) was re-amended (Day 27 and Day 59), further dechlorination of MCB to benzene did not occur. This is consistent with the observations gathered for this culture during years of sequential transfers and maintenance (Figure S1, Supplementary Information). For the second feeding event (Day 93), the added mass of lindane

was 1.8 times greater than the first feeding. As expected, corresponding stoichiometric amounts

of MCB and benzene were produced. After 13 days (Day 106), 53% of the re-amended

theoretical mass of lindane was dechlorinated at an average dechlorination rate of 3.9 ± 0.3

µmol/L/d (1.1 ± 0.1 mg/L/d). Electron donor limitation became evident after sampling on Day

121, so ethanol was re-amended (Day 123). All lindane was dechlorinated afterwards. At the end

of phase I, the average molar ratio of benzene to MCB was 0.32 ± 0.06. In the control bottles

with autoclaved inoculum, only trace amounts of benzene and MCB were detected (**Error!**

Reference source not found.B), likely due to abiotic reductive dechlorination.

The qPCR-determined 16S rRNA gene copies per mL of culture of *Dehalobacter*,

Desulfuromomadales/*Geobacter*, and total Bacteria are shown in **Error! Reference source not**

found.C and D. There was a significant increase in *Dehalobacter* and total Bacteria copy

numbers from Day 93 (prior to the second lindane feeding) to Day 148. *Dehalobacter* copy

numbers increased by 4.3-fold. Desulfuromomadales/*Geobacter* copy numbers did not increase.

This indicates that the growth of *Dehalobacter* is linked to the dechlorination of lindane to MCB

and benzene, and that bacteria belonging to the Desulfuromomadales/*Geobacter* are likely not

directly implicated. These findings are consistent with observations reported earlier for culture

I.[21] Since some *Geobacter* spp. are known to dechlorinate organochlorines[36-38], we had

hypothesized that they could be involved in lindane dechlorination. *Geobacter* in culture I are

potentially using ethanol (supplied during culture maintenance) or other bacterial metabolites,

e.g. acetate or H_2 as electron donor. The use of ethanol as electron donor is a known feature of

some *Geobacter* spp. such as *G. chapellei* strain 172T, *G. metallireducens* GS-15T, *G.*

hydrogenophilus H-2T, and *G. grbiciae* TACP-2T[37, 39]. In addition, *G. hydrogenophilus* H-2T, *G.*

grbiciae TACP-2 T, and *G. sulfurreducens* can use use H_2 as electron donor[39] and require acetate

as a carbon source for growth with H_2. *Geobacter* spp. can use a wide range of electron

acceptors, e.g. elemental sulfur, nitrate, malate, Mn (IV), Fe (III), and some organochlorines, and

can also transfer electrons directly to other microbes[40]. Bacteria belonging to the

Desulfuromomadales could also be fermenting organic acids.

Interestingly, *Dehalobacter* but not Desulfuromomadales/*Geobacter* copy numbers were

quantifiable via qPCR in the bottles with autoclaved inoculum. *Dehalobacter* gene copies were

also detected in heat-treated experimental control bottles from a previous study with

Dehalobacter-enriched cultures.[26] *Dehalobacter* spp. are members of the Firmicutes which are

known to produce heat-resistant endospores.[41] Endospores may survive the autoclaving process

and thus result in cell viability and DNA preservation. *Dehalobacter* genomes do contain

sporulation genes[42], yet the *Dehalobacter restrictus* type strain, DSM 9455, formerly known as

PER-K23, was reported as not spore-forming.[43] Additional studies on *Dehalobacter* spp. are

required to determine if spore formation is a feature of select strains.

Phase I: Dechlorination of lindane after addition of culture I

Error! Reference source not found.

Phase II: dechlorination of MCB to benzene

On Day 148, six of the active bottles (experimental bottles 1 to 6) from phase I were
bioaugmented with culture II. Experimental bottles 7, 8, 9 were not bioaugmented with culture
II, and remained as controls. **Error! Reference source not found.** shows the steady
dechlorination of MCB to benzene in five out of the six bioaugmented bottles; one of the
bioaugmented bottles did not initially exhibit significant MCB dechlorination, and is not
included in the graph. A slight reduction in dechlorination rates was observed for the time period

between Day 237 and 254 with some bottles exhibiting significant deviations from the mean

dechlorination rate. As a result, a second bioaugmentation event, in which culture II was again

added to experimental bottles 1 to 6, was performed on Day 275. Interestingly, after this

bioaugmentation event, MCB dechlorination was kick-started in the single bottle in which no

significant MCB dechlorination had been observed resulting in detectable MCB to benzene

transformation in all the bioaugmented bottles. The mean MCB to benzene dechlorination rate

for phase III was 0.3 ± 0.2 µmol/L/day (0.03 ± 0.02 mg/L/day, n = 5). This rate is about 5 times

lower than the average rate measured during routine cultivation of culture II (0.03 vs 0.16

mg/L/day). One of the factors that may have contributed to the rate reduction is the increased

competition for electron donor that the MCB-dechlorinating population likely faced in these

experimental bottles. As shown in **Error! Reference source not found.**, culture I (lindane

enrichment) is a mixed enrichment culture with organisms known to be able to use methanol, e.g.

for methane production in certain members of the *Methanosarcinales*, or potentially as electron

donor. Several amplicon sequence variants in culture I were classified to the genus

Desulfosporosinus. Some *Desulfosporosinus* spp. such as *D. meridie*, a homoacetogenic

bacterium, use methanol as electron donor during sulfate reduction.[44]

Benzene and MCB amounts in the control bottles (non-bioaugmented with culture II) are shown

in **Error! Reference source not found.**B. As expected, benzene did not increase in these bottles.

As phase II progressed, there was a slight decline in the amount of MCB and benzene measured

in the control and active bottles. This was due to addition of inoculum (10 mL) to the active

bottles during the second bioaugmentation event, and to the fact that all the experimental bottles

were topped up to 90 mL on Day 275. As shown in **Error! Reference source not found.**C, the

qPCR-measured Bacteria and *Dehalobacter* gene copies per mL of culture did not increase in the

active bottles over the 127-day period following inoculation. However, there was a measurable

decline (14-fold) of *Dehalobacter* gene copies per mL of culture in the control (non-

bioaugmented) bottles (**Error! Reference source not found.**D). This finding provides additional

evidence for the role of the *Dehalobacter* population in culture I as the lindane-dechlorinating

population. In the active bottles, the *Dehalobacter* gene copies did not decline because the

lindane-dechlorinating *Dehalobacter* population was being replaced by the MCB-dechlorinating

Dehalobacter population native to culture II. This will be further discussed later. In the controls

(non-bioaugmented with culture II, Desulfuromomadales/*Geobacter* gene copies did not decline

suggesting that these organisms were metabolizing some of the available exogenous or

endogenous electron donor (e.g. acetate) that likely was still present in the bottles; ethanol was

added to these controls in two occasions (Day 198 and 218).

On Day 337, before proceeding with the addition of culture III, some MCB still remained in the

culture II-bioaugmented bottles. For complete MCB dechlorination to have occurred, additional

bioaugmentation events, likely with a higher inoculum density, would have been required. As

will be discussed later, the *Dehalobacter* population in culture II is usually in the order of 10^7

cells/mL of culture, not very dense compared to other dechlorinating enrichment cultures, and its

calculated yield is lower than that of other *Dehalobacter* populations respiring different

chlorinated substrates.

Phase II: Dechlorination of MCB to benzene after addition of culture II

Error! Reference source not found.

Phase III: benzene biodegradation

After 190 days into phase II (Day 337), we proceeded with phase III, i.e. the bioaugmentation

with the benzene-degrading culture (culture III). At this point, $67 \pm 7\%$ of the mass of MCB had

been converted to benzene following the bioaugmentation with culture II. Culture III was added

to three of the active bottles from phase II (experimental bottles 4, 5, and 6); the remaining three

bottles from phase II served as controls. As shown in **Error! Reference source not found.**,

benzene was biodegraded at a rate of 1.3 ± 0.1 µmol/L/day (0.1 ± 0.01 mg/L/day) in two out of

the three bioaugmented bottles. In the remaining bioaugmented bottle, benzene degradation

occurred as well but at a lower rate. In the controls (not bioaugmented with culture III), benzene

degradation did not occur (**Error! Reference source not found.**B).

After the bioaugmentation with culture III, the resulting gene copies per mL of

Deltaproteobacterial candidate Sva0485 (previously referred to as ORM2), were in the order of

10^8 and remained at 10^7 after benzene degradation (**Error! Reference source not found.**C).

Recall that ORM2 is the bacterium responsible for the initial attack on the benzene ring. The

average amount of benzene produced in the experimental bottles (6.5 mg/L) was likely not

enough to maintain the population of ORM2 at such high density (10^8 ORM2 gene copies per

mL); culture III is generally maintained at higher concentrations, ranging from 30 to 150 mg/L,

in order to support high cell density. The gene copies per mL for

Desulfuromomadales/*Geobacter* remained at 10^7 for the active bottles and declined to 10^6 in the

control bottles (**Error! Reference source not found.**D). In the active bottles, endogenous

electron donors may have been available to sustain the population of

Desulfuromomadales/*Geobacter*. In the control bottles, electron donor was not added during

phase III, and thus the microbial community derived from phase I likely faced electron donor

shortage. The decline in Desulfuromomadales/*Geobacter* copy numbers in these control bottles

strongly suggests that these organisms were metabolizing some of the available electron donor

which was likely exhausted during the transition from phase II to phase III (methanol was last

supplied on Day 296).

Neat benzene (not produced from lindane dechlorination) was subsequently added to the three

experimental bottles on Day 441 to confirm that the benzene-degrading microbial community

remained active. As shown in Figure S4, benzene was indeed biodegraded afterwards. The

bottles that were not bioaugmented with culture II or culture III did not exhibit further

dechlorination or benzene degradation.

Error! Reference source not found.

MCB is dechlorinated to benzene in culture II by a native KB-1-derived *Dehalobacter*

population distinct from the native *Dehalobacter* population in culture I

As mentioned earlier, culture II originated from a KB-1-derived enrichment culture (referred to

in the lab as 1,2,4TCB/M_2014) that dechlorinates 1,2,4-trichlorobenzene (1,2,4-TCB) to a

mixture of dichlorobenzene, MCB, and benzene.[26] Culture II (500 mL scale-up) had been

maintained on MCB as the sole electron acceptor for about 2 years. To clearly demonstrate

Dehalobacter population growth during MCB dechlorination to benzene, the culture was

sampled frequently over a period of 127 days and MCB/Benzene and *Dehalobacter* abundances

were analyzed. As shown in Figure S5, during the dechlorination of $\sim$ 100 μmol of MCB, the

relative abundance of *Dehalobacter* increased from 2 to 17% which corresponds to a 12.5-fold

change in its population. Assuming one 16S rRNA gene copy per *Dehalobacter* cell, the

resulting *Dehalobacter* yield is 2.8 ± 0.05 x 10^{10} cells per mole of Cl⁻ released. This yield is two

orders of magnitude lower than that reported for the population grown using 1,2,4-TCB as

electron acceptor ($4.6x10^{12}$ to $1.8x10^{13}$ cells per mole Cl⁻ released).[26] The lower yield may be

related to the fact that only two electron equivalents can be transferred per mole of MCB in

contrast to four or six electron equivalents per mole of TCB depending on the resulting

dechlorination products.

As mentioned earlier, in the bottles bioaugmented with culture II, the qPCR-determined

Dehalobacter gene copies (**Error! Reference source not found.**C) did not decline because the

lindane-dechlorinating *Dehalobacter* population was replaced by the MCB-dechlorinating

Dehalobacter population native to culture II. We were able to demonstrate this after a close

examination of the 16S rRNA gene amplicon sequencing data. The *Dehalobacter* in culture I are

represented by a single amplicon sequence variant (ASV), while *Dehalobacter* in culture II are

represented by two dominant ASVs. The absolute abundance of *Dehalobacter* ASVs in cultures I

and II, as well as in the active experimental bottles before and after inoculation with culture II

are shown in **Error! Reference source not found.**. The sum of the absolute abundance of all

Dehalobacter ASVs, which were estimated from the product of the relative ASV abundance and

the qPCR-determined Bacteria gene copies, closely matches the qPCR-determined abundance of

Dehalobacter (Figure S6); this result adds confidence to our approach to estimate the absolute

abundance of each ASV of interest.

Dehalobacter ASV1 belongs to culture I, while *Dehalobacter* ASV2 and ASV3 belong to culture

II (**Error! Reference source not found.**A). There are thirteen nucleotide differences between

ASV1 and ASV2, twelve differences between ASV1 and ASV3, and only one nucleotide

difference between ASV2 and ASV3 (see Figure S7; ASVs can be found in the Supplementary

Excel file). After addition of culture II, the population represented by ASV1 declined and was

replaced by that of ASV2 and ASV3 which further confirms that the dechlorination of MCB to

benzene was carried out by *Dehalobacter* populations originating from culture II. ASV1, the

single dominant ASV in culture I is a 100% match to one of the *Dehalobacter* operational

taxonomic units (OTU 1134628) reported in Qiao et al.[21] for the HCH-dechlorinating culture set

that includes culture I.

A

Dehalobacter ASVs in culture inoculum

B

Dehalobacter ASVs in active experimental bottles

Error! Reference source not found.

After the second bioaugmentation with culture II, ASV2 remained in the biaugmented bottles

and ASV3 was found below detection. The differences observed in the absolute abundance of

ASV2 and ASV3 in the different samples indicates that these ASVs represent different

Dehalobacter populations in culture II. After an examination of historical 16S rRNA gene

amplicon sequencing data, we were able to determine that ASV2 is in fact the dominant

Dehalobacter ASV in the KB-1-derived enrichment cultures that dechlorinate chlorinated

benzenes, i.e. this ASV was the most abundant *Dehalobacter* sequence, and representative

operational taxonomic unit, in previous 16S rRNA gene amplicon analyses of the KB-1-derived

chlorinated benzene-fed cultures. The 468 bp sequence of ASV2 is a 100% match to the

corresponding region of the partial 16S rRNA gene (1508 bp, GI: 1221108851) of *Dehalobacter*

sp. TCB1 (referred to as KB1_124TCB1 in Puentes Jácome and Edwards [26]). Two additional

ASVs, at a relative abundance lower than 0.1%, were also detected in the bottles inoculated with

culture II, yet these ASVs were not consistently found among all inoculated bottles. These ASVs

might represent other *Dehalobacter* populations that exist in culture II at very low abundances.

The mother culture of culture II contains multiple *Dehalobacter* strains,[26, 45] so it is not unusual

to detect other ASVs that likely belong to low-abundance populations.

Implications for HCH, MCB, and benzene contamination at field sites

In this study, we have demonstrated for the first time that lindane (γ-hexachlorocyclohexane) can

be anaerobically biotransformed by sequential bioaugmentation using specialized microbial

enrichment cultures. The sequential treatment resulted in the biotransformation of $73 \pm 3\%$ of the

added lindane all the way to the non-toxic end products. Not only lindane, but also toxic MCB

and carcinogenic benzene were biodegraded by the anaerobic enrichment cultures. Although this

study focused on the γ isomer of HCH, this approach is applicable to other HCH isomers since

enrichment cultures that dechlorinate α-, β-, and δ-HCH to MCB and benzene are also currently

studied in the lab.[21] The success of the different biotransformation steps in this lab scale setting

was aided by the sequential nature of the treatment. Keeping the three transformation steps

separate could possibly have minimized any detrimental microbial interactions between the

different cultures, such as competition for the different electron donors or acceptors present. The

enrichment cultures described here have been studied extensively in the lab and in-house

techniques for culture growth and maintenance have improved during several years of

cultivation. Specifically, benzene-degrading culture III, has recently been evaluated as an

inoculum with different field site materials (soil and groundwater) and is undergoing field scale

pilot tests. At the field scale, HCH remediation could be tested sequentially, analogous to the lab

scale, taking into account the spatial and temporal gradients at the contaminated site and

different rates of transformation. This study further establishes the key role of *Dehalobacter* spp.

as chlorobenzene- and HCH-organohalide-respiring bacteria and demonstrates that sequential

treatment with specialized anaerobic cultures may be explored at field sites in order to address

legacy soil and groundwater contamination with HCH.

Acknowledgements

The authors would like to thank: Shen Guo for conducting GC sampling and DNA extraction

during phase III of the biotransformation experiment; Dr. Courtney Toth for performing the

preliminary steps in the QIIME 2 pipeline for the analysis of the 16S rRNA amplicon sequencing

data; and Suly Rambinaising for performing the qPCR runs with the

Desulfuromonadales/*Geobacter* primer. L. P. J. was supported by the NSERC CREATE

RENEW program [180804567], the Government of Ontario through the Ontario Graduate

Scholarship program, the Ontario Research Fund INTEGRATE project [ORF-RE05-WR-01],

and the Genomic Applications Partnership Program GAPP [OGI-102].

References

1.   Santos, A.; Fernandez, J.; Guadano, J.; Lorenzo, D.; Romero, A., Chlorinated organic compounds in liquid wastes (DNAPL) from lindane production dumped in landfills in Sabinanigo (Spain). *Environ. Pollut.* **2018,** *242*, (Pt B), 1616-1624.

2.   Fernández, J.; Arjol, M.; Cacho, C., POP-contaminated sites from HCH production in Sabiñánigo, Spain. *Environmental Science and Pollution Research* **2013,** *20*, (4), 1937-1950.

3.   Heidrich, S.; Schirmer, M.; Weiss, H.; Wycisk, P.; Grossmann, J.; Kaschl, A., Regionally contaminated aquifers - toxicological relevance and remediation options (Bitterfeld case study). *Toxicology* **2004,** *205*, (3), 143-155.

4.   Zhang, G.; Parker, A.; House, A.; Mai, B.; Li, X.; Kang, Y.; Wang, Z., Sedimentary Records of DDT and HCH in the Pearl River Delta, South China. *Environ. Sci. Technol.* **2002,** *36*, (17), 3671-3677.

5.   Laquitaine, L.; Durimel, A.; de Alencastro, L. F.; Jean-Marius, C.; Gros, O.; Gaspard, S., Biodegradability of HCH in agricultural soils from Guadeloupe (French West Indies): identification of the lin genes involved in the HCH degradation pathway. *Environmental Science and Pollution Research* **2016,** *23*, (1), 120-127.

6.   Vijgen, J.; Weber, R.; Lichtensteiger, W.; Schlumpf, M., The legacy of pesticides and POPs stockpiles-a threat to health and the environment. *Environmental Science and Pollution Research* **2018,** *25*, (32), 31793-31798.

7.   Vijgen, J.; Aliyeva, G.; Weber, R., The Forum of the International HCH and Pesticides Association-a platform for international cooperation. *Environmental Science and Pollution Research* **2013,** *20*, (4), 2081-2086.

8.   Vijgen, J.; Abhilash, P. C.; Li, Y. F.; Lal, R.; Forter, M.; Torres, J.; Singh, N.; Yunus, M.; Tian, C. G.; Schaffer, A.; Weber, R., Hexachlorocyclohexane (HCH) as new Stockholm Convention POPs-a global perspective on the management of Lindane and its waste isomers. *Environmental Science and Pollution Research* **2011,** *18*, (2), 152-162.

9.   Lindane Compound Summary. https://pubchem.ncbi.nlm.nih.gov/compound/727 (October 14, 2020).

10.   Nantel, A. J. Lindane. http://www.inchem.org/documents/pims/chemical/pim859.htm#3.%20PHYSICO-CHEMICAL%20PROPERTIES (October 10, 2020).

11.   Lindane Pesticide Information Profile http://pmep.cce.cornell.edu/profiles/extoxnet/haloxyfop-methylparathion/lindane-ext.html (October 14, 2020).

12.   Sang, S. P., S.; Cuddeford, V. Lindane – A review of toxicity and environmental fate. http://chm.pops.int/Portals/0/docs/from_old_website/documents/meetings/poprc/submissions/Comments_2006/wwf/WWF%20canada.pdf (September 16, 2020).

13.   Bhatt, P.; Kumar, M. S.; Chakrabarti, T., Fate and Degradation of POP-Hexachlorocyclohexane. *Crit. Rev. Environ. Sci. Technol.* **2009,** *39*, (8), 655-695.

14.   Hung, H.; Halsall, C. J.; Blanchard, P.; Li, H. H.; Fellin, P.; Stern, G.; Rosenberg, B.,
Temporal trends of organochlorine pesticides in the Canadian Arctic atmosphere. *Environ. Sci.*
Technol. **2002,** *36*, (5), 862-8.

15.   Harner, T.; Shoeib, M.; Kozma, M.; Gobas, F. A. P. C.; Li, S. M.,
Hexachlorocyclohexanes and Endosulfans in Urban, Rural, and High Altitude Air Samples in the
Fraser Valley, British Columbia: Evidence for Trans-Pacific Transport. *Environ. Sci. Technol.*
2005, *39*, (3), 724-731.

16.   Langenhoff, A. A.; Staps, S. J.; Pijls, C.; Rijnaarts, H. H., Stimulation of
hexachlorocyclohexane (HCH) biodegradation in a full scale in situ bioscreen. *Environ. Sci.*
Technol. **2013,** *47*, (19), 11182-8.

17.   Phillips, T. M.; Seech, A. G.; Lee, H.; Trevors, J. T., Biodegradation of
hexachlorocyclohexane (HCH) by microorganisms. *Biodegradation* **2005,** *16*, (4), 363-392.

18.   Boyle, A. W.; Haggblom, M. M.; Young, L. Y., Dehalogenation of lindane (gamma-
hexachlorocyclohexane) by anaerobic bacteria from marine sediments and by sulfate-reducing
bacteria. *FEMS Microbiol. Ecol.* **1999,** *29*, (4), 379-387.

19.   Van Doesburg, W.; Van Eekert, M. H. A.; Middeldorp, P. J. M.; Balk, M.; Schraa, G.;
Stams, A. J. M., Reductive dechlorination of ß-hexachlorocyclohexane (ß-HCH) by a
Dehalobacter species in coculture with a Sedimentibacter sp. *FEMS Microbiol. Ecol.* **2005,** *54*,
(1), 87-95.

20.   Elango, V.; Kurtz, H. D., Jr.; Anderson, C.; Freedman, D. L., Use of gamma-
hexachlorocyclohexane as a terminal electron acceptor by an anaerobic enrichment culture. *J.*
Hazard. Mater. **2011,** *197*, 204-210.

21.   Qiao, W.; Puentes Jacome, L. A.; Tang, X.; Lomheim, L.; Yang, M. I.; Gaspard, S.;
Avanzi, I. R.; Wu, J.; Ye, S.; Edwards, E. A., Microbial Communities Associated with Sustained
Anaerobic Reductive Dechlorination of alpha-, beta-, gamma-, and delta-Hexachlorocyclohexane
Isomers to Monochlorobenzene and Benzene. *Environ. Sci. Technol.* **2019**.

22.   Maphosa, F.; van Passel, M. W. J.; de Vos, W. M.; Smidt, H., Metagenome analysis
reveals yet unexplored reductive dechlorinating potential of Dehalobacter sp. E1 growing in co-
culture with Sedimentibacter sp. *Environmental Microbiology Reports* **2012,** *4*, (6), 604-616.

23.   Bashir, S.; Kuntze, K.; Vogt, C.; Nijenhuis, I., Anaerobic biotransformation of
hexachlorocyclohexane isomers by Dehalococcoides species and an enrichment culture.
Biodegradation **2018,** *29*, (4), 409-418.

24.   Kaufhold, T.; Schmidt, M.; Cichocka, D.; Nikolausz, M.; Nijenhuis, I., Dehalogenation
of diverse halogenated substrates by a highly enriched Dehalococcoides-containing culture
derived from the contaminated mega-site in Bitterfeld. *FEMS Microbiol. Ecol.* **2013,** *83*, (1),
176-188.

25.   Liang, X.; Devine, C. E.; Nelson, J.; Lollar, B. S.; Zinder, S.; Edwards, E. A., Anaerobic
Conversion of Chlorobenzene and Benzene to CH4 and CO2 in Bioaugmented Microcosms.
Environ. Sci. Technol. **2013,** *47*, (5), 2378-2385.

26.   Puentes Jácome, L. A.; Edwards, E. A., A switch of chlorinated substrate causes
emergence of a previously undetected native Dehalobacter population in an established

Dehalococcoides-dominated chloroethene-dechlorinating enrichment culture. *FEMS Microbiol.*
Ecol. **2017,** *93*, (12).

27.   Nelson, J. L.; Jiang, J. D.; Zinder, S. H., Dehalogenation of Chlorobenzenes,
Dichlorotoluenes, and Tetrachloroethene by Three *Dehalobacter* spp. *Environ. Sci. Technol.*
2014, *48*, (7), 3776-3782.

28.   Nelson, J. L.; Fung, J. M.; Cadillo-Quiroz, H.; Cheng, X.; Zinder, S. H., A Role for
Dehalobacter spp. in the Reductive Dehalogenation of Dichlorobenzenes and
Monochlorobenzene. *Environ. Sci. Technol.* **2011,** *45*, (16), 6806-6813.

29.   Luo, F.; Devine, C. E.; Edwards, E. A., Cultivating microbial dark matter in benzene-
degrading methanogenic consortia. *Environ. Microbiol.* **2016,** *18*, (9), 2923-36.

30.   Devine, C. E.; University of Toronto (Canada). Chemical Engineering and Applied
Chemistry. Identification of key organisms, genes and pathways in benzene-degrading
methanogenic cultures. 2013.

31.   Bolyen, E.; Rideout, J. R.; Dillon, M. R.; Bokulich, N. A.; Abnet, C.; Al-Ghalith, G. A.;
Alexander, H.; Alm, E. J.; Arumugam, M.; Asnicar, F.; Bai, Y.; Bisanz, J. E.; Bittinger, K.;
Brejnrod, A.; Brislawn, C. J.; Brown, C. T.; Callahan, B. J.; Caraballo-Rodríguez, A. M.; Chase,
J.; Cope, E.; Da Silva, R.; Dorrestein, P. C.; Douglas, G. M.; Durall, D. M.; Duvallet, C.;
Edwardson, C. F.; Ernst, M.; Estaki, M.; Fouquier, J.; Gauglitz, J. M.; Gibson, D. L.; Gonzalez,
A.; Gorlick, K.; Guo, J.; Hillmann, B.; Holmes, S.; Holste, H.; Huttenhower, C.; Huttley, G.;
Janssen, S.; Jarmusch, A. K.; Jiang, L.; Kaehler, B.; Kang, K. B.; Keefe, C. R.; Keim, P.; Kelley,
S. T.; Knights, D.; Koester, I.; Kosciolek, T.; Kreps, J.; Langille, M. G. I.; Lee, J.; Ley, R.; Liu,
Y.-X.; Loftfield, E.; Lozupone, C.; Maher, M.; Marotz, C.; Martin, B. D.; McDonald, D.;
McIver, L. J.; Melnik, A. V.; Metcalf, J. L.; Morgan, S. C.; Morton, J.; Naimey, A. T.; Navas-
Molina, J. A.; Nothias, L. F.; Orchanian, S. B.; Pearson, T.; Peoples, S. L.; Petras, D.; Preuss, M.
L.; Pruesse, E.; Rasmussen, L. B.; Rivers, A.; Robeson, I. I. M. S.; Rosenthal, P.; Segata, N.;
Shaffer, M.; Shiffer, A.; Sinha, R.; Song, S. J.; Spear, J. R.; Swafford, A. D.; Thompson, L. R.;
Torres, P. J.; Trinh, P.; Tripathi, A.; Turnbaugh, P. J.; Ul-Hasan, S.; van der Hooft, J. J. J.;
Vargas, F.; Vázquez-Baeza, Y.; Vogtmann, E.; von Hippel, M.; Walters, W.; Wan, Y.; Wang,
M.; Warren, J.; Weber, K. C.; Williamson, C. H. D.; Willis, A. D.; Xu, Z. Z.; Zaneveld, J. R.;
Zhang, Y.; Zhu, Q.; Knight, R.; Caporaso, J. G., QIIME 2: Reproducible, interactive, scalable,
and extensible microbiome data science. *PeerJ Preprints* **2018,** *6*, e27295v2.

32.   Martin, M., Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data AnalysisDO -
10.14806/ej.17.1.200 **2011**.

33.   Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S.
P., DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **2016,**
13, (7), 581-+.

34.   Bokulich, N. A.; Kaehler, B. D.; Rideout, J. R.; Dillon, M.; Bolyen, E.; Knight, R.;
Huttley, G. A.; Gregory Caporaso, J., Optimizing taxonomic classification of marker-gene
amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **2018,** *6*, (1), 90.

35.   Ulrich, A. C.; Edwards, E. A., Physiological and molecular characterization of anaerobic
benzene-degrading mixed cultures. *Environ. Microbiol.* **2003,** *5*, (2), 92-102.

36. Wagner, D. D.; Hug, L. A.; Hatt, J. K.; Spitzmiller, M. R.; Padilla-Crespo, E.; Ritalahti,
K. M.; Edwards, E. A.; Konstantinidis, K. T.; Löffler, F. E., Genomic determinants of
organohalide-respiration in Geobacter lovleyi, an unusual member of the Geobacteraceae. *BMC*
Genomics **2012,** *13*.

37. Nevin, K. P.; Holmes, D. E.; Woodard, T. L.; Covalla, S. F.; Lovley, D. R.,
Reclassification of Trichlorobacter thiogenes as Geobacter thiogenes comb. nov. *Int. J. Syst.*
Evol. Microbiol. **2007,** *57*, 463-466.

38. Sung, Y.; Fletcher, K. F.; Ritalaliti, K. M.; Apkarian, R. P.; Ramos-Hernandez, N.;
Sanford, R. A.; Mesbah, N. M.; Loffler, F. E., Geobacter lovleyi sp nov strain SZ, a novel metal-
reducing and tetrachloroethene-dechlorinating bacterium. *Appl. Environ. Microbiol.* **2006,** *72*,
(4), 2775-2782.

39. Coates, J. D.; Bhupathiraju, V. K.; Achenbach, L. A.; McInerney, M. J.; Lovley, D. R.,
Geobacter hydrogenophilus, Geobacter chapellei and Geobacter grbiciae, three new, strictly
anaerobic, dissimilatory Fe(III)-reducers. *Int. J. Syst. Evol. Microbiol.* **2001,** *51*, 581-588.

40. Röling, W. F. M., The Family Geobacteraceae. In *The Prokaryotes: Deltaproteobacteria*
and Epsilonproteobacteria, Rosenberg, E.; DeLong, E. F.; Lory, S.; Stackebrandt, E.;
Thompson, F., Eds. Springer Berlin Heidelberg: Berlin, Heidelberg, 2014; pp 157-172.

41. Galperin, M. Y., Genome Diversity of Spore-Forming Firmicutes. *Microbiol Spectr* **2013,**
1, (2), TBS-0015-2012.

42. Tang, S. Q.; Gong, Y. C.; Edwards, E. A., Semi-Automatic In Silico Gap Closure
Enabled De Novo Assembly of Two Dehalobacter Genomes from Metagenomic Data. *Plos One*
2012, *7*, (12).

43. Holliger, C.; Hahn, D.; Harmsen, H.; Ludwig, W.; Schumacher, W.; Tindall, B.;
Vazquez, F.; Weiss, N.; Zehnder, A. J. B., Dehalobacter restrictus gen. nov. and sp. nov., a
strictly anaerobic bacterium that reductively dechlorinates tetra- and trichloroethene in an
anaerobic respiration. *Arch. Microbiol.* **1998,** *169*, (4), 313-321.

44. Robertson, W. J.; Bowman, J. P.; Franzmann, P. D.; Mee, B. J., Desulfosporosinus
meridiei sp. nov., a spore-forming sulfate-reducing bacterium isolated from gasolene-
contaminated groundwater. *Int. J. Syst. Evol. Microbiol.* **2001,** *51*, (Pt 1), 133-40.

45. Puentes Jácome, L. A. Anaerobic Biodegradation of Chlorinated Benzenes and
Hexachlorocyclohexane by Mixed Microbial Cultures Derived from Contaminated Field Sites.
University of Toronto, Toronto, 2019.