Nitric Oxide Regulates the *Ralstonia solanacearum* Type 3 Secretion System

Connor G. Hendrich[1], Alicia N. Truchon[1], Beth L. Dalsing[1,2], and Caitilyn Allen[1*]

[1] Department of Plant Pathology, University of Wisconsin – Madison, Madison,

Wisconsin, USA

[2] Present address: Syngenta, Durham, NC

[*] Corresponding author: Caitilyn Allen, E-mail: callen@wisc.edu

Keywords: *Ralstonia solanacearum*, Type III Secretion, nitric oxide, bacterial wilt,

vascular wilt, tomato

Funding: National Sciences Foundation grant IOS 1258082.

Abstract

Ralstonia solancearum causes bacterial wilt disease on diverse plant hosts. *R.*

solanacearum cells enter a host from soil or infested water through the roots, then

multiply and spread in the water-transporting xylem vessels. Despite the low nutrient

content of xylem sap, *R. solanacearum* grows very well inside the host, using

denitrification to respire in this hypoxic environment. *R. solanacearum* growth *in planta*

also depends on the successful deployment of protein effectors into host cells via a

Type III Secretion System (T3SS). The T3SS is absolutely required for *R. solanacearum*

virulence, but it is metabolically costly and can trigger host defenses. Thus, the

pathogen's success depends on optimized regulation of the T3SS. We found that a

byproduct of denitrification, the toxic free-radical nitric oxide (NO), positively regulates

the *R. solanacearum* T3SS both *in vitro* and *in planta*. Using chemical treatments and

R. solanacearum mutants with altered NO levels, we show that the expression of a key

T3SS regulator is induced by NO in culture. Analyzing the transcriptome of *R.*

solanacearum responding to varying levels of NO both in culture and *in planta* revealed

that the T3SS and effectors were broadly upregulated with increasing levels of NO. This

regulation was specific to the T3SS and was not shared by other stressors. Our results

suggest that *R. solanacearum* experiences an NO-rich environment in the plant host

and may use this NO as a signal to activate T3SS during infection.

Keywords: *Ralstonia solanacearum*, Type III Secretion, nitric oxide, bacterial wilt,

vascular wilt, tomato

Introduction

Ralstonia solancearum (*Rs*) is a Gram-negative soil-borne betaproteobacterium

that causes bacterial wilt disease on a wide range of plant hosts, including important

crops like tomato, potato, and banana (Elphinstone 2005). The bacteria survive in and

are spread through infected soil and water, infecting host roots through wounds and

natural openings (Vasse et al. 1994). Once inside, *R. solanacearum* cells move to the

xylem, where they can grow and spread throughout the plant. Although this bacterium is

slow growing and sensitive to stress when grown *in vitro*, in plant xylem vessels *R.*

solanacearum is a formidable force. *R. solanacearum* quickly grows to high cell

densities in xylem even though this habitat is low in oxygen and nutrients, and

accessible to host defenses (Genin and Denny 2012). As it grows, *R. solanacearum*

produces an arsenal of virulence factors, including highly mucoid extracellular polymeric

substances (EPS), a consortium of cell wall degrading enzymes, and dozens of protein

effectors that are injected into host cells via the Type III Secretion System (T3SS)

(Genin and Denny 2012). Water transport in infected plants is eventually blocked by a

combination of host-produced gels and tyloses, degradation of vessel walls, EPS, and

the sheer mass of bacterial cells. Without sufficient water, plants wilt and die while *R.*

solanacearum cells escape back into the soil, where they can find another host

(Swanson et al. 2007).

The lethality of bacterial wilt disease combined with its wide host range and a

lack of effective treatments make *R. solanacearum* an important agricultural problem

around the world (Yuliar et al. 2015; Elphinstone 2005). Successful disease

management depends on understanding *R. solanacearum* biology, but the pathogen

behaves very differently in culture than it does in its natural environment (Jacobs et al.

2012). In particular, the T3SS, which is absolutely essential for virulence and subject to

a complex regulatory cascade, is affected by the plant environment in ways that remain

poorly understood (Poueymiro and Genin 2009). The PhcA quorum sensing system

upregulates T3 secretion in culture, but not *in planta* (Génin et al. 2005; Jacobs et al.

2012; Monteiro et al. 2012a; Khokhani et al. 2017). An unidentified 'diffusible factor'

increases T3SS expression *in planta* via the HrpG global regulator (Genin and Denny

2012).

We have discovered that the diffusible free radical nitric oxide (NO) plays a role

in *R. solanacearum* life inside plants. Xylem sap is rich in nitrate, which *R.*

solanacearum uses as an alternate electron acceptor to respire in the hypoxic xylem

environment (Dalsing et al. 2015). The *R. solanacearum* strain GMI1000 reduces nitrate

completely to dinitrogen gas by means of denitrifying respiration. The genes required for

this process are highly expressed during infection (Jacobs et al. 2012; Khokhani et al.

2017). However, denitrifying respiration produces NO as an intermediate. In large

quantities, NO is toxic and strongly inhibits *R. solanacearum* growth. The pathogen

manages this threat by detoxifying NO with the nitric oxide reductase NorB and the

flavohemoglobin HmpX, which are both highly expressed *in planta* and required for full

virulence (Dalsing et al. 2015; Jacobs et al. 2012). Plants use NO extensively for both

signaling and defense, and can produce NO through multiple different pathways

(Scheler et al. 2013; Delledonne et al. 1998; Simontacchi et al. 2015; Wilson et al. 2008;

Leitner et al. 2009). Thus, *R. solanacearum* likely encounters NO during plant

pathogenesis.

In this study, we identify a link between NO and regulation of the *R.*

solanacearum T3SS. We first show that T3SS gene expression increases when *R.*

solanacearum is actively denitrifying. We then show that this effect depends specifically

on the presence of NO. Based on these results, we conducted a transcriptomics

experiment to determine the effects of NO on *R. solanacearum* gene expression in

culture and *in planta* and show that levels of many transcripts, including 107 T3SS-

related genes, are altered by changes in NO levels.

Results

Nitric oxide changes expression of the *R. solanacearum* T3SS

Regulation of the T3SS in *R. solanacearum* is multi-layered and incompletely

understood (Fig 1.A). Because *R. solanacearum* actively denitrifies *in planta*, we

wondered if this pathway or its products affect T3SS expression during plant infection

(Dalsing et al. 2015; Jacobs et al. 2012). To test this hypothesis, we used an *R.*

solanacearum reporter strain carrying a fusion of the *lux* operon to the promoter of *hrpB*,

which encodes a key T3SS regulator (Genin and Denny 2012). Transcription of *hrpB*

increased >3-fold when the bacterium was denitrifying in culture (Fig. 1B).

An intermediate product of denitrification is the highly reactive free radical NO.

Because NO is a common signaling molecule in eukaryotes, we decided to determine if

it was responsible for *hrpB* activation observed during denitrification. To block the

effects of NO, we used the NO-scavenging chemical cPTIO (Pfeiffer et al. 1997). While

cPTIO is commonly used to both detect and deplete NO from solutions, it can also be

reduced or inactivated by factors other than NO (D'Alessandro et al. 2013). To mitigate

cPTIO effects unrelated to NO, we tested its ability to affect *hrpB* expression in two

conditions. First, we incubated *R. solanacearum* in denitrification-promoting conditions

with and without cPTIO. We measured *hrpB* expression in RNA extracted from these

cultures at 16 hpi, near the peak of *hrpB::lux* expression. In these conditions, NO

scavenging by cPTIO slightly but significantly reduced *hrpB* expression (Fig. 1C). We

then tested the effect of a host on the interaction between *hrpB* expression and NO. We

incubated a cell suspension of R. solanacearum *hrpB::lux* with tomato seedlings for 24

hours with or without cPTIO. As expected, the presence of a host plant activated *hrpB*

gene expression. However, adding cPTIO to the mixture completely reversed this effect

(Fig. 1D). Together, these results indicated that the denitrifying respiration-dependent

increase in *hrpB* expression observed in culture is specifically triggered by NO. Further,

this result suggests that NO could be the unknown soluble signal that increases *hrp*

regulon expression *in planta*.

Fig. 1. Nitric oxide affects expression of the *R. solanacearum* T3SS. **A,** Summary of

known T3SS regulation in *R. solanacearum* strain GMI1000. Arrows indicate positive

regulation, barred lines indicate repression, dashed lines indicate an unknown or

uncharacterized mechanism, green text indicates plant-derived signals, and blue text

indicates metabolites. T3SS genes, such as the T3 effector *ripX*, are controlled by the

transcriptional regulator HrpB. HrpB is transcriptionally activated by HrpG and PrhG,

which are themselves controlled by the quorum-sensing regulator PhcA, but only *in*

vitro. PrhG activity also depends on three uncharacterized proteins, PrhK, PrhL, and

PrhM, through an unknown mechanism. HrpG is activated by the contact-dependent

plant cell wall sensor PrhA as well as by an unknown soluble signal present during plant

infection. **B,** Expression of a bacterial luciferase transcriptional reporter of HrpB. *R.*

solanacearum strain GMI1000 *hrpB::lux* was grown in denitrification-inducing conditions

with or without 30 mM nitrate. Both culture density (OD_{600}) and luminescence were

measured every 30 min, and luminescence was normalized to culture density. Error

bars represent standard error of the mean. Cultures grown with nitrate had significantly

increased *hrpB* expression (2-way ANOVA, *P*=0.0003). **C,** The NO scavenger cPTIO

reduces *hrpB* expression in *R. solanacearum* cells grown in denitrifying conditions.

Cultures were grown as in B with and without 1 mM cPTIO and harvested at 16 hpi.

hrpB expression was measured using qRT-PCR. Expression of *hrpB* was reduced in

cPTIO-treated cultures (*P*<0.0001, students T test) Error bars represent standard error

of the mean. **D,** The NO scavenger cPTIO decreased T3SS activity in the presence of a

host. Luminescence of *R. solanacearum hrpB::lux* was measured after 24 h incubation

in water with or without five day-old tomato seedlings, and with or without 1 mM cPTIO,

an NO-scavenger. Exposure to tomato seedlings induced *hrpB::lux* expression

(*P*=0.004, students T test). *R. solanacearum* cells exposed to both tomato seedlings

and cPTIO had luminescence indistinguishable from that of cells without tomato

seedlings. Error bars represent standard error of the mean.

Denitrifying metabolism causes broad changes in *R. solanacearum* gene

expression.

While the above results showed that NO can affect *hrpB* expression, many other

genes are involved in regulation, biosynthesis, and function of the *R. solanacearum*

T3SS. To better understand the broader effects of NO on pathogen biology and T3SS

gene expression, we sequenced the transcriptomes of *R. solanacearum* cells grown

with varying levels of NO both in culture and in infected tomato plants (Fig. 2A). In

culture, we grew the cells in 0.1% O_2 in a modified Van den Mooter medium (VDM),

which promotes denitrifying respiration (Dalsing et al. 2015). We manipulated

intracellular *R. solanacearum* NO levels genetically by using mutants lacking either the

NarG nitrate reductase (the *ΔnarG* mutant cannot denitrify and does not produce NO),

or the NorB nitric oxide reductase (the *ΔnorB* mutant accumulates NO, see

Supplementary Fig. 1). Exogenous NO levels were manipulated by treating cells with

CysNO, a nitrosothiol NO donor. Because high NO levels are toxic to *R. solanacearum*,

we controlled for general stress responses by also profiling transcriptomes of cultures

treated with H_2O_2, a non-NO source of stress. After about 18 h, growth of the *ΔnorB*

mutant was inhibited by accumulating NO, so we sampled cultures 16 hpi, when *ΔnarG*,

ΔnorB, and the wild-type strain were at similar densities (Supplementary Fig. 1). CysNO

or H_2O_2 treatments were added 4 h before sampling, at the highest concentrations that

did not significantly alter culture growth (Supplementary Fig 2A). Total RNA was

extracted from bacteria *in planta* 3 days after tomato plants were inoculated with ~2000

CFU of *R. solanacearum* through a cut petiole (Khokhani et al. 2017). This method

synchronized the infection process better than root inoculation. Bacterial colonization in

sampled tissue was standardized by extracting RNA only from stems containing

between 5×10^7 and 10^9 CFU/g stem, as determined by dilution plating a ground stem

section (Supplementary Fig. 2B). Each of the three biological replicates contained

pooled RNA from 4 or 5 plants.

B

Fig. 2. Nitric oxide causes broad changes in *R. solanacearum* gene expression. **A,**

Design of RNA-seq experiment. We extracted RNA from *R. solanacearum* grown with

altered NO levels from stems of infected tomato plants (*in planta)* and from denitrifying

VDM medium (in culture) and NO levels were increased chemically by adding the NO

donor CysNO to wild type (WT) and increased genetically by using a NO-accumulating

*Δ*norB mutant. The *ΔnarG* mutant has reduced levels of NO. H_2O_2 -treated WT was

included as a control for non-nitrosative general stress responses. **B,** Whole-

transcriptome datasets were used for principal component analysis; plot created in R

with the package Vegan (Oksanen et al. 2019). Each data point represents one

biological replicate, with three replicates per condition. Ellipses represent a 95%

confidence interval for each condition (some ellipses are so small that they appear as

lines).

Around 97% of RNA-seq reads from *in vitro* samples mapped to the *R.*

solanacearum genome (Supplementary Fig. 3A). Because tomato stem slices also

included host RNA, only 9.2% of the *in planta* RNA reads mapped to the *R.*

solanacearum genome. However, the absolute number of bacterial transcripts from the

in planta samples still exceeded 10^6 reads per replicate. Principal component analysis

clustered the *in vitro* samples together based on treatment, and the *in planta* samples

were even more tightly clustered, reflecting less variation in gene expression than the

culture samples (Fig. 2B). Genes were considered differentially expressed if they had

an adjusted P-value <0.05. A full list of gene expression values can be found in the

supplementary materials.

We validated our transcriptomic results by using qRT-PCR to measure the

expression of two regulators of T3SS, *hrpB* and *hrpG*, and one well-expressed T3E,

ripX. Expression levels of each of these genes in the *ΔnarG* and *ΔnorB* mutants

generally matched those obtained from RNA-seq analysis (Fig. 3). As with *hrpB*

expression, *hrpG* and *ripX* were both activated by NO, whether it was produced

endogenously or exogenously.

Fig. 3. Expression analysis of selected T3SS-related genes by qRT-PCR validated

RNA-seq results. RNA was extracted from *R. solanacearum* cultured in VDM media at

0.1% oxygen for 16 hours for both qRT-PCR and RNA-seq analysis. *R. solanacearum*

samples were: wild type strain GMI1000 (WT, untreated control reference), non-

denitrifying low-NO mutant *(ΔnarG*) and NO-accumulating mutant (*ΔnorB*), and WT

treated for 4h with exogenous NO (CysNO). Gene expression levels in all samples were

normalized using *serC* (Jacobs et al. 2012). Error bars indicate standard error of the

mean. Black bars, log$_2$fold-change expression of T3SS regulatory gene *hrpB;* green

bars*,* log$_2$ fold-change expression of T3SS regulatory gene *hrpG;* blue bars*,* log$_2$ fold-

change expression of Type 3-secreted effector gene *ripX. * *P*<0.05, for qRT, student's

t-test compared to gene expression level in untreated WT bacteria; for RNA-seq, FDR

using Benjamini-Hochberg.

Denitrifying conditions in culture do not replicate *in planta* conditions

Because xylem fluid contains relatively small amounts of nutrients, minimal

media has often been considered to mimic the xylem environment (Arlat et al. 1992;

Lowe-Power et al. 2018a). Because VDM is a minimal medium and *R. solanacearum*

experiences low-oxygen conditions and uses denitrifying respiration during tomato

infection, we asked if anaerobic growth in VDM would better mimic the tomato xylem

environment than rich media (Dalsing et al. 2015). However, when we examined

expression of *R. solanacearum* genes in tomato stems compared to in VDM, we found a

total of 1038 genes were differentially regulated in these conditions (Supplementary

Table 2). Most genes involved in denitrification were expressed at a higher level in VDM

than *in planta*. Other broad categories that were altered include sugar metabolism,

flagellar motility, amino acid transporters, iron uptake genes, and ribosomal proteins. In

line with previous results, T3SS-related genes were among the most highly upregulated

genes *in planta* (Jacobs et al. 2012). Overall, growth in VDM did not closely replicate

the *in planta* environment and the bacterium seemed more reliant on denitrifying

metabolism in VDM.

Altering denitrification and NO levels changes expression of denitrification and

virulence-related genes *in planta*

To determine if the effects of NO on the T3SS were specific or a part of a non-

specific general response, we examined genes differentially regulated the non-

denitrifying *ΔnarG* and NO-accumulating *ΔnorB in planta*. In the stem, 187 genes were

significantly altered by mutating *narG* and thus blocking denitrification and NO

production, while only 49 genes were altered in the NO accumulating *ΔnorB* mutant. In

ΔnarG, nine of the most significantly upregulated genes were directly involved in nitrate

sensing and uptake or were components of the nitrate reductase protein. In both *ΔnarG*

and *ΔnorB*, genes involved in motility and sulfur metabolism were decreased compared

to WT (Supplementary Table 2). Overall, this muted response highlights the robust

regulatory state of *R. solanacearum* cells growing *in planta* and shows that they can

tolerate even changes in fundamental aspects of its biology (Peyraud et al. 2018).

Consistent with the observation that the entire transcriptome of WT *R.*

solanacearum was very different *in planta* than in culture (Fig. 2aA) the transcriptomic

profiles of the denitrification mutants grown in culture were much more divergent from

that of WT, with 3,145 significantly altered genes in *ΔnarG* and 3,976 altered in *ΔnorB*.

In the non-denitrifying *ΔnarG*, the bacterium showed decreased expression in many

KEGG pathways involved in growth and metabolic activity such as purine and

pyrimidine metabolism, oxidative phosphorylation, and tRNA synthetases

(Supplementary Table 3).

In culture, genes up-regulated in the NO-accumulating *ΔnorB* mutant were

enriched in the KEGG pathways of flagellar assembly, sulfur relay system and fatty acid

degradation (Supplementary Table 3). Downregulated genes were enriched in central

metabolic KEGG pathways like oxidative phosphorylation, pyrimidine metabolism,

biosynthesis of amino acids, and fatty acid biosynthesis. In contrast to the activation of

the T3SS, CysNO-treated *R. solanacearum* cells and *ΔnorB* saw reduction in

expression of Types I, II, and VI secretion systems. Overall, the transcriptomic

responses of *Rs* cells to either increased or decreased NO did not align well with their

responses *in planta*. This is consistent with previous studies showing that cultured *Rs*

cells do not reflect biology of the pathogen in its natural habitat (Jacobs et al. 2012;

Perrier et al. 2018).

R. solanacearum encounters more nitrosative stress in VDM than *in planta*

To estimate the degree of nitrosative stress induced by our treatments, we

looked at the expression of *norB* and *hmpX*, which directly detoxify NO in *R.*

solanacearum (Farr and Kogoma 1991; Dalsing et al. 2015). Compared with *R.*

solanacearum growing in denitrifying culture conditions, *norB* and *hmpX* expression

decreased 5-fold and 20-fold *in planta*, suggesting that in this environment the

bacterium experiences lower nitrosative stress levels (Supplementary Table 2). In

striking contrast, expression of the oxidative stress-response catalase gene *katE* was

increased 35-fold *in planta*. *In planta*, neither the non-denitrifying *ΔnarG* strain nor the

NO overproducing *ΔnorB* strain showed significant changes *norB* or *hmpX* expression.

Conversely, the *ΔnorB* mutant grown in culture had an 11-fold up-regulation of the NO-

detoxifying *hmpX*. In culture, our oxidative stress control treatment with H_2O_2 increased

expression of *katE*, as expected, but also had increased expression of NO detoxifying

genes *hmpX* and *norB*. Overall, these results showed some overlap in the response to

ROS and RNS stress in culture but suggested that *R. solanacearum* cells experienced

less nitrosative stress in tomato xylem than when denitrifying in VDM culture.

Nitric oxide induced expression of T3SS-related genes both *in vitro* and *in planta*.

To determine if in addition to *hrpB*, other T3SS-associated genes are induced by

NO in culture, we measured relative expression of all 107 known *R. solanacearum*

T3SS genes under each tested condition. These genes encode T3SS structural

elements, regulators, secreted effectors, and chaperones (Fig. 4) (Cunnac et al. 2004;

Peeters et al. 2013; Van Gijsegem et al. 1995; Plener et al. 2010; Lonjon et al. 2017;

Zhang et al. 2015b, 2018, 2011; Brito et al. 1999; Marc Marenda et al. 1998; Lonjon et

al. 2015; Génin et al. 2005). Cells of the NO-accumulating *ΔnorB* mutant grown in

culture had broadly increased T3SS gene expression relative to wild type, with nine of

sixteen structural elements significantly increased. Only one structural element, *hrpF*,

trended non-significantly towards reduced expression (Fig. 4A). Five out of thirteen

known regulators of T3SS were significantly upregulated in *ΔnorB*, although expression

of the *prhA*, *prhG*, *prhO*, and the *prhKLM* operon were all reduced. Two out of the four

T3E chaperones had significantly elevated expression in *ΔnorB*, as did most T3Es. The

differential expression of effectors did not seem to correlate with their known

interactions with chaperones (Supplementary Table 4). The expression of T3SS genes

in CysNO-treated cells largely mirrored that of *norB*, with 80% of tested genes having a

similar trend. However, the magnitude of the effect was lower. T3SS gene expression in

ΔnarG was more varied, with fewer significantly differentially regulated genes than

ΔnorB. In contrast to *ΔnorB*, *ΔnarG*, and CysNO-treated cultures, H_2O_2-treated cultures

had only eight significantly changed T3SS genes (Fig. 4A). Overall, we saw a greater

induction of T3SS genes in culture by our high-NO conditions than low-NO conditions.

Fig. 4. Nitric oxide alters expression of T3SS-related genes both *in vitro* and *in planta*.

A, Heatmap showing relative expression levels of 107 T3SS-associated genes in *R.*

solanacearum cells growing in rich medium with varying NO levels. All fold-change

numbers are compared to levels in untreated wild-type (WT) cells. Gene annotations,

fold change, and significance levels can be found in the supplementary materials. NO

levels were modified genetically using mutants lacking *narG* (deleting this nitrate

reductase reduces endogenous NO) and *norB* (deleting this nitric oxide reductase

increases endogenous NO). NO levels were increased chemically by adding 1 mM

CysNO. Cells treated with 100 μM hydrogen peroxide (H_2O_2) were a non-NO control to

show general stress responses. **B,** Relative expression levels of 107 T3SS-associated

genes in *R. solanacearum* cells growing *in planta*. About 2000 CFU of WT *R.*

solanacearum, ΔnarG, or *ΔnorB* were introduced into 17-day old Bonny Best tomato

plants through a cut petiole and RNA was harvested from a stem slice 3 days after

inoculation. **C,** Expression of most T3SS-associated genes in *R. solanacearum* strain

GMI1000 was enhanced when the bacterium grew in tomato plants relative to when *R.*

solanacearum cells multiplied *in vitro* under denitrifying conditions (VDM medium, 0.1%

O_2).

To determine if altering NO levels affects *R. solanacearum* T3SS gene

expression in a biologically relevant environment with all natural T3SS regulating inputs,

we examined changes in T3SS gene expression in *ΔnarG* and *ΔnorB in planta*.

Compared to *R. solanacearum* grown in culture, wild type *R. solanacearum in planta*

had much higher expression of almost every T3SS-associated gene (Fig. 3C). When

growing in tomato stems, *ΔnorB* grown *in planta* had no significantly differentially

regulated T3SS genes, possibly because relatively high levels of NO are already

present in the plant environment (Fig. 3B). In contrast, the non-denitrifying *ΔnarG*

mutant showed an overall trend of lower T3SS gene expression, with expression of nine

genes significantly reduced (Fig. 3B). These included two components of the T3SS

structure, *hrpF* and *hrcN*, the regulators *hrpB* and *prhG*, the effector chaperone *hpaG*,

and four T3SS effectors. Even in the highly T3SS-inducing *in planta* environment,

reducing *R. solanacearum*-produced NO was sufficient to reduce T3SS gene

expression in tomato xylem.

Discussion

The T3SS is essential for *R. solanacearum* virulence, as it is for many Gram-

negative pathogenic bacteria. Involving at least 16 structural proteins, this secretion

system is metabolically expensive to synthesize and operate, so the bacterium is under

strong selection pressure to express it only when it increases fitness (Genin et al. 1992;

Boucher et al. 1985). Many factors contribute to the regulation of T3SS in *R.*

solanacearum, including the presence of undefined host cell wall materials, unknown

soluble factors, unknown metabolic cues, and (in culture) quorum sensing (Zuluaga et

al. 2013; Brito et al. 1999; Aldon et al. 2000; Genin and Denny 2012; Zhang et al. 2018,

2011; Senuma et al. 2020). While many T3SS regulators have been described, open

questions remain, such as why T3SS is repressed in rich media and whether specific

host cell wall components activate PrhA. In addition, many genes that affect T3SS gene

expression do so via unknown mechanisms (Zhang et al. 2011, 2015b, 2018). The

results presented here suggest that NO participates in this complex regulatory network.

Despite its toxicity at high concentrations, NO is widely used as a signaling

molecule in all domains of life (Asgher et al. 2017; Astuti et al. 2018; Simmonds et al.

2014). Many effects of NO are mediated by NO-dependent protein post-translational

modifications, including binding transition metals, S-nitrosylation of cysteine, and

tyrosine nitration (Simmonds et al. 2014; Cooper 1999; Leitner et al. 2009; Wünsche et

al. 2011; Kolbert et al. 2017; Zhang et al. 2015a; Russwurm and Koesling 2004). While

most well-characterized mechanisms of NO signaling have been described in

eukaryotes, examples of NO signaling in bacteria have also been identified. Species in

Legionella, *Shewanella*, *Vibrio*, and *Silicibacter* have been shown to use H-NOX

proteins to sense NO and regulate biofilm formation (Rao et al. 2015; Henares et al.

2012; Liu et al. 2012; Network et al. 2012; Carlson et al. 2010). NO also contributes to

biofilm regulation in *Pseudomonas aeruginosa* via the protein NosP (Hossain and Boon

2017). Both H-NOX proteins and NosP bind NO using iron centers, but no orthologues

of these genes are apparent in the *R. solanacearum* genome. The first example of S-

nitrosylation in bacteria was the oxidative stress response protein OxyR in *E. coli* (Seth

et al. 2012). OxyR regulates a distinct set of genes when a key cysteine is nitrosylated

rather than oxidized. In addition to this, nitrosative stress damage triggers S-

nitrosylation of the *Salmonella enterica* redox sensor SsrB and affects the S-nitroso

proteome of *Mycobacterium tuberculosis* (Rhee et al. 2005; Husain et al. 2010).

Although we show NO activates a key virulence factor in *R. solanacearum*, its specific

mechanism of action is not yet known. Using *in silico* predictions of NO-dependent post-

translational modifications, we could identify and selectively modify potentially important

residues in T3SS regulatory proteins to determine the mechanism of NO-dependent

T3SS activation (Xue et al. 2010; Liu et al. 2011).

We found that expression of T3SS-associated genes was enhanced by the

presence of a plant host and by exogenous NO released by the NO donor CysNO, and

the plant-mediated induction was suppressed by the NO scavenger cPTIO (Fig. 1C).

T3SS gene expression also increased in response to nitric oxide produced by *R.*

solanacearum itself, both at natural levels generated by denitrifying respiration and at

enhanced levels caused by deleting *norB*. As previously reported (Monteiro et al.

2012a; Jacobs et al. 2012; Khokhani et al. 2017), *R. solanacearum* cells infecting

tomato plants highly expressed most T3SS genes, but reducing *R. solanacearum*-

produced NO by preventing denitrifying respiration with a Δ*narG* mutation significantly

decreased expression of some T3SS-related genes *in planta*. These included genes

encoding structural components, HrpF and HrcN, key regulators HrpB and PrhG, the

chaperone HpaG, and four T3-secreted effectors (Fig. 4B). This was especially striking

because overall gene expression in Δ*narG* was otherwise very similar to that of wild

type. These results raise the possibility that *R. solanacearum* uses the presence of NO

as an indirect indicator of the host xylem environment. There was no increase in T3SS

gene expression in the NO-overproducing Δ*norB* mutant, which may indicate that T3SS

genes are already so highly expressed *in planta* that the addition of more NO does not

further increase this elevated level of expression.

408   Besides its effects on the T3SS, altering NO concentrations in culture elicited

broad changes in gene expression. We reduced NO levels using the non-denitrifying

mutant *ΔnarG*. However, preventing the cells from using the denitrification pathway has

significant effects on the biology and metabolic state of the cells, making the

interpretation of these results challenging. For example, the *ΔnarG* cells grown in

culture showed evidence of decreases in growth rate and metabolic activity, indicated

by decreased expression in the biosynthetic pathways of nucleotides, reduced

expression of the oxidative phosphorylation system, and decreased production of many

tRNAs. Treating the cells with NO, either with exogenously applied CysNO or with the

NO-overproducing *ΔnorB*, caused similar changes in many metabolic pathways in

culture. However, a primary force in culture was the stress caused by increased NO

concentrations. This manifested in decreased expression of denitrification and nitrate

uptake functions and an increased expression of the NO-detoxifying enzyme *hmpX*.

This stress was associated with decreased expression of most *R. solanacearum*

secretion systems. However, expression of the T3SS showed the opposite trend,

indicating the specificity of NO-dependent T3SS regulation.

424   When denitrification mutants grew *in planta* they had far fewer alterations in gene

expression relative to wild type than when the mutants grew in culture. This suggests

that inputs from the plant host help *R. solanacearum* control its metabolism and

behaviors. Overall, the cells grown *in planta* seem to encounter less nitrosative stress

than cells grown in denitrifying conditions in culture. This could mean that NO stress is

not as important *in planta* than in culture, but it does not take into account differences in

the microenvironments of the xylem environment. Bacteria grown in culture experience

a uniform environment with consistent NO stress. In contrast, *R. solanacearum* cells

colonizing plant stems occupy multiple different microenvironments, which likely contain

varying NO levels.

A previous study compared gene expression between *R. solanacearum* grown in

rich media and *R. solanacearum* living in tomato xylem (Jacobs et al. 2012). Although

there were methodological differences between that experiment and this one, we saw

many similarities in the DEGs present between in culture and *in planta R.*

solanacearum. In both experiments, many of the most highly upregulated genes are

involved in the T3SS. Of the genes that were upregulated *in planta* in both experiments,

31 out of 141 (~22%) encoded structural components, regulators, or effectors of the

T3SS. Three exceptions to this trend were *prhKLM*. Mutating any of these genes leads

to a loss of T3SS gene expression, yet all three were highly downregulated *in planta*

compared to either CPG or VDM (Supplementary Table 4) (Zhang et al. 2011; Jacobs et

al. 2012). Several other genes that were highly upregulated *in planta* in both

experiments encode metabolism of sucrose or myo-inositol, two carbon sources

important for *R. solanacearum* during infection (Lowe-Power et al. 2018b; Hamilton et

al. 2020; Xian et al. 2020). Additionally, genes involved in flagellar motility were

upregulated in both experiments. A similar number of genes (137) were downregulated

in planta in both experiments. These included several genes predicted to be involved in

amino acid transport and metabolism and the superoxide dismutase genes *sodB* and

sodC.

Components of the denitrification pathway were among the most highly

upregulated *R. solanacearum* genes *in planta* compared to in rich medium. Consistent

with this observation, *R. solanacearum* mutants unable to denitrify were reduced in

growth *in planta* and in bacterial wilt virulence (Dalsing et al. 2015). However, the same

genes were around 20-fold more highly expressed in denitrifying culture than *in planta*.

Further, the RNS-responsive NO-detoxifying genes *norB* and *hmpX* were upregulated *in*

planta compared to rich media but were expresses much less *in planta* than in

denitrifying conditions in culture. This indicates that *Rs* experiences more nitrosative

stress when growing in denitrifying culture conditions than when growing in the plant

host, which in turn is more nitrosatively stressful than aerobic growth in rich media.

These data add to the strong evidence that *R. solanacearum* gene expression and

metabolism are very different in the biologically realistic host environment than in

culture. This may be due to the complex structure of the xylem environment, where host

inputs modulate *R. solanacearum* gene expression, the ability to form defensive biofilms

and multiple microenvironments with different nutrient or oxygen concentrations, or

because the xylem is a flowing environment, constantly bringing new nutrients into the

bacterial habitat. The bacterium's behavior in culture should be regarded as artifactual

unless proven otherwise.

We attempted to directly measure the concentrations of NO in healthy and

infected tomato xylem collected from fluid pooling on a cut stem using the NO-reactive

fluorescent dye DAF-FM, electro paramagnetic resonance (EPR), and an NO-sensitive

microsensor (Namin et al. 2013; Kleschyov et al. 2007; Yoshioka et al. 1996; Schreiber

et al. 2008). While a fluorescent signal was observed when DAF-FM was added to

xylem sap, no signal was detected through EPR using the NO spin trap cPTIO or with

the microsensor (Unisense, Aarhus, Denmark), suggesting that the DAF-FM

fluorescence was not specific to NO. These results highlight the need to measure NO

using multiple methods in biological systems (Gupta and Igamberdiev 2013). This

failure to detect NO in tomato xylem may not reflect true levels of NO in an intact plant.

We collected xylem sap by de-topping tomato plants just above the cotyledons and

allowing root pressure to force xylem fluid to pool on the cut stem. Any material exuded

in the first two minutes after cutting was removed in an attempt to limit the amount of

cellular debris and phloem in the collected material. This method, while useful for

obtaining sap to analyze for stable metabolites may not work with a transient metabolite

like NO (Lowe-Power et al. 2018a). Cutting the stems stimulates production of reactive

oxygen species, which quickly react with NO to form other reactive nitrogen species,

potentially diluting any NO signal (Milling et al. 2011; Beckman and Koppenol 1996; Van

Faassen and Vanin 2007). Furthermore, this approach can only measure NO levels in

bulk xylem sap. The xylem is a dynamic and physically heterogenous environment.

Xylem flow, xylem structure, and *R. solanacearum*-produced biofilms may all create

gradients or microenvironments with altered NO concentrations in the xylem of intact

plants (Zimmermann, 1983; Kim *et al.*, 2016; Tran *et al.*, 2016; Lowe-Power, Khokhani

and Allen, 2018). These factors complicate direct measurement of xylem NO. Confocal

microscopy of intact infected stems using fluorescent markers to visualize *R.*

solanacearum, biofilms, and NO may eventually give a more complete understanding of

this challenging landscape.

Overall, these results reveal a strong connection between NO levels and

expression of the *R. solanacearum* T3SS. They suggest that *R. solanacearum* regulates

virulence in part by using denitrifying metabolism and the resulting NO as a signal of the

hypoxic high-nitrogen host xylem environment. Ongoing work is exploring how *R.*

solanacearum manages nitrosative stress and identify additional effects of NO on

biology of both host plant and bacterial pathogen.

Materials and Methods

Bacterial strains and culture conditions.

All strains, plasmids, and primers used in this study are listed in Supplementary

Table 1. *R. solanacearum* was cultured in either CPG or a modified VDM medium at

28°C (Hendrick and Sequeira 1984; Dalsing et al. 2015). *E. coli* was grown in Luria-

Bertani (LB) media at 37°C. As needed, cultures were supplemented with antibiotics: 25

μg/mL kanamycin or 15 μg/mL gentamicin.

Strain construction.

An *R. solanacearum* GMI1000 mutant lacking *hrpB* (RSp0873) was created

using a modified *sacB*-dependent positive selection vector, pUFR80 (Castañeda et al.

2005). The regions directly upstream and downstream of *hrpB* were amplified using the

ΔhrpBup/dwn primer pairs. These fragments were then inserted into pUFR80 digested

with *Hind*III using Gibson assembly (Sievers et al. 2011). The resulting plasmid,

pUFR80-ΔhrpB was transformed into *R. solanacearum* GMI1000 and successful

plasmid integrations were selected for kanamycin resistance. The resulting colonies

were then counter selected on CPG + 5% w/v sucrose. Successful deletions were

confirmed using PCR, sequencing, and a functional screen for the ability to induce HR

on the incompatible host *N. tabacum* (Poueymiro et al. 2009).

The *ΔhrpB* mutation was complemented by inserting a DNA fragment encoding

hrpB and *hrcC* with the ~600 bp upstream containing their native promoter into the

selectively neutral *att* site in the *R. solanacearum* chromosome using pRCK (Monteiro

et al. 2012b). The fragment was amplified using hrpBcomp_F/R and inserted into

AvRII/XbaI-digested pRCK using Gibson assembly. This vector was then transformed

into *ΔhrpB* and selected on CPG+kan.

Plant growth conditions.

Wilt-susceptible tomato cv. Bonny Best were germinated and grown in BM2 all-

purpose germination and propagation mix (Berger, Saint-Modeste, QC). Plants were

grown at 28˚C with a 12 hr photoperiod. Seedlings were transplanted after 14 days.

After transplanting, plants were watered on alternate days with 1/2 strength Hoagland

solution. Tomato seeds for axenic seedling production were sterilized by washing with

10% bleach for ten minutes, followed by two five-minute washes in 70% ethanol, then

rinsed five times with water and germinated on water agar plates.

Lux reporter assays.

To test the effect of denitrification on T3SS gene expression, *R. solanacearum*

strain GMI1000 carrying a *hrpB::lux* reporter gene fusion was grown in denitrifying

conditions as follows (Monteiro et al. 2012a). Overnight cultures grown in CPG were

resuspended in VDM with or without 30 mM potassium nitrate to an initial OD_{600} of

0.001. These cultures were grown in 200 μL volumes in clear-bottomed white-walled 96

well plates at 28°C in 0.1% O_2 with shaking. Every half hour, the Abs_{600} and

luminescence was measured using a BioTek SynergyHT microplate reader (BioTek,

Winooski, VT, USA). To determine the effects of the NO scavenger cPTIO on T3SS

gene expression, *hrpB::lux* cultures were resuspended in 24-well plates in 1 mL of

water, either with or without two five day-old sterile tomato seedlings. As appropriate,

cPTIO was added to a final concentration of 1 mM. The plates were incubated at 28°C

with shaking for ~24 hours. The pigmented liquid was removed from the seedlings and

transferred to 1.5 mL Eppendorf tubes, after which the bacteria were pelleted by

centrifugation. The pigmented supernatant was removed, and the pellets resuspended

in 200 μL of water. These suspensions were transferred to a new plate and the Abs_{600}

and luminescence was measured as detailed above.

RNA extraction for sequencing.

To profile gene expression of *R. solanacearum* at various levels of NO stress, we

extracted RNA from *R. solanacearum* grown in culture and *in planta*. For each in

condition, we used *R. solanacearum* from three separate overnight cultures as

biological replicates. For the *in vitro* samples, overnight cultures grown in CPG were

resuspended to a final OD_{600} of 0.01 in 50 mL of VDM + 30 mM potassium nitrate in

conical tubes. These cultures were incubated without shaking for 12 hours at 28°C in

0.1% O_2. Four hours before harvesting, CysNO or H_2O_2 were added to the relevant

cultures to a final concentration of 1 mM and 100 μM, respectively. CysNO was

synthesized by adding HCl to a solution containing 200 mM L-cysteine and 200 mM

sodium nitrite. The solution was incubated in the dark for ten minutes before it was

neutralized by adding sodium hydroxide to a final concentration of 200 mM. The

molarity of the solution was determined using the absorbance at 334 nm and the

extinction coefficient 90/M*cm. After 16h total incubation, sub-samples were collected

for dilution plating to determine CFU/ml and the tubes were capped and centrifuged at

room temperature for 5 min at 8000 rpm. The supernatant was removed and pellets

were frozen in liquid nitrogen. RNA extractions were carried out using a modified

version of the Quick-RNA™ MiniPrep kit (Zymo Research, Irvine, CA, USA), as follows.

Pellets were resuspended in 400 μL of ice cold TE pH 8 with 1 mg/mL lysozyme, 0.25

μL Superase Inhibitor (Ambion, Austin, TX, USA), and 80 μL of 10% SDS, vortexed for

10s, transferred to a new 2 mL tube, and shaken at ~300 rpm for 2 min. 800 μL of RNA-

Lysis lysis buffer was added, the tubes were vortexed again for 10 s, cleaned according

to the kit manufacturer's instructions, and eluted in 100 μL of water. Nucleic acid

concentrations were estimated using a nanodrop, normalized to 200 ng/uL, and cleaned

using the DNA-free DNAse kit (Invitrogen, Carlsbad, CA, USA). Samples were

incubated at 37C for 1 hour, with 2 μL more of DNAse added at 30 min. After DNAse

inactivation, samples were further cleaned by chloroform extraction, then precipitated

overnight at -20°C with 100 μM Sodium Acetate pH 5.5 and 66 % ethanol. Samples

were checked for concentration on a Nanodrop, for DNA contamination by PCR using

the qRT-PCR primers *serC*_F/R, and for RNA integrity (RIN) using an Agilent

Bioanalyzer 21000 (Agilent, Santa Clara, CA, USA). All samples had RIN values above

7.3.

591   The *in planta* samples were harvested after 21 day old Bonny Best tomatoes

were inoculated with ~2000 CFU of each bacteria strain through the cut petiole of the

first true leaf. At 3 dpi, approximately 0.1 g of stem tissue was collected from the site of

inoculation, immediately frozen in liquid nitrogen, and stored at -80°C. Another ~0.1 g of

tissue was collected from directly below the inoculation site and was ground in bead

beater tubes using a PowerLyzer (Qiagen, Hilden, Germany) for two cycles of 2200 rpm

for 90 s in each with a 4 min rest between cycles. This material was then dilution plated

to measure bacterial colonization. From each biological replicate, we chose six plants

with the most similar level of colonization. RNA was extracted from these using a

modified hot phenol-chloroform method (Jacobs et al. 2012). Between 4 and 5 individual

plants were pooled per biological replicate. Nucleic acid sample quality was checked

using a nanodrop, Agilent bioanalyzer, and qRT-PCR primers Actin_F/R. All samples

had RIN values of above 7.2.

RNA sequencing and data analysis.

All RNA was sent to Novogene (Beijing, China) for library preparation,

sequencing, and analysis. rRNA depletion, fragmentation, and library construction were

done using NEBNext® μLtra™ Directional RNA Library Prep Kit for Illumina® (NEB,

Ipswitch, MA, USA) following the manufacturer's recommendation and starting with 3 μg

of RNA per sample. After the libraries were constructed, their quality was assessed

using an Agilent Bioanalyzer 21000 system and sequenced using an Illumina platform

to generate paired-end reads. Between 19,000,000 and 40,000,000 reads were

produced per sample. Raw reads were converted to FASTQ files using Illumina

CASAVA v1.8, and then filtered for quality, removing reads with adaptor contamination,

greater than 10% uncertain nucleotides, or more than 50% of nucleotides with a Q_{pred}

less than or equal to 5. Over 95% of reads were of good quality, and were mapped to

the *R. solanacearum* GMI1000 genome

(https://www.ncbi.nlm.nih.gov/assembly/GCF_000009125.1) using Bowtie2 -2.2.3

(Langmead et al. 2009). From this, gene expression was calculated using HTseq v0.6.1

and differential gene expression was calculated using DESeq 1.18.0. P-values

calculated by DESeq were adjusted to control for the false discovery rate (FDR) using

the Benjamini-Hochberg approach. Gene Ontology enrichment was analyzed using

GOseq Release2.12 and KEGG enrichment was calculated using KOBAS v2.0

software.

Measuring T3SS expression with qRT-PCR.

Expression of T3SS genes was measured in denitrification mutants grown in 40

mL of VDM + 30 mM nitrate grown in GasPak™ EZ microaerobic pouches (BD, Franklin

Lakes, NJ, USA) at 28˚C with shaking for 18 hours starting from an OD_{600} of 0.001.

RNA extractions were carried out using a hot phenol-chloroform method (Jacobs et al.

2012). Extractions were carried out using the Quick-RNA™ MiniPrep kit (Zymo

Research, Irvine, CA, USA). cDNA synthesis was done using the Superscript VILO

cDNA synthesis kit (Invitrogen, Carlsbad, CA, USA). qPCR reactions were carried out in

10 μL volumes using 5 ng of total template using PowerUp Syber Green Master) in a

QuantStudio 5 Real-Time PCR System Mix (Applied Biosystems, Foster City, CA,

USA).

Data Availability

The data supporting this research are openly available in the Gene Expression

Omnibus at https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE160024, GEO

accession: GSE160024.

Acknowledgments

The authors would like to thank Jon Jacobs for discussion and advice throughout the

project and Max Miao for his expertise and discussion about statistics and R. CGH and

BLD were supported by NSF predoctoral fellowships. This material is based upon work

supported by the National Science Foundation Graduate Research Fellowship Program

under Grant No. DGE-1747503. Any opinions, findings, and conclusions or

recommendations expressed in this material are those of the author(s) and do not

necessarily reflect the views of the National Science Foundation. The authors declare

no competing financial interests

References

Aldon, D., Brito, B., Boucher, C., and Genin, S. 2000. A bacterial sensor of plant cell
contact controls the transcriptional induction of *Ralstonia solanacearum*
pathogenicity genes. Embo J. 19:2304–2314

Arlat, M., Gough, C. L., Zischek, C., Barberis, P. a, Trigalet, A., and Boucher, C. a.
1992. Transcriptional organization and expression of the large hrp gene cluster of
Pseudomonas solanacearum. Mol. Plant. Microbe. Interact. 5:187–193

Asgher, M., Per, T. S., Masood, A., Fatma, M., Freschi, L., Corpas, F. J., and Khan, N.
A. 2017. Nitric oxide signaling and its crosstalk with other plant growth regulators in
plant responses to abiotic stress. Environ. Sci. Pollut. Res. 24:2273–2285

Astuti, R. I., Nasuno, R., and Takagi, H. 2018. Nitric Oxide Signalling in Yeast. Adv.
Microb. Physiol. 72:29–63

Beckman, J. S., and Koppenol, W. H. 1996. Nitric oxide, superoxide, and peroxynitrite:
the good, the bad, and ugly. Am. J. Physiol. 271:C1424-37

Boucher, C. A., Barberis, P. A., Trigalet, A. P., and Demery, D. A. 1985. Transposon
mutagenesis of *Pseudomonas solanacearum*: Isolation of Tn5-induced avirulent
mutants. J. Gen. Microbiol. 131:2449–2457

Brito, B., Marenda, M., and Barberis, P. 1999. *prhJ* and *hrpG*, two new components of

the plant signal-dependent regulatory cascade controlled by PrhA in *Ralstonia*
solanacearum. Mol. Microbiol. 31:237–251

Carlson, H. K., Vance, R. E., and Marletta, M. A. 2010. H-NOX Regulation of c-di-GMP
Metabolism and Biofilm Formation in *Legionella pneumophila*. Mol. Microbiol.
77:930–942

Castañeda, A., Reddy, J., El-Yacoubi, B., and Gabriel, D. 2005. Mutagenesis of all eight
avr genes in *Xanthomonas campestris* pv. *campestris* had no detected effect on
pathogenicity, but one *avr* gene affected race specificity. Mol. Plant-Microbe
Interact. 18:1306–1317

Cooper, C. E. 1999. Nitric oxide and iron proteins. Biochim. Biophys. Acta - Bioenerg.
1411:290–309

Cunnac, S., Occhialini, A., Barberis, P., Boucher, C., and Genin, S. 2004. Inventory and
functional analysis of the large Hrp regulon in *Ralstonia solanacearum*:
identification of novel effector proteins translocated to plant host cells through the
type III secretion system. Mol. Microbiol. 53:115–128

D'Alessandro, S., Posocco, B., Costa, A., Zahariou, G., Schiavo, F. Lo, Carbonera, D.,
and Zottini, M. 2013. Limits in the use of cPTIO as nitric oxide scavenger and EPR
probe in plant cells and seedlings. Front. Plant Sci. 4:340

Dalsing, B. L., Truchon, A. N., Gonzalez-Orta, E. T., Milling, A. S., and Allen, C. 2015.
Ralstonia solanacearum uses inorganic nitrogen metabolism for virulence, ATP
production, and detoxification in the oxygen-limited host xylem environment. MBio.
6:1–13

Delledonne, M., Xia, Y., Dixon, R. a, and Lamb, C. 1998. Nitric oxide functions as a
signal in plant disease resistance. Nature. 394:585–588

Elphinstone, J. 2005. The current bacterial wilt situation: a global overview. Pages 9–28
in: Bacterial Wilt: The Disease and the *Ralstonia solanacearum* Species Complex.,
C. Allen, P. Prior, and A.C. Hayward, eds. Society, American Phytopathological, St
Paul, MN.

Van Faassen, E., and Vanin, A. F. 2007. *Nitric oxide radicals and their reactions*.
Elsevier B.V.

Farr, S. B., and Kogoma, T. 1991. Oxidative stress responses in *Escherichia coli* and
Salmonella typhimurium. Microbiol. Rev. 55:561–585

Ge, S. X., Son, E. W., and Yao, R. 2018. iDEP: An integrated web application for
differential expression and pathway analysis of RNA-Seq data. BMC
Bioinformatics. 19:1–24

Génin, S., Brito, B., Denny, T. P., and Boucher, C. 2005. Control of the *Ralstonia*
solanacearum Type III secretion system (Hrp) genes by the global virulence
regulator PhcA. FEBS Lett. 579:2077–2081

Genin, S., and Denny, T. P. 2012. Pathogenomics of the *Ralstonia solanacearum*
species complex. Annu. Rev. Phytopathol. 50:67–89

Genin, S., Gough, C. L., Zischek, C., and Boucher, C. A. 1992. Evidence that the *hrpB*
gene encodes a positive regulator of pathogenicity genes from *Pseudomonas*
solanacearum. Mol. Microbiol. 6:3065–3076

Van Gijsegem, F., Gough, C., Zischek, C., Niqueux, E., Arlat, M., Genin, S., Barberis,
P., German, S., Castello, P., and Boucher, C. 1995. The hrp gene locus of
Pseudomonas solanacearum, which controls the production of a type III secretion
system, encodes eight proteins related to components of the bacterial flagellar
biogenesis complex. Mol. Microbiol. 15:1095–1114

Gupta, K. J., and Igamberdiev, A. U. 2013. Recommendations of using at least two
different methods for measuring NO. Front. Plant Sci. 4:2–5

Hamilton, C. D., Steidl, O. R., Macintyre, A. M., Hendrich, C. G., and Allen, C. 2020.
Ralstonia solanacearum depends on catabolism of myo-inositol, sucrose, and
trehalose for virulence in an infection stage-dependent manner. MPMI.

Hanahan, D. 1983. Studies on Transformation of *Escherichia coli* with Plasmids. J. Mol.
Biol. 166:557–580

Henares, B. M., Higgins, K. E., and Boon, E. M. 2012. Discovery of a Nitric Oxide
Responsive Quorum Sensing Circuit in *Vibrio harveyi*. ACS Chem. Biol. 7:1331–
1336

Hendrick, C. A., and Sequeira, L. 1984. Lipopolysaccharide-defective mutants of the wilt
pathogen *Pseudomonas solanacearum*. Appl. Environ. Microbiol. 48:94–101

Hossain, S., and Boon, E. M. 2017. Discovery of a Novel Nitric Oxide Binding Protein
and Nitric-Oxide- Responsive Signaling Pathway in *Pseudomonas aeruginosa*.
ACS Infect. Dis. 3:454–461

Husain, M., Jones-carson, J., Song, M., Mccollister, B. D., and Bourret, T. J. 2010.
Redox sensor SsrB Cys 203 enhances *Salmonella* fitness against nitric oxide
generated in the host immune response to oral infection. Proc Natl Acad Sci U S A.
107:4–13

Jacobs, J. M., Babujee, L., Meng, F., Milling, A. S., and Allen, C. 2012. The *in planta*
transcriptome of *Ralstonia solanacearum*: conserved physiological and virulence
strategies during bacterial wilt of tomato. MBio. 3:e00114-12

Khokhani, D., Lowe-Power, T. M., Tran, T. M., and Allen, C. 2017. A Single Regulator
Mediates Strategic Switching between Attachment / Spread and Growth / Virulence
in the Plant Pathogen *Ralstonia solanacearum*. MBio. 8:1–20

Kim, M. K., Ingremeau, F., Zhao, A., Bassler, B. L., and Stone, H. A. 2016. Local and
global consequences of flow on bacterial quorum sensing. Nat. Microbiol. 1:15005

Kleschyov, A. L., Wenzel, P., and Munzel, T. 2007. Electron paramagnetic resonance
(EPR) spin trapping of biological nitric oxide. J. Chromatogr. B Anal. Technol.
Biomed. Life Sci.

Kolbert, Z., Feigl, G., Bordé, Á., Molnár, Á., and Erdei, L. 2017. Protein tyrosine nitration
in plants: Present knowledge, computational prediction and future perspectives.
Plant Physiol. Biochem. 113:56–63

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S. L. 2009. Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol.
10

Leitner, M., Vandelle, E., Gaupels, F., Bellin, D., and Delledonne, M. 2009. NO signals
in the haze. Nitric oxide signalling in plant defence. Curr. Opin. Plant Biol. 12:451–
458

Liu, N., Xu, Y., Hossain, S., Huang, N., Coursolle, D., and Gralnick, J. A. 2012. Nitric
Oxide Regulation of Cyclic di-GMP Synthesis and Hydrolysis in *Shewanella*
woodyi. Biochemistry. 51:2087–2099

Liu, Z., Cao, J., Ma, Q., Gao, X., Ren, J., and Xue, Y. 2011. GPS-YNO2: computational
prediction of tyrosine nitration sites in proteins. Mol. Biosyst. 7:1197

Lonjon, F., Lohou, D., Cazalé, A. C., Büttner, D., Ribeiro, B. G., Péanne, C., Genin, S.,
and Vailleau, F. 2017. HpaB-Dependent Secretion of Type III Effectors in the Plant
Pathogens *Ralstonia solanacearum* and *Xanthomonas campestris* pv. *vesicatoria*.
Sci. Rep. 7:1–14

Lonjon, F., Turner, M., Henry, C., Rengel, D., Lohou, D., van de Kerkhove, Q., Cazale,
A.-C., Peeters, N., Genin, S., and Vailleau, F. 2015. Comparative Secretome
Analysis of *Ralstonia solanacearum* Type 3 Secretion-Associated Mutants Reveals
a Fine Control of Effector Delivery, Essential for Bacterial Pathogenicity. Mol. Cell.
Proteomics.

Lowe-Power, T. M., Hendrich, C. G., von Roepenack-Lahaye, E., Li, B., Wu, D., Mitra,
R., Dalsing, B. L., Ricca, P., Naidoo, J., Cook, D., Jancewicz, A., Masson, P.,
Thomma, B., Lahaye, T., Michael, A. J., and Allen, C. 2018a. Metabolomics of
tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum* produces
abundant putrescine, a metabolite that accelerates wilt disease. Environ. Microbiol.

20:1330–1349

Lowe-Power, T. M., Khokhani, D., and Allen, C. 2018b. How *Ralstonia solanacearum*
Exploits and Thrives in the Flowing Plant Xylem Environment. Trends Microbiol.
26:929–942

Marc Marenda, Callard, B. B. D., Genin, S., Barberis, P., Boucher, C., and Arlat, M.
1998. PrhA controls a novel regulatory pathway required for the specific induction
of *Ralstonia solanacearum hrp* genes in the presence of plant cells. Mol. Microbiol.
27:437–453

Milling, A. S., Babujee, L., and Allen, C. 2011. *Ralstonia solanacearum* extracellular
polysaccharide is a specific elicitor of defense responses in wilt-resistant tomato
plants. PLoS One. 6

Monteiro, F., Genin, S., van Dijk, I., and Valls, M. 2012a. A luminescent reporter
evidences active expression of *Ralstonia solanacearum* type III secretion system
genes throughout plant infection. Microbiol. (United Kingdom). 158:2107–2116

Monteiro, F., Solé, M., van Dijk, I., and Valls, M. 2012b. A chromosomal insertion
toolbox for promoter probing, mutant complementation, and pathogenicity studies in
Ralstonia solanacearum. Mol. Plant. Microbe. Interact. 25:557–68

Namin, S. M., Nofallah, S., Joshi, M. S., Kavallieratos, K., and Tsoukias, N. M. 2013.
Kinetic analysis of DAF-FM activation by NO: Toward calibration of a NO-sensitive
fluorescent dye. Nitric Oxide - Biol. Chem. 28:39–46

Network, C. S., Plate, L., and Marletta, M. A. 2012. Article Nitric Oxide Modulates
Bacterial Biofilm Formation through a Multicomponent. Mol. Cell. 46:449–460

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E.,
and Wagner, H. 2019. vegan: Community Ecology Package. R package version
2.5-6. Available at: https://cran.r-project.org/package=vegan.

Peeters, N., Carrère, S., Anisimova, M., Plener, L., Cazalé, A.-C., and Genin, S. 2013.
Repertoire, unified nomenclature and evolution of the Type III effector gene set in
the *Ralstonia solanacearum* species complex. BMC Genomics. 14:859

Perrier, A., Barlet, X., Peyraud, R., Rengel, D., Guidot, A., and Genin, S. 2018.
Comparative transcriptomic studies identify specific expression patterns of
virulence factors under the control of the master regulator PhcA in the *Ralstonia*
solanacearum species complex. Microb. Pathog. 116:273–278

Peyraud, R., Cottret, L., Marmiesse, L., and Genin, S. 2018. Control of primary
metabolism by a virulence regulatory network promotes robustness in a plant

pathogen. Nat. Commun. 9:418

Pfeiffer, S., Leopold, E., Hemmens, B., Schmidt, K., Werner, E. R., and Mayer, B. 1997.
Interference of carboxy-PTIO with nitric oxide-and peroxynitrite- mediated
reactions. Free Radic. Biol. Med. 22:787–794

Plener, L., Manfredi, P., Valls, M., and Genin, S. 2010. PrhG, a transcriptional regulator
responding to growth conditions, is involved in the control of the type III secretion
system regulon in *Ralstonia solanacearum*. J. Bacteriol. 192:1011–1019

Poueymiro, M., Cunnac, S., Barberis, P., Deslandes, L., Peeters, N., Cazale-Noel, A.-
C., Boucher, C., and Genin, S. 2009. Two Type III Secretion System Effectors from
Ralstonia solanacearum GMI1000 Determine Host-Range Specificity on Tobacco.
Mol. Plant-Microbe Interact. 22:538–550

Poueymiro, M., and Genin, S. 2009. Secreted proteins from *Ralstonia solanacearum*: a
hundred tricks to kill a plant. Curr. Opin. Microbiol. 12:44–52

Rao, M., Smith, B. C., and Marletta, M. A. 2015. Nitric Oxide mediates biofilm formation
and symbiosis in *Silicibacter* sp. strain TrichCH4B. MBio. 6:1–10

Rhee, K. Y., Erdjument-Bromage, H., Tempst, P., and Nathan, C. F. 2005. S-nitroso
proteome of *Mycobacterium tuberculosis*: Enzymes of intermediary metabolism and
antioxidant defense. Proc. Natl. Acad. Sci. U. S. A. 102:467–72

Russwurm, M., and Koesling, D. 2004. NO activation of guanylyl cyclase. EMBO J.
23:4443–4450

Scheler, C., Durner, J., and Astier, J. 2013. Nitric oxide and reactive oxygen species in
plant biotic interactions. Curr. Opin. Plant Biol. 16:534–539

Schreiber, F., Polerecky, L., and De Beer, D. 2008. Nitric oxide microsensor for high
spatial resolution measurements in biofilms and sediments. Anal. Chem. 80:1152–
1158

Senuma, W., Takemura, C., Hayashi, K., Ishikawa, S., Kiba, A., Ohnishi, K., Kai, K., and
Hikichi, Y. 2020. The putative sensor histidine kinase PhcK is required for the full
expression of *phcA* encoding the global transcriptional regulator to drive the
quorum-sensing circuit of *Ralstonia solanacearum* strain OE1-1. Mol. Plant Pathol.
:1–15

Seth, D., Hausladen, A., Wang, Y.-J., and Stamler, J. S. 2012. Endogenous Protein S-
Nitrosylation in *E. coli*: Regulation by OxyR. Science (80-.). 336:470–3

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R.,
McWilliam, H., Remmert, M., Soding, J., Thompson, J. D., and Higgins, D. G. 2011.

Fast, scalable generation of high-quality protein multiple sequence alignments
using Clustal Omega. Mol. Syst. Biol. 7:539–545

Simmonds, M. J., Detterich, J. A., and Connes, P. 2014. Nitric oxide, vasodilation and
the red blood cell. Biorheology. 51:121–134

Simontacchi, M., Galatro, A., Ramos-Artuso, F., and Santa-María, G. E. 2015. Plant
Survival in a Changing Environment: The Role of Nitric Oxide in Plant Responses
to Abiotic Stress. Front. Plant Sci. 6:977

Swanson, J. K., Montes, L., Mejia, L., and Allen, C. 2007. Detection of latent infections
of *Ralstonia solanacearum* race 3 biovar 2 in geranium. Plant Dis. 91:828–834

Tran, T., MacIntyre, A., Khokhani, D., Hawes, M. C., and Allen, C. 2016. Extracellular
DNases of *Ralstonia solanacearum* modulate biofilms and facilitate bacterial wilt
virulence. Environ. Microbiol. Environ. Microbiol. Reports. 00:1–15

Vasse, J., Frey, P., and Trigalet, A. 1994. Microscopic studies of intercellular infection
and protoxylem invasion of tomato roots by *Pseudomonas solanacearum*. MPMI.
8:241–251

Wilson, I. D., Neill, S. J., and Hancock, J. T. 2008. Nitric oxide synthesis and signalling
in plants. Plant, Cell Environ. 31:622–631

Wünsche, H., Baldwin, I. T., and Wu, J. 2011. S-Nitrosoglutathione reductase (GSNOR)
mediates the biosynthesis of jasmonic acid and ethylene induced by feeding of the
insect herbivore *Manduca sexta* and is important for jasmonate-elicited responses
in *Nicotiana attenuata*. J. Exp. Bot. 62:4605–4616

Xian, L., Yu, G., Wei, Y., Rufian, J. S., Li, Y., Zhuang, H., Xue, H., Morcillo, R. J. L., and
Macho, A. P. 2020. A Bacterial Effector Protein Hijacks Plant Metabolism to
Support Pathogen Nutrition. Cell Host Microbe. :1–10

Xue, Y., Liu, Z., Gao, X., Jin, C., Wen, L., Yao, X., and Ren, J. 2010. GPS-SNO:
Computational prediction of protein s-nitrosylation sites with a modified GPS
algorithm. PLoS One. 5:1–7

Yoshioka, T., Iwamoto, N., and Ito, K. 1996. An application of electron paramagnetic
resonance to evaluate nitric oxide and its quenchers. J. Am. Soc. Nephrol. 7:961–
965

Yuliar, Nion, Y. A., and Toyota, K. 2015. Recent trends in control methods for bacterial
wilt diseases caused by *Ralstonia solanacearum*. Microbes Environ. 30:1–11

Zhang, R., Hess, D. T., Qian, Z., Hausladen, A., Fonseca, F., Chaube, R., Reynolds, J.
D., and Stamler, J. S. 2015a. Hemoglobin βCys93 is essential for cardiovascular

function and integrated response to hypoxia. Proc. Natl. Acad. Sci. U. S. A.
112:6425–6430

Zhang, Y., Kiba, A., Hikichi, Y., and Ohnishi, K. 2011. prhKLM genes of *Ralstonia*
solanacearum encode novel activators of hrp regulon and are required for
pathogenesis in tomato. FEMS Microbiol. Lett. 317:75–82

Zhang, Y., Li, J., Zhang, W., Shi, H., Luo, F., Hikichi, Y., Shi, X., and Ohnishi, K. 2018.
A putative LysR-type transcriptional regulator PrhO positively regulates the type III
secretion system and contributes to the virulence of *Ralstonia solanacearum*. Mol.
Plant Pathol. :1–12

Zhang, Y., Luo, F., Wu, D., Hikichi, Y., Kiba, A., Igarashi, Y., Ding, W., and Ohnishi, K.
2015b. PrhN, a putative marR family transcriptional regulator, is involved in positive
regulation of type III secretion system and full virulence of *Ralstonia solanacearum*.
Front. Microbiol. 6:1–12

Zimmermann, M. H. 1983. *Xylem Structure and the Ascent of Sap*. Springer-Verlag,
Berlin.

Zuluaga, A. P., Puigvert, M., and Valls, M. 2013. Novel plant inputs influencing
Ralstonia solanacearum during infection. Front. Microbiol. 4:1–7

Supplementary Table 1. Bacterial strains, plasmids, and primers

Strains	Relevant characteristics	Reference
R. solanacearum		
GMI1000	Phylotype I seq 18 strain isolated from tomato in French Guyana	(Boucher et al. 1985)
hrpB::lux	GMI1000 expressing the bacterial *lux* reporter controlled by the *hrpB* promoter, Kan[R]	(Monteiro et al. 2012 a)
ΔnarG	GMI1000 lacking *narG*, a nitrate reductase. Cannot denitrify and does not produce NO, Gm[R]	(Dalsing et al. 2015)
ΔnorB	GMI1000 lacking *norB*, a nitric oxide reductase. Can denitrify, but accumulates NO, Gm[R]	(Dalsing et al. 2015)
ΔhrpB	GMI1000 lacking *hrpB*. Avirulent and does not express a T3SS	This study
hrpB comp	*ΔhrpB* complemented.,Kan[R]	This study
E. coli		
DH5α	dlacZ, Delta M15 Delta(lacZYA-argF) U169 recA1 endA1 hsdR17(rK-mK+) supE44 thi-1 gyrA96 relA1	(Hanahan 1983)
Plasmid	**Description**	**Reference**
pUFR80	Positive sucrose selection vector, Suc[S], Kan[R]	(Castañeda et al. 2005)
pRCK-GWY	Gateway destination vector that integrates into the GMI1000 genome at the neutral *att* site; Kan[R]	(Monteiro et al. 2012 b)
pUFR80-ΔhrpB	Vector used to make unmarked *R. solanacearum ΔripB*, Suc[S], Kan[R]	This study
pRCK-hrpBco	Complementation vector containing WT *hrpB* and *hrcC*, Kan[R]	This study

mp		
Cloning Primers		
Primer	**Sequence[a]**	**Reference**
ΔhrpBup_F	5' – cgacggccagtgccaGTTCGATACCAGCGAAAGC target locus: RSp0873	This study
ΔhrpBup_R	5' – cggagggcGGCAATGCTCCTGAAGCG target locus: RSp0873	This study
ΔhrpBdwn_F	5' – gcattgccGCCCTCCGGCGGACCGCC target locus: RSp0873	This study
ΔhrpBdwn_R	5' – acctgcaggcatgcaGCTGTTGCCGACGTATTGACCGGTGC target locus: RSp0873	This study
hrpBcomp_F	5' – tgcgcgagcagggggaattgcGACAACGTCTCCGGGTTC target locus: RSp0873-RSp0874	This study
hrpBcomp_R	5' – cgaccctagtctaagatcttCTTCGGCATTCGGCAATG target locus: RSp0873-RSp0874	This study
qRT-PCR Primers		
Primer	**Sequence**	**Reference**
hrpB_F	5' – CCTGCCCGAATACGCAAATG target locus: RSp0873	This study
hrpB_R	5' – CGAATGGCGGATCAGGCGCT target locus: RSp0873	This study
hrpG_F	5' – GCAGAACGTGGAATGTGTCC target locus: R. solanacearump0852	This study
hrpG_R	5' – AGACTGAATGCCTCGGTACG target locus: RSp0852	This study
ripAA_F	5' – GGAATACAGCAACTGCGTGC target locus: RSc0608	This study
ripAA_R	5' – TTGTAGTTGCGGACCCTCAGtarget locus: RSc0608	This study
serC_F	5' – CGCGCAAATACGGTGAAGTG target locus: R. solanacearumc0903	This study
serC_R	5' – GTGCACAGATGCACGTAAGC target locus: RSc0903	This study
Actin_F	5' – TCAGCAACTGGGATGATATG target locus: BT013524	(Milling et al. 2011)

Actin_R	5' – TTAGGGTTGAGAGGTGCTTC target locus: BT013524	(Milling et al. 2011)

^a Capitalized bp anneal to the target locus, non-capitalized bp anneal to other fragments

in a Gibson assembly reaction

Supplementary Fig. 1. Growth and NO levels of *R. solanacearum* strains under

denitrifying conditions in culture. Cultures were started at a concentration of 10^6

CFU/mL in VDM containing the NO-detecting fluorescent dye DAF-FM DA at 10 μM.

Cultures were incubated in in clear-bottomed, white-walled 96-well plates at 30C for 24

h at [O$_2$] =0.1%. Culture growth (dotted lines) is shown as absorbance at 600 nm. NO

concentration (solid lines) was measured as DAF-FM fluorescence at 495/515 nm. The

ΔnorB mutant strain accumulates NO, which significantly impairs its growth.

A

B

Supplementary Fig. 2. In vitro cell densities and tomato stem colonization of samples

used in RNA-seq. **A,** Cell densities of culture samples as measured by dilution plating.

One H_2O_2-treated plate was uncountable due to plate contamination. OD_{600} values of

each culture were also collected and had comparable values. **B,** Colonization of tomato

plants used for transcriptome sequencing. Roughly 0.1 g of stem tissue was collected

from directly below the sample used for sequencing. The tissue was ground and dilution

plated to enumerate CFU/g stem.

A

B

Supplementary Fig. 3. General RNA-seq quality **A,** Total and mapped reads per RNA-

seq sample. The *in planta* samples had lower percentage of mapped reads due to the

presence of host RNA. **B,** Hierarchical clustering and expression heat map of the 1000

most variable genes in the dataset. Blue indicates higher expression and brown

indicates lower expression. All genes were normalized to the mean gene expression

across all conditions. Samples collected from *R. solanacearum* grown *in planta* are

denoted with the prefix "p". Heatmap created using iDEP.90 (Ge et al. 2018).

Supplementary Table 2 Complete list of gene expression and $\log_2$ fold change values

for all conditions included in this study. All fold-change numbers are compared to levels

in untreated wild-type (WT) cells. NO levels were modified genetically using mutants

lacking *narG* (deleting this nitrate reductase reduces endogenous NO) and *norB*

(deleting this nitric oxide reductase increases endogenous NO). NO levels were

increased chemically by adding 1 mM CysNO. Cells treated with 100 μM hydrogen

peroxide (H_2O_2) were a non-NO control to identify general stress responses. *P*-values

were calculated by DESeq were adjusted to control for the false discovery rate (FDR)

using the Benjamini-Hochberg approach.

Supplementary Table 3 KEGG enrichment of genes altered in each condition.

enrichment was calculated using KOBAS v2.0 software. *P* values were calculated using

a hypergeometric test and adjusted using the Benjamini and Hochberg FDR correction

method.

Supplementary Table 4 Annotations, Log$_2$ fold change relative to wild-type and

adjusted p-values for *R. solanacearum* T3SS related genes in response to changes in

NO concentration both *in planta* and in culture (see Figure 3 for heatmap of these data).

All fold-change numbers are compared to levels in untreated wild-type (WT) cells. *in*

planta samples are denoted by a 'p' in front of the treatment (e.g. *pnorB*). Genes

organized by their category, including structural genes, regulatory genes, chaperones,

and effectors. Effectors are subgrouped based on their known association with

chaperones (Lonjon et al. 2015).