# Interpreting the dynamic pathogenesis of Parkinson's disease by longitudinal blood transcriptome analysis

Gang Xue[1,4], Gang Wang[1,2,4], Qianqian Shi[1], Hui Wang[1], Bo-Min Lv[1], Min Gao[1,3], Xiaohui Niu[1*] and Hong-Yu Zhang[1*]

[1]Hubei Key Laboratory of Agricultural Bioinformatics, College of Informatics, Huazhong Agricultural University, Wuhan, China

[2]Department of Basic Medical Laboratory, General Hospital of the Central Command Theater of PLA, Wuhan, 430015, PR China

[3]Lab of Epigenetics and Advanced Health Technology, Space Science and Technology Institute (Shenzhen), Shenzhen 518117

[4]These authors contributed equally

*Corresponding authors: Hong-Yu Zhang (zhy630@mail.hzau.edu.cn) and Xiaohui Niu (niuxiaoh@mail.hzau.edu.cn)

**Abstract**

Achieving an improved understanding of the temporal sequence of factors involved in

Parkinson's disease (PD) pathogenesis may accelerate drug discovery. In this study,

we performed a longitudinal transcriptome analysis to identify associated genes

underlying the pathogenesis of PD at three temporal phases. We firstly found that

multiple initiator genes, which are related to processes of olfactory transduction and

stem cell pluripotency, indicate PD risk to those subjects at the prodromal phase. And

many facilitator genes involved in calcium signaling and stem cell pluripotency

contribute to PD onset. We next identified 325 aggravator genes whose expression

could lead to disease progression through damage to dopaminergic synapses and

ferroptosis via an integrative analysis with DNA methylation. Last, we made a

systematic comparison of gene expression patterns across PD development and

accordingly provided candidate drugs at different phases in an attempt to prevent the

neurodegeneration process.

## Introduction

Parkinson's disease (PD) is a common chronic neurodegenerative disorder characterized by tremor and postural instability in motor symptoms along with non-motor symptoms, including hyposmia, sleeping disorder and cognitive decline[1,2]. The primary pathological features of PD are aggregations of alpha-synuclein (α-syn) in the cytoplasm, as well as the elimination of dopaminergic neurons in the substantia nigra pars compacta (SNc)[1].

Currently, the pathogenesis of PD has not been thoroughly elucidated, and many factors have been identified to contribute to the neurodegenerative process of PD. Recently, a new conceptual model of PD pathogenesis was proposed, in which these multifaceted factors can be divided into three temporal phases, namely, triggers, facilitators, and aggravators, and they may play distinct roles at different phases of PD[3]. Triggers are agents, such as environmental toxins and viral infections, that initiate the disease process. Accompanying or following triggers, facilitators spread the pathology to the central nervous system, leading to the onset of PD motor symptoms. Facilitators are proposed to include systemic inflammation, mitochondrial dysfunction, and genetic mutations. After PD manifests, aggravators, such as impaired autophagy and neuroinflammation, can directly contribute to disease progression. This conceptual model provides a new definition for PD pathogenesis, but the framework requires further investigation and validation at the molecular level. Elucidating the temporal sequence of pathogenesis in the disease process may be helpful for applying appropriate therapeutic agents when the related factor is most

active[3].

Transcriptome sequencing is a useful technique that is employed to quantitatively

investigate genes responsible for the underlying pathogenesis. Previous transcriptomic

studies of PD examined gene expression at a single time point[4,5,6], hindering the

observation of expression changes in disease development. By monitoring the same

individual at different stages or time points, longitudinal transcriptomics analysis can

capture the gene expression patterns during the temporal phases of disease

and therefore has the potential to be employed to study the systemic evolution of

pathogenesis[7]. The Parkinson's Progression Markers Initiative (PPMI) is a landmark

longitudinal observational study to comprehensively evaluate PD and identify

biomarkers of disease progression[8]. In 2019, the PPMI reported the longitudinal RNA

resource of whole blood samples, which enabled us to determine the sequence of

pathogenesis in PD development.

In this study, we performed a longitudinal transcriptome analysis with blood

samples from the PPMI to study the dynamic pathogenesis of PD (Fig. 1). By

analyzing the subjects who developed from prodromal to PD, we first identified

initiator genes that indicated the risk of disease by survival analysis. Next, facilitator

genes were determined to contribute to the onset of PD through differential expression

analysis. Moreover, aggravator genes were identified in 252 *de novo* PD patients

observed from baseline to the third year. Causal inference and methylome-wide

association study (MWAS) provided further validation and insight into aggravation.

Based on the distinct gene expression patterns observed in these three phases, we

suggested potential therapies to be employed at the appropriate stages to prevent

neurodegeneration in PD. Notably, we demonstrated that withaferin A has the ability

to promote the differentiation of neural stem cells into dopaminergic neurons *in vitro*

experiment. The results suggest great potential of withaferin A to treat prodromal or

early stage PD patients.

**Results**

**Overview of study**

In this study, two cohorts from PPMI were utilized to interpret the dynamic

pathogenesis of PD (Fig. 1). Sixty-four prodromal subjects who had a diagnosis of

hyposmia or rapid eye movement sleep behavior disorder (RBD) or at least one motor

symptom of PD were followed from 2013 to 2019, some of whom had converted to

PD during longitudinal observation. The conversion process was employed to infer

the pathogenesis of initiation and facilitation. To investigate the pathogenesis of

disease aggravation, 252 *de novo*, untreated idiopathic PD patients were selected at

baseline and subsequently observed at 1 year, 2 years, and 3 years. In this instance,

the Movement Disorder Society-Unified Parkinson's Disease Rating Scale

(MDS-UPDRS) total score was applied to measure the clinical severity of PD.

It is difficult to obtain brain transcriptomes from PD patients, while blood

transcriptomes are more accessible and less invasive to obtain from patients, making

it easy to dynamically monitor cellular injuries that occur in PD. RNA-seq data were

obtained from whole blood samples under the stringent quality controls applied by

PPMI (Methods). Notably, ~60-70% of genes expressed in the blood are also

expressed in brain tissues, as determined by examining mRNA only, suggesting that

the blood may serve as a surrogate tissue for brain transcriptome analysis and

facilitate the identification of PD-related molecular signatures.

**Identification of initiators**

In the previous conceptual model of PD pathogenesis, triggers are agents or events

that initiate disease pathogenesis during prodromal PD. These triggers, such as

pathogens and environmental toxins, are exogenous factors that initiate the disease

process[3]. However, we attempted to infer endogenous factors at the molecular level

by examining initiator genes, which might indicate the risk of disease and are thought

to act in the prodromal stage. To identify these genes, we defined prodromal

conversion to PD as an "event" and defined time interval of conversion as the

"survival time" to perform survival analysis (Methods). Next, genes that were

significantly associated with the risk of conversion were considered initiator genes of

PD (2,714 genes, log-rank test $P \leq 0.05$, Supplementary Table 1).

Many initiator genes that influence the risk of PD onset were involved in signaling

pathways regulating pluripotency of stem cells, olfactory transduction, and hepatitis C

(Fig. 2a). In line with the triggers proposed, hepatitis C, which is an infectious

pathogen, is associated with a higher risk of developing PD[9]. In addition, the

dysfunction of olfactory transduction is strongly related to hyposmia, which is one of

the earliest non-motor symptoms of PD[10] and is consistent with the clinical symptoms

of prodromal subjects, as previously described. Together with the evidence of α-syn

aggregates in olfactory[11], these findings indicated that the olfactory system may be the

major initiator site in PD pathogenesis. Notably, *OR8D1*, one of the key genes to help

maintain normal olfactory function[12], ranks first among the initiator genes (log-rank

test, $P$ = 3.2e-05). Prodromal subjects with lower expression levels of *OR8D1* were at

high risk of converting to PD (Fig. 2b). Thus, the expression of *OR8D1* could be

employed to help early diagnosis. In PD, ~50% of nigral neurons are lost before the

onset of clinical motor symptoms[13]. According to our results, we hypothesize that the

lost neurons might not be renewed due to the damaged pluripotency of stem cells,

which can influence the neurogenesis process[14,15].

**Identification of facilitators**

Facilitators are proposed to help initiators spread PD pathology and damage the

central nervous system[3]. Unlike initiator genes, facilitator genes are thought to

contribute to the onset of PD and show different expression patterns in prodromal and

early-stage PD. To find the facilitator genes, we identified differentially expressed

genes (DEGs) by comparing the expression profiles from prodromal to confirmed PD

patients in a discovery dataset, which contained 17 RNA-seq samples of five subjects

during their natural conversion (Method, Fig. 3a). We identified 1,326 DEGs whose

changes in expression during conversion may facilitate the onset of PD ($P \leq 0.05$,

Supplementary Table 2). To assess the robustness of our results, a similar analysis was

performed in a validation dataset with 57 prodromal subjects and 390 PD subjects at

baseline. Although these two cohorts have no conversion relationship between each

other, the large-scale samples can largely eliminate the influence of differences

deriving from genetic background in our comparative research. The DEGs in the

discovery and validation datasets (false discovery rate (FDR) $\leq$ 0.01, Supplementary

Table 3) had a significant overlap (Fig. 3b), supporting the results in the discovery

dataset with limited samples. Therefore, these 1,326 DEGs refer to facilitator genes of

PD.

The facilitator genes were mostly expressed in the cerebral cortex, suggesting that

the central nervous system might have been damaged (Fig. 3c). In addition, the

facilitator genes were mainly downregulated in the calcium signaling pathway and

signaling pathways regulating pluripotency of stem cells during conversion (Fig. 3d).

Evidence has shown that changes in calcium ($Ca^{2+}$) levels may trigger a series

of events to increase SNc neuronal death sensitivity[16], including mitochondrial

dysfunction, enhanced ROS generation and damaged ATP production. $Ca^{2+}$

homeostasis is also regulated by PD-related proteins, such as α-syn, PINK-1/parkin,

leucine-rich repeat kinase 2 (LRRK2), and DJ-1 protein[16]. These findings confirmed

the facilitators of a previously conceptual model[3] and indicated that dysfunction in

$Ca^{2+}$ signaling is one of the earliest events in the pathogenesis of PD[16]. We also noted

that the pathway of stem cells pluripotency was also captured by initiator genes; thus,

the neurogenesis process was influenced in both the prodromal and early stages of PD,

which may also imply its causative role in the onset of PD. Taken together, these

findings indicated that the dysfunctions in calcium signaling and stem cell

pluripotency might be considered facilitators of PD pathogenesis.

**Identification of aggravators**

Once PD manifests, aggravators can directly promote the neurodegenerative process

and contribute to disease progression. In fact, although there are some interventions

for PD, at present, there is no agent that can halt PD progression. Most of the patients

showed aggravation within a few years after diagnosis[17]. In addition, consistent with

our observation (Supplementary Fig. 1a-b), the clinical manifestations of patients

differ greatly in the trajectory of aggravation[18], hindering efforts to identify a suitable

treatment. Although molecular mechanism underlying PD aggravation has critical

importance in the long-term prevention and treatment of this disease, it has not been

elucidated to date. Therefore, we searched for potential aggravator genes to determine

the underlying mechanisms of PD.

With the longitudinal blood transcriptome of 252 patients at four serial time points

(baseline, 1 year, 2 years, and 3 years), we constructed linear mixed effects models

(Methods) to characterize the association between gene expression change and

clinical severity during disease aggravation. Using each subject's baseline samples as

their own control, this analysis reduces the noise derived from the genetic background

of patients and improves the precision. We identified 325 genes significantly

associated with PD aggravation (FDR $\leq$ 0.05; Supplementary Table 4), 307 of which

were consistently downregulated and 18 of which were upregulated, across the entire

aggravation process (Fig. 4a). Taking downregulated aggravator genes as an example,

at early stages (baseline and 1 year), these genes were expressed at a relatively high

level, but the gene expression level decreased significantly in many patients after 2-3

years. Such an expression trend was also observed in an independent dataset from PD

patients (Supplementary Fig. 2), suggesting that the dynamic molecular functions

dominating disease aggravation may help the development of drugs to intervene in the

process. Moreover, the downregulated aggravator genes were mainly enriched in the

dopaminergic synapse pathway, cGMP-PKG signaling pathway, and ferroptosis (Fig.

4b). It is clear that the suppressed function of dopaminergic synapses can damage

dopaminergic neurons. Ferroptosis is an iron-dependent autophagic cell death

process[19], and prior research has elucidated its mechanism in PD neurodegeneration.

For example, as a key component of ferroptosis, iron generates hydroxyl radicals and

induces the oxidation of dopamine, which may lead to an oxidative environment and

thus contribute to the death of dopaminergic neurons in the SNc[20].

What remained to be confirmed was whether the changes in the expression levels

of aggravator genes were the consequences or causes of PD progression. Therefore,

we next performed cross-lagged panel analyses (Methods, Fig. 4c) to investigate the

cause-effect relationship between the 325 aggravator genes and clinical severity (i.e.,

MDS-UPDRS total score). Ten aggravator genes showed significant path coefficients

($\beta_4$) from baseline gene expression to follow-up MDS-UPDRS total score (the third

year) but did not show significant path coefficients ($\beta_2$) from baseline MDS-UPDRS

total score to follow-up gene expression, demonstrating that the changes in expression

exhibited by these genes may be the causes, rather than the consequences of PD

aggravation (Supplementary Table 5). Based on the unidirectional relationship from

baseline gene expression to follow-up clinical severity at ten genes, we further

examined the association between weighted gene expression risk score (ERS) at

baseline and MDS-UPDRS total score at follow-up and found a significant positive

correlation between them (Methods), which confirmed that these genes drive the

process of aggravation and could be utilized for disease prediction (Fig. 4d).

**Integrative analysis of aggravator genes and DNA methylation**

As is well-known, the alterations of gene expression are affected by comprehensive

factors, such as genetic variation and epigenetic changes. In this study, we further

integrated the baseline blood methylation profiles to elucidate the epigenetic

regulation related to disease aggravation. Methylome-wide association study (MWAS)

were performed to distinguish the disease signals from CpG methylation (Method).

Since methylation alterations may achieve long-term regulation of disease

development and do not cause clinical manifestations immediately[21,22]

(Supplementary Fig. 3), we constructed a linear model with the baseline methylome

and the rate of PD aggravation. A total of 1,416 CpG sites ($P \leq 0.001$, Supplementary

Table 6) were found to be associated with the rate of aggravation (Fig. 5a), and

methylation score derived from these CpG sites accounted for approximately 19% of

its variance (Method, Fig. 5b). In addition, previous research identified 138 CpGs that

significantly changed over time in PD patients with longitudinal genome-wide

methylation data[23]. Although there is no overlap between our aggravator

rate-associated CpGs and the previously identified 138 CpGs, the genes mapped by

CpGs showed significant overlap ($P$ = 7.8e-06, Fig. 5c), further indicating that these

1,416 CpGs identified by us were related to disease aggravation. Moreover, the

aberrant methylation events that occurred in many PD-related pathways overlapped

with those pathways enriched by aggravator genes, e.g., dopaminergic synapse and

ferroptosis, which led us to investigate their functional associations.

We subsequently explored the associations between our aggravator genes and these

aggravator rate-associated CpGs (Method) and found that the changes in expression in

all aggravator genes exhibited significant relationships with these CpGs ($P \leq 0.05$). In

particular, cg13298827 was positively correlated with expression changes of

*PCDHA11* located within +/-10 kb (Fig. 5d). *PCDHA11* may be involved in the

creation and maintenance of specific neuronal connections in the brain, and its

dysfunction may influence the signaling and communication between neural cells,

thereby leading to the aggravation of PD. Since some regulations may have

long-range interactions, we next integrated high-throughput chromosome

conformation capture (Hi-C) data from the brain dorsolateral prefrontal cortex[24]

(http://resource.psychencode.org) and identified 18 CpG-gene interactions in the

topologically associating domains (TADs) (Supplementary Table 7). For example,

cg17512517 may influence disease aggravation by negatively regulating expression

changes of *CCDC50* (Fig. 5d). Taken together, our results indicated that DNA

methylation may regulate the changes in expression levels of aggravator genes, which

may modulate PD progression.

**Inferred cellular changes associated with aggravation**

In addition to identifying genes associated with aggravation, we also examined which cell types were changed along with PD aggravation. In this study, we applied the xCell algorithm[25] to infer the abundance scores of 64 immune and stromal cell types from the RNA-seq expression profiles of each PD sample at baseline and after 1-3 years. Analogous to our aggravator gene identification flow, 14 cell types were found to be significantly associated with PD aggravation (FDR $\leq$ 0.05, Supplementary Fig. 4). Remarkably, neurons, although slightly increased in the third year, significantly decreased overall with disease aggravation, which is in keeping with the loss of neurons observed in the SNc of PD patients[1]. In addition, some of the gene signatures of neurons were also downregulated aggravator genes, such as *NDRG4*, *PKIA*, *SHISA6*, *STX16*, *ZNF14*, and *ZNF606*, which confirmed our previous results. Nonetheless, it has not been determined whether changes in cellular proportions take place before transcriptomic changes or after them. Moreover, a recent study provided more detailed evidence regarding the cell types underlying the pathogenesis of PD based on single-cell transcriptomic data, demonstrating that cholinergic neurons, monoaminergic neurons, enteric neurons, and oligodendrocytes were related to PD and that these cells changed at the earliest stages of disease progression[26].

**Distinct expression patterns between initiator, facilitator and aggravator genes**

We next investigated how PD develops by tracking the trajectory of gene expression

patterns. We first observed that the initiator, facilitator and aggravator genes have

little overlap (Fig. 6a), which is reminiscent of the distinct biological processes of

these three temporal phases, as previously discussed. When further investigating their

connection in the protein-protein interaction (PPI) network, we observed that the

facilitator genes possess more degrees than initiator genes and tend to be located in

the hub of the network (Fig. 6b-c), while aggravator genes are located in the hub

compared to the facilitator genes (Fig. 6d-e). Since the expression changes in initiator

genes precede the facilitator genes and the changes in facilitator genes precede the

aggravator genes, it appears that the dysfunction first occurred in the signature genes

at the early phase, which would further spread the disease disturbance to spur

neurodegeneration by giving way to new signature genes, exhibiting dynamic

pathogenesis in disease development and implying that although distinct biological

functions are presented by the three temporal phase genes, they can connect to one

another in various ways. This complex evolution of pathogenesis may account for the

difficulties in halting or slowing PD development. Taken together, these results

confirmed the difference in pathogenesis among the three temporal phases of PD

while also demonstrating the connections among them.

**Identification of potential drugs for different phases of PD**

After elucidating the factors involved in PD pathogenesis in three temporal phases,

we next sought to identify potential therapies to reverse or slow the development of

PD. Currently, pharmacological treatment and deep brain stimulation are the main

therapies to alleviate the symptoms of PD, but they are unable to prevent the

neurodegenerative process[27,28]. Another promising therapy is cell-based treatments,

which can replace lost neurons with new and healthy neurons to reconstruct disrupted

circuits[29,30,31]. We propose that these therapies should be applied at the appropriate

phase of PD to maximize the chances of modifying the neurodegeneration process.

As mentioned above, the stem cell pluripotency was influenced in both the

prodromal and early stages of PD, and analysis of facilitator genes indicated that the

signaling pathways regulating the pluripotency of stem cells (KEGG ID: hsa04550)

were downregulated when prodromal subjects were converted to PD patients (Fig. 7a).

This result motivated us to seek novel candidate agonists to treat prodromal or early

stage PD patients by recovering the function of stem cells and promoting their

revitalization of neurons. To this end, L1000CDS² analysis was performed to identify

potential agents by upregulating the expression of DEGs in the hsa04550 pathway

(Methods, Supplementary Fig. 5a). Next, we obtained three candidate compounds, i.e.,

atorvastatin, curcumin and withaferin A (WFA), the former two of which have been

reported to promote the differentiation of neurons in neural stem cells (NSCs)[32,33].

Moreover, similar results were also obtained in the validation dataset using the same

approach (Supplementary Fig. 5a), indicating that WFA is of high confidence for

promoting NSCs differentiation into neurons.

To verify our inferences, we first evaluated the effects of WFA on the proliferation

and differentiation of NSCs *in vitro* (Methods). The optimum concentration for NSCs

differentiation was identified by the CCK8 test (Fig. 7b). Since NSCs will start to

differentiate spontaneously in the absence of basic fibroblast growth factor (bFGF)[34],

we applied WFA or DMSO (control) to treat NSCs without bFGF, and we assessed

differentiation by immunofluorescence staining. We observed that WFA can promote

NSCs to differentiate more neurons (Tuj$^+$ cells), fewer astrocytes (GFAP$^+$ cells) and

oligodendrocytes (GalC$^+$ cells) compared with the DMSO-treated control group (Fig.

7c-d), which demonstrated that WFA can promote NSCs to differentiate into neurons.

Given that the loss of dopaminergic neurons is an important pathological change in

PD[1], we performed further experiments and observed that the proportion of

dopaminergic neurons (TH$^+$Tuj$^+$ cells) in neurons (Tuj$^+$ cells) under WFA treatment

was significantly higher than that in the control (Fig. 7e-f), suggesting that WFA can

effectively promote NSCs to differentiate into dopaminergic neurons. Moreover, five

WFA-regulated genes were identified from the DEGs (Supplementary Fig. 5a) and

further validated by quantitative RT-PCR (Supplementary Table 8, Fig. 7g).

Specifically, *FGF2*, *ID1*, and *TCF3* were upregulated, which is consistent with our

hypothesis that NSCs differentiation is promoted by upregulating the DEGs. While

*WNT6* and *WNT11* were downregulated, we speculated that the difference between

blood tissue and NSCs may lead to this inconsistency. Notably, *TCF3* is a

transcriptional regulator involved in the maintenance of neural stem cells[35]; thus, its

functional recovery probably promotes the revitalization of neurons. *TCF3* is also a

hub node in the network of initiator-facilitator genes (Fig. 6c). Taken together, these

results strongly indicated that WFA has considerable potential to cure prodromal and

early-stage PD by promoting the differentiation of NSCs into dopaminergic neurons.

With the aggravation of PD, we found that biological terms, such as dopaminergic

synapse and the cGMP-PKG signaling pathway, were also enriched by the drug

targets of PD, implying their connections in a biological context. Among our

aggravator genes, six downregulated aggravator genes were also drug targets of PD,

and three of them have agonists in the DrugBank database, which might be considered

viable drugs in attempts to slow PD aggravation (Table 1). For example, by

upregulating *DRD3* in the dopaminergic synapse pathway, levodopa may be suitable

to treat patients with relatively more severe symptoms. In fact, levodopa is effective

in controlling most motor symptoms and remains the gold standard in the drug

treatment of PD[36]. Moreover, to yield more effective disease-modifying treatments,

some of these agonists should be combined to target the various pathogeneses in

aggravation. In addition, although *SLC18A1* is not a target of PD to date, it is located

in the dopaminergic synapse pathway and connected to known drug targets by

dopamine transduction. The downregulation of *SLC18A1* expression may lead to

dysfunction of the synaptic vesicle circle, thereby accelerating the severity of the

disease (Supplementary Fig. 5b); therefore it may be reasonable to consider *SLC18A1*

as a potential target for the design of therapeutic interventions in the future.

**Discussion**

In this study, we identified distinct molecular patterns in the development of PD with

a longitudinal blood transcriptomic analysis and provided a modified conceptual

model for PD pathogenesis. The results of this study helped us to identify appropriate

therapies for PD patients at different stages of the disease.

At the prodromal stage of PD, we found that olfactory-associated genes were first

reflected in the initiation of the disease and subsequently provided a simple and

effective method to help early diagnosis by examining the expression of the olfactory

receptor *OR8D1* in blood samples. Since hyposmia is also a clinical feature of

early-stage patients[10], our results could be coupled with clinical observations.

Nonetheless, the validity of these results should be confirmed in larger prospective

experiments. In addition, the expression of the initiator genes also supported that

hepatitis C is a trigger to spark the disease[3,9].

After the prodromal subjects were diagnosed as PD patients, the expression of the

facilitator genes indicated that dysfunction of calcium regulation might contribute to

neuronal death, and the neurons were unable to be renewed due to the decreased

pluripotency of stem cells. Changes in calcium levels can induce a series of events,

including mitochondrial dysfunction, enhanced ROS generation, and damaged ATP

production[16]. Calcium is also regulated by PD-related proteins, such as α-syn and

LRRK2[16]. These findings confirmed the facilitators of the previously proposed

conceptual model[3].

With the progression of PD, the expression of the aggravator genes showed that the

dysfunction of dopaminergic synapses, cGMP-PKG signaling, and ferroptosis may

represent the main factors of the neurodegeneration process. Ferroptosis is an

autophagic cell death process that contributes to autophagy activation[19]. In line with

the role played by aggravators in a previous conceptual model, these findings

highlight the importance of autophagy in disease aggravation[3]. Other aggravators

identified by this study need to be confirmed through further experiments. In addition,

we noted that although the expression levels of aggravator genes are significantly

changed with progression, the magnitude of these expression level changes was not

dramatic ($|\log2$ (third year/baseline)$| \leq 1$). One of the reasons for this finding may be

that PD injury occurs in brain tissue; thus, signal intensity may be reduced when it

reaches the bloodstream. The blood environment may also possess biological signals

from the internal environment or other organs, which may interfere with the

extraction of disease signals. In addition, PD is a chronic disease; therefore, gene

expression does not change dramatically, which also highlights the necessity of

conducting longitudinal studies that can capture this moderate dynamic change.

Interestingly, by integrating DNA methylation, we observed that methylation

alterations at baseline did not cause clinical manifestations immediately but may have

influenced the PD aggravation rate by regulating the changes in expression levels of

aggravator genes. A similar phenomenon was also reported by other chronic

conditions[21], making it possible to detect and monitor PD aggravation earlier.

However, it has not been determined how these CpGs regulate gene expression

changes and how long they are sustained.

According to the pathogenesis involved at different stages of PD, we further

inferred and demonstrated that WFA has the ability to promote the differentiation of

neural stem cells into dopaminergic neurons with strong potential to treat prodromal

or early stage PD patients (Fig. 7b-g). WFA is a major active component of *Withania*

*somnifera*, which has been demonstrated to protect dopaminergic neurons in

Parkinsonian mouse models and has exhibited a strong antioxidative effect[37].

Moreover, *Withania somnifera* was also reported to treat other disorders related to the

central nervous system[37]; thus, the future research should investigate whether WFA

has a similar mode of action in the treatment of other diseases, such as Alzheimer's

disease. For advanced PD patients, we proposed a list of drugs, such as levodopa,

droxidopa, and dextromethorphan, that may be employed to slow PD aggravation.

The appropriate drug combination addressing multiple targets and pathogenesis would

benefit severe patients. Moreover, our results may also lead to the reconsideration of

previously failed clinical trials, which may have been performed in nonspecific PD

patients[3].

**Methods**

**Clinical data analysis of PD patients**

In this study, the Movement Disorder Society-Unified Parkinson's Disease Rating

Scale (MDS-UPDRS) total score, which relates to various aspects, including

non-motor and motor symptoms, as well as motor complications[38], was applied to

measure the severity of PD patients. A higher score indicates an individual with a

more serious condition. For the use of PD medications, there are seven medical

treatments defined in PPMI, including unmedicated, levodopa, and levodopa plus

dopamine agonist. In this study, we simply classified patients as unmedicated and

medicated for subsequent analyses; for age characteristics, patients were classified

into three groups adopted from PPMI: group 1 (< 56 years), group 2 (56-65 years),

group 3 (> 65 years). Missing values of demographic and clinical information were

imputed using the median value.

**Omics data and quality control**

For RNA-seq data of whole blood samples, quality control and quantification of gene

or transcript expression were applied by PPMI. Briefly, FASTQ files were merged and

aligned to GRCh37 (hs37d5) by STAR (v2.4K)[39]. The read counts of genes or

transcripts were calculated by FeatureCounts[40] and abundance estimates (transcripts

per million, TPM) via salmon[41]. Samples with obvious quality control issues, such as

gender mismatch, principal component analysis (PCA) outliers and low read counts,

were filtered. After quality control, most samples have a mean Spearman correlation (>

0.87) to all other samples. To reduce the influence of technical noise, we removed

genes with counts/TPM of 0 in more than 80% of the samples in the subsequent

analysis. Importantly, according to PPMI analysis, ~60-70% of genes in whole blood

are also expressed in brain tissue by examining mRNA expression only. Detailed

information regarding the RNA sequencing and quality control can be found on the

PPMI website (http://ppmi-info.org/).

For DNA methylation (DNAm) data of whole blood samples, PPMI applied the

Illumina Human Methylation EPIC array (alias 850K methylation array) to assess

methylation status over 850,000 methylation sites across the whole genome. The β

values for each probe, which ranged from 0 (no methylation) to 1 (100% methylation),

were utilized to measure the methylation level of the targeted CpG site. Quality

control included removing low-quality samples and probes, as well as normalizing

blood cell composition. On that basis, we also applied the following filtering criteria

in this study[42]: removal of probes targeting X and Y chromosomes ($N$ = 19,701),

removal of cross-reactive probes on the EPIC array ($N$ = 43,254), and removal of

probes overlapping genetic variants at targeted CpG sites ($N$ = 12,510), leaving

792,150 CpG sites for the subsequent analysis. CpGs mapped to genes according to

GPL21145    from    Gene    Expression    Omnibus    (GEO,

http://www.ncbi.nlm.nih.gov/geo/).

**Identifying initiator genes**

Sixty-four prodromal subjects were recruited at baseline with a longitudinal

observation beginning in 2013, and 18 prodromal subjects converted to PD by 2019.

After matching RNA-seq samples and removing subjects without longitudinal

observation, 53 prodromal subjects were kept for the downstream analysis. To identify

initiator genes of PD, we defined prodromal conversion to PD as an "event" and

defined the time interval from prodromal to PD as the "survival time" to perform

survival analysis. For each gene, baseline prodromal subjects were divided into two

groups based on the median expression of the gene. Next, survival curves were

constructed by Kaplan-Meier survival curves, and the log-rank test was applied to

compare whether there was a significant difference in the overall conversion between

the two groups using the R package survival.

**Identifying facilitator genes**

To identify facilitator genes, we first performed differential expression analysis in a discovery dataset, which contained 17 RNA-seq samples of five subjects after matching clinical status and RNA-seq samples. Specifically, the heterogeneity between subjects is an important confounding factor identified by PCA (Fig. 3a); thus, we applied a multi-factor design that included the subjects and their condition (prodromal and PD) information in the design formula to identify DEGs using the R package/Bioconductor DESeq2[43]. This design will account for differences between the subjects when estimating the effect due to the condition. The validation dataset contained 57 prodromal and 390 PD RNA-seq samples at baseline without a conversion relationship between each other; therefore, the DEGs of the validation dataset were identified only by estimating the effect due to the condition.

**Identifying aggravator genes**

In this study, 252 PD patients both included clinical and RNA-seq data at four serial time points (baseline, 1-3 years). We first filtered the genes whose expression did not significantly change during the four time points based on repeated measures ANOVA with the R package ez. The *P*-value was adjusted by sphericity corrections followed by multiple testing using FDR. Genes with FDR $\leq$ 0.01 were kept for subsequent analyses (3435 genes). The linear mixed-effects model was constructed to identify which gene was associated with PD aggravation using the *lme* function in the R

package nlme. Each model included the MDS-UPDRS total score as the dependent

variable; gene expression, age, and time (i.e., baseline, 1-3 years), and medication

(unmedicated or medicated) as fixed effects; Each subject variation was included as a

random effect and considered their different response to medical treatment, therefore

each model was constructed as follows:

MDS-UPDRS total score ~ Gene expression + Age + Time + Medication, random =

~1+ Medication | Subject.

Genes with FDR $\leq$ 0.05 were defined as aggravator genes (325 genes, Supplementary

Table 4). The $\beta$ coefficient of the gene in the model lower than 0 indicates that it was

downregulated with PD aggravation; otherwise, it was upregulated. The R package

Pheatmap was subsequently used to plot the heatmap of aggravator genes. Notably,

some PD samples were abandoned in the above analysis because they did not possess

RNA-seq data for four serial years. In fact, most of them were measured three times,

involving 425 RNA-seq samples, which were regarded as an independent dataset to

assess the robustness of the aggravator genes.

**Cross-lagged panel analysis**

Cross-lagged panel analysis was used to investigate cause-effect relationship between

the expression of aggravator genes and clinical severity (Fig. 4c). Briefly, baseline

gene expression and MDS-UPDRS total score were adjusted for age by regression

residual analyses and then standardized with Z-transformation (mean = 0, SD = 1).

The same process was performed in the follow-up year (third year) with additional

adjustment for the times of medical treatment. Then, two following equations were

built:

$$X_{follow-up} = \beta_1 X_{baseline} + \beta_2 Y_{baseline} + \varepsilon_X$$

$$Y_{follow-up} = \beta_3 Y_{baseline} + \beta_4 X_{baseline} + \varepsilon_Y$$

Here, *X* is aggravator gene and *Y* is MDS-UPDRS total score. The linear regression

coeffcients $\beta_1$ and $\beta_3$ describe the autoregressive effects,  and the regression coeffcients

$\beta_2$ and $\beta_4$ represent cross-lagged effects. All parameters and statistics were estimated

by the *sem* function in the R package lavaan based on these two equations. Next, the

weighted gene expression risk score (ERS) was calculated as follows:

$\text{Weighted } ERS = \frac{1}{n}(\beta_1 \times Gene_1 + \beta_2 \times Gene_2 + \beta_3 \times Gene_3 + ... + \beta_{n-1} \times Gene_{n-1} + \beta_n \times Gene_n)$

where $\beta$ is the cross-lagged path coefficient ($\beta_4$) for each gene, and $n$ is the number of

genes included.

**MWAS analysis**

A linear regression model was constructed to assess the associations between each

CpG's methylation level (β value) and clinical traits of PD at baseline ($N = 330$). The

model used for each CpG was as follows:

MDS-UPDRS total score ~ Methylation β value + Age

Since variation of DNA methylation (DNAm) may arise prior to disease[21,22], and we

hypothesized that DNAm may influence the aggravation of PD in a long-term manner,

we applied the rate of PD aggravation, i.e., the change of MDS-UPDRS total score

from baseline to third years, as clinical traits of PD to identify the aggravation

rate-associated CpGs. After matching longitudinal clinical and DNAm data, 268 PD

patients were retained for the subsequent analysis. Using the *lm* function in R, the

linear regression model used for each CpG was as follows:

541            Rate of PD aggravation ~ Methylation β value + Age + Medication times

The quantile-quantile (Q-Q) plot was used to diagnose the above MWAS results

using the R package qqman. To estimate how much of the rate of PD aggravation

could be explained by blood DNAm, the top association CpGs were selected first ($P \leq$

0.001, Supplementary Table 6). Next, PCA was performed on these CpGs, and each

patient was scored by the first principal component using the R package psych. A

linear model was constructed to measure the relationship between the score calculated

by these CpGs and the rate of PD aggravation.

**Functional analysis**

KEGG pathway enrichment and gene set enrichment analysis (GSEA) analysis (2000

total permutations) of gene sets were performed by the R/Bioconductor package

clusterProfiler[44]. Tissue enrichment analysis of DEGs between prodromal and PD was

investigated by R package/Bioconductor TissueEnrich[45].

**Integrating transcriptome and methylome on PD aggravation**

The RNA-seq profiles and DNA methylation array data were pairwise integrated ($N =$

191) to study the regulatory relationship between each aggravator gene and each

aggravation rate-associated CpG site. Each linear regression model included delta

gene expression level change from baseline to third years as the dependent variable

(log2 (third year/baseline)), β value of a CpG site at baseline as the independent

variable, and age and medication times as covariates:

Δ Gene expression level ~ Methylation β value + Age + Medication times

**xCell analysis**

xCell[25] was used to estimate the abundance scores of 64 immune and stromal cell

types from the RNA-seq profiles of each PD patent over time (i.e., baseline, 1, 2, 3

years), the abundance scores was then standardized with Z-transformation (mean = 0,

SD = 1)  Similar to our aggravator genes analysis, linear mixed-effects model was

used to inferred which cell type was associated with PD aggravation:

MDS-UPDRS total score ~ Cell score + Age + Medication, random = ~1+

572           Medication | Subject

**Observation of initiator, facilitator and aggravator genes in the PPI network**

The interactions between initiator, facilitator and aggravator genes were examined in

a protein-protein interaction (PPI) network. The PPI network was derived from the

STRING database (http://www.string-db.org), and the interaction with experimental

score higher than 400 was considered reliable. Notably, since it difficult to observe the

network relationship between aggravator genes and facilitator genes identified from

the discovery dataset, we merged the facilitator genes both from the discovery and

validation datasets to observe their connections with aggravator genes in the PPI

network. The gene networks were subsequently visualized using Cytoscape[46].

**Drug discovery by L1000CDS² analysis**

The Library of Integrated Network-based Cellular Signatures (LINCS) program has

generated more than 1 million transcriptional profiles to elaborate the human cell

expression change under different perturbations of agents[47]. The L1000 dataset is part

of LINCS, which is based on the expression of only ~1000 genes[48]. L1000CDS² is a

LINCS L1000 characteristic direction signature search engine that links drugs, genes,

and diseases by comparing the input signatures to the background

(http://amp.pharm.mssm.edu/L1000CDS2/#/index). In this study, we applied

L1000CDS² to predict potential small molecular compounds that may reverse the

expression of DEGs between prodromal and PD subjects.

In the discovery dataset, 50 agents were returned by L1000CDS² according to eight

DEGs ($P \leq 0.05$) in pathways of regulating pluripotency of stem cells (KEGG ID:

hsa04550). We further selected three candidate compounds, i.e., atorvastatin,

curcumin and withaferin A (WFA), which were used to treat patients with mental

disorders in clinical trials (https://clinicaltrials.gov/), indicating the ability of delivery

across the blood brain barrier. Using the same approach, WFA was also identified in

the validation dataset (57 prodromal and 390 PD RNA-seq samples). To identify

potential genes regulated by WFA, we further identified six DEGs in the discovery

dataset that overlapped with background genes of WFA in the L1000 dataset, and five

of these six potential WFA-regulated genes in the discovery dataset were confirmed

by the validation dataset (Supplementary Fig. 5a).

**Isolation and culture of rat fetal neural stem cells**

Rat fetal NSCs were isolated and cultured from brain tissues of Sprague-Dawley rats

at 14−18 days post-coitum. Briefly, brain tissues were cut into small blocks, digested

by pre-warmed StemPro® Accutase (A11105-01, Gibco, NY, USA), filtered by a

70-µm cell strainer (352350, BD Biosciences, NY, USA) and centrifuged. The cell

pellet was resuspended in pre-warmed StemPro® NSC SFM complete medium

(consisting of KnockOut D-MEM/F-12 with 2% StemPro® Neural Supplement, 20

ng/mL bFGF, 20 ng/mL EGF, and 1% GlutaMAX-I; A1050901, Gibco, NY, USA),

inoculated in a dish at $1 \times 10^7$/mL and incubated at 37°C, 5% $CO_2$ and 90% humidity.

The NSCs that developed into neurospheres were passaged after 5-7 days

(Supplementary Fig. 6a). In subculture, the NSCs were digested by pre-warmed

StemPro® Accutase and plated in dishes precoated with 20 µg/mL fibronectin

(33016015, Gibco, NY, USA) to adherent culture at 37°C, 5% $CO_2$ and 90% humidity

(Supplementary Fig. 6b). The NSCs were passaged again when 75-90% confluent.

The cells were identified by immunofluorescence staining of Nestin at passage 2 (P2)

(Supplementary Fig. 6c-d). All WFA effects research experiments were carried out on

cells at passages 2 to 3 (P2~3) to minimize the experimental deviations.

**Analysis of WFA on NSCs proliferation by CCK8 assay**

WFA (HY-N2065, MCE, NJ, USA) was dissolved in DMSO to make a 1 mmol/L

stock solution, and subsequently, we made culture media containing different

concentrations of WFA with StemPro® NSC SFM complete medium. The working

concentrations of WFA were 0, 2, 10, 50, 250, 1000, 5000, and 25000 nmol/L. After

that step, rat fetal NSCs at P2 were collected and resuspended in culture media

containing WFA at $5 \times 10^4$/mL and then seeded at 100 μL/well in 96-well plates

immediately. To minimize the experimental deviations, culture medium containing

WFA but without NSCs was used as a blank control for each concentration. After

incubating the cells at 37°C, 5% $CO_2$ and 90% humidity for 48 hours, 10 μL CCK8

reagent (HY-K0301, MCE, NJ, USA) was added to each well and incubated for 4

hours. The absorbance was measured at 450 nm using a microplate reader (Multiskan

MK3, Thermo Fisher Scientific, MA, USA). The working concentration of WFA with

the highest average absorbance was used to analyze the effect of WFA on NSCs

differentiation.

**Intervention of NSCs differentiation by WFA**

Rat fetal NSCs at P3 were plated as an adherent culture on chamber slides (177380,

Thermo Fisher Scientific, MA, USA) or 6-well plates pre-coated with 20 μg/mL

fibronectin for 1 hour. To investigate the effect of WFA on NSCs differentiation, we

withdrew bFGF from StemPro® NSC SFM complete medium and added WFA stock

solution (1 mmol/L in DMSO, 1:4000; according to the experimental results of CCK8

assay, the work concentration is 250 nmol/L) or DMSO (D2438, 1:4000, Sigma, MO,

USA) into the media. Then, the cells were incubated at 37°C, 5% $CO_2$ and 90%

humidity until subsequent immunofluorescence and quantitative RT-PCR detection.

**Analysis of WFA on NSCs differentiation by immunofluorescence staining**

After intervention by WFA or DMSO for 48 hours, NSCs undergoing differentiation

were fixed, permeabilized and blocked and then incubated with primary mouse

anti-Tuj antibody (ab78078, 1:1000, Abcam, Cambridge, UK), primary mouse

anti-GALC antibody (MAB342, 1:100, Millipore, MA, USA) or primary rabbit

anti-GFAP antibody (16825-1-AP, 1:2000, ProteinTech, Wuhan, China) overnight at

4°C. After washing with PBS, cells were incubated with Alexa Fluor 594 conjugated

goat anti-mouse IgG (H+L) cross-adsorbed secondary antibody (A-11005, 1:200,

Thermo Fisher Scientific, MA, USA) or Alexa Fluor 488 conjugated goat anti-rabbit

IgG (H+L) cross-adsorbed secondary antibody (A-11008, 1:200, Thermo Fisher

Scientific, MA, USA) at 37°C for 30 min. Finally, cells were washed with PBS, dyed

with DAPI (Roche, Basel, Switzerland), mounted with 50% glycerol/PBS and

photographed under a fluorescence microscope (BX51, Olympus, Tokyo, Japan).

After treating NSCs with WFA or DMSO for seven days, double

immunofluorescence staining of Tuj and tyrosine hydroxylase (TH) was used to

analyze whether WFA could further affect NSCs differentiation into dopaminergic

neurons. Briefly, after fixation, permeabilization and blocking, cells were incubated

with both primary mouse anti-Tuj antibody and primary rabbit anti-TH antibody

(ab112, 1:1000, Abcam, Cambridge, UK) overnight at 4°C. Then, the cells were

incubated with mixed secondary antibodies of Alexa Fluor 594-conjugated goat

anti-mouse IgG and Alexa Fluor 488-conjugated goat anti-rabbit IgG for 30 min.

Finally, the cells were dyed with DAPI, mounted with 50% glycerol/PBS and

photographed under a BX51 fluorescence microscope.

**Quantitative RT-PCR**

Total RNA was extracted from NSCs treated with WFA or DMSO for 48 hours with

TRIzol reagent (15596026, Thermo Fisher Scientific, MA, USA) and reverse

transcribed into cDNA using the RevertAid RT Reverse Transcription Kit (K1691,

Thermo Fisher Scientific, MA, USA). Then, qRT-PCR mixtures were prepared with

the KAPA SYBR® FAST Universal qPCR Kit (KK4601, Sigma, MO, USA)

according to the manufacturer's instructions and run in LightCycler 96 (Roche,

Switzerland). The mRNA expression of genes regulating the pluripotency of stem

cells was normalized to the expression of GAPDH. The results were compared by the

$2-\Delta\Delta Ct$ method. The specific primers (Supplementary Table 8) were synthesized by

Tsingke Biological Technology (Wuhan, China).

**Statistical analysis of NSCs experiments**

Image-Pro Plus 6.0 software (Media Cybernetics, MA, USA) was used for analysis

and cell counting of the fluorescence images. The results from different groups were

compared using Student's *t*-test or one-way ANOVA followed by Tukey's multiple

comparison tests. *P*-value $\leq$ 0.05 was considered to be significant. All statistical

analyses were performed with GraphPad Prism 6 software (GraphPad Software, CA,

USA).

**PD drugs and targets collection**

695    The drugs for PD were obtained from clinical trials (release 2019.4.22, at least phase

696    2 clinical trial drugs, https://clinicaltrials.gov/), and the drug targets were extracted

697    from DrugBank[49], DGIdb[50] and TTD[51]. We finally collected 104 PD drugs

698    corresponding to 363 target genes (Supplementary Table 9).

**Data availability**

701    Data used in this study were obtained from the Parkinson's Progression Markers

702    Initiative (PPMI) database (www.ppmiinfo.org/data). For up-to-date information on

703    the study, please visit www.ppmiinfo.org.

**Acknowledgments**

We would like to acknowledge PPMI - a public-private partnership, which is funded

by the Michael J. Fox Foundation for Parkinson's Research and funding partners,

including Abbvie, Allergan, Amathus Therapeutics, Avid Radiopharmaceuticals,

Biogen, BioLegend, Bristol-Myers Squibb, Calico, Celgene, Denali, GE Healthcare,

Genetech, GlaxoSmithKline, Golub Capital, Handl Therapeutics, Insitro, Janssen

Neuroscience, Lilly, Lundbeck, Merck, Meso Scale Discovery, Pfizer, Piramal,

Prevail, Roche, Sanofi Genzyme, Servier, Takeda, Teva, UCB, Verily, and Voyager

Therapeutics. We also thank the National Natural Science Foundation of China (grant

31670779) for supporting this study.

**Author contributions**

717 H.Y.Z. and X.H.N. conceived the project; G.X. collected and performed data analysis

718 with help from X.H.N., Q.Q.S., M.G., and H.W; G.W. performed the WFA

719 experiment. G.X., G.W., Q.Q.S., H.Y.Z., X.H.N., H.W., and B.L wrote and edited the

720 manuscript.

**Competing interests**

723 The authors declare no competing interests.

**References**

1.    Dauer W, Przedborski S. Parkinson's disease: mechanisms and models. *Neuron* **39**, 889-909 (2003).

2.    Chaudhuri KR, Healy DG, Schapira AH. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol.* **5**, 235-245 (2006).

3.    Johnson ME, Stecher B, Labrie V, Brundin L, Brundin P. Triggers, facilitators, and aggravators: Redefining Parkinson's disease pathogenesis. *Trends Neurosci.* **42**, 4-13 (2019).

4.    Shamir R*, et al.* Analysis of blood-based gene expression in idiopathic Parkinson disease. *Neurology* **89**, 1676-1683 (2017).

5.    Chen-Plotkin AS. Blood transcriptomics for Parkinson disease? *Nat. Rev. Neurol.* **14**, 5-6 (2018).

6.    Fan Y, Xiao S. Progression rate associated peripheral blood biomarkers of Parkinson's disease. *J. Mol. Neurosci.* **65**, 312-318 (2018).

7.    Androulakis IP, Yang E, Almon RR. Analysis of time-series gene expression data: methods, challenges, and opportunities. *Annu. Rev. Biomed. Eng.* **9**, 205-228 (2007).

8.    Marek K*, et al.* The parkinson progression marker initiative (PPMI). *Prog. Neurobiol.* **95**, 629-635 (2011).

9.    Kim J-M*, et al.* Association between hepatitis C virus infection and Parkinson's disease. *Mov. Disord.* **31**, 1584-1585 (2016).

10.   Doty RL. Olfaction in Parkinson's disease and related disorders. *Neurobiol.*

*Dis.* **46**, 527-552 (2012).

11.     Braak H, Del Tredici K, Rüb U, De Vos RA, Steur ENJ, Braak E. Staging of

brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging.* **24**,

197-211 (2003).

12.     Mainland JD*, et al.* The missense of smell: functional variability in the human

odorant receptor repertoire. *Nat. Neurosci.* **17**, 114 (2014).

13.     Ross GW*, et al.* Parkinsonian signs and substantia nigra neuron density in

decendents elders without PD. *Ann. Neurol.* **56**, 532-539 (2004).

14.     Wichterle H, Przedborski S. What can pluripotent stem cells teach us about

neurodegenerative diseases? *Nat. Neurosci.* **13**, 800-804 (2010).

15.     Le Grand JN, Gonzalez-Cano L, Pavlou MA, Schwamborn JC. Neural stem

cells in Parkinson's disease: a role for neurogenesis defects in onset and

progression. *Cell. Mol. Life Sci.* **72**, 773-797 (2015).

16.     Calì T, Ottolini D, Brini M. Calcium signaling in Parkinson's disease. *Cell*

*Tissue Res.* **357**, 439-454 (2014).

17.     Poewe W, Mahlknecht P. The clinical progression of Parkinson's disease.

*Parkinsonism Relat. Disord.* **15**, S28-S32 (2009).

18.     Chenplotkin A*, et al.* Finding useful biomarkers for Parkinson's disease. *Sci.*

*Transl. Med.* **10**, eaam6003 (2018).

19.     Gao M, Monian P, Pan Q, Zhang W, Xiang J, Jiang X. Ferroptosis is an

autophagic cell death process. *Cell Res.* **26**, 1021-1032 (2016).

20.     Guiney SJ, Adlard PA, Bush AI, Finkelstein DI, Ayton S. Ferroptosis and cell

death mechanisms in Parkinson's disease. *Neurochem. Int.* **104**, 34-48 (2017).

21.    Chen R*, et al.* Longitudinal personal DNA methylome dynamics in a human

with a chronic condition. *Nat. Med.* **24**, 1930-1939 (2018).

22.    Rakyan VK, Down TA, Balding DJ, Beck S. Epigenome-wide association

studies for common human diseases. *Nat. Rev. Genet.* **12**, 529-541 (2011).

23.    Henderson-Smith A*, et al.* DNA methylation changes associated with

Parkinson's disease progression: outcomes from the first longitudinal

genome-wide methylation analysis in blood. *Epigenetics* **14**, 365-382 (2019).

24.    Wang D*, et al.* Comprehensive functional genomic resource and integrative

model for the human brain. *Science* **362**,    (2018).

25.    Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular

heterogeneity landscape. *Genome Biol.* **18**, 220 (2017).

26.    Bryois J*, et al.* Genetic identification of cell types underlying brain complex

traits yields insights into the etiology of Parkinson's disease. *Nat. Genet.* **52**,

482-493 (2020).

27.    Connolly BS, Lang AE. Pharmacological treatment of Parkinson disease: a

review. *JAMA* **311**, 1670-1683 (2014).

28.    Benabid AL, Chabardes S, Mitrofanis J, Pollak P. Deep brain stimulation of

the subthalamic nucleus for the treatment of Parkinson's disease. *Lancet*

*Neurol.* **8**, 67-81 (2009).

29.    Barker RA, Drouin-Ouellet J, Parmar M. Cell-based therapies for Parkinson

disease—past insights and future potential. *Nat. Rev. Neurol.* **11**, 492 (2015).

30.  Reddy AP, Ravichandran J, Carkaci-Salli N. Neural regeneration therapies for

Alzheimer's and Parkinson's disease-related disorders. *Biochim. Biophys. Acta*

*Mol. Basis. Dis.* **1866**, 165506 (2020).

31.  Qian H*, et al.* Reversing a model of Parkinson's disease with in situ converted

nigral neurons. *Nature* **582**, 550-556 (2020).

32.  Choi N-Y*, et al.* Atorvastatin rejuvenates neural stem cells injured by

oxygen-glucose deprivation and induces neuronal differentiation through

activating the PI3K/Akt and ERK pathways. *Mol. Neurobiol.* **56**, 2964-2977

(2019).

33.  Chen F*, et al.* Curcumin increased the differentiation rate of neurons in neural

stem cells via wnt signaling in vitro study. *J. Surg. Res.* **192**, 298-304 (2014).

34.  Gage FH. Mammalian neural stem cells. *Science* **287**, 1433-1438 (2000).

35.  Ohtsuka T*, et al.* Gene expression profiling of neural stem cells and

identification of regulators of neural differentiation during cortical

development. *Stem Cells* **29**, 1817-1828 (2011).

36.  Poewe W, Antonini A, Zijlmans JC, Burkhard PR, Vingerhoets F. Levodopa in

the treatment of Parkinson's disease: an old drug still going strong. *Clin. Interv.*

*Aging* **5**, 229 (2010).

37.  Singh N, Rai SN, Singh D, Singh SP. Withania somnifera shows ability to

counter Parkinson's Disease: An Update. *SOJ Neurol.* **2**, 1-4 (2015).

38.  Goetz CG*, et al.* Movement Disorder Society-sponsored revision of the

Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation

and clinimetric testing results. *Mov. Disord.* **23**, 2129-2170 (2008).

39.    Dobin A*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,

15-21 (2013).

40.    Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**,

923-930 (2014).

41.    Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast

and bias-aware quantification of transcript expression. *Nat. Methods.* **14**,

417-419 (2017).

42.    Pidsley R*, et al.* Critical evaluation of the Illumina MethylationEPIC

BeadChip microarray for whole-genome DNA methylation profiling. *Genome*

*Biol.* **17**, 208 (2016).

43.    Love MI, Huber W, Anders S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

44.    Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS.* **16**, 284-287 (2012).

45.    Jain A, Tuteja G. TissueEnrich: Tissue-specific gene enrichment analysis.

*Bioinformatics* **35**, 1966-1967 (2019).

46.    Shannon P*, et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res.* **13**, 2498-2504 (2003).

47.    Keenan AB*, et al.* The library of integrated network-based cellular signatures

NIH program: system-level cataloging of human cells response to

834           perturbations. *Cell Syst.* **6**, 13-24 (2018).

835    48.    Subramanian A*, et al.* A next generation connectivity map: L1000 platform

836           and the first 1,000,000 profiles. *Cell* **171**, 1437-1452. e1417 (2017).

837    49.    Law V*, et al.* DrugBank 4.0: shedding new light on drug metabolism. *Nucleic*

838           *Acids. Res.* **42**, D1091-D1097 (2014).

839    50.    Wagner AH*, et al.* DGIdb 2.0: mining clinically relevant drug–gene

840           interactions. *Nucleic Acids. Res.* **44**, D1036-D1044 (2016).

841    51.    Qin C*, et al.* Therapeutic target database update 2014: a resource for targeted

842           therapeutics. *Nucleic Acids. Res.* **42**, D1118-D1123 (2014).

**Figure**

**Fig. 1 Study workflow.** Longitudinal RNA-seq of whole blood samples and

corresponding clinical assessment were obtained from Parkinson's Progression

Markers Initiative (PPMI) database (https://www.ppmi-info.org/data). (up) Flowchart

of analysis in the conversion process. (right) Flowchart of analysis in the aggravation

process.

**Fig. 2 Pathway enrichment and survival analysis of initiator genes. a** KEGG

pathway enrichment of initiator genes by gene set enrichment analysis (GSEA) (genes

ranked by -log10($P$-value)). **b** Kaplan-Meier curves of conversion in prodromal

subjects ($N$ = 53). The magenta line indicates prodromal subjects with higher

expression of *OR8D1*, while the blue line indicates prodromal subjects with lower

expression of *OR8D1*, which are at high risk of conversion.

**Fig. 3 Identifying DEGs between prodromal and PD. a** PCA clustering plot of

RNA-seq data between prodromal and PD. **b** Overlaps of DEGs in the discovery and

validation datasets ($P$ = 4.75e-09). **c** Tissue enrichment of DEGs between prodromal

and PD (*$P \leq 0.05$). **d** KEGG pathway enrichment of DEGs by GSEA (genes ranked

by log2FoldChange).

**Fig. 4 Functional analysis of aggravator genes and causal inference. a** Heatmap of

aggravator genes in longitudinal PD patients ($N = 252$). **b** KEGG pathway enrichment

of downregulated aggravator genes. **c** Causal inference based on the cross-lagged path

analysis model. $\beta_2$ and $\beta_4$ are cross-lagged path coefficients. Genes with significant

unidirectional paths ($\beta_4$, $P \leq 0.05$) from baseline gene expression to follow-up

MDS-UPDRS total score (the third year), while those without significant path

coefficients ($\beta_2$) from baseline MDS-UPDRS total score to follow-up gene

expression are considered to be causing genes in disease aggravation. **d** Correlation

between the weighted gene expression risk score (ERS) at baseline and MDS-UPDRS

total score at follow-up (the third year).

**Fig. 5 Assessment of the blood methylome and investigation of their regulatory**

**relationship with aggravator genes. a** Quantile-quantile (Q-Q) plot of MWAS result

using rate of aggravation as clinical traits ($N = 268$). **b** Linear relationship between

DNA methylation score and rate of aggravation of PD patients. **c** Overlaps of our

aggravator rate associated-genes with literature-reported PD aggravator-associated

genes (CpGs map to genes). **d** (upside) The associations between cg13298827,

*PCDHA11* and aggravation; (downside) The associations between cg17512517,

*CCDC50* and aggravation. Positive (+) and negative (-) associations are shown in the

figure.

**Fig. 6 Relationship between initiator, facilitator and aggravator genes. a** Initiator,

facilitator and aggravator genes do not show significant overlaps with each other ($P >$

0.05). **b** Boxplot of the connected degrees of initiator and facilitator genes in the PPI

network. Facilitator genes showed more connections than initiator genes ($P$ = 2e-04).

**c** Network diagram depicting the interaction between initiator and facilitator genes.

The facilitator genes (yellow) tend to be located in the hub of the network. **d** Boxplot

of the connected degrees of facilitator and aggravator genes in the PPI network.

Aggravator genes showed more connections than facilitator genes ($P$ = 4.2e-09). **e**

Network diagram depicting the interaction between facilitator and aggravator genes.

The aggravator genes (pink) tend to be located in the hub of the network.

**Fig. 7 WFA promotes the proliferation and differentiation of NFCs. a** GSEA plot

showing downregulated signaling pathways regulating pluripotency of stem cells

when prodromal conversion to PD. **b** Identification of the optimum concentration of

WFA on the proliferation of NFCs by the CCK8 test. WFA significantly promoted

NSCs proliferation at 50, 250, and 1000 nmol/L (*$P \leq$ 0.05), and the proliferative

effects at 250 nmol/L were significantly higher than those at 50 nmol/L ($^{\#}P \leq$ 0.05),

suggesting that WFA has a dose-dependent effect, whereas no significant difference

was observed between 250 nmol/L and 1000 nmol/L, indicating that 250 nmol/L may

reach the plateau stage of a dose-dependent effect; thus, 250 nmol/L was adopted to

evaluate the effects of WFA on the next NSCs differentiation experiment. **c** Detection

of NFCs differentiation by immunofluorescence staining (scale bar = 100 μm). **d**

Statistical analysis of neurons (Tuj$^+$ cells), astrocytes (GFAP$^+$ cells) and

oligodendrocytes (GalC$^+$ cells) between the WFA- and DMSO-treated control groups

(*$P \leq 0.05$). **e** Detection of dopaminergic neurons derived from NSCs using

immunofluorescence staining (Scale bar = 100 μm). **f** Statistical analysis of the

proportion of dopaminergic neurons (TH$^+$ Tuj$^+$ cells) in neurons (Tuj$^+$ cells) between

the WFA and control groups (*$P \leq 0.05$). **g** Assessment of the mRNA expression of

genes regulating the pluripotency of stem cells by quantitative RT-PCR (*$P \leq 0.05$).

**Table 1 Downregulated aggravator genes and the involved agonists**

| Aggravator genes | Agonists (DrugBank) |
| --- | --- |
| *ADRA2C* | Brimonidine, Norepinephrine, Droxidopa, Clonidine, Bromocriptine, Apomorphine, Paliperidone, Ephedra sinica root, Metamfetamine, Xylometazoline, Agmatine, Oxymetazoline, Racepinephrine |
| *DRD3* | Ropinirole, Pramipexole, Apomorphine, Cariprazine, Pergolide, Bromocriptine, Lisuride, Cabergoline, Rotigotine, Dopamine, Levodopa, Captodiame, Dihydro-alpha-ergocryptine, Tianeptine |
| *OPRD1* | Codeine, Dextropropoxyphene, Hydrocodone, Morphine, Butorphanol, Fentanyl, Etorphine, 3-Methylthiofentanyl, Dimethylthiambutene, Loperamide, 3-Methylfentanyl, Oxycodone, Diamorphine, Tramadol, Methadone, Sufentanil, Carfentanil, Amitriptyline, Diphenoxylate, Levorphanol, Dihydromorphine, Remifentanil, Ketobemidone, Dextromethorphan, Opium, Loxicodegol, Butyrfentanyl, Metenkefalin, Hydromorphone |

**Supplementary Figure**

**Supplementary Fig. 1 Clinical aggravation of PD patients. a** The overall

aggravation of PD patients ($N = 252$). Although most patients are under drug

treatment after enrollment, the disease condition is still in progression (Repeated

Measures ANOVA $P = 9.15\text{e-}20$). **b** The aggravation trajectory for each PD patient.

**Supplementary Fig. 2 Heatmap of aggravator genes in independent PD patients**

**(425 RNA-seq samples).**

**Supplementary Fig. 3 QQ plot of MWAS using baseline MDS-UPDRS total score**

**as a clinical trait.** As shown in the plot, almost all the CpGs on the diagonal of the

plot and seldom the $P$-value of CpGs were smaller than expected, which indicates that

little CpG was found to be significantly associated with this clinical trait.

**Supplementary Fig. 4 Heatmap of cell types associated with PD aggravation**.

Abbreviations: Th1 cells, Type 1 T-helper cells; aDC, Activated dendritic cells;

CD4+Tem, CD4+ effector memory T-cells; CD8+ Tcm, CD8+ central memory T cells;

MSC, Mesenchymal stem cells; Th2 cells, Type 2 T-helper cells; HSC, Hematopoietic

stem cells.

**Supplementary Fig. 5 Identifying drugs and potential drug targets for PD. a**

L1000CDS² analysis for drug discovery. DEGs in the pathway of regulating

pluripotency of stem cells (KEGG ID: hsa04550) were submitted to L1000CDS² to

predict potential agents. DEGs overlapping with background genes of withaferin A in

L1000 signatures were considered WFA-regulated genes. Five WFA-regulated genes

were confirmed by the validation datasets. **b** Downregulated aggravator genes and PD

drug targets in the neuron presynaptic terminal of dopaminergic synapse. The green

color indicates drug targets of PD, while the magenta color indicates aggravator gene.

**Supplementary Fig. 6 Morphology and immunofluorescent identification of**

**NSCs. a** Bright field image of suspended neurospheres at P1 (passage 1) that was

cultured in complete StemPro® NSC SFM for 6 days (Scale bar = 100 μm). **b** Bright

field image of adherent NSCs at P2 (passage 2) that were cultured in complete

StemPro® NSC SFM (scale bar = 100 μm). **c** Immunofluorescent identification of

NSCs with primary mouse anti-Nestin monoclonal antibody (ab6142, 1:100, Abcam,

Cambridge, UK) (Scale bar = 100 μm). **d** Merged fluorescence image of nestin and

DAPI (scale bar = 100 μm).

Longitudinal RNA-seq of whole blood samples

Longitudinal clinical assessment

Prodromal

Conversion

PD

Initiator genes

Facilitator genes

Drug discovery

Validation in vitro experiment

NSCs

Agent

Control

Neurons

Astrocytes

Oligodendrocytes

Function analysis
&
Comparison

De Novo PD

Baseline, 1-3 years

Aggravator genes

Drug discovery

Causal inference

Integration with DNA methylation

Advanced PD

**a**

**b**

**a**

**b**

**c**

**d**

**a**

Age

Medical treatment

MDS-UPDRS total score

Time

↑ Upregulated

Downregulated ↓

**Age**
<56
56-65
>65

**Medical treatment**
Unmedicated
Medicated

**MDS-UPDRS total score**
100

**Time**
Baseline
Year 1
Year 2
Year 3

**b**

Dopaminergic synapse
Adrenergic signaling in cardiomyocytes
cGMP−PKG signaling pathway
Amphetamine addiction
Ferroptosis
Metabolism of xenobiotics by cytochrome P450
Selenocompound metabolism
Chemical carcinogenesis
Fatty acid biosynthesis
mRNA surveillance pathway
Estrogen signaling pathway
Inflammatory mediator regulation of TRP channels

$-\log 10\ (P)$

**c**

Baseline gene expression $\xrightarrow{\beta_1}$ Follow-up gene expression

✗ $\beta_2$

√ $\beta_4$

Baseline MDS-UPDRS total score $\xrightarrow{\beta_3}$ Follow-up MDS-UPDRS total score

**d**

Adjusted R² = 0.054 , $P$ = 0.00052

Follow-up MDS-UPDRS total score

ERS

**a**

**b**

**c**

**d**

**a**

Running Enrichment Score

signaling pathways regulating
pluripotency of stem cells

PD vs Prodromal
NES = -1.560
Adjusted *P* = 0.004

Ranked list metric

Rank in Ordered Dataset

**b**

OD450

WFA (nmol/L)

**c**

DMSO    WFA

Tuj

GFAP

GALC

**e**

DMSO    WFA

Tuj        Tuj

TH         TH

DAPI      DAPI

Merge     Merge

**d**

% of total cells

Tuj    GFAP    GALC

**f**

% of total cells

TH

**g**

Rative mRNA expression

■ DMSO  ■ WFA

FGF2    ID1    TCF3    WNT6    WNT11