Pseudomonas aeruginosa PA14 biofilms produce R-bodies, extendable protein polymers with roles in host colonization and virulence

Bryan Wang[1], Yu-Cheng Lin[1], Jeanyoung Jo[1], Alexa Price-Whelan[1], Shujuan Tao McDonald[1,2],

5   Lewis M. Brown[1,2], and Lars E.P. Dietrich[1]*

1Department of Biological Sciences, Columbia University, New York, NY, USA
2Quantitative Proteomics and Metabolomics Center, Columbia University, New York, NY, USA

10   *email: LDietrich@columbia.edu

reb genes code for R-bodies: large, extendable polymers that are known for their roles in obligate endosymbioses. In the non-endosymbiotic pathogen *Pseudomonas aeruginosa*,

15 **reb homologues are part of a cluster found in virulent strains. Here, we demonstrate that R-bodies are produced in abundance by *P. aeruginosa* PA14 subpopulations during biofilm growth, identify regulators of reb gene expression, and show that reb genes are required for full colonization and virulence in host models.**

20 R-bodies are enigmatic biological machines that have been characterized most extensively in *Caedibacter taeniospiralis*, an obligate endosymbiont of paramecia. These structures are composed of ~ 12 kDa monomers that polymerize to form hypercoiled ribbons ~ 0.5 μm in diameter [1]. In response to changes in parameters such as pH, temperature or ionic strength [2–4], R-body coils extend into 10-20-μm-long needle-like structures (i.e., ~ 10X the length of the

25 producing cell), lysing the bacterium [4,5]. In the *Caedibacter-Paramecium* endosymbiosis, R-body-mediated lysis is thought to release an unidentified factor that is toxic to other paramecia. R-body-producing endosymbionts therefore confer onto their host cell a "killing" phenotype and a competitive advantage against naive peer cells, a phenomenon that was first described by Tracy M. Sonneborn in 1938 [6].

Diverse proteobacterial genomes contain homologues to genes for R-bodies (i.e., reb genes) [7]. In the opportunistic pathogen *P. aeruginosa*, reb gene homologues are part of a cluster that is conserved in a substantial fraction (147/241) of complete strain genomes, including those of the virulent strains PA14 and LESB58, but that is not present in the popular and notably less-

35 virulent model strain PAO1 [8,9] (Supplementary File 1). We will refer to genes in the *P. aeruginosa* PA14 reb cluster using names shown in **Figure 1a**, which are based on a prior study [10] and phylogenetic analyses conducted using the *C. taeniospiralis* structural genes for R-body production (**Supplementary Figure 1**). Included in this conserved locus are two regulatory

genes, *rebZ* and *fecI2*, which code for a transcription factor and sigma factor, respectively, and

40 which in a prior study were required for full virulence in a *C. elegans* killing assay [9] (**Figure 1a**).

Via bioinformatic analyses, we found that though *reb* structural gene homologues are scattered

across the phylogenetic tree of the genus *Pseudomonas* **(Supplementary Figure 2)**,

homologues of the regulatory genes show strong conservation in strains that contain *reb*

structural genes, both within the species *P. aeruginosa* (in 146/147 *reb* homologue-containing

45 genomes; **Supplementary Table 1**) and among other pseudomonads (**Figure 1a,**

Supplementary Figure 2). Though all of these observations hinted at a potential role for R-

bodies in *P. aeruginosa*-host associations, R-body production by *P. aeruginosa* biofilms and

during host colonization had not been investigated.

50 To examine whether PA14 produces R-bodies, we applied a modified R-body enrichment

protocol [11] to biofilms formed by the WT and by Δ*pel*, a mutant that is unable to produce the

major PA14 exopolysaccharide and yields biofilms that are more amenable to disruption. We

performed protein mass spectrometry (MS) and scanning-electron microscopy (SEM) on the

isolated SDS-insoluble fractions (**Figure 1b**). Products of the *reb* cluster were detected by MS.

55 Three products—PA14_27680, RebP1, and RebP2-- were detected at similar levels and were

among the 12 most abundant proteins in our sample (**Figure 1c**). RebP1 and RebP2 have 40-

42% and 34-40% homology, respectively, to RebB and RebA from *C. taeniospiralis,* which

constitute the main structural components of R-bodies in that organism [12]. *PA14_27680* is

conserved in all pseudomonads that contain at least one *reb* structural gene, suggesting that it

60 may encode another R-body structural component. One other uncharacterized protein coded for

by the *reb* gene cluster, PA14_27675, was also detected in the SDS-insoluble fractions

(**Supplementary File 2 and Figure 1c**). Consistent with the MS results, our SEM analyses

revealed the presence of R-bodies in preparations from WT PA14 biofilms (**Figure 1d**). We

detected R-bodies in various states of extension ranging from hypercoiled (~ 450 nm long) to

65   fully extended (~ 5.5 μm long) (**Figure 1d, e**). These structures were not observed in a deletion

mutant lacking the *rebP1P2* operon (Δ*rebP1P2*) (**Supplementary Figure 3**). Taken together,

these data indicate that R-bodies are produced by PA14 biofilms.

Based on the conservation of *rebZ* and *fecI2* in strains containing R-body structural gene

70   homologues (**Figure 1a; Supplementary File 1; Supplementary Figure 1**), we reasoned that

one or both of these genes control expression of *rebP1P2*. To test this, we generated constructs

that report expression from the *rebP1P2* promoter (P$_{rebP1}$) as red (mScarlet) or green (GFP)

fluorescence and inserted these into a neutral site on the chromosome in WT PA14 and

regulatory gene mutants. We observed P$_{rebP1}$ activity in WT colony biofilms that was dependent

75   on the presence of both *rebZ* and *fecI2* (**Figure 1f-h** and **Supplementary Figure 3**). Because

biofilm development leads to the formation of steep chemical gradients that can affect gene

expression patterns along the z-axis [13], we wanted to visualize P$_{rebP1}$ activity across depth. We

employed a thin sectioning protocol to prepare vertical sections taken from the center region of

three-day-old colony biofilms [14]. Confocal microscopy of P$_{rebP1}$-*mScarlet* biofilm sections

80   revealed that, strikingly, cells expressing *rebP1P2* are present in thin striations such that both

bright and dark cells are situated at the same depth. For comparison, we prepared sections of

biofilms in which *gfp* expression was driven by a constitutive promoter (P$_{PA1/04/03}$-*gfp*) and

observed fluorescence that is more consistent between cells at the same depth

(**Supplementary Figure 3**). Our results suggest that P$_{rebP1}$-driven expression is stochastic and

85   that expression status is heritable during vertical growth in the the biofilm.

The fact that homologues of the *C. taeniospiralis* R-body structural genes are found in diverse

free-living bacteria suggests that their roles extend beyond endosymbiotic interactions [7]

(**Supplementary Figure 1**). *P. aeruginosa* forms biofilms and acts as an extracellular pathogen

90   during colonization of a range of hosts including plant and nematode models [15,16]. To examine

whether *rebP1P2* is expressed during interactions with hosts, we employed the P$_{rebP1}$-*mScarlet* reporter strain in *Arabidopsis thaliana* seedling and *Caenorhabditis elegans* gut colonization assays. In both hosts, as in biofilms, *rebP1P2* expression patterns were distinct from those exhibited by a promoter driving constitutive expression of mScarlet (P$_{PA1/04/03}$-*mScarlet*). While

95 constitutive mScarlet production was uniform and fuzzy across colonized regions of hosts (**Figure 2a, b**), *rebP1P2* expression was punctate and restricted to a subset of PA14 cells (**Figure 2a**, **b**). To rule out the possibility that constitutive mScarlet production represented non-specific, background fluorescence, we expressed mScarlet without a promoter (MCS-*mScarlet*) in these hosts. Indeed, promoterless mScarlet expression was not observed in either *A. thaliana*

100 or *C. elegans* (**Figure 2a, b**). The distinct *rebP1P2* expression patterns we observed in the colony biofilm and host contexts may indicate that there is an advantage to restricting R-body production to a subset of cells in the population.

In *C. taeniospiralis* and *Azorhizobium caulinodans*, R-body production has been linked to host-

105 associated phenotypes that confer toxicity [4,17,18]. We hypothesized that R-bodies contribute to PA14 host colonization and/or pathogenicity and therefore performed established assays to test such contributions of PA14 R-body production in our two host systems. In *A. thaliana*, we quantified the extent to which PA14 WT and mutant strains colonized seedlings after inoculation and five days of plant growth. For the *C. elegans* virulence assay, we quantified percent survival

110 for up to four days after synchronized populations of worms were exposed to PA14 WT and mutant strains. When compared to WT PA14, Δ*rebP1P2* displayed attenuated host colonization (**Figure 2e**) and virulence (**Figure 2f**); both phenotypes were restored to WT levels upon complementation of *rebP1P2*.

115 In many hosts and various infection sites in humans, biofilm formation is critical to bacterial colonization and virulence and requires production of matrix [19]. Matrix secretion is a defining

feature of biofilms; the matrix functions as the "glue" that holds bacterial cells together and can facilitate attachment to the host [20]. To test whether matrix production is affected by a defect in R-body production, we examined biofilm development using a standardized colony morphology

120   assay, in which the dye Congo red is provided in the medium and binds to biofilm matrix. We found that matrix production and biofilm development were unaltered in the Δ*rebP1P2* mutant (**Figure 2c**). Furthermore, Δ*rebP1P2* showed no fitness disadvantage when grown in mixed-strain colony biofilms with the WT parent (**Figure 2d**). We conclude the *A. thaliana* colonization and *C. elegans* killing defects observed in the Δ*rebP1P2* mutant are not mediated via effects on

125   biofilm formation and that the benefit of R-body production is specific to the host-associated lifestyle.

We have shown here the production of R-bodies by PA14 cells growing in biofilms and demonstrated a role for these structures during host infection. Expression of the PA14 R-body

130   structural genes is stochastic and can be observed both in independent biofilms and during association with hosts. However, an advantage to R-body production is detectable only during host colonization, suggesting that R-body function is specific to infection. *Caedibacter* R-body extension is triggered by low pH and occurs in phagosomes of naive paramecia, prompting *Caedibacter* lysis and the release of other toxins within *Caedibacter* cells and thereby killing the

135   eukaryotic predator [4,5]. We speculate that PA14 R-body extension is similarly triggered in a host-specific manner and that cells in the R-body-producing subpopulation act as "kamikazes" to release other bacterial products that facilitate colonization and enhance virulence. Our findings identify R-bodies as novel *P. aeruginosa* virulence factors and raise the possibility that R-body production contributes to the behaviors of diverse free-living bacteria during interactions

140   with other organisms.

METHODS

Bacterial strains and growth conditions. Strains used in this study are listed in

145   Supplementary Table 1. Liquid cultures of *Pseudomonas aeruginosa* strain UCBPP-PA14

(PA14) were grown in lysogeny broth (LB) at 37°C with shaking at 250 rpm. Biological replicates

were inoculated from distinct clonal-source colonies grown on LB + 1.5% agar plates. Overnight

precultures were grown for 14-16 h and subcultures were prepared by diluting precultures 1:100

in LB and growing until mid-exponential phase (OD at 500 nm ~ 0.5).

Construction of PA14 mutant strains. Markerless deletion, complementation, and

constitutive-expression strains were made as described previously [21]. Briefly, flanking

sequences containing the target locus were amplified using the primers listed in Supplementary

Table 2 and inserted into pMQ30 through gap repair cloning in *Saccharomyces cerevisiae*

155   InvSc1 [22]. Each plasmid, listed in Supplementary Table 3, was moved into PA14 and resolved

out using selective media. PA14 double recombinants (markerless strains) were confirmed by

PCR.

Construction of PA14 reporter strains. The desired promoter region (500 bp upstream of the

160   *rebP1P2* operon for P$_{rebP1P2}$-*mScarlet*; the synthetic *lac* promoter for P$_{PA1/03/04}$-*mScarlet*) was

amplified using the primers listed in Supplementary Table 1 and ligated into pLD3208, upstream

of *mScarlet*. Resulting plasmids were integrated into a neutral site in the PA14 genome using

biparental conjugation and the plasmid backbone was resolved out via FLP-FRT recombination.

Reporter constructs containing *mScarlet* were isolated using selective media and confirmed by

165   PCR.

Colony biofilm morphology assays. Ten µL of liquid subcultures were spotted on 60 mL of

colony morphology medium (1% tryptone, 1% agar, 40 µg/mL Congo red dye, 20 µg/mL

Coomassie blue dye) and incubated for up to five days in the dark at 25°C with >90% humidity

170   (Percival CU-22L) and imaged daily with an Expression 11000XL scanner (Epson). Images

shown are representative of at least six biological replicates.

Competition assays. Fluorescent, mScarlet-expressing and non-fluorescent cells were mixed

together in a 1:1 ratio from subcultures and grown as mixed-strain biofilms as described above.

175   After three days, biofilms were resuspended in 1 mL of 1% tryptone and homogenized in a bead

mill homogenizer (Omni Bead Ruptor 12) for 99 s on the "high" setting. Serial dilutions of

homogenized cells were plated onto 1% tryptone + 1.5% agar plates for colony forming unit

(CFU) determination. Fluorescent CFUs were determined by imaging with a Typhoon FLA7000

fluorescent scanner (GE Healthcare).

Purification of SDS-insoluble biofilm fraction. Twenty, five-day-old PA14 Δ*pelB-G* colony

biofilms were resuspended in 80 mL of sterile phosphate-buffered saline (PBS) and sonicated

for 2 x 10 s on ice with the microtip of a Sonifier 250 (Branson). SDS and β-mercaptoethanol

were added to final concentrations of 2% and 5%, respectively. After nutating on a Nutating

185   Mixer (Fisher Scientific) at room temperature (RT; ~ 25°C) for 60 min, the sample was aliquoted

into 1.5-mL microfuge tubes and the insoluble fraction was spun down at 16,873 x g for 5 min.

The supernatant was discarded and pellets were pooled together.

Mass spectrometry of SDS-insoluble proteins. The SDS-insoluble pellet was washed three

190   times with Optima water (Fisher Scientific), solubilized in 100 μL of 98% formic acid at room

temperature for 1 h. spin-vacuum dried for 1.5 h, washed with 50% methanol and water, and

then dissolved in 2% RapiGest (Waters Corp.) with 6 mM DTT. Samples were then sonicated,

boiled, cooled, cysteines reduced and alkylated and proteins digested with trypsin as described

previously [23]. Liquid Chromatography mass/ spectrometry (120 min runs) was performed with

195   the Synapt G2 (Waters Corp.) in positive resolution/ion mobility mode with proteins identified

with ProteinLynx Global Server (PLGS) as described previously [24]. Proteins were also identified

and semi-quantitatively measured with a Q Exactive HF (Orbitrap, Thermo Scientific) with

Mascot V. 2.5 as described previously [25]. Identification of the most abundant proteins in these

preparations by the Q Exactive HF were confirmed by the qualitative orthogonal method (Synapt

200   G2 ion mobility/PLGS analysis). The reference proteome UP000000653 for *Pseudomonas*

aeruginosa strain PA14 from UniProt (Release 2015_10, 11/10/2015) was used for all database

searches. All raw MS data files will be publicly available at the MassIVE data repository

(https://massive.ucsd.edu).

205   **Scanning-electron microscopy.** The SDS-insoluble pellet was resuspended in pre-fixative

solution (2% paraformaldehyde, 2.5% glutaraldehyde, 0.0075% L-lysine in PBS), nutated at RT

for 30 min in the dark, washed in PBS, and fixed in 2.5% glutaraldehyde in PBS at RT for 30

min in the dark. Fixed pellets were washed twice in PBS, dehydrated through a series of 60-min

ethanol washes (25%, 50%, 75%, 95%, 3 x 100%) and visualized with a Helios NanoLab

210   DualBeam 660 (FEI).

Fluorescence visualization in *P. aeruginosa* colony biofilms. For whole-colony fluorescence

imaging, colony biofilms were grown as described for morphology assays but on growth medium

without dyes. After three days, bright field images and fluorescence images were visualized with

215   a Zeiss Axio Zoom.V16 fluorescence stereo zoom microscope. Thin sections of colony biofilms

(n ≥ 3 biological replicates) were prepared as described previously [14]. Differential interference

contrast (DIC) and fluorescent confocal images were captured using an LSM800 confocal

microscope (Zeiss).

220 ***A. thaliana* colonization assays.** *Arabidopsis thaliana* ecotype Columbia (Col-0) seeds were

sterilized using standard bleach protocols [26]. Washed, surface-sterilized seeds were

resuspended in 0.1% agar and subjected to 24 h of cold treatment at 4˚C, then plated and

germinated on half-strength MS agar medium containing Gamborg vitamins [27]. Seeds were

incubated at 22˚C with a 12-h-light/12-h-dark photoperiod (100-150 µE/m^2/s; Percival CU-22L)

225 for three weeks. *A. thaliana* seedlings were then inoculated with PA14 using a flood inoculation

assay [28] and incubated under the same light/dark conditions for five days. For visualization of

PA14-infected *A. thaliana*, leaves were excised and *mScarlet* fluorescence was visualized with

a Zeiss Axio Zoom.V16 fluorescence stereo zoom microscope. For quantification of PA14

colonization of *A. thaliana*, seedlings were harvested, weighed, and surface-sterilized with 1 mL

230 of 5% H_2O_2 in water for 3 min. These seedlings were washed five times with sterile PBS and

homogenized in 1 mL of PBS in a bead mill homogenizer on the "high" setting for 99 sec. Serial

dilutions of homogenized tissue were plated onto LB agar plates containing 4 µg/ml tetracycline

and grown overnight at 37°C to select for *P. aeruginosa*, and CFU counts were quantified.

235 ***C. elegans* pathogenicity assays.** Ten µL of overnight PA14 cultures were spotted onto NGM

agar plates [29] and incubated at 24 h at 37˚C, followed by 48 h at 25˚C. Thirty to 35 larval stage 4

(L4) worms were picked onto the PA14-seeded plates and incubated at 25˚C. For visualization of

PA14-infected worms, worms were exposed to PA14 for three days, immobilized in 10 mM

levamisole in water, mounted on a 2% agarose pad on a glass slide, and imaged at 40x on a

240 Zeiss Axio Imager D1 epifluorescence microscope with an AxioCam Mrm. For pathogenicity killing

assays, which were modified from Tan 1999, live/dead worms were counted for up to four days

after plating onto PA14-seeded plates. *unc-44*(*e362*) worms, which exhibit body movement

deficits, were used instead of wild-type to prevent worms from crawling off the plates.

245 **Phylogenetic analysis**. Trees in Supplementary Figures 1 and 2 were generated with Geneious

Prime 2020.2.4 using Muscle for sequence alignment and the neighbor-joining method for tree

building (genetic distance model: Jukes-Cantor).

250 **ACKNOWLEDGEMENTS**

This study was supported by NIH/NIAID grant R01AI103369, an NSF CAREER

award to L.E.P.D., and NSF GRFP grant DGE - 1644869 to B.W. Mass spectrometer acquisition

and operations were funded by New York State Stem Cell Science Board (NYSTEM contracts

CO2361 and C029159) with matching funds from Columbia University and the Columbia Stem

255 Cell Initiative (LMB). We thank Jason Reed (UNC Chapel Hill) for providing *A. thaliana* Col-0

seeds, Hillary Callahan (Barnard College) for technical support in growing seedlings, Martin

Chalfie for provision of *C. elegans unc-44(e362)*, and Iva Greenwald for technical support with

imaging of *C. elegans*. We also thank Zarina Akbary for technical assistance in creating the

$\Delta rebZ\Delta fecI2$ P_{rebP1}-*gfp* strain.

REFERENCES

1. Quackenbush, R. L. & Burbach, J. A. Cloning and expression of DNA sequences associated with the killer trait of Paramecium tetraurelia stock 47. *Proc. Natl. Acad. Sci. U. S. A.* **80**, 250–254 (1983).

2. Gibson, I. A comparison of the refractile bodies (R-bodies) of certain bacteria—III. Nucleotide sequence homologies and R-body function. *Micron and Microscopica Acta* **15**, 253–259 (1984).

3. Gibson, I., Bedingfield, G., Dobbs, H. & Shackleton, J. Effects of various agents on the R bodies of certain bacteria. *Micron and Microscopica Acta* **18**, 71–75 (1987).

4. Pond, F. R., Gibson, I., Lalucat, J. & Quackenbush, R. L. R-body-producing bacteria. *Microbiol. Rev.* **53**, 25–67 (1989).

5. Jurand, A., Preer, J. R. & Rudman, B. M. Further investigations on the prelethal effects of the killing action of kappa containing killer stocks ofParamecium aurelia. *J. Exp. Zool.* **206**, 25–47 (1978).

6. Sonneborn, T. M. Mating Types in Paramecium Aurelia: Diverse Conditions for Mating in Different Stocks; Occurrence, Number and Interrelations of the Types. *Proc. Am. Philos. Soc.* **79**, 411–434 (1938).

7. Raymann, K., Bobay, L.-M., Doak, T. G., Lynch, M. & Gribaldo, S. A genomic survey of Reb homologs suggests widespread occurrence of R-bodies in proteobacteria. *G3* **3**, 505–516 (2013).

8. Winsor, G. L. *et al.* Enhanced annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database. *Nucleic Acids Res.* **44**, D646-53 (2016).

9. Feinbaum, R. L. *et al.* Genome-wide identification of Pseudomonas aeruginosa virulence-related genes using a Caenorhabditis elegans infection model. *PLoS Pathog.* **8**, e1002813

(2012).

10. Schulz, S. *et al.* Elucidation of sigma factor-associated networks in Pseudomonas

    aeruginosa reveals a modular architecture with limited and function-specific crosstalk. *PLoS*

290   *Pathog.* **11**, e1004744 (2015).

11. Polka, J. K. & Silver, P. A. A Tunable Protein Piston That Breaks Membranes to Release

    Encapsulated Cargo. *ACS Synth. Biol.* **5**, 303–311 (2016).

12. Heruth, D. P., Pond, F. R., Dilts, t. J. A. & Quackenbush, R. L. Characterization of Genetic

    Determinants for R Body Synthesis and Assembly in Caedibacter taeniospiralis 47 and 116.

295   *J. Bacteriol.* **176**, 3559–3567 (1994).

13. Stewart, P. S. & Franklin, M. J. Physiological heterogeneity in biofilms. *Nat. Rev. Microbiol.*

    6, 199–210 (2008).

14. Cornell, W. C. *et al.* Paraffin Embedding and Thin Sectioning of Microbial Colony Biofilms

    for Microscopic Analysis. *J. Vis. Exp.* (2018) doi:10.3791/57196.

300 15. Tan, M. W., Mahajan-Miklos, S. & Ausubel, F. M. Killing of Caenorhabditis elegans by

    Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proc. Natl.*

    Acad. Sci. U. S. A. **96**, 715–720 (1999).

16. Walker, T. S. *et al.* Pseudomonas aeruginosa-plant root interactions. Pathogenicity, biofilm

    formation, and root exudation. *Plant Physiol.* **134**, 320–331 (2004).

305 17. Preer, J. R., Jr, Preer, L. B. & Jurand, A. Kappa and other endosymbionts in Paramecium

    aurelia. *Bacteriol. Rev.* **38**, 113–163 (1974).

18. Matsuoka, J.-I. *et al.* A putative TetR-type transcription factor AZC_3265 from the legume

    symbiont Azorhizobium caulinodans represses the production of R-bodies that are toxic to

    eukaryotic cells. *Soil Sci. Plant Nutr.* **63**, 452–459 (2017).

310 19. Flemming, H.-C. *et al.* Biofilms: an emergent form of bacterial life. *Nat. Rev. Microbiol.* **14**,

    563–575 (2016).

20. Donlan, R. M. Biofilms: microbial life on surfaces. *Emerg. Infect. Dis.* **8**, 881–890 (2002).

21. Okegbe, C. *et al.* Electron-shuttling antibiotics structure bacterial communities by modulating cellular levels of c-di-GMP. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5236–E5245 (2017).

22. Shanks, R. M. Q., Caiazza, N. C., Hinsa, S. M., Toutain, C. M. & O'Toole, G. A. Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes from gram-negative bacteria. *Appl. Environ. Microbiol.* **72**, 5027–5036 (2006).

23. Oswald, E. S., Brown, L. M., Bulinski, J. C. & Hung, C. T. Label-free protein profiling of adipose-derived human stem cells under hyperosmotic treatment. *J. Proteome Res.* **10**, 3050–3059 (2011).

24. Yang, W. S. *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156**, 317–331 (2014).

25. Wobma, H. M. *et al.* The influence of hypoxia and IFN-γ on the proteome and metabolome of therapeutic mesenchymal stem cells. *Biomaterials* **167**, 226–234 (2018).

26. Clough, S. J. & Bent, A. F. Floral dip: a simplified method forAgrobacterium-mediated transformation ofArabidopsis thaliana. *The Plant Journal* vol. 16 735–743 (1998).

27. Murashige, T. & Skoog, F. A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiologia Plantarum* vol. 15 473–497 (1962).

28. Ishiga, Y., Ishiga, T., Uppalapati, S. R. & Mysore, K. S. Arabidopsis seedling flood-inoculation technique: a rapid and reliable assay for studying plant-bacterial interactions. *Plant Methods* **7**, 32 (2011).

29. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

30. Rahme, L. G. *et al.* Use of model plant hosts to identify Pseudomonas aeruginosa virulence factors. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 13245–13250 (1997).

a

b

colony biofilms → strong sonication → 2% SDS → pellet insoluble fraction → Mass Spectrometry / Scanning-electron Microscopy

c

Protein Name	Locus (PA14/ PAO1)	Semiquantitative abundance	
		WT	Δ*pel*
HsiB2	PA14_43040/ PA1657	0.7	11.0
OprF	PA14_41570/ PA1777	6.8	10.4
OprL	PA14_51710/ PA0973	4.3	9.4
-	PA14_06990/ PA0537	1.5	8.4
HsiB3	PA14_34070/ PA2365	0.4	5.0
HsiC2	PA14_43030/ PA1658	0.5	4.9
HsiC3	PA14_34050/ PA2366	1.7	3.8
-	PA14_50740/ PA1053	1.9	2.6
-	**PA14_27680**	**0.6**	**2.1**
RebP2	**PA14_27630**	**0.8**	**1.4**
RebP1	**PA14_27640**	**0.8**	**1.4**
AlgP	PA14_69370/ PA5253	0.2	1.3

d

e f g h

Fig.1. Components of the *reb* gene cluster, which are found in diverse pseudomonads, support R-body production by *Pseudomonas aeruginosa* PA14 biofilms. a, Chromosomal arrangement of genes associated with R-body production in representative *Pseudomonas* species and strains. Regulatory genes are shaded yellow and orange. Genes shaded blue are homologous to the R-body structural genes from *C. taeniospiralis*. *PA14_27680* and its homologues, which may code for a novel R-body component, are shaded green. **b,** Schematic showing method for preparation of SDS-insoluble material from PA14 biofilms. **c,** Top 12 most abundant proteins (semi-quantitative) detected in the SDS-insoluble formic acid solubilized fraction of WT (one replicate) and Δ*pel* (two biological replicates) by mass spectrometry. **d,** Left panel: Scanning-electron micrograph (SEM) image of R-bodies (false-colored turquoise) in the insoluble fraction. Scale bar = 1 μm. Right: SEM images, cartoon representations, and dimensions of individual selected R-bodies in fully coiled and fully extended states. Scale bars = 250 nm. **e,** SEM image of a single, partially extended R-body. Scale bar = 400 nm. **f,** Schematic indicating planes of view for panels **g** and **h**. **g,** Bright-field and fluorescence microscopy images of three-day-old biofilms formed by the indicated genotypes, containing the P$_{rebP1}$-*mScarlet* construct. Dotted lines demarcate the edge of the biofilm. Scale bar = 25 mm. **h,** Representative images of thin sections prepared from three-day-old biofilms formed by the indicated genotypes, containing the P$_{rebP1}$-*mScarlet* construct. Dotted lines denote the interfaces with air (top) and agar (bottom). Scale bar = 10 μm.

Fig. 2. *rebP1P2* is expressed during host association and contributes to host colonization and killing, but does not confer an advantage during host-independent growth. a, Fluorescent micrographs of *A. thaliana* leaves infected with WT PA14 expressing mScarlet under the control of the *rebP1* promoter (P_{rebP1}-*mScarlet*), no promoter (MCS-*mScarlet*), or a constitutive promoter ($P_{PA1/04/03}$-*mScarlet*). Smaller panels show images of the whole seedling and the imaged leaf indicated by the white box. Images are representative of nine biological replicates from three experiments. Scale bars = 1 mm. **b,** Images of *C. elegans* fed WT PA14 containing the indicated reporter constructs. Top panels, Nomarski; middle panels, fluorescence; bottom panels, overlay. White boxes indicate regions shown in close-up images. Images are representative of 15 replicates from three experiments. Scale bars = 10 µm. **c,** Three-day-old WT and Δ*rebP1P2* biofilms grown on colony morphology medium. Images are representative of nine replicates from three experiments. Scale bar = 4 mm. **d,** Representation of each strain in colony forming units (CFUs) obtained from mixed-strain colony biofilms grown for three days. Error bars represent standard deviation of biological triplicates. **e,** CFUs obtained from *A. thaliana* seedlings inoculated with the indicated genotypes and incubated for five days, normalized to seedling weight. Each circle represents a biological replicate, with color intensity indicating replicates from the same experiment; colored lines indicate means. p-values were calculated using unpaired, two-tailed t-tests. **f,** Killing kinetics of *C. elegans* following exposure to various PA14 genotypes. The role of *gacA* in *C. elegans* pathogenicity has been described previously [30] and *gacA::Tn* therefore serves as a control for defective killing in this assay. Error bars denote standard deviation and p-values were calculated using unpaired, two-tailed t-tests. n ≥ 4 biological replicates from three experiments.