Pseudomonas aeruginosa PA14 produces R-bodies, extendable protein polymers with roles in host colonization and virulence

Bryan Wang[1], Yu-Cheng Lin[1], Jeanyoung Jo[1], Alexa Price-Whelan[1], Shujuan Tao McDonald [1,2],

5   Lewis M. Brown[1,2], and Lars E.P. Dietrich[1]*

[1]Department of Biological Sciences, Columbia University, New York, NY, USA
[2]Quantitative Proteomics and Metabolomics Center, Columbia University, New York, NY, USA

10   *email: LDietrich@columbia.edu

Abstract

Pseudomonas aeruginosa causes biofilm-based infections in diverse hosts. A majority of sequenced *P. aeruginosa* genomes, including that of the popular model strain UCBPP-PA14 ("PA14"), contain a gene cluster with the potential to confer production of large, extendable protein polymers called R-bodies. R-bodies are best known for their role in killing of paramecia by the bacterium *Caedibacter taeniospiralis*, and genes in PA14's R-body-associated cluster had previously been linked to its virulence. However, the ability to produce R-bodies had not been investigated in this strain. Here we show that R-bodies are among the most abundant proteins present in SDS-insoluble preparations from PA14 biofilms and that two adjacent regulatory genes, including the gene coding for the sigma factor FecI2, are required for expression of R-body structural genes. Subpopulations of PA14 cells express R-body structural genes during biofilm growth and during colonization of plant and animal hosts. Moreover, R-body production is required for wild-type levels of colonization of *A. thaliana* seedlings and full virulence in a *C. elegans* killing assay. Our results suggest that R-body production may contribute to the observed differences in virulence among *P. aeruginosa* isolates.

Introduction

R-bodies are enigmatic biological machines that have been characterized most extensively in *Caedibacter taeniospiralis*, an obligate endosymbiont of paramecia. These structures are composed of ~ 12 kDa monomers that polymerize to form hypercoiled ribbons ~ 0.5 μm in diameter [1]. In response to changes in parameters such as pH, temperature, or ionic strength [2–4], R-body coils extend into 10-20-μm-long needle-like structures (i.e., ~ 10X the length of the producing cell), lysing the bacterium [4,5]. In the *Caedibacter-Paramecium* endosymbiosis, tolerant host paramecia shed *C. taeniospiralis* into the environment, where the bacteria can be taken up by neighboring, sensitive paramecia. R-body-mediated lysis is triggered by lysosomal conditions and thought to release an unidentified factor that is toxic to sensitive paramecia. R-body-producing endosymbionts therefore confer onto their host cell a "killing" phenotype and a competitive advantage in a phenomenon that was first described by Tracy M. Sonneborn in 1938 [6]. Subsequent genetic studies have implicated three genes--*rebA, B,* and *D*--in *C. taeniospiralis* R-body production. Oddly, RebA, B, and D are all homologous to each other, but RebA and RebB constitute the primary *C. taeniospiralis* R-body components [7–9].

The opportunistic pathogen *Pseudomonas aeruginosa* is a major cause of biofilm-based
infections. A substantial fraction (147/241) of *P. aeruginosa* complete strain genomes, including
that of strain UCBPP-PA14 ("PA14"), contain *reb* gene homologs as part of a cluster that
includes the gene for the sigma factor FecI2 and targets of FecI2 [10]; we refer to this region as
the "*reb* cluster". Genes in the *reb* cluster have been identified as contributing to increased
virulence when distinct *P. aeruginosa* isolates are compared [11,12]. Notably, the popular model
strain PAO1 does not contain a *reb* cluster and this is consistent with its relatively low virulence
in a *Caenorhabditis elegans* infection assay. These observations, combined with the known
contributions of R-bodies to host damage in other systems [4,13,14], led us to hypothesize that
PA14 produces R-bodies that contribute to pathogenicity.

In this report we detail characteristics of the PA14 *reb* cluster. We provide evidence of R-body
production by PA14 biofilms and show that PA14 *rebP1P2*, an operon encoding *C.
taeniospiralis reb* gene homologs, is required for this. We also find that cells in PA14 liquid
cultures and biofilms exhibit stochastic P_{rebP1} expression that is dependent on local regulatory
genes and that this promoter is active during interactions with *Arabidopsis thaliana* and
Caenorhabditis elegans hosts. Finally, we show that *rebP1P2* is necessary for wild type-like
levels of colonization of *A. thaliana* seedlings and virulence in *C. elegans*. Together, these
findings raise the possibility that R-body production is a component of the marked strain-
dependent variation in pathogenicity observed for *P. aeruginosa* [15].

Results

Phylogenetic analysis of pseudomonad R-body-associated genes and description of the PA14
reb *cluster*

A previous analysis showed that homologs of *C. taeniospiralis reb* genes are broadly distributed
throughout the proteobacterial clade [16]. To investigate the phylogenetic relationships between
Reb homologs from proteobacterial genomes, we BLASTed PA14_27640, which is the PA14
protein most similar to *C. taeniospiralis* RebB*,* against all complete bacterial genomes used for
the prior analyses conducted by [16] plus 14 representative pseudomonads. We generated a tree
that shows the relationships of all Reb homologs from these proteobacteria (**Figure 1a**). In this
tree, *C. taeniospiralis* RebA and RebB form a distinct grouping and are closely related to Reb
homologs from non-pseudomonad gammaproteobacteria, such as those in the genera

Shewanella and *Allivibrio* (**Supplementary Figure 1**). Reb homologs from pseudomonads fall
into two groups: (i) one that contains *C. taeniospiralis* RebD and PA14_27685 and for which we
have therefore designated its members as "RebD"; and (ii) one that is distinct from the *C.
taeniospiralis* Rebs but that includes representatives from many *reb* cluster-containing
pseudomonads and chromobacteria, and for which we have therefore designated its members
as "RebP". The two *rebP* homologs present in PA14, *PA14_27640* and *PA14_27630*, are
predicted to form an operon that we refer to as *rebP1P2*.

We examined the chromosomal regions of representative pseudomonad strains that contain
rebD and *rebP1P2* and refer to the strain-specific groups of genes in these loci as "*reb* clusters".
In pseudomonads that contain *rebD* and *rebP1P2*, there are three additional ORFs that are well-
conserved. These include two regulatory genes, *PA14_27700* and *fecI2*, which code for a
transcription factor and sigma factor, respectively, and which in a prior study were required for
full virulence in a *C. elegans* killing assay [17] (**Figure 1b**). We have given *PA14_27700* the
designation "*rcgA*" (for <u>r</u>eb <u>c</u>luster <u>g</u>ene <u>A</u>). The third gene, *PA14_27680*, was previously shown
(together with *PA14_27700*) to contribute to PA14's enhanced virulence when it was compared
to *P. aeruginosa* strain PAO1 in *C. elegans* infection studies [11]. *PA14_27680* is conserved in all
pseudomonads that contain at least one *reb* structural gene, suggesting that it may be
associated with R-body function and we have therefore named it "*rapA*" (<u>R</u>-body-<u>a</u>ssociated
<u>p</u>rotein <u>A</u>). Although *rebP* homologs are scattered across the phylogenetic tree of the genus
Pseudomonas (**Supplementary Figure 2**), homologs of the regulatory genes show strong
conservation in strains that contain *rebP* genes, both within the species *P. aeruginosa* (in
146/147 *rebP*-containing genomes; **Supplementary File 1**) and among other pseudomonads
(**Figure 1b**).

PA14 cells growing in biofilms produce R-bodies

For most bacteria that contain *reb* genes, the ability to produce R-bodies has not been
investigated. To determine whether PA14 produces R-bodies, we applied a modified R-body
enrichment protocol [18] to biofilms formed by the WT and by Δ*pel*, a mutant that is unable to
produce the major PA14 exopolysaccharide and that therefore yields biofilms that are more
amenable to disruption. We performed protein mass spectrometry (MS) and scanning-electron
microscopy (SEM) on the isolated SDS-insoluble fractions (**Figure 2a**). Strikingly, three
products of the *reb* cluster—RapA, RebP1, and RebP2–were detected by MS at similar levels

and were among the 12 most abundant proteins in our sample (**Figure 2b**). One other
uncharacterized protein coded for by the *reb* gene cluster, PA14_27675, was also detected
specifically in the SDS-insoluble fractions prepared from Δ*pel* biofilms and at levels two times
lower than RebP1 and RebP2 (PA14_27675 was not detected in preparations from WT biofilms)
(**Figure 2b**). Consistent with the MS results, our SEM analyses revealed the presence of R-
bodies in preparations from WT PA14 biofilms (**Figure 2c**). We detected R-bodies in various
states ranging from hypercoiled (~ 450 nm long) to extended (~ 5.5 µm long) (**Figure 2c, d, e**).
These structures were not observed in a deletion mutant lacking the *rebP1P2* operon
(Δ*rebP1P2*) (**Supplementary Figure 3**). Taken together, our data indicate that R-bodies are
produced by PA14 biofilms.

rebP1P2 *expression is stochastic and dependent on RcgA and FecI2, in both liquid cultures and*
biofilms

Having observed that *rebP1P2* is required for R-body production by PA14 biofilms, we sought to
characterize the expression of this locus. We generated a construct that reports expression from
the *rebP1P2* promoter (P$_{rebP1}$) as red (mScarlet) fluorescence and inserted it into a neutral site
on the chromosome in WT PA14. We also generated similar strains with mScarlet driven by a
constitutive promoter (P$_{PA1/04/03}$-*mScarlet*) or a promoterless sequence in the WT background to
use as controls. We grew the WT P$_{rebP1}$ and control reporter strains in shaken liquid batch
cultures to examine expression over the growth curve. We found that P$_{rebP1}$ showed modest
activity compared to that of the constitutive promoter and that it was expressed specifically in
the stationary phase of growth (**Figure 3a**). We then created a dual reporter strain containing
P$_{rebP1}$-*mScarlet* and P$_{PA1/04/03}$-*gfp* and grew it to stationary phase in liquid culture before
examination via fluorescence microscopy. This imaging revealed that the low relative
fluorescence of the P$_{rebP1}$-*mScarlet* strain (**Figure 3a**) was not due to low uniform expression but
instead to stochastic expression of *rebP1P2*, with mScarlet fluorescence visible in just 0.26% ±
0.006% of the cell population (**Figure 3b**).

Based on the conservation of *rcgA* and *fecI* in *Pseudomonas reb* clusters (**Figure 1b;**
Supplementary Figure 1; Supplementary File 1), we reasoned that the encoded
transcriptional regulators control expression of *rebP1P2*. To test this, we moved the P$_{rebP1}$-
mScarlet construct into the PA14 Δ*rcgA* and Δ*fecI2* mutant backgrounds. We observed P$_{rebP1}$

activity in WT liquid cultures and colony biofilms that was dependent on the presence of *rcgA* and *fecI2* (**Figure 3c, d, Supplementary Figure 4**). Deletion of one or both of these putative regulatory genes resulted in abrogation of *rebP1P2* expression, indicating that RcgA and FecI2 are both required for P*rebP1* activity. Complementation of these genes back into their endogenous loci restored P*rebP1* expression (**Figure 3c, d and Supplementary Figure 4**).

Because biofilm development leads to the formation of steep chemical gradients that can affect gene expression patterns along the z-axis [19], we chose to visualize P*rebP1* activity across colony depth. To do this, we employed a thin sectioning protocol to prepare vertical sections taken from the center region of three-day-old colony biofilms [20]. Confocal microscopy of the dual reporter (P*rebP1*-*mScarlet* P*PA1/04/03*-*gfp*) biofilm sections revealed that cells expressing *rebP1P2* are present in thin striations such that both bright and dark cells are situated at the same depth (**Figure 3d**). The constitutive reporter (P*PA1/04/03*-*gfp*) showed fluorescence that is more consistent between cells at the same depth (**Figure 3c, d**). In agreement with our findings for liquid culture, P*rebP1*-driven expression in biofilms appears to be stochastic. The vertical arrangement of P*rebP1*-active cells may indicate that expression status is heritable during vertical growth in the biofilm.

In many hosts and various infection sites in humans, biofilm formation by pathogens such as *P. aeruginosa* is critical to bacterial colonization and virulence. Biofilm formation also exacerbates the challenge of treating resistant infections. Matrix secretion is a defining feature of biofilms; the matrix functions as the "glue" that holds bacterial cells together and can facilitate attachment to the host [21,22]. Having observed R-body production by PA14 biofilms, we sought to examine whether this activity contributes to biofilm development, matrix production, or fitness during biofilm growth. We examined biofilm development using a standardized colony morphology assay, in which the dye Congo red is provided in the medium and binds to biofilm matrix. We found that matrix production and biofilm development were unaltered in the Δ*rebP1P2* mutant (**Figure 3e**). Furthermore, Δ*rebP1P2* showed no fitness disadvantage when grown in mixed-strain colony biofilms with the WT parent (**Figure 3f**).

rebP1P2 *is expressed and contributes to PA14 pathogenicity in plant and nematode hosts*

The fact that homologs of the *C. taeniospiralis* R-body structural genes are found in diverse free-living bacteria suggests that their roles extend beyond endosymbiotic interactions [16]. To

examine whether *rebP1P2* is expressed during interactions with hosts, we employed the P*rebP1*-*mScarlet* reporter strain in *Arabidopsis thaliana* seedling and *Caenorhabditis elegans* gut colonization assays. In both hosts, as in colony biofilms, *rebP1P2* expression patterns were distinct from those exhibited by a promoter driving constitutive expression of mScarlet (P*PA1/04/03*-

185   *mScarlet*). While constitutive mScarlet production was uniform and fuzzy across colonized regions of hosts (**Figure 4a, b**), *rebP1P2* expression was punctate and restricted to a subset of PA14 cells (**Figure 4a**, **b**). The distinct *rebP1P2* expression patterns we observed in the colony biofilm and host contexts may indicate that there is an advantage to restricting R-body production to a subset of cells in the population.

In *C. taeniospiralis* and the plant symbiont *Azorhizobium caulinodans*, R-body production has been linked to host toxicity [4,13,14], and *P. aeruginosa* forms biofilms and acts as an extracellular pathogen during colonization of a range of hosts including plant and nematode models [23,24]. We hypothesized that R-bodies contribute to PA14 host colonization and/or pathogenicity and

195   therefore performed established assays to test such contributions of PA14 R-body production in our two host systems. In *A. thaliana*, we quantified the extent to which PA14 WT and mutant strains colonized seedlings after inoculation and five days of plant growth. For the *C. elegans* virulence assay, we quantified percent survival for up to four days after synchronized populations of worms were exposed to PA14 WT and mutant strains. When compared to WT

200   PA14, Δ*rebP1P2* displayed attenuated host colonization (**Figure 4c**) and virulence (**Figure 4d**); both phenotypes were restored to WT levels upon complementation of *rebP1P2* (**Figure 4c, d**). In combination with the results of our biofilm competition assays (**Figure 3f**), these results indicate that the effects of R-body production are specific to the host-associated lifestyle.

205   **Discussion**

We have identified genes required for R-body production by *P. aeruginosa* PA14. In *C. taeniospiralis*, the proteins RebA, RebB, and RebD have been implicated in R-body production. While these proteins are homologous to each other, RebD is an outlier in that it is shorter than

210   RebA and RebB and in that its expression is not required for R-body production. PA14 contains one homolog to RebD, and two RebB homologs that we have designated as "RebP1" and "RebP2" due to their distinct phylogenetic grouping. Similar to the case in *C. taeniospiralis*, RebP1 and RebP2 are longer than PA14 RebD, and RebP1 and RebP2 were both detected in SDS-insoluble preparations from PA14 biofilms, while RebD was not. Expression of the putative

215 *rebP1P2* operon is regulated by the transcription factor RcgA and sigma factor FecI2. RcgA and FecI2 had previously been associated with virulence [12,17], but their exact roles have eluded definition. We have also provided evidence that *rebP1P2* expression is stochastic and that it can be observed both in independent biofilms and during association with hosts. While expression patterns between these two systems appear to be similar, an advantage to R-body production is

220 detectable only during host colonization, suggesting that R-body function is specific to infection.

Though R-body production was initially thought to be limited to parasitic bacteria [25], it has since been shown that *reb* gene homologs are found in diverse proteobacteria, including many non-symbionts [16]. Our results indicate that R-bodies made by *P. aeruginosa* PA14 contribute to

225 colonization of a plant host and to virulence in a *C. elegans* model of infection. In a recent study, the regulator RcgA was found to be necessary for *C. elegans* ribosomal injury that occurs during PA14 infection [26]. Together with our findings, the results of this study hint at a role for R-bodies in releasing a bacterial enzyme, or activating a host enzyme, that cleaves *C. elegans* rRNA. Aside from *C. taeniospiralis*, and now *P. aeruginosa*, to our knowledge a role for R-bodies in

230 host damage has been observed only for the plant symbiont *A. caulinodans.* The *reb* genes in this plant symbiont are repressed by the regulatory protein PraR under both free-living and host-associated conditions, but deletion of PraR allows R-body production and leads to killing of host plant cells [14]. Because R-body production has not been observed in wild type *A. caulinodans*, its significance in the context of bacterium-plant symbiosis is unclear [14]. Nevertheless, the fact that

235 the damaging effects of R-bodies have now been described for three divergent bacterial species hints at a general role for these structures in interactions between proteobacteria and eukaryotes.

Caedibacter R-body extension is triggered by low pH and occurs in phagosomes of naive

240 paramecia, prompting *Caedibacter* lysis and the release of other toxins within *Caedibacter* cells and thereby killing the eukaryotic predator [4,5]. We speculate that PA14 R-body extension is similarly triggered in a host-specific manner and that cells in the R-body-producing subpopulation act as "kamikazes" to release other bacterial products that facilitate colonization and enhance virulence. Our findings raise the possibility that R-body production contributes to

245 the behaviors of diverse free-living bacteria via analagous mechanisms during association with other organisms.

Methods

Bacterial strains and growth conditions. Strains used in this study are listed in **Supplementary Table 1**. Cultures of *Pseudomonas aeruginosa* strain UCBPP-PA14 (PA14; [27] were grown in lysogeny broth (LB; 1% tryptone, 1% NaCl, 0.5% yeast extract [28] in 13 mm x 100 mm culture tubes at 37°C with shaking at 250 rpm unless otherwise indicated. Biological replicates were

255   inoculated from distinct clonal-source colonies grown on LB + 1.5% agar plates. Overnight precultures were grown for 14-16 h and subcultures were prepared by diluting precultures 1:100 in LB in 13 mm x 100 mm culture tubes and growing at 37°C with shaking at 250 rpm until mid-exponential phase (OD at 500 nm ~ 0.5). Subcultures were used in experiments unless otherwise noted.

Construction of mutant strains. Markerless deletion and complementation strains were made as described previously [29]. Briefly, ~ 1 kb of flanking sequence from each side of the target locus were amplified using the primers listed in **Supplementary Table 2** and inserted into pMQ30 through gap repair cloning in *Saccharomyces cerevisiae* InvSc1 [30]. Each plasmid, listed in

265   **Supplementary Table 2**, was transformed into *Escherichia coli* strain UQ950, verified by restriction digests and sequencing, and moved into *P. aeruginosa* PA14 using biparental conjugation. PA14 single recombinants were selected on LB agar plates containing 100 μg/ml gentamicin. Double recombinants (markerless mutants) were selected on a modified LB medium (containing 10% sucrose and lacking NaCl) and genotypes were confirmed by PCR.

270   Combinatorial mutants were constructed by using single mutants as hosts for biparental conjugation as indicated in **Supplementary Table 1**.

Phylogenetic analysis. All trees were generated with Geneious Prime 2020.2.4 using Muscle for sequence alignment and the neighbor-joining method for tree building (genetic distance model:

275   Jukes-Cantor (**Supplementary Figure 1**); Tamura-Nei (**Supplementary Figure 2**)). Trees in Figure 1a and **Supplementary Figure 2** were displayed using the online Interactive Tree Of Life (iTOL) tool [31].

Construction of PA14 reporter strains. A transcriptional reporter for the *rebP1P2* operon was

280   constructed using primers listed in **Supplementary Table 2** to amplify the promoter region (500 bp upstream of the *rebP1P2* operon), adding an SpeI digest site to the 5' end and an EcoRI digest site to the 3' end of the promoter. Purified PCR products were digested and ligated into the

multiple cloning site of the pLD3208 vector, upstream of the *mScarlet* coding sequence. This

plasmid (pLD3210) was transformed into *E. coli* UQ950, verified by sequencing, and integrated

285   into a neutral site in the PA14 genome using biparental conjugation with *E. coli* S17. PA14 single

recombinants were selected on M9 minimal medium agar plates (47.8 mM $Na_2HPO_4 \cdot 7H_2O$, 22

mM KH_2PO_4, 8.6 mM NaCl, 18.6 mM NH_4Cl, 1 mM $MgSO_4$, 0.1 mM $CaCl_2$, 20 mM sodium citrate,

1.5% agar) containing 100 µg/ml gentamicin. The plasmid backbone was resolved out of PA14

using FLP-FRT recombination by introduction of the pFLP2 plasmid [32] and selected on M9

290   minimal medium agar plates containing 300 µg/ml carbenicillin and confirmed on LB agar plates

without NaCl and modified to contain 10% sucrose. The presence of *mScarlet* in the final clones

was confirmed by PCR. To create the strain that constitutively expresses *mScarlet*, the

constitutively expressing synthetic *lac* promoter ($P_{PA1/03/04}$) was ligated into pLD3208 and mating

the resulting plasmid, pLD3433, into the *glmS* locus of the PA14 genome via triparental mating

295   with *E. coli* donor strain BW29427 and *E. coli* helper strain ß2155.

Colony biofilm morphology assays. Ten µl of liquid subcultures were spotted onto 60 mL of

colony morphology medium (1% tryptone, 1% agar containing 40 µg/mL Congo red dye and 20

µg/mL Coomassie blue dye) in 100mm x 100mm x 15mm square Petri dishes (LDP D210-16).

300   Plates were incubated for up to five days in the dark at 25°C with >90% humidity (Percival CU-

22L) and imaged daily with an Epson Expression 11000XL scanner. Images shown are

representative of at least six biological replicates.

Competition mixing assays. OD at 500 nm of subcultures were determined in a Synergy 4 plate

305   reader (BioTek) and subcultures were mixed together in a 1:1 ratio of fluorescent, mScarlet-

expressing and non-fluorescent cells. Ten µl of this mixture were spotted onto colony morphology

medium and grown as described above. After three days, biofilms were harvested, resuspended

in 1 mL of 1% tryptone, and homogenized in an Omni Bead Ruptor 12 bead mill homogenizer for

99 s on the "high" setting. Serial dilutions of homogenized cells were plated onto 1% tryptone +

310   1.5% agar plates and grown overnight at 37˚C and colony forming units (CFU) were determined.

Fluorescent CFUs were determined by imaging with a Typhoon FLA7000 fluorescent scanner

(GE Healthcare).

Purification of SDS-insoluble biofilm fraction. To isolate the sodium dodecyl sulfate-insoluble

315   (SDS-insoluble) fraction of biofilms, we used a PA14 strain defective in producing the main

exopolysaccharide Pel (Δ*pelB-G*). Biological replicates of five-day-old colony biofilms (n = 20,

grown on morphology agar as described above) were harvested from the growth medium with a sterile pipette tip, resuspended in 80 mL of sterile phosphate-buffered saline (PBS), and sonicated for 2 x 10 s on ice with the microtip of a Sonifier 250 (Branson). SDS and β-mercaptoethanol

320 were added to final concentrations of 2% and 5%, respectively, and the sample was nutated on a Nutating Mixer (Fisher Scientific 88 861 043) at room temperature (RT; ~ 25°C) for 60 min. The sample was then transferred into 1.5-mL microfuge tubes and the insoluble fraction spun down at 16,873 x g for 5 min. The supernatant was discarded and the pellets (i.e., the Congo red-binding, SDS-insoluble fraction) were pooled together.

Mass spectrometry of SDS-insoluble proteins. The SDS-insoluble pellet was washed three times with Optima water (Fisher Scientific), solubilized in 100 µL of 98% formic acid at room temperature for 1 h. spin-vacuum dried for 1.5 h, washed with 50% methanol and water, and then dissolved in 2% RapiGest (Waters Corp.) with 6 mM DTT. Samples were then sonicated,

330 boiled, cooled, cysteines reduced and alkylated and proteins digested with trypsin as described previously [33]. Liquid Chromatography mass/ spectrometry (120 min runs) was performed with the Synapt G2 (Waters Corp.) in positive resolution/ion mobility mode with proteins identified with ProteinLynx Global Server (PLGS) as described previously [34]. Proteins were also identified and semi-quantitatively measured with a Q Exactive HF (Orbitrap, Thermo Scientific) with

335 Mascot V. 2.5 as described previously [35]. Identification of the most abundant proteins in these preparations by the Q Exactive HF were confirmed by the qualitative orthogonal method (Synapt G2 ion mobility/PLGS analysis). The reference proteome UP000000653 for *Pseudomonas aeruginosa* strain PA14 from UniProt (Release 2015_10, 11/10/2015) was used for all database searches. All raw MS data files will be publicly available at the MassIVE data repository

340 (https://massive.ucsd.edu).

Scanning-electron microscopy. The SDS-insoluble pellet was resuspended in pre-fixative solution (2% paraformaldehyde, 2.5% glutaraldehyde, 0.0075% L-lysine in PBS) and nutated at RT for 30 min in the dark. The pellet was then washed in sterile PBS and fixed in 2.5%

345 glutaraldehyde in PBS at RT for 30 min in the dark. Fixed pellets were washed twice in PBS and dehydrated through a series of 60-min ethanol washes (25%, 50%, 75%, 95%, 3 x 100% ethanol). Samples were visualized with a Helios NanoLab DualBeam 660 (FEI).

Liquid culture growth assays. Overnight precultures were diluted 1:100 in 200 µl of 1% tryptone

350 in a flat bottom, polystyrene, 96-well plate (Greiner Bio-One 655001) and incubated at 25°C with

continuous shaking on the medium setting in a Biotek Synergy 4 or Biotek Synergy H1 plate reader. The expression of mScarlet was assessed by taking fluorescence readings at excitation and emission wavelengths of 569 nm and 599 nm, respectively, every 30 minutes for up to 24 h. Growth was assessed by taking OD readings at 500 nm simultaneously with

355   the fluorescence readings.

Quantification of cells expressing P$_{rebP1}$-mScarlet. Five µl of overnight PA14 cultures were mounted on a 2% agarose pad on a glass slide, and imaged at 63x on a Zeiss Axio Imager D1 epifluorescence microscope with an AxioCam MRm. Five independent fields of views were

360   captured at random for each biological replicate. Number of cells in each image was determined using the MicrobeJ plugin in Fiji [36,37].

Fluorescence visualization in *P. aeruginosa* colony biofilms. For whole-colony fluorescence imaging, ten µl of liquid subcultures were spotted onto 1% tryptone, 1% agar and colony biofilms

365   were grown in the dark at 25°C with >90% humidity (Percival CU-22L). After three days, bright field images and fluorescence images were visualized with a Zeiss Axio Zoom.V16 fluorescence stereo zoom microscope (excitation, 545 nm; emission, 605 nm for imaging of mScarlet; excitation, 488 nm; emission, 509 nm for imaging of GFP).

370   Thin sections of PA14 colony biofilms were prepared as described previously [20]. Briefly, after three days of growth as described above, biofilms were overlaid with 1% agar and sandwiched biofilms were lifted from the bottom layer and fixed overnight in 4% paraformaldehyde in PBS at 25°C for 24 h in the dark. Fixed biofilms were washed twice in PBS and dehydrated through a series of 60-min ethanol washes (25%, 50%, 70%, 95%, 3 x 100% ethanol) and cleared via three

375   60-min incubations in Histoclear-II (National Diagnostics); these steps were performed using an STP120 Tissue Processor (Thermo Fisher Scientific). Biofilms were then infiltrated with wax via two separate 2-h washes of 100% paraffin wax (Paraplast Xtra) at 55°C, and allowed to polymerize overnight at 4°C. Trimmed blocks were sectioned in 10-µm-thick sections perpendicular to the plane of the biofilm, floated onto water at 45°C, and collected onto slides.

380   Slides were air-dried overnight, heat-fixed on a hotplate for 1 h at 45°C, and rehydrated in the reverse order of processing. Rehydrated colonies were immediately mounted in TRIS-buffered DAPI:Fluorogel (Electron Microscopy Sciences) and overlaid with a coverslip. Differential interference contrast (DIC) and fluorescent confocal images were captured using an LSM800

confocal microscope (Zeiss). At least three biological replicates of each strain were prepared in
385   this manner.

A. thaliana colonization assays. _Arabidopsis thaliana_ ecotype Columbia (Col-0) seeds were
sterilized using standard bleach protocols [38]. Washed, surface-sterilized seeds were
resuspended in 0.1% agar and subjected to 24 h of cold treatment at 4°C, then plated and
390   germinated on half-strength MS agar medium containing Gamborg vitamins [39]. Seeds were
incubated at 22°C with a 12-h-light/12-h-dark photoperiod (100-150 μE/m^2/s; Percival CU-22L)
for three weeks. _A. thaliana_ seedlings were then inoculated with PA14 using a flood inoculation
assay [40] and incubated under the same light/dark conditions for five days. For visualization of
PA14-infected _A. thaliana_, leaves were excised and _mScarlet_ fluorescence was visualized with
395   a Zeiss Axio Zoom.V16 fluorescence stereo zoom microscope. For quantification of PA14
colonization of _A. thaliana_, seedlings were harvested, weighed, and surface-sterilized with 1 mL
of 5% H_2O_2 in water for 3 min. These seedlings were washed five times with sterile PBS and
homogenized in 1 mL of PBS in a bead mill homogenizer on the "high" setting for 99 sec. Serial
dilutions of homogenized tissue were plated onto LB agar plates containing 4 μg/ml tetracycline
400   and grown overnight at 37°C to select for _P. aeruginosa_, and CFU counts were quantified.

C. elegans pathogenicity assays. Ten μl of overnight PA14 cultures were spotted onto NGM
agar plates [41] and incubated at 24 h at 37°C, followed by 48 h at 25°C. Thirty to 35 larval stage 4
(L4) worms were picked onto the PA14-seeded plates and incubated at 25°C. For visualization of
405   PA14-infected worms, worms were exposed to PA14 for three days, immobilized in 10 mM
levamisole in water, mounted on a 2% agarose pad on a glass slide, and imaged at 20x and 63x
on a Zeiss Axio Imager D1 epifluorescence microscope with an AxioCam MRm. For pathogenicity
killing assays, which were modified from [23], live/dead worms were counted for up to four days
after plating onto PA14-seeded plates. _unc-44_(_e362_) worms, which exhibit body movement
410   deficits, were used instead of wild-type to prevent worms from crawling off the plates.

ACKNOWLEDGEMENTS
This study was supported by NIH/NIAID grant R01AI103369, an NSF CAREER
415   award to L.E.P.D., and NSF GRFP grant DGE - 1644869 to B.W. Mass spectrometer acquisition
and operations were funded by New York State Stem Cell Science Board (NYSTEM contracts
CO2361 and C029159) with matching funds from Columbia University and the Columbia Stem

Cell Initiative (LMB). We thank Jason Reed (UNC Chapel Hill) for providing *A. thaliana* Col-0 seeds, Hillary Callahan (Barnard College) for technical support in growing seedlings, Martin Chalfie for provision of *C. elegans unc-44(e362)*, and Iva Greenwald for technical support with imaging of *C. elegans*. We also thank Zarina Akbary for technical assistance in creating strain LD3137.

Fig.1. Components of the *reb* gene cluster are found in diverse pseudomonads.

a, Phylogenetic tree of proteins from 38 bacterial strains that are homologous to *C. taeniospiralis* RebB. The bacterial strains included are those with complete genomes from Figure 6 of Raymann *et al*. [16] plus 14 representative pseudomonads (see Figure S1 for the complete annotated tree). *C. taeniospiralis* RebA and RebB are shown in purple. The cluster containing *C. taeniospiralis* RebD and its *Pseudomonas* homologs is shown in orange. We have assigned the designation "RebP" to Reb homologs that cluster with *P. aeruginosa* PA14_27630 and PA14_27640 (shown in blue). **b**, Chromosomal arrangement of genes associated with R-body production in a set of strains that represent all arrangements found in pseudomonads. Regulatory genes are shaded yellow. Genes shaded blue are homologs of *rebP* while genes shaded orange are homologous to *C. taeniospiralis rebD*. *rapA* (*PA14_27680*) and its homologs, which may code for a novel R-body component, are shaded green. The asterisk denotes a gene that is annotated in BioCyc (PA14_ RS11205) [42], but not in the Pseudomonas Genome DB [43].

a

colony biofilms → strong sonication → 2% SDS → pellet insoluble fraction → MS / SEM

b

Protein Name	Locus (PA14/ PAO1)	Semiquantitative abundance	
		WT	Δ*pel*
HsiB2	PA14_43040/ PA1657	0.7	11.0
OprF	PA14_41570/ PA1777	6.8	10.4
OprL	PA14_51710/ PA0973	4.3	9.4
-	PA14_06990/ PA0537	1.5	8.4
HsiB3	PA14_34070/ PA2365	0.4	5.0
HsiC2	PA14_43030/ PA1658	0.5	4.9
HsiC3	PA14_34050/ PA2366	1.7	3.8
RplT	PA14_28680/ PA2741	0.0	2.9
-	PA14_50740/ PA1053	1.9	2.6
-	**PA14_27680**	**0.6**	**2.1**
RebP2	**PA14_27630**	**0.8**	**1.4**
RebP1	**PA14_27640**	**0.8**	**1.4**
AlgP	PA14_69370/ PA5253	0.2	1.3
OprM	PA14_05550/ PA0427	0.0	1.2
Pnp	PA14_62710/ PA4740	0.0	1.1
-	PA14_70400/ PA5333	0.3	1.0
PchI	PA14_09320/ PA4222	0.5	0.8
FusA1	PA14_08820/ PA4266	0.0	0.8
-	**PA14_27675**	**0.0**	**0.7**
LecB	PA14_20610/ PA3361	0.0	0.5

c

e

d

445 nm | 415 nm | 5.4 µm | 210 nm

Fig 2. R-bodies are produced by *Pseudomonas aeruginosa* PA14 biofilms. a, Schematic showing method for preparation of SDS-insoluble material from PA14 biofilms. **b**, Top 20 most abundant proteins detected in the SDS-insoluble, formic acid-solubilized fraction of WT (one replicate) and Δ*pel* (two biological replicates) biofilms by mass spectrometry (MS). **c**, Scanning electron micrograph (SEM) image of R-bodies (false-colored turquoise) in the WT sample prepared as shown in **a**. Scale bar is 1 µm. **d**, SEM images, cartoon representations, and dimensions of individual selected R-bodies in fully coiled and fully extended states. Scale bars is 250 nm. **e**, SEM image of a single, partially extended R-body. Scale bar is 400 nm.

a

b P_rebP1-*mScarlet* P_PA1/04/03-*gfp*

P_rebP1 expression: 0.26%

OD_500 / reporter

- P_PA1/04/03-*mScarlet (consititutive)*
- P_rebP1-*mScarlet*
- MCS-*mScarlet (promoterless)*

c P_rebP1-*mScarlet* P_PA1/04/03-*gfp*

WT   ΔrcgAfecI2   ΔrcgA   ΔfecI2   ΔrcgAfecI2::rcgAfecI2

biofilm (top view)

d

biofilm (thin section)

air

biofilm

agar

e WT   ΔrebP1P2

f Colony forming units

0   50   100%

WT : **WT**

WT : **ΔrebP1P2**

ΔrebP1P2 : **WT**

Fig 3. Expression of PA14 R-body genes is stochastic and controlled by RcgA and FecI2*, and does not confer a competitive advantage during growth in biofilms.* a, Reporter strain growth and expression of mScarlet under the control of the *rebP1P2* promoter (P*rebP1*-*mScarlet*), no promoter (MCS-*mScarlet*), or a constitutive promoter (P*PA1/04/03*-*mScarlet*) in shaken 1% tryptone liquid cultures incubated at 25 °C. **b**, Fluorescent image of PA14 cells containing P*rebP1* transcriptional reporter construct (P*rebP1*-*mScarlet*) and a constitutive promoter driving *gfp* (P*PA1/04/03*-*gfp*) in shaken 1% tryptone liquid cultures grown at 25 °C for 16 h. Image is representative of 6 replicates from three experiments. Scale bar is 20 µm. **c**, Left: Schematic indicating plane of view for right-hand panels. Right: Bright-field and fluorescence microscopy images of three-day-old biofilms formed by the indicated genotypes, containing the P*rebP1* transcriptional reporter construct (P*rebP1*-*mScarlet*) and constitutively expressing GFP (via P*PA1/04/03*-*gfp*). Dotted lines demarcate the edges of biofilms. Scale bar is 2 mm. **d**, Left: Schematic indicating plane of view for right-hand panels. Right: Representative images of thin sections prepared from three-day-old biofilms formed by the strains as indicated for **c**. Dotted lines denote the interfaces with air (top) and agar (bottom). Scale bar is 20 µm. **e**, Three-day-old WT and Δ*rebP1P2* biofilms grown on colony morphology medium. Images are representative of nine replicates from three experiments. Scale bar is 4 mm. **f**, Relative percentages of colony forming units (CFUs) of each strain obtained from mixed-strain colony biofilms grown for three days. Each mixed biofilm contained one unlabeled strain and one constitutively expressing mScarlet. Error bars represent standard deviation of biological triplicates.

Fig. 4. *rebP1P2* is expressed during host association and contributes to host colonization and killing.
a, Fluorescent micrographs of *A. thaliana* leaves infected with WT PA14 expressing mScarlet under the control of the *rebP1P2* promoter (P$_{rebP1}$-*mScarlet*), no promoter (MCS-*mScarlet*), or a constitutive promoter (P$_{PA1/04/03}$-*mScarlet*). Smaller panels show images of the whole seedling and the imaged leaf indicated by the white box. Images are representative of nine biological replicates from three experiments. Scale bars are 1 mm.
b, Images of *C. elegans* fed WT PA14 containing the indicated reporter constructs. Top panels, Nomarski; middle panels, fluorescence; bottom panels, overlay. White boxes indicate regions shown in close-up images. Images are representative of 15 replicates from three experiments. Scale bars are 10 µm. **c**, CFUs obtained from *A. thaliana* seedlings inoculated with the indicated genotypes and incubated for five days, normalized to seedling weight. Each circle represents a biological replicate, with color intensity indicating replicates from the same experiment; colored lines indicate means. p-values were calculated using unpaired, two- tailed t-tests.
d, Killing kinetics of *C. elegans* following exposure to various PA14 genotypes. The role of *gacA* in *C. elegans* pathogenicity has been described previously [27] and *gacA::Tn* therefore serves as a control for defective killing in this assay. Error bars denote standard deviation and p-values were calculated using unpaired, two-tailed t-tests. n ≥ 4 biological replicates from three experiments.

425   **REFERENCES**

1.   Quackenbush, R. L. & Burbach, J. A. Cloning and expression of DNA sequences
      associated with the killer trait of Paramecium tetraurelia stock 47. *Proc. Natl. Acad. Sci. U.
      S. A.* **80**, 250–254 (1983).

430   2.   Gibson, I. A comparison of the refractile bodies (R-bodies) of certain bacteria—III.
      Nucleotide sequence homologies and R-body function. *Micron and Microscopica Acta* **15**,
      253–259 (1984).

3.   Gibson, I., Bedingfield, G., Dobbs, H. & Shackleton, J. Effects of various agents on the R
      bodies of certain bacteria. *Micron and Microscopica Acta* **18**, 71–75 (1987).

435   4.   Pond, F. R., Gibson, I., Lalucat, J. & Quackenbush, R. L. R-body-producing bacteria.
      Microbiol. Rev. **53**, 25–67 (1989).

5.   Jurand, A., Preer, J. R. & Rudman, B. M. Further investigations on the prelethal effects of
      the killing action of kappa containing killer stocks of Paramecium aurelia. *J. Exp. Zool.* **206**,
      25–47 (1978).

440   6.   Sonneborn, T. M. Mating Types in Paramecium Aurelia: Diverse Conditions for Mating in
      Different Stocks; Occurrence, Number and Interrelations of the Types. *Proc. Am. Philos.
      Soc.* **79**, 411–434 (1938).

7.   Heruth, D. P., Pond, F. R., Dilts, t. J. A. & Quackenbush, R. L. Characterization of Genetic
      Determinants for R Body Synthesis and Assembly in Caedibacter taeniospiralis 47 and
445   116. *J. Bacteriol.* **176**, 3559–3567 (1994).

8.   Kanabrocki, J. A., Quackenbush, R. L. & Pond, F. R. Organization and expression of
      genetic determinants for synthesis and assembly of type 51 R bodies. *J. Bacteriol.* **168**,
      40–48 (1986).

9.   Pirritano, M. *et al.* Dual-Seq reveals genome and transcriptome of Caedibacter
450   taeniospiralis, obligate endosymbiont of Paramecium. *Sci. Rep.* **10**, 9727 (2020).

10.   Schulz, S. *et al.* Elucidation of sigma factor-associated networks in Pseudomonas
      aeruginosa reveals a modular architecture with limited and function-specific crosstalk.
      PLoS Pathog. **11**, e1004744 (2015).

11.   Lee, D. G. *et al.* Genomic analysis reveals that Pseudomonas aeruginosa virulence is
455   combinatorial. *Genome Biol.* **7**, R90 (2006).

12. Vasquez-Rifo, A., Veksler-Lublinsky, I., Cheng, Z., Ausubel, F. M. & Ambros, V. The Pseudomonas aeruginosa accessory genome elements influence virulence towards Caenorhabditis elegans. *Genome Biol.* **20**, 270 (2019).

13. Preer, J. R., Jr, Preer, L. B. & Jurand, A. Kappa and other endosymbionts in Paramecium aurelia. *Bacteriol. Rev.* **38**, 113–163 (1974).

14. Matsuoka, J.-I. *et al.* A putative TetR-type transcription factor AZC_3265 from the legume symbiont Azorhizobium caulinodans represses the production of R-bodies that are toxic to eukaryotic cells. *Soil Sci. Plant Nutr.* **63**, 452–459 (2017).

15. Mikkelsen, H., McMullan, R. & Filloux, A. The Pseudomonas aeruginosa reference strain PA14 displays increased virulence due to a mutation in ladS. *PLoS One* **6**, e29113 (2011).

16. Raymann, K., Bobay, L.-M., Doak, T. G., Lynch, M. & Gribaldo, S. A genomic survey of Reb homologs suggests widespread occurrence of R-bodies in proteobacteria. *G3* **3**, 505–516 (2013).

17. Feinbaum, R. L. *et al.* Genome-wide identification of Pseudomonas aeruginosa virulence-related genes using a Caenorhabditis elegans infection model. *PLoS Pathog.* **8**, e1002813 (2012).

18. Polka, J. K. & Silver, P. A. A Tunable Protein Piston That Breaks Membranes to Release Encapsulated Cargo. *ACS Synth. Biol.* **5**, 303–311 (2016).

19. Stewart, P. S. & Franklin, M. J. Physiological heterogeneity in biofilms. *Nat. Rev. Microbiol.* **6**, 199–210 (2008).

20. Cornell, W. C. *et al.* Paraffin Embedding and Thin Sectioning of Microbial Colony Biofilms for Microscopic Analysis. *J. Vis. Exp.* (2018) doi:10.3791/57196.

21. Donlan, R. M. Biofilms: microbial life on surfaces. *Emerg. Infect. Dis.* **8**, 881–890 (2002).

22. Flemming, H.-C. *et al.* Biofilms: an emergent form of bacterial life. *Nat. Rev. Microbiol.* **14**, 563–575 (2016).

23. Tan, M. W., Mahajan-Miklos, S. & Ausubel, F. M. Killing of Caenorhabditis elegans by Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 715–720 (1999).

24. Walker, T. S. *et al.* Pseudomonas aeruginosa-plant root interactions. Pathogenicity, biofilm formation, and root exudation. *Plant Physiol.* **134**, 320–331 (2004).

25. Heckmann, K. & Görtz, H.-D. Prokaryotic Symbionts of Ciliates. in *The Prokaryotes: A Handbook on the Biology of Bacteria: Ecophysiology, Isolation, Identification, Applications* (eds. Balows, A., Trüper, H. G., Dworkin, M., Harder, W. & Schleifer, K.-H.) 3865–3890 (Springer New York, 1992).

490   26.   Vasquez-Rifo, A., Ricci, E. P. & Ambros, V. Pseudomonas aeruginosa cleaves the decoding center of Caenorhabditis elegans ribosomes. *PLoS Biol.* **18**, e3000969 (2020).

27. Rahme, L. G. *et al.* Use of model plant hosts to identify Pseudomonas aeruginosa virulence factors. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 13245–13250 (1997).

28. Bertani, G. Lysogeny at mid-twentieth century: P1, P2, and other experimental systems. *J.*
495   *Bacteriol.* **186**, 595–600 (2004).

29. Okegbe, C. *et al.* Electron-shuttling antibiotics structure bacterial communities by modulating cellular levels of c-di-GMP. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5236–E5245 (2017).

30. Shanks, R. M. Q., Caiazza, N. C., Hinsa, S. M., Toutain, C. M. & O'Toole, G. A.
500   Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes from gram-negative bacteria. *Appl. Environ. Microbiol.* **72**, 5027–5036 (2006).

31. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).

32. Hoang, T. T., Karkhoff-Schweizer, R. R., Kutchma, A. J. & Schweizer, H. P. A broad-host-
505   range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa mutants. *Gene* **212**, 77–86 (1998).

33. Oswald, E. S., Brown, L. M., Bulinski, J. C. & Hung, C. T. Label-free protein profiling of adipose-derived human stem cells under hyperosmotic treatment. *J. Proteome Res.* **10**,
510   3050–3059 (2011).

34. Yang, W. S. *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156**, 317–331 (2014).

35. Wobma, H. M. *et al.* The influence of hypoxia and IFN-γ on the proteome and metabolome of therapeutic mesenchymal stem cells. *Biomaterials* **167**, 226–234 (2018).

515   36. Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077 (2016).

37. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

38. Clough, S. J. & Bent, A. F. Floral dip: a simplified method forAgrobacterium-mediated
520   transformation ofArabidopsis thaliana. *The Plant Journal* vol. 16 735–743 (1998).

39. Murashige, T. & Skoog, F. A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiologia Plantarum* vol. 15 473–497 (1962).

40. Ishiga, Y., Ishiga, T., Uppalapati, S. R. & Mysore, K. S. Arabidopsis seedling flood-

inoculation technique: a rapid and reliable assay for studying plant-bacterial interactions.

525 *Plant Methods* **7**, 32 (2011).

41. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

42. Karp, P. D. *et al.* The BioCyc collection of microbial genomes and metabolic pathways. *Brief. Bioinform.* **20**, 1085–1093 (2019).

43. Winsor, G. L. *et al.* Enhanced annotations and features for comparing thousands of

530 *Pseudomonas* genomes in the Pseudomonas genome database. *Nucleic Acids Res.* **44**, D646–53 (2016).