Hidden genetic variation in plasticity increases the potential to adapt to novel environments

Greg M. Walter[1,2]*, James Clark[1,3], Delia Terranova[4], Salvatore Cozzolino[5], Antonia Cristaudo[4],

Simon J. Hiscock[3] and Jon R. Bridle[1,6]

[1]University of Bristol, School of Biological Sciences, Bristol BS8 1TQ, UK

[2]Current address: Monash University, School of Biological Sciences, Melbourne 3800, Australia

[3]University of Oxford, Department of Plant Sciences, Oxford, OX1 3RB, UK

[4]University of Catania, Department of Biological, Geological and Environmental Sciences,

Catania 95128, Italy

[5]University of Naples Federico II, Department of Biology, Naples 80126, Italy

[6]Current address: University College London, Department of Genetics, Evolution and

Environment, London WC1E 6BT, UK

* Corresponding Author: Greg M. Walter

15   Email: greg.walter@monash.edu

Abstract

Adaptive plasticity increases population persistence, but can slow adaptation to changing environments by hiding the effects of different alleles on fitness. However, if plastic responses are no longer adaptive in novel environments, then differences among alleles can emerge and increase genetic variation in fitness that allows rapid adaptation. We tested this hypothesis by transplanting cuttings and seeds of a Sicilian daisy within and outside its native range, and quantifying variation in morphology, physiology, gene expression and fitness. We show that genetic variance in plasticity increases the potential for rapid adaptation to novel environments. Genetic variation in fitness was low across native environments where plasticity effectively tracked familiar environments. In the novel environment however, genetic variation in fitness increased threefold, and correlated with genetic variation in plasticity. Furthermore, genetic variation that can increase fitness in the novel environment had the lowest fitness at the native site, suggesting that adaptation to novel environments relies on genetic variation in plasticity that is selected against in native environments.

Introduction

Understanding how populations can respond to novel environments is vital for predicting the resilience of ecological communities to environmental change (Parmesan 2006; Shaw and Etterson 2012). Populations persist in novel environments either by genotypes adjusting their phenotype to match the new conditions (adaptive plasticity) (Via et al. 1995; Charmantier et al. 2008), and/or by increases in the frequency of alleles that increase fitness in novel environments, causing adaptation (termed 'evolutionary rescue') (Gomulkiewicz and Holt 1995; Bell and Gonzalez 2009). However, if adaptation to novel environments relies on alleles that increase fitness via beneficial plastic responses, then selection on plasticity can promote evolutionary rescue (Lande 2009; Chevin and Lande 2011).

Given that plasticity can only evolve to buffer environmental variation encountered within its native range, existing plasticity should only maintain fitness in previously experienced conditions (Chevin et al. 2013), which reduces genetic variation in fitness in familiar environments (Bradshaw 1991). However, the buffering effect of adaptive plasticity should become less effective in novel environments, exposing genetic variation to selection because genotypes will vary more in their phenotype and fitness depending on their (previously untested) sensitivity to the new conditions (Hermisson and Wagner 2004; Lande 2009; Chevin et al. 2010; Chevin and Lande 2011; Ashander et al. 2016). This exposure of genetic variation in novel environments could allow rapid evolutionary rescue in novel conditions (Nussey et al. 2005; Lande 2009). If so, estimates of genetic variation in fitness within a species' current geographical range are likely to underestimate the potential for evolutionary rescue in novel environments, and therefore the resilience of populations and ecological communities to environmental change.

The amount of additive genetic variance in relative fitness (i.e. the amount of genetic variation

relative to mean fitness) predicts the increase in mean fitness in the subsequent generation

(Fisher 1930; Bonnet et al. 2019), and therefore the potential for evolutionary rescue (Shaw

2018). Although genetic variation in fitness is known to vary depending on the environment

(Sheth et al. 2018), changes in the amount of genetic variation in fitness is rarely characterized in

novel environments, and has not been linked to genetic variation in plasticity in natural

environments.

In this study, we transplanted multiple clones of 314 genotypes (produced by a quantitative

genetics breeding design) across three environments to test: (1) In familiar environments,

adaptive plasticity maintains high mean fitness but low genetic variance in fitness because all

genotypes can track the environment. However, in a novel environment, the ability for plasticity

to allow genotypes to track their environment will be limited, which will reduce mean fitness,

but increase genetic variation in fitness that can facilitate evolutionary rescue. (2) Genetic

variation in fitness in the novel environment will be correlated with genetic variation in plasticity

for morphology, physiology and gene expression.

We use a short-lived perennial Sicilian daisy, *Senecio chrysanthemifolius* (Asteraceae), that

grows in disturbed habitats at low altitudes (c.400-1,000 m.a.s.l [metres above sea level]) on

Mount Etna, an active volcano, and throughout lowland Sicily (**Fig. 1A**). We mated naturally-

occurring *S. chrysanthemifolius* genotypes in a quantitative genetic full-sibling, half-sibling

breeding design (Lynch and Walsh 1998) to produce 314 offspring genotypes (**Fig. 1B**). We then

used multiple clones of each genotype to estimate the fitness, morphology, physiology and gene

expression of each genotype at three elevations, both within and outside the species' native range

(**Fig. 1B**). Our breeding design allowed us to calculate the among-sire variance, which represents

the additive genetic variance for all phenotypic and fitness measurements recorded.

Results

(1) Reduced mean fitness but increased genetic variance in fitness in the novel environment

Mortality of cuttings was low, even at the novel elevation (12% across all elevations), allowing

us to assay fitness for 6-8 cuttings per genotype at each transplant site. In September-October,

after 5 months of growth, we estimated fitness by counting all flower heads produced by each

plant. This trait correlates closely with total seeds produced per plant (*see Supplementary*

Material and **Fig. S1**), and has been used routinely to assay fitness in other short-lived perennials

(e.g. Pujol et al. 2014). Fitness (number of flowers) was consistently high within the native range

(500m and 1,500m), but decreased by 87% at 2,000m (**Fig. 1C**). Additive genetic variance in

relative fitness was near zero within the native range (500m and 1,500m), but more than three

times greater at 2,000m where it predicts a 13.8% increase in mean fitness in the subsequent

generation (**Fig. 1D**; **Table S2**). Low genetic variance in relative fitness within the native range

suggests that adaptive plasticity allows all genotypes to track phenotypic optima and maintain

high mean fitness in familiar environments. By contrast, the plastic responses of genotypes at the

novel elevation fail (to differing extents) to maintain high mean fitness, creating high levels of

genetic variation in relative fitness, and increasing the potential for evolutionary rescue in

unfamiliar environments (Shaw 2018). Furthermore, seven of the nine sires with the highest

fitness in the novel environment were collected from two sites (**Table S3**), suggesting that

genetic variation present at certain sites within the native range might be key to promoting

evolutionary rescue.

Fig. 1. The experimental design (**A-B**) and the reduction in mean fitness, but increase in genetic variance in fitness at the novel elevation (**C-D**). (**A**) *Senecio chrysanthemifolius* in its natural habitat. Inset map presents the location of the study system in relation to Europe. **(B)** Schematic of Mount Etna showing the experimental design. We sampled individuals from five sites in the foothills of Mount Etna, which we crossed in the glasshouse and transplanted cuttings of their offspring at three elevations. We mated 36 sires each to three dams; we grew three offspring per cross in the glasshouse from which we sampled cuttings that were transplanted at the three elevations. Inset leaves show the average change in leaf morphology with elevation observed for a single genotype. (**C**) Mean fitness dropped outside the native range. Credible intervals represent 95% Highest Posterior Density (HPD) intervals of the mean. (**D**) Posterior distributions for the estimates of additive genetic variance in fitness. Genetic variance in fitness was significantly greater at the novel elevation (2,000m).

(2) Fitness in the novel environment correlates with genetic variance in plasticity

Quantifying phenotypes across elevation: We measured leaf morphology and leaf pigment

content by sampling 3-4 young, but fully expanded leaves from each plant. We scanned the

leaves and quantified morphology using the morphometric software '*Lamina*' (Bylesjo et al.

2008), which produced five leaf morphology traits: leaf area, leaf complexity ($\frac{leaf\ perimeter^2}{leaf\ area}$),

the number of leaf indents standardized by the perimeter, the width of leaf indents and Specific

Leaf Area ($SLA = \frac{leaf\ area}{leaf\ weight}$). These leaf measurements represent leaf morphology and

investment traits that show plastic responses to the abiotic environment in *S. chrysanthemifolius*

(Walter et al. 2020b), and in other plant systems, including sunflowers (Royer et al. 2009). We

also used a Dualex instrument (Force-A) to measure the chlorophyll, flavonol and anthocyanin

pigment content of the leaf. Flavonols and anthocyanins are secondary metabolites that combat

oxidative stress created by stressful abiotic (e.g. light and temperature) and biotic (e.g. herbivore)

conditions (Mierziak et al. 2014). Within each transplant site, we observed a significant

association between phenotype and fitness (**Table S5** and **Fig. S2**), and a significant change in

multivariate phenotype across elevation (MANOVA $F_{2,23} = 35.647$, $P<0.0001$; *see*

Supplementary Material), suggesting that these traits are ecologically important and show

significant plasticity across elevation (**Fig. 2A**).

Fig. 2. When compared to the home site (500m), the magnitude of plasticity was greater at the
novel elevation (2,000m) compared to the edge of the range (1,500m). (**A**) The change in raw
trait values across elevation. Black lines and circles represent the overall mean (±1 SE), with all
the genotypes from the breeding design in gray. (**B**) Frequency distribution for plasticity in each
trait, which was calculated as the degree to which each genotype changed trait values from the
home site (vertical red line) to the edge of the range (light blue) and to the novel elevation (dark
blue). Values of one represent no change from the home site. Values above and below one
represent plasticity as an increase and decrease in the trait value, respectively.

Identifying genetic variance in plasticity: To estimate genetic variation in plasticity, we first

quantified plasticity as the phenotypic change from the home site to the edge of the range (500-

1,500m), and separately, to the novel elevation (500m-2,000m). We calculated plasticity

separately for each of the 314 genotypes by dividing the trait values of each genotype at 1,500m

and 2,000m by the mean trait value of that genotype at the home site. This standardization

calculates the trait values at 1,500m and 2,000m relative to the home site for each genotype.

Values of one reflect no change in phenotype between sites, while values above and below one

respectively reflect increases and decreases in trait values from the home site (**Fig. 2B**). We

estimated plasticity for all traits and then used multivariate linear mixed effects models

(including fitness and plasticity in all eight traits) to partition the total variance into sire variance,

representing additive genetic variance in plasticity for each trait. We tested whether our estimates

of genetic variance in plasticity were statistically significant by comparing our observed

estimates to a null distribution (*see Methods*).

Our observed estimates of genetic variance in plasticity to the edge of the range (500 to 1,500m)

were within the null distribution for four traits, while four traits showed greater plasticity than

the null expectation (**Fig. 3**). By contrast, our observed estimates exceeded the null distribution

for all eight traits at 2,000m (**Fig. 3**), suggesting more genetic variance in plasticity at the novel

elevation. Genetic variation in plasticity was created by genotype-specific responses to the novel

environment (**Fig. S3**). Such genetic variance in plasticity could promote evolutionary rescue if

correlated with fitness variation at the novel elevation.

Fig. 3. Greater additive genetic variance in plasticity is observed at the novel elevation than at the range edge. Credible intervals represent 95% HPD for the null distribution. Where the observed estimate exceeds the random distribution indicates statistically significant estimates of genetic variance in plasticity. Significant genetic variance in plasticity was observed for all traits at the novel elevation, compared to only four traits at the range edge. Traits: AR=leaf area, LC=leaf complexity, NI=number of indents, IW=indent width, SLA=specific leaf area, CH=chlorophyll content, FL=flavonol content and AN=anthocyanin content.

Genetic correlations between leaf plasticity and fitness: We detected significant genetic

correlations (>70% of the posterior distribution did not overlap with zero) between the

magnitude of trait plasticity and fitness at the novel elevation for four traits: leaf complexity (-

0.25), number of indents (-0.31), SLA (0.4) and flavonol content (0.39) (**Fig. 4A**). Therefore,

genetic variance in plasticity for four leaf traits was correlated with genetic variation in fitness,

and so could cause evolutionary rescue in novel environments. Furthermore, depending on the

trait, genetic variation in fitness was associated both with increases and decreases in the

magnitude of plastic responses. For leaf complexity and the number of indents, *ER* genotypes

show greater plasticity than *HR* genotypes, which was created by stronger reductions in trait

values between 500m and 2,000m (**Fig. 4B)**. By contrast, for SLA and flavonol content, *ER*

genotypes show less plasticity than the *HR* genotypes (**Fig. 4B**).

Fig. 4. Genetic variance in plasticity was significantly correlated with fitness at the novel
elevation. (**A**) Genetic correlations between plasticity and fitness were moderately strong for leaf
complexity, the number of indents, specific leaf area and flavonol content. Numbers inset
represent genetic correlations with the 95% HPD interval in parentheses. Gray shading
represents the 95% HPD intervals for phenotypic correlations with fitness. (**B**) Histogram for the
sire genetic values (that underlie the genetic correlations from **A**), which represent the magnitude
of plastic change in trait means from the home site (red vertical line) to 2,000m. Shown are the
25% of sires with the highest (*ER* genotypes in blue) and lowest (*HR* genotypes in green) fitness
at 2,000m. Greater fitness at 2,000m is associated with plasticity as a greater reduction in leaf
complexity and number of indents, but with a lesser change in SLA and flavonol content.

Variation in gene expression: Overall, a larger number of genes were differentially expressed

with increasing distance from the home site. To the edge of the range (500m-1,500m), the *ER*

genotypes showed significant changes (adj. *P*<0.01) in more genes than the *HR* genotypes (1,376

in *ER* vs 514 genes in *HR*), suggesting that the *HR* genotypes adjusted fewer genes to maintain

fitness within their range (**Fig. 5A**). However, the number of genes that showed expression

changes tripled outside the native range (500m-2,000m) for both classes of genotypes (*ER*=4,876

genes vs *HR*=4,720) with the *ER* genotypes again showing a greater magnitude (1.2-fold mean

difference) of differential expression per gene than *HR* genotypes (**Fig. 5B**). For both classes of

genotypes, the mean magnitude of expression change was greater at the novel elevation

(*ER*=1.41, *HR*=1.21) when compared to the edge of the range (*ER*=1.19, *HR*=1.04). Compared to

the *HR* genotypes, *ER* genotypes showed twice as many underexpressed genes at the novel

elevation, but half as many overexpressed genes (**Fig. 5C**). Distinct patterns of gene expression

were therefore associated with genotypic differences in fitness at the novel elevation.

Gene ontology: To identify the function of differentially expressed genes, we tested for

functional enrichment of Gene Ontology (GO) terms. *ER* and *HR* genotypes differed in the

functional categories of genes that varied in expression at the novel elevation (**Fig. 5D**). *ER*

genotypes showed differential expression of genes relating to biosynthesis, while differentially

expressed genes in the *HR* genotypes were related to photosystems. Comparing the enriched GO

terms to those of a 2017 transplant experiment (Walter et al. 2020b) that included the high

elevation sister species (*S. aethnensis*), we found that four (of 17) GO terms were enriched in

S.aethnensis as well as in the *ER* genotypes, compared to only one (of 20) GO term shared by *S.*

aethnensis and the *HR* genotypes (**Fig. 5D**). Therefore, genotypes of *S. chrysanthemifolius* that

showed the highest fitness at the novel elevation also showed a more similar gene expression

response to the species native to that elevation.

GO terms that were significantly enriched at 2,000m included light harvesting in photosystem I,

cell wall organization, and responses to cold (**Table S8**). To illustrate the contrasting patterns of

expression, we selected the five most differentially expressed genes from each GO term and

compared the reaction of the *ER* and *HR* genotypes across elevation. For genes involved in

photosystem I (**Fig. 5E**), *ER* genotypes showed stronger patterns of underexpression than *HR*

genotypes at the novel elevation. For genes involved in cold responses (**Fig. 5F**), *ER* genotypes

showed a stronger response than *HR* genotypes, which included greater overexpression or

underexpression depending on the gene.

Fig. 5. *ER* and *HR* genotypes differ in gene expression at the novel elevation. **(A)** Within the native range, more genes were differentially expressed in the *ER* genotypes. **(B)** Outside the native range, more genes were differentially expressed, and more genes in the *ER* genotypes showed a greater magnitude of differential gene expression. **(C)** Numbers of differentially expressed genes (adjusted *P*<0.01 and log-fold change <2 or >-2 for over- and underexpression) for the *ER* and *HR* genotypes between the home site (500m) and outside the range (2,000m). *ER* genotypes underexpressed more genes than *HR* genotypes. **(D)** Enriched Gene Ontology (GO) terms for differentially expressed genes for both genotypes and for a high elevation sister species (*S. aethnensis*) transplanted along the same elevation gradient in 2017 (Walter et al. 2020b). Only the three most significant terms are shown. *ER* genotypes shared more GO terms with *S. aethnensis,* suggesting that high fitness is associated with similar genetic pathways to the native species. **(E)** Comparing elevational changes in mean expression for five genes (associated with light harvesting) for the *ER* (unfilled shapes, dashed lines) and *HR* (filled shapes, solid lines) genotypes. Each gene is represented by a different shape and credible intervals represent 95% confidence intervals. *ER* genotypes showed stronger patterns of underexpression at 2,000m. **(F)** Elevational changes in mean expression for five genes associated with responses to cold. *ER* genotypes show stronger under- and overexpression, but the pattern was gene-dependent.

Selection at the native environment hides genetic variation that could promote evolutionary

rescue in novel environments

Surprisingly, genetic variance in fitness at the novel environment was uncorrelated with fitness at

the native environment (**Table S2**; genetic correlation of -0.1 [-0.79,0.6 HPD]). **Fig 6A** shows

that the sires with the highest fitness at the novel elevation had the lowest fitness at the home site

and *vice versa,* suggesting that genetic variance in relative fitness not only increases in novel

environments, but that a genetic trade-off exists between the native and novel elevations.

Therefore, although evolutionary rescue may be facilitated by genetic variation that is present in

natural populations, these genotypes are selected against within the native range and only have

positive effects on fitness in novel environments.

Comparing fitness of seeds vs. cuttings: The 314 genotypes used in this study were generated by

crossing among adult genotypes sampled from the field. They were then grown from seed in the

greenhouse and transplanted into the field as cuttings. This means they did not experience

selection in the field at early life history stages. To test whether genetic variation that increases

fitness in novel environments is selected against at the home site during early life history, we

transplanted seeds from all 104 full-sibling families at the same elevations in spring (April

2019).

We transplanted 25 seeds of each full-sibling family at each transplant site (*n*=2,600 seeds/site;

total *N*=8,000), and measured the number of seedlings that emerged and tracked their survival to

autumn (October, 2019). We then estimated the genetic covariance in performance between

seeds and cuttings, and between the home site and the novel elevation (*see Supplementary*

Material for the analysis between 500 to 1,500m). Using a multivariate generalized mixed

model, we calculated a 4×4 covariance matrix with the diagonal representing genetic variance in

relative fitness as seeds and cuttings, and the off diagonal representing the genetic covariance

(and correlation) between seeds and cuttings both within and between elevations (**Fig. 6B**).

If genetic variance that can increase fitness at novel elevations is selected against at the home

site, sires that performed well as cuttings at the novel elevation (2,000m) should perform

similarly well as seedlings at 2,000m. However, the same sires that performed best at 2,000m

should show lower performances at the home site (500m). **Fig. 6B-D** presents seedling

emergence as the fitness component for early life history, but results were similar for survival

(**Fig. S8**). As predicted, performance as seedlings and cuttings at 2,000m were positively

correlated (0.19; -0.55,0.88 HPD; **Fig. 6B**), suggesting genetic variation in relative fitness at the

novel elevation was common to both life history stages. However, consistent with the lower

performance of *ER* genotypes as seedlings at the home site, seedling fitness at a given elevation

(500m or 2,000m) showed weak negative genetic correlations with fitness as cuttings at the other

elevation (-0.15 [-0.85,0.6 HPD] and -0.04 [-0.67,0.6 HPD]; **Fig. 6B**). The fittest sires at the

novel elevation therefore had the lowest fitness at the native site, while the sires with the lowest

fitness at the novel elevation showed the highest fitness at the native site (**Fig. 6C**).

To confirm that selection on seedlings at the home site reduces genetic variation suitable for

evolutionary rescue at the novel elevation, we used matrix decomposition to describe the matrix

in **Fig. 6B**. This method, analogous to principal components analysis, identifies axes that

describe genetic variation in fitness that is shared between elevations and life history stages. The

first axis represents genetic variance in seedling emergence common to both elevations (**Fig.
6D**), largely because genetic variance in seedling performance was much greater (**Fig. 6B**).

However, the second axis, representing 25% of total genetic variance in fitness, describes genetic

290   variation in fitness that differed between the native and novel elevations because the 500m and

291   2,000m sites possess loadings with high values that have different signs (**Fig. 6D**). Genetic

292   variation that increased fitness at the novel elevation showed lower fitness at the native

293   elevation, suggesting that selection on seedlings within the native range reduces the viability of

294   genotypes that would allow evolutionary rescue in novel environments.

Fig. 6. Genetic variation that could promote evolutionary rescue shows lower performance at the
home site. **(A)** Sire fitness values as cuttings at the home site and novel elevation show that the
ER genotypes had lower fitness than the *HR* genotypes at the home site. **(B)** The 4×4 covariance
matrix representing genetic variation for fitness as seedlings and cuttings at the home site and
novel elevation. Gray shading represents genetic variance in fitness along the diagonal, with the
lower diagonal presenting genetic correlations, and the upper diagonal the genetic covariances.
(**C**) Sires that performed well at the novel elevation (as both seedlings and cuttings) showed
lower fitness than the *HR* genotypes at the native elevation as both seedlings (left panel) and
cuttings (right panel). **(D)** Axes of genetic variation in fitness describing the matrix in (**B**).
Numbers in bold emphasize trait loadings greater than 0.2 to aid interpretation. The second axis
describes genetic variation underlying differences in fitness between the home site and the novel
elevation.

Discussion

This study provides evidence that adaptive plasticity is restricted to native environments, and

fails to sustain high mean fitness in a novel environment, exposing genetic variation in fitness

that could promote evolutionary rescue by allowing rapid changes in allele frequencies.

Increased genetic variation in fitness of *S. chrysanthemifolius* at the novel elevation was

correlated with genetic variation for greater plasticity that reduced the trait values of leaf

complexity and leaf indentation, but lower plasticity that maintained the trait values of specific

leaf area and flavonol content across elevation. Our data also suggest that genetic variation that

would increase fitness in novel environments showed lower fitness in the native environment.

Therefore, genetic variation important for evolutionary rescue is likely to be maintained at low

frequency by selection within the native range (Brennan et al. 2019). Studies estimating genetic

variation in fitness *in situ* are therefore likely to underestimate the potential for evolutionary

rescue because the effects of alleles important for evolutionary rescue only emerge in novel

environments (Sheth et al. 2018; Kulbaba et al. 2019).

Gene expression responses are determined through complex genetic and epigenetic regulation,

which are likely to determine plasticity in phenotype, and how environmental sensitivity can

evolve and promote resilience in novel environments (Oostra et al. 2018). Interestingly, the gene

expression data revealed that, compared to the low fitness genotypes at 2,000m, genotypes with

high fitness differentially expressed more genes, which was associated with greater

underexpression. Furthermore, differential expression in the high fitness genotypes occurred in

functional categories associated with plasticity in the closely related high elevation (2,000m+)

sister species. This suggests that adaptation to novel environments requires genotypes that create

the most beneficial gene expression profiles (Wang and Althoff 2019), which contrasts with

previous evidence that adaptation to novel environments involves non-adaptive plasticity in gene

expression (Ghalambor et al. 2015), or that a lack of genetic variance in gene expression will

prevent beneficial responses to novel environments (Oostra et al. 2018).

Overall, our results provide strong evidence that evolutionary rescue depends on genetic

variation that increases fitness by selecting for the most beneficial plastic responses in novel

environments (Lande 2009; Chevin and Lande 2011; Chevin and Hoffmann 2017), created by a

fine-tuning of gene-products and trait values in novel environments. Our results also support

studies that show that genetic variance in fitness depends on the environment in which it is

quantified (Sheth et al. 2018; Kulbaba et al. 2019). Genetic variation for sensitivity to the

environment is likely to be unevenly distributed across a species' range (Stratton 1994; Colautti

and Barrett 2013; Sheth and Angert 2016; Walter et al. 2020a), with local adaptation determining

which populations provide the potential to promote evolutionary rescue (Hargreaves and Eckert

2019). To quantify rates of adaptation to novel environmental regimes, research should therefore

assess genetic variation in environmental sensitivity for multiple populations exposed to a range

of novel environments. It is also important to determine how and when the expression of genetic

variation likely to cause evolutionary rescue is masked by existing environmental conditions.

Methods

Genotype sampling and crossing design: To establish the parental generation, in June 2017 we

collected cuttings from 72 individuals located at five geographically separate sites on the

foothills of Mt. Etna (Table S1; also see Fig. 1 in Walter et al. 2020b). Where possible, we

sampled individuals that were at least ten meters apart from each other to minimize chances of

sampling close relatives. We removed whole branches from mature plants, which we cut into 4-

5cm segments at the glasshouse (Giarre, Italy), dipped them in a rooting plant growth regulator

for softwood cuttings (Germon® Bew., Der. NAA 0.5%, L. Gobbi, Italy) and placed each cutting

in one cell of an 84-cell tray containing a compressed mix of 1:1 perlite and coconut coir. For

three weeks, we kept cuttings on benches covered by plastic to maintain humidity and encourage

root growth. For each individual sampled in the field, we placed one cutting in a 30cm diameter

pot with standard potting mix, which we watered regularly and left to grow. To encourage

growth, we suspended 25W LED tubes (TSA Technology, Italy) 1m above the bench. Once

plants produced buds, we covered flowering branches with perforated bread bags in order to

prevent pollinators from entering while allowing airflow. We randomly designated each

individual as a dam or sire and grouped them into 12 blocks, each containing three sires ($n = 36$)

and three dams ($n = 36$), see **Table S1**. Within each block we mated all sires to all dams to

produce nine full-sibling families per block ($n = 104$ total full-sibling families, with four crosses

failing to produce seeds).

Six seeds from each full-sibling family were germinated by cutting the top off each seed (<1mm)

and placing them on moistened filter paper in a plastic Petri dish. Petri dishes were kept in the

dark for two days, and then transferred to growth cabinets maintained at 22°C with a 12h:12h

light:dark photoperiod. After one-week we transferred seedlings to the glasshouse where one

individual was transplanted into one cell of an 84-cell tray containing a mix of 1:1 perlite and

coconut coir. After growing 3-4 true leaves, we transferred seedlings to 14cm diameter pots

containing standard potting mix. We transferred three seedlings (in three separate pots) for each

full-sibling family, which we arranged randomly in the greenhouse. When the main stem of each

seedling reached c.12cm of growth, we cut the main stem c.4cm above ground level to promote

lateral branching and generate enough cuttings to replicate each genotype in the field transplant.

This meant that each of the 312 individuals grew 4-5 side branches, each long enough to take 5-6

cuttings.

Field transplant of the breeding design as cuttings: When branches started producing buds, we
took cuttings from all 312 individuals (hereafter, genotypes), cutting each branch from each plant
into smaller segments each 4-5cm long with 2-3 leaf nodes. For almost all genotypes we were
able to take 21 cuttings (transplant 1). Each cutting was treated as described above and left for
two weeks to produce roots. To produce a second round of cuttings, all plants were left to regrow
for three weeks, at which point we repeated the process to generate another 14 cuttings per plant
(transplant 2). Due to the deaths of two genotypes in the glasshouse after taking the initial
cuttings, we replaced these genotypes with siblings for the second round of cuttings, which led to
two extra genotypes (n=314 rather than 312). Cuttings were collected as above and stored in
high-humidity tunnels for two weeks until they produced roots and were ready to transplant.

Transplant 1, May 23-25 2018: To prepare the cuttings, for each transplant site we randomized
six cuttings per genotype into 24 84-cell trays. We split the 24 trays into two sets of 12, each
containing three cuttings for each genotype, which provide two replicate environmental blocks at
each site, each containing three cuttings for each of the 312 genotypes (transplant 1: $n = 936$
cuttings per block; $n = 1,872$ cuttings per transplant site; total N = 5,634 cuttings).

Transplant 2, 10-11 July: We used all remaining cuttings available to transplant two additional
cuttings/genotype at each field site in two additional blocks, as well as transplanting extra
cuttings to replace cuttings lost due to c.10% mortality at 500m and 2,000m due to a short spell
of intense heat in late June (transplant 2: $n = 321-525$ cuttings per block; $n = 718-927$ cuttings
per transplant site; total N = 2,515 cuttings). In total, we transplanted 8,149 cuttings for the 314
genotypes.

In total, we transplanted 7-10 cuttings from each of the 314 genotypes at each of three transplant

sites along an elevational gradient (n=c.2,700 cuttings/site; total N=8,149 cuttings) that

represented typical conditions within the species' native range (500m.a.s.l), the edge of its range

(1,500m.a.s.l), and a novel elevation (2,000m.a.s.l). The 500m site was located on abandoned

land, the 1,500m site in an apple and pear orchard, and the 2,000m site on a lava flow from

c.1980. Elevations under 1,000m are exposed to high temperatures that regularly exceed 40°C

during summer, and above 1,000m the temperature regularly drops below 0°C during winter.

Soil at lower elevations soil is characterized as a sandy loam, but changes to volcanic sand above

1,500m (Walter et al. 2020b).

To prepare experimental plots, the soil surface was cleared of plants and the soil turned (30cm

deep) immediately prior to transplanting the cuttings. We then transplanted cuttings into large

grids of different sizes, spacing each plant 30cm apart. Transplanted cuttings were maintained

moist until they established, after which we reduced watering to minimal amounts at regular

intervals, sufficient to maintain survival for all genotypes while having a minimal effect on

growth or flower production.

Bayesian implementation of statistical tests: We used Bayesian models to estimate genetic

variance in fitness and phenotypic plasticity across elevation. For each equation, we included a

substantial burn-in and thinning interval to allow model convergence. We checked

autocorrelation by confirming that the effective sample sizes exceeded 85% of the number of

samples that were saved. We used uninformative parameter-expanded priors, which we checked

by changing the scale parameter and ensuring that there was no effect on the posterior

distribution. From each implementation, we extracted 2,000 samples, which provided the

posterior distribution for each parameter estimated.

Quantifying genetic variance in fitness: To quantify genetic variance in fitness, we used the R

(v3.6.1; R Core Team 2019) package '*MCMCglmm*' (Hadfield 2010) to implement the

generalized linear mixed effects model

$$y_{ijklmn} = T_i + s_{j(k)} + d_{k(j)} + g_{l(jk)} + b_{m(i)} + e_{n(ijklm)} , \qquad (1)$$

where the only fixed effect represented transplant site (T_i). The random effects $s_{j(k)}$ represented

the *j*th sire, $d_{k(j)}$ the *k*th dam and $g_{l(jk)}$ the *l*th individual of the breeding design nested within

dam and sire. Additional random effects included block within transplant site ($b_{m(i)}$), which

removed variance due to differences between the two transplants and between the experimental

blocks. $e_{n(ijklm)}$ represented the error variance. For the $s_{j(k)}$, $d_{k(j)}$ and $g_{l(jk)}$ components we

were interested in estimating the variance among genotypes within each transplant site, as well

as the covariance among sites. To do this, we estimated random intercepts for each transplant

site, and random slopes between transplant sites, resulting in a 3×3 covariance matrix for each

random effect. We implemented equation 1 while estimating residual variance at each transplant

site, but not the covariance among sites because plants could not be at both sites at once. Fitness

was included as a univariate, poisson-distributed response variable (y_{ijklmn}), which meant that

for each random component our estimates of variance represented variance in relative fitness

(Bonnet et al. 2019). We implemented equation 1 with 1 million iterations, a burn-in of 100,000

iterations and thinning interval of 500 iterations. The additive genetic (co)variance matrix for

fitness is presented in **Table S2**.

Quantifying genetic variance in plasticity: For the subsequent analyses, for all traits we

calculated the average across all leaves sampled from each plant. We calculated plastic changes

in phenotype across elevation by standardizing all clones of each genotype (at 1,500m and

2,000m) by the mean value of that genotype at the home site (500m). This standardization

calculated the trait values for each genotype at the edge of the range, and outside the range,

relative to their trait value at the home site (see **Fig. 2**). We estimated plasticity separately for

each transplant date to account for differences between transplants in the linear mixed model

below. To quantify genetic variance in plasticity, and the genetic correlation between phenotypic

plasticity and fitness, we used a multivariate linear mixed effects model

452
$$y_{ijklm} = s_{i(j)} + d_{j(i)} + g_{k(ij)} + b_l + e_{m(ijkl)}, \qquad (2)$$

where the random effects $s_{i(j)}$ represented the ith sire, $d_{j(i)}$ the jth dam and $g_{k(ij)}$ the kth

individual of the breeding design nested within dam and sire. Additional random effects included

block within transplant site (b_l), which removed variance due to differences between the two

transplants and between the experimental blocks. The error variance is represented by $e_{m(ijkl)}$.

We applied equation 2 to plasticity in all eight phenotypic traits as a multivariate response

variable (y_{ijklm}). Fitness was included as the ninth trait as a poisson-distributed response

variable. We applied equation 2 separately for the edge of the range (1,500m) and outside the

range (2,000m). We implemented equation 2 with 2 million iterations, a burn-in of 200,000

iterations and thinning interval of 1,000 iterations.

Extracting the sire component from the output of equation 3 and multiplying it by four (Lynch

and Walsh 1998) quantified the 9×9 additive genetic variance-covariance matrix for plasticity,

separately for the 1,500m and 2,000m elevations. In **Table S6** we present the estimates for

genetic variance in plasticity for each trait, and each comparison. To quantify the genetic

correlation between plasticity and fitness we calculated the correlation matrix and extracted

genetic correlations between trait plasticity and fitness, which are presented in **Fig. 4**.

In a Bayesian framework, estimates of genetic variance in plasticity are constrained to be

positive, making comparisons to zero uninformative. To test whether we detected biologically

meaningful observed estimates of genetic variance in plasticity, we created a null distribution by

randomizing the parentage of all individuals and re-implementing the same equation. To prevent

the fixed effect of the two transplants and the experimental blocks having undue influence on the

randomizations, we randomized the pedigree within each experimental block. We randomized

and re-implemented equation 2 1,000 times to create a suitable null distribution. To reduce

computing time, we reduced the total number of iterations while retaining the same burn-in and

thinning interval to ensure the same mixing properties of the chain (Walter et al. 2018). The

convergence of our observed models showed that 100 iterations were required to obtain a stable

estimate of the mean for each variance parameter. We calculated the mean of each model

implemented on the randomized data, which gave us the null distribution of genetic variance in

plasticity expected under random sampling. If our observed estimates of genetic variance

exceeded our null distribution, this is evidence for significant genetic variance in plasticity.

Gene expression analyses: To prioritize genotypes for gene expression analyses, we quantified

changes in fitness between the home site (500m) and the novel elevation (2,000m). We chose six

genotypes that showed the greatest increase in relative fitness from the home site to outside the

range (*see Supplementary Material*), representing genotypes that could promote evolutionary

rescue in the novel environment (hereafter, '*ER*' genotypes). We also identified six genotypes

with the greatest decrease in relative fitness outside their range, suggesting a higher level of

specialization to their native range (hereafter, '*HR*' genotypes).

For each of the chosen genotypes, we sampled leaves from three clones chosen at random at each

transplant site. From each plant, we sampled 3-4 young leaves (c.15mm long), which we

immediately submerged in RNAlater and stored at 4°C for 24 hours and then at -80°C before

RNA extraction. To control for variation due to weather and diurnal effects, sampling was

constrained to three consecutive days between 9am and 11am under similar weather conditions.

We extracted RNA for the leaves of each plant and quantified gene expression at each elevation

using 3' RNAseq (QuantSeq). See *Supplementary Material* for RNA extraction protocols and the

reference transcriptome assembly.

Detecting Differentially Expressed Genes: 3' reads were mapped to the reference transcriptome

using Salmon v1.1.0 (Patro et al. 2017). On average 79.15% of reads were mapped (72.6-87.3%).

Transcript abundance estimates were imported into R using the 'txImport' package (Soneson et

al. 2015), with estimates normalized according to library size but not by transcript length, as the

3' sequencing method removes this bias. We visualized the broad patterns of variation in gene

expression among samples by first normalizing the transcript abundance using the variance

stabilizing transformation function in DESeq2 and then subjecting the transcript abundance

matrix to a principal components analysis **(Fig. S5)**. Differential expression of transcripts was

then calculated using non-normalized transcript counts between transplant sites for each set of

genotypes (*ER* and *HR*) separately using DESeq2 (Love et al. 2014). Differentially expressed

genes were defined as those showing both significant (adjusted p-value < 0.01) and strong (log2-

fold change > 2 or < -2) changes in expression between treatments.

We tested for enrichment of gene ontology (GO) terms among differentially expressed genes

between each transplant site for both sets of genotypes (**Table S8**). Enrichment tests were

performed using both a Kolmogorov-Smirnoff test with the 'weight' algorithm and Fisher's

exact test in the topGO package (Alexa and Rahnenführer 2019).

Estimating seedling performance across elevation: To test the performance of the same sires at

early life history stages, we planted siblings from all full-sibling families as seeds at the same

sites as the cuttings. At each transplant site, we transplanted 25 seeds from each family

randomized into five replicate blocks at each transplant site (family $n = 104$, $n = 520$ seeds per

block, $n = 2,600$ seeds per transplant site, total $N = 8,000$ seeds). We used a small drop of non-

drip superglue to attach each seed to the middle of a toothpick, which we randomized into five

environmental blocks at each site (Walter et al. 2016). We then placed each toothpick into a

randomized grid (replicating the experimental grids in the field) made of 3cm thick polystyrene

glued to plywood (**Fig. S6A**). To construct the experimental blocks in the field, we cleared the

ground and attached 4x4cm grid fishing net to the ground using 20cm steel u-shaped pegs.

Seeds in natural populations germinate in autumn and spring with available water. To replicate

natural germination conditions, we initiated the transplant in spring (7-8 May 2019), suspended

80% shade-cloth 20cm above each block and kept the seeds moist for three weeks, after which

we reduced watering. During this period it rained often, suggesting the experiment matched

natural germination timing. Shortly after reducing water, and once seedling emergence

stabilized, we replaced the shade-cloth with a lighter shade-cloth (40%) that represented light

shading common in natural populations (5 weeks at 500m; 7 weeks for 1,500m; 8 weeks for

2,000m). We removed all shade-cloth once seedlings reached their first two true leaves (after 9

weeks at 500m, and after 10 weeks for 1,500m and 2,000m). The differences in timing were

necessary because the lower temperatures at higher elevations slowed the rate of emergence and

growth. To make conditions as similar as possible for the four sites, and as natural as possible,

we removed shade-cloth and ceased watering at the same average developmental point at each

site. We collected data weekly by recording whether seedlings emerged and then tracking their

survival. **Fig. S6B** presents a photo of a seedling block at 2,000m, two months after the initial transplant.

Seedling emergence was high (500m=86.8%; 1,5000m=76.5%; 2,000m=90.0%). Compared to the home site, mortality after emergence was higher at the edge of the range and outside the range (**Fig. S7**). For the analyses of genetic variance below, we used seedling emergence and survival as measures of seedling performance. For survival, we used the data collected at the end of summer (24 September, 139 days from commencement of the experiment), at which point mortality stabilized and 67.8% of the plants that emerged still remained across all transplant sites (**Fig. S7**).

Our *a priori* hypothesis focused on describing the genetic correlation between seedlings and cuttings across the elevational extremes (i.e. between the home site and the novel elevation). Consequently, we focus on the 500m and 2,000m elevations, but we also provide the analogous analysis with the intermediate (1,500m) elevation in the *Supplementary Material*. To connect fitness for the same sires measured as seedlings and cuttings, we re-implemented equation 1 with the combined seedling and cutting data. To account for the difference among blocks in the seed experiment we included an additional random term ($r_{p(i)}$) representing the pth experimental seed block within the ith transplant site. To estimate the genetic covariance between fitness as seeds and as cuttings, we implemented equation 1 with seed and cutting fitness included as a bivariate response variable. Both random intercepts and random slopes were estimated for each transplant site and life history stage, resulting in a 4×4 genetic covariance matrix ($\mathbf{G}_m$) representing genetic variance in fitness as both cuttings and seeds within each transplant site, as well as the genetic covariance among transplant sites for seedlings and cuttings. We implemented equation 1 for

emergence and separately for survival. In the main text we present seedling emergence, but

results for seedling survival were consistent (**Fig. S8**).

To identify genetic variation common to both seed and cuttings experiments, and common

among transplant sites, we decomposed $\mathbf{G}_m$, which quantified the orthogonal axes of genetic

variation underlying total genetic variation in fitness. We predicted that if genetic variance

underlying evolutionary rescue was removed at early life history stages, we would observe an

axis of genetic variance with trait loadings opposite for the home site and the site outside the

home range. To identify whether our axes of genetic variation described biologically meaningful

genetic variance in fitness, we created a suitable null distribution for which to compare our

observed matrix decomposition. We constructed our random distribution using the same method

of randomizing the pedigree information and re-implementing the same model. To prevent

random and fixed effects having undue influence on the randomizations, we randomized the

pedigree within each experimental block (separately for seedlings and cuttings).

We found that all axes describing genetic variance in fitness represented greater genetic variance

in fitness than expected under random sampling (**Fig. S9**). This means that the second

eigenvector that described differences in fitness between the home site and the site outside the

range also described greater genetic variance than expected under random sampling.

References

Alexa, A., and J. Rahnenführer. 2019. Gene set enrichment analysis with topGO. R package

version 2.38.1.

Ashander, J., L. M. Chevin, and M. L. Baskett. 2016. Predicting evolutionary rescue via evolving

580   plasticity in stochastic environments. Proceedings of the Royal Society B-Biological

581   Sciences 283.

582   Bell, G., and A. Gonzalez. 2009. Evolutionary rescue can prevent extinction following

583   environmental change. Ecology Letters 12:942-948.

584   Bonnet, T., M. B. Morrissey, and L. E. B. Kruuk. 2019. Estimation of genetic variance in fitness,

585   and inference of adaptation, when fitness follows a log-normal distribution. Journal of

586   Heredity 110:383-395.

587   Bradshaw, A. D. 1991. The Croonian Lecture, 1991 - Genostasis and the Limits to Evolution.

588   Philosophical Transactions of the Royal Society B-Biological Sciences 333:289-305.

589   Brennan, R. S., A. D. Garrett, K. E. Huber, H. Hargarten, and M. H. Pespeni. 2019. Rare genetic

590   variation and balanced polymorphisms are important for survival in global change

591   conditions. Proceedings of the Royal Society B-Biological Sciences 286:20190943.

592   Bylesjo, M., V. Segura, R. Y. Soolanayakanahally, A. M. Rae, J. Trygg, P. Gustafsson, S.

593   Jansson et al. 2008. LAMINA: a tool for rapid quantification of leaf size and shape

594   parameters. BMC Plant Biology 8.

595   Charmantier, A., R. H. McCleery, L. R. Cole, C. Perrins, L. E. B. Kruuk, and B. C. Sheldon.

596   2008. Adaptive phenotypic plasticity in response to climate change in a wild bird

597   population. Science 320:800-803.

598   Chevin, L. M., R. Gallet, R. Gomulkiewicz, R. D. Holt, and S. Fellous. 2013. Phenotypic

599   plasticity in evolutionary rescue experiments. Philosophical Transactions of the Royal

600   Society of London Series B-Biological Sciences 368:1-12.

Chevin, L. M., and A. A. Hoffmann. 2017. Evolution of phenotypic plasticity in extreme
    environments. Philosophical Transactions of the Royal Society of London Series B-
    Biological Sciences 372:20160138.

Chevin, L. M., and R. Lande. 2011. Adaptation to marginal habitats by evolution of increased
    phenotypic plasticity. Journal of Evolutionary Biology 24:1462–1476.

Chevin, L. M., R. Lande, and G. M. Mace. 2010. Adaptation, plasticity, and extinction in a
    changing environment: towards a predictive theory. PLoS Biology 8:e1000357.

Colautti, R. I., and S. C. Barrett. 2013. Rapid adaptation to climate facilitates range expansion of
    an invasive plant. Science 342:364-366.

Fisher, R. A. 1930, The genetical theory of natural selection. Oxford, Oxford University Press.

Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, and K. A. Hughes.
    2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
    nature. Nature 525:372-375.

Gomulkiewicz, R., and R. D. Holt. 1995. When does evolution by natural selection prevent
    extinction? Evolution 49:201-207.

Hadfield, J. D. 2010. MCMC Methods for multi-response generalized linear mixed models: The
    MCMCglmm R package. Journal of Statistical Software 33:1-22.

Hargreaves, A. L., and C. G. Eckert. 2019. Local adaptation primes cold-edge populations for
    range expansion but not warming-induced range shifts. Ecology Letters 22:78-88.

Hermisson, J., and G. P. Wagner. 2004. The population genetic theory of hidden variation and
    genetic robustness. Genetics 168:2271-2284.

Kulbaba, M. W., S. N. Sheth, R. E. Pain, V. M. Eckhart, and R. G. Shaw. 2019. Additive genetic

variance for lifetime fitness and the capacity for adaptation in an annual plant. Evolution

73:1746-1758.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. Journal of Evolutionary Biology 22:1435-1446.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome biology 15:550.

Lynch, M., and B. Walsh. 1998, Genetics and analysis of quantitative traits. Sunderland, Sinauer

Associates, Inc.

Mierziak, J., K. Kostyn, and A. Kulma. 2014. Flavonoids as important molecules of plant

interactions with the environment. Molecules 19:16240-16265.

Nussey, D. H., E. Postma, P. Gienapp, and M. E. Visser. 2005. Selection on heritable phenotypic

plasticity in a wild bird population. Science 310:304-306.

Oostra, V., M. Saastamoinen, B. J. Zwaan, and C. W. Wheat. 2018. Strong phenotypic plasticity

limits potential for evolutionary responses to climate change. Nature Communications

9:1-11.

Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. Annual

Review of Ecology, Evolution, and Systematics 37:637-669.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. Salmon provides fast

and bias-aware quantification of transcript expression. Nature methods 14:417.

Pujol, B., P. Marrot, and J. R. Pannell. 2014. A quantitative genetic signature of senescence in a

short-lived perennial plant. Current Biology 24:744-747.

R Core Team. 2019 R: A language and environment for statistical computing, version 3.6.1. R

Foundation for Statistical Computing, Vienna, Austria.

Royer, D. L., L. A. Meyerson, K. M. Robertson, and J. M. Adams. 2009. Phenotypic plasticity of

leaf shape along a temperature gradient in *Acer rubrum*. PLoS One 4:e7653.

Shaw, R. G. 2018. From the Past to the Future: Considering the Value and Limits of

Evolutionary Prediction. The American Naturalist 193:1-10.

Shaw, R. G., and J. R. Etterson. 2012. Rapid climate change and the rate of adaptation: insight

from experimental quantitative genetics. New Phytologist 195:752-765.

Sheth, S. N., and A. L. Angert. 2016. Artificial Selection Reveals High Genetic Variation in

Phenology at the Trailing Edge of a Species Range. The American Naturalist 187:182-

193.

Sheth, S. N., M. W. Kulbaba, R. E. Pain, and R. G. Shaw. 2018. Expression of additive genetic

variance for fitness in a population of partridge pea in two field sites. Evolution 72:2537-

2545.

Soneson, C., M. I. Love, and M. D. Robinson. 2015. Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. F1000Research 4.

Stratton, D. A. 1994. Genotype-by-environment interactions for fitness of *Erigeron annuus* show

fine-scale selective heterogeneity. Evolution 48:1607-1618.

Via, S., R. Gomulkiewicz, G. De Jong, S. M. Scheiner, C. D. Schlichting, and P. H. Van

Tienderen. 1995. Adaptive phenotypic plasticity: consensus and controversy. Trends in

Ecology & Evolution 10:212-217.

Walter, G. M., J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018. Evolution of genetic

variance during adaptive radiation. The American Naturalist 191:E108–E128.

Walter, G. M., S. Catara, J. R. Bridle, and A. Cristaudo. 2020a. Population variation in early

development can determine ecological resilience in response to environmental change.

New Phytologist 226:1312-1324.

Walter, G. M., J. Clark, A. Cristaudo, B. Nevado, S. Catara, M. Paunov, V. Velikova et al.

2020b. Adaptive divergence generates distinct plastic responses in two closely related

Senecio species. bioRxiv.

Walter, G. M., M. J. Wilkinson, M. E. James, T. J. Richards, J. D. Aguirre, and D. Ortiz-

Barrientos. 2016. Diversification across a heterogeneous landscape. Evolution 70:1979-

1992.

Wang, S. P., and D. M. Althoff. 2019. Phenotypic plasticity facilitates initial colonization of a

novel environment. Evolution 73:303-316.

Acknowledgements: We thank J.D. Aguirre, S. Barrett, R. Dugand and A. Hoffmann for

comments on previous versions. We are very grateful to those who helped with fieldwork: A.

Barbato, O. Brayley, G. Burstein, M. Castrogiovanni, S. du Plessis, C. Impelluso, E. la Spina,

M.G. Majorana, J. Menzies, M. Millen, G. Pepe and D. Ward, and to Piante Faro for providing

glasshouse resources. This work was carried out using the computational facilities of the

Advanced Computing Research Centre, University of Bristol. **Funding:** This work was

supported by joint NERC grants NE/P001793/1 and NE/P002145/1 awarded to JB and SH.

Author contributions: GW, SH and JB designed the experiment with input from SC and AC.

GW, AC and DT conducted the experiments. JC collected, extracted and processed RNA

samples, and analyzed the transcriptome data. GW analyzed the data, and wrote the manuscript

with help from JC, JB and SH, and input from all other authors.