Identification and functional prediction of anthocyanin biosynthesis

regulatory long non-coding RNAs (lncRNAs) in carrot

Constanza Chialva[1#], Thomas Blein[2#], Martin Crespi[2*] & Diego Lijavetzky[1*]

[1]Instituto de Biología Agrícola de Mendoza (IBAM), UNCuyo, CONICET, Facultad de

Ciencias Agrarias, Almirante Brown 500, M5528AHB, Chacras de Coria, Mendoza,

Argentina. [2]Institute of Plant Sciences Paris-Saclay (IPS2), CNRS, INRA, University

Paris-Saclay and University of Paris Batiment 630, Gif Sur Yvette, France.

[#]Constanza Chialva and Thomas Blein contributed equally to this work.

[*]Correspondence and requests for materials should be addressed to D.L. (email:

dlijavetzky@conicet.gov.ar) or M.C. (email: martin.crespi@ips2.universite-paris-

saclay.fr)

ABSTRACT

Carrot (*Daucus carota* L.) is one of the most cultivated vegetable in the world and of

great importance in the human diet. Its storage organs can accumulate large quantities of

anthocyanins, metabolites that confer the purple pigmentation to carrot tissues and

whose biosynthesis is well characterized. Long non-coding RNAs (lncRNAs) play

critical roles in regulating gene expression of various biological processes in plants. In

this study, we used a high throughput stranded RNA-seq to identify and analyze the

expression profiles of lncRNAs in phloem and xylem root samples using two genotypes

with a strong difference in anthocyanin production. We identified 639 differentially

expressed lncRNAs between genotypes, and certain were specifically associated with a

particular tissue. We then established regulatory correlations between lncRNAs and

anthocyanin biosynthesis genes in order to identify a molecular framework for the

differential expression of the pathway between genotypes. A specific natural antisense

transcript (NAT) linked to the *DcMYB7* key anthocyanin biosynthetic transcription

factor suggested how the regulation of this pathway may have evolved between

genotypes.

INTRODUCTION

Anthocyanins are flavonoids, a class of phenolic compounds synthesized via the

phenylpropanoid pathway, a late branch of the shikimic acid pathway[1]. They are

secondary metabolites that confer purple, red, and blue pigmentation to several organs

and tissues of many plant species[2]. These water-soluble pigments serve in various roles

in the plant, including attracting pollinators to flowers and seed dispersers to fruits,

protection against UV radiation, amelioration of different abiotic and biotic stresses,

such as drought, wounding, cold temperatures, and pathogen attacks[3,4], as well as

participation in physiological processes such as leaf senescence[5,6]. As dietary

components, anthocyanins possess various health-promoting effects, mainly due to their

antioxidant and anti-inflammatory properties, including protection against cancer,

strokes and other chronic human disorders[7].

Carrot (*Daucus carota* subsp. *carota* L.; $2n = 2x = 18$) is a globally important root crop

with yellow and purple as the first documented colors for domesticated carrot in Central

Asia approximately 1,100 years ago[8]. Orange carrots were not reliably reported until the

sixteenth century in Europe[9,10], where its popularity was fortuitous for modern

consumers because the orange pigmentation results from high quantities of α- and β-

carotene, making carrots the richest source of provitamin A in the US diet[11].

Additionally, with its great nutrition and economic value, carrot has been well known as

a nice model plant for genetic and molecular studies[11]. Carrot is one of the crops that

can accumulate large quantities of anthocyanins in its storage roots (up to 17–18

mg/100 g fresh weight)[12]. Purple carrots accumulate almost exclusively derivatives of

cyanidin glycosides with five cyanidin pigments reported in most studies[13,14]. The root

content of these five anthocyanin pigments vary across carrot genetic backgrounds[12,15].

In addition, anthocyanin pigmentation also varies between root tissues, ranging from

fully pigmented roots (i.e., purple color in the root phloem and xylem) to pigmentation

only in the outer-most layer of the phloem[16,17].

Regardless of the plant species, at least two classes of genes are involved in anthocyanin

biosynthesis: structural genes encoding the enzymes that directly catalyze the

production of anthocyanins, and regulatory genes that control the transcription of

structural genes[18,19]. In most cases, the anthocyanin biosynthetic structural genes are

regulated by transcription factors (TFs) belonging to the R2R3–MYB, basic helix-loop-

helix (bHLH) and WD-repeat protein families, in the form of the 'MBW' complex[19,20].

Recent reports pointed out that gene regulation by TFs may play a key role controlling

anthocyanin pigmentation in purple carrots[17,21,22]. Moreover, the broad variation

observed among purple carrot root genotypes, regarding both anthocyanin concentration

and pigment distribution in the phloem and xylem tissues, suggests independent genetic

regulation in these two root tissues[23]. In this sense, Xu et al.[16] found that the expression

pattern of a R2R3–MYB TF, *DcMYB6*, is correlated with anthocyanins production in

carrot roots and that the overexpression of this gene in *Arabidopsis thaliana* enhanced

anthocyanins accumulation in vegetative and reproductive tissues in this heterologous

system. Similarly, Kodama et al.[24] found that a total of 10 *MYB*, *bHLH* and *WD40* genes

were consistently up- or downregulated in a purple color-specific manner, including

DcMYB6. Iorizzo et al.[25] identified a cluster of MYB TFs, with *DcMYB7* as a candidate

gene for root and petiole pigmentation, and *DcMYB11* as a candidate gene for petiole

pigmentation. Bannoud et al.[23] showed that *DcMYB7* and *DcMYB6* participate in the

regulation of phloem pigmentation in purple-rooted samples. Finally, Xu et al.[26], by

means of lost- and gain-of-function mutation experiments, demonstrated that *DcMYB7*

is the main determinant that controls purple pigmentation in carrot roots.

Non-coding RNAs with a length higher than 200 nucleotides are defined as long

noncoding RNAs (lncRNAs). They were originally considered to be transcriptional

byproducts, or transcriptional 'noise', and were often dismissed in transcriptome

analyses due to their low expression and low sequence conservation compared with

protein-coding mRNAs. However, specific lncRNAs were shown to be involved in

chromatin modification, epigenetic regulation, genomic imprinting, transcriptional

control as well as pre- and post-translational mRNA processing in diverse biological

processes in plants[27–30]. Certain lncRNAs can be precursors of small interfering RNA

(siRNA) or microRNA (miRNAs), triggering the repression of protein-coding genes at

the transcription level (transcriptional gene silencing or TGS) or at post-transcriptional

level (PTGS)[27,31]. Additionally, other lncRNAs can act as endogenous target mimics of

miRNAs, to fine-tune the miRNA-dependent regulation of target genes[32,33]. It has been

suggested that lncRNAs can regulate gene expression in both the *cis*- and *trans*-acting

mode[34]. The *cis*-acting lncRNAs can be classified by their relative position to annotated

genes[27,35,36] and notably include   long noncoding natural antisense (lncNATs)

transcribed in opposite strand of a coding gene, overlapping with at least one of its

exons[37,38]. Other so-called intronic lncRNAs are transcribed within introns of a protein-

coding gene[39] whereas long intergenic ncRNAs (lincRNAs) are transcripts located

farther than 1 kb from protein-coding genes[27,35,36]. Among these cis-lncRNAs, NATs

are of special interest as they have been shown to provide a mechanism for locally

regulating the transcription or translation of the target gene on the other strand,

providing novel mechanisms involved in the regulation of key biological processes[40]

and environmentally dependent gene expression[37,38].

As mentioned above, several differential expression analyses have been performed

between purple and non-purple carrot roots allowing the identification of the main

structural genes and TFs involved in anthocyanin biosynthesis in whole roots and/or

phloem tissues[16,21,23–26]. However, the identification and functional prediction of

lncRNA in carrot or putatively involved in carrot anthocyanin biosynthesis regulation

has not yet been reported. In the present study, we combined a high throughput stranded

RNA-Seq based approach with a dedicated bioinformatic pipeline, to annotate lncRNAs

and analyze the expression profiles of lncNATs putatively associated to the carrot root

anthocyanin biosynthesis regulation. In addition, we individually analyzed the gene

expression patterns in phloem and xylem root of purple and orange *D. carota*

genotypes. Our findings point for a role of antisense transcription in the anthocyanin

biosynthesis regulation in the carrot root at a tissue-specific level.

RESULTS

RNA-seq data mining, identification and annotation of anthocyanin-related

lncRNAs

In order to thoroughly identify and annotate lncRNAs related to anthocyanin

biosynthesis regulation in carrot roots, we performed a whole transcriptome RNA-seq

analysis of the carrot genotypes 'Nightbird' (purple phloem and xylem) and 'Musica'

(orange phloem and xylem) that had been studied in a tissue specific manner

(Supplementary Figure S1). We generated an average of 51.4 million of reads per

sample from the 12 carrot root samples (i.e., two phenotypes x two tissues x three

biological replicates), ranging from 43.5 million to 60.3 million. The average GC

content (%) was 44.8% and the average ratio of bases that have phred[41] quality score of

over 30 (Q30) was 94.1%. The average mapping rate to the carrot genome was 90.9%

(Supplementary Table S1). We identified and annotated 8484 new transcripts, including

2095 new protein-coding transcripts and 6373 noncoding transcripts (1521 lncNATs,

4852 lincRNAs and 16 structural transcripts) (Supplementary Table S2 and

Supplementary File S1). Those were added to the 34263 known carrot transcripts[42] to

conform the final set of 42747 transcripts used for this work. The set contains 34204

coding transcripts and 7288 noncoding transcripts (1521 lncNATs, 5767 lincRNAs) and

1255 structural transcripts (Fig 1A and Supplementary Table S3). Most noncoding

transcripts presented less than 1000 bp long, being 400-800 bp the most frequent length

class. Coding transcripts between 500-1000 bp long were the most frequent, while most

structural transcripts presented less than 200 bp (Fig 1B). Noncoding transcripts

predominantly presented one exon and unexpectedly[43], only one exon was also the most

frequent class for coding transcripts (Fig 1C). Finally, we found no particular bias for

the distribution of the noncoding transcripts along the nine carrot chromosomes (Fig

1D). As expected, the average expression level of the ncRNAs, measured in Transcripts

Per Kilobase Million (TPMs), was lower than that of the mRNAs (Fig. 1E).

Variation in coding and noncoding expression was mainly explained by the

anthocyanin-pigmentation phenotype difference between orange and purple

carrots

We sampled orange and purple carrot genotypes in two different tissues, phloem and

xylem (Supplementary Figure S1). Considering the global gene variation of the 12

evaluated libraries (i.e., three for each phenotype/tissue combination), the color

phenotype was clearly the main source of variation (PC1, 49 %), while the tissue

specificity factor was also important (PC2, 18%), but significantly lower than the root

color phenotype effect (Fig. 2A).

We then assessed the variation in mRNA and ncRNA gene expression between purple

and orange carrot roots in our RNA-seq analysis. A total of 3567 genes were

differentially expressed (DEG) between purple and orange carrots (Bonferroni's

adjusted p-value < 0.01), divided in 2928 mRNA and 639 lncRNAs (Fig. 2B) and

representing 10% and 15% of the mRNA and lncRNA expressed genes, respectively.

Within the 3567 DEGs, we found 1664 downregulated and 1907 upregulated

transcripts. In turn, the downregulated transcripts were distributed into 1343 coding and

319 noncoding transcripts, while the upregulated were divided into 1585 and 320

coding and noncoding transcripts, respectively (Fig. 2B). All information concerning

the differentially expressed analysis and gene annotation is detailed in Supplementary

Table S4.

As expected, we identify several differentially expressed genes (DEG) between the two

genotypes known to be involved in carrot root anthocyanin biosynthesis[21,23–26]. Most of

the known genes of the pathway and their main regulators were differentially expressed

between the two genotypes (Supplementary Table S4). Several genes were induced in

purple tissues and they mainly comprised genes representing: i) the early step in the

flavonoid/anthocyanin pathway, like chalcone synthase (*DcCHS1*/DCAR_030786);

chalcone isomerase (*DcCHI1*/DCAR_027694) and (*DcCHIL*/DCAR_019805);

flavanone 3-hydroxylase (*DcF3H1*/DCAR_009483), and flavonoid 3'-hydroxylase

(*DcF3'H1*/DCAR_014032); ii) cytochrome P450 (CYP450) proteins, putatively related

to the flavonoid and isoflavonoid biosynthesis pathways[23,44]; iii) ATP-binding cassette

(ABC) transporters, potentially related to anthocyanin transport[45,46]; and iv) genes from

the late steps of the pathway, like dihydro-flavonol 4-reductase

(*DcDFR1*/DCAR_021485), leucoanthocyanidin dioxygenase (*DcLDOX1*/

DCAR_006772), and UDP-glycosyltransferase (*DcUFGT*/DCAR_009823) and the

recently described *DcUCGXT1*/DCAR_021269 and *DcSAT1*/MSTRG.8365, which

were confirmed to be responsible for anthocyanins glycosylation and acylation,

respectively[26,47]. Finally, most significant regulatory genes of the pathway, belonging to

the MYB, bHLH and WD40 TF gene families[21,23–26] were also differentially expressed

between purple and orange genotypes (Supplementary Table S4). We further analyzed

the tissue differential expression distribution of those 26 'MBW' TFs and found that

DcMYB6 and *DcMYB7,* the two most studied TFs associated with anthocyanin

biosynthesis regulation[23–26], were differentially expressed between purple and orange

carrots, both in phloem and xylem tissues (Supplementary Figure S2). Interestingly,

three genes recently described to be regulated by *DcMYB7*[26] (i.e. *DcbHLH3,*

DcUCGXT1 and *DcSAT1*) also displayed no tissue specificity. *DcbHLH3* was described

as a co-regulator in anthocyanin biosynthesis, while *DcUCGXT1* and *DcSAT1*

participate in anthocyanin glycosylation and acylation, respectively[26,47]. Additionally,

seven TFs showed xylem preferential expression-specificity, while only one was

preferentially expressed specifically in phloem. Finally, differential expression of 11

TFs was just detected when the 12 libraries were jointly analyzed, presumably by

presenting significant but low expression differences (Supplementary Figure S2).

Putative regulation of anthocyanin-related genes by carrot antisense lncRNAs

In order to investigate the putative involvement of carrot lncRNAs in the regulation of

the anthocyanin biosynthesis in different carrot root tissues, we predicted the potential

targets of lncRNAs in *cis*-regulatory relationship, particularly those classified as natural

antisense transcripts (lncNATs). The selection of such lncRNAs was based in three

assumptions: i) both, the lncRNA and the putative target were differentially expressed

between purple and orange tissues (Supplementary Table S4); ii) the lncRNAs were

antisense of the target genes; and iii) the Pearson and Spearman correlation coefficients

between the expression levels of these genes were ≥ 0.70 or ≤ -0.70, and $p < 0.01$.

According to these criteria, we found 19 differentially expressed lncNATs, since the

lncRNAs were located in the antisense orientation (in the opposite strand) to a target

mRNA, being most of them fully overlapping pairs (Supplementary Table S4 and S5).

About 79% of those lncNATs were expressed in concordance with sense strand

transcript, while five out of the 19 presented discordant expression (i.e. when the

lncNAT expression increase, the sense strand transcript was repressed) (Supplementary

Table S4 and S5). Interestingly, we detected two lncNATs (MSTRG.27767/asDcMyb6

and MSTRG.9120/asDcMyb7) in antisense relationship to *DcMYB6* and *DcMYB7*,

respectively, with concordant expression correlation (Fig. 3). *DcMYB6* showed a log2

fold-change of 7.6 with an adjusted *p*-value of 4.5 10^{-30}, while *DcMYB7* presented a

log2 fold-change of 11.7 with an adjusted *p*-value of 3.8 10^{-37}. Accordingly, the two

detected antisense lncRNAs also presented significant differential expression, where

asDcMYB6 displayed a log2 fold-change of 6.5 with an adjusted *p*-value of 2.1 10^{-13}

and *asDcMYB7* presented a log2 fold-change of 6.1 with an adjusted *p*-value of 1.3 10^{-}

^{04} (Supplementary Table S4). Finally, the Pearson and Spearman correlation coefficients

between the expression levels of each sense/antisense pair were ≥ 0.79 and *p*-value $<$

0.01 (Supplementary Table S5). On the other hand, as also detailed in Supplementary

Table S4, two out of the four lncNATs showing discordant expression were found in the

antisense relationship with disease resistance-like related genes (a predicted Catalase,

and probable disease resistance protein At5g63020).

DISCUSION

The presence of color in flowers, fruits and other organs and tissues, plays several

biological functions mostly driven by the adaptive behavior of plants in response to the

environment[2,20,48,49]. But in turn, plant organ pigmentation have served as natural

genetic markers since the early works of Mendel[50,51]. Anthocyanins are flavonoid

pigments that accumulate in plant cell vacuoles[52] and are mainly responsible for most

tissue and organ coloration[19,20,48]. Genetic analyses using model plant species like

Arabidopsis, petunia and maize allowed the identification of most structural genes in the

anthocyanin biosynthesis pathway as well as the main regulatory genes controlling

pigment synthesis. In carrot, anthocyanin pigmentation is responsible for the purple

phenotype[9,53]. Two main genes, *P₁ and P₃*, have been identified in chromosome 3 and

suggested to be responsible for the two independent mutations underlying the

domestication of purple carrots[17]. Despite several carrot structural genes from the

anthocyanin biosynthesis pathway have shown expression correlation with the purple

phenotype[21,22], none of them co-localize with *P₁ and P₃*. Similar situation occurred in

other plants like grapevine, where accumulation of anthocyanins correlated with the

expression of several structural genes of the pathway but none of them co-localized with

the 'color locus' in chromosome 2[54,55]. Finally, this discrepancy was solved by a study

describing an insertion mutation in the promotor of a R2R3–MYB TF (i.e. *VviMybA1*)[56]

explaining the lack of color of white grapevine cultivars. In the same direction, several

recent works[16,23–25,47] focused on the role of carrot TFs putatively involved in the

regulation of anthocyanin biosynthesis in purple genotypes, particularly those belonging

to the 'MBW' complex (i.e., R2R3–<u>M</u>YB, basic helix-loop-helix -<u>b</u>HLH- and <u>W</u>D-

repeat TFs). Two recent reports showed that three R2R3–MYB TFs are involved in the

P₁ and P₃ loci: *DcMYB113* has been suggested to correspond to *P₁*[47], while *DcMYB6*

and *DcMYB7* were proposed as the two main candidate TFs underlying the carrot root

anthocyanin pigmentation in the *P₃* locus[25]. However, knockdown and overexpression

functional analyses demonstrated that *DcMYB7* (but not *DcMYB6*) is the *P₃* gene

controlling purple pigmentation in carrot roots[26]. Likewise described for the grapevine

VviMybA1 gene[56], non-purple carrot genotypes seems to arise by an insertion mutation

in the promoter region of *DcMYB7*[26], yet the authors imply the existence of an

additional genetic factor suppressing the expression of *DcMYB7* in non-purple

pigmented peridermal carrot root tissues.

In this work, we performed a thorough transcriptomic analysis by comparing two carrot

hybrids with contrasted anthocyanin pigmentation phenotypes (i.e. purple vs. orange),

both in phloem and xylem tissues. The study corroborates the involvement of the

principal reported structural genes of the anthocyanin biosynthesis pathway[21,22], but

mostly, the key TF genes reported as the main regulators explain the carrot purple

phenotype (i.e. *DcMYB6* and *DcMYB7*)[16,25,26]. Interestingly, the performed dissection

between phloem and xylem purple samples, allowed us to show that there is no tissue-

specific expression of such key genes, contrary to previously suggested for *DcMYB6*

and *DcMYB7*[16,23,25].

We showed here the first whole genome identification and annotation of lncRNAs in

carrot by combining a high throughput stranded RNA-Seq based approach with a

focused bioinformatic pipeline. Through this process, we identified 6373 novel

lncRNAs, as compared to the 915 sequences annotated in the original carrot genome

assembly[42]. Moreover, 10% of them (641 genes) can be defined as anthocyanin

biosynthesis-related lncRNAs since we found them differentially expressed between

purple and orange carrots. For the functional annotation of such lncRNAs, we focused

on those showing an antisense relationship to differentially expressed protein coding

genes, known (or putatively) involved in carrot anthocyanin biosynthesis and depicted

in the precedent paragraph. Additionally, the selected lncNATs had to present a

statistically significant Pearson and Spearman correlation with their putative targets.

This led us to identify 19 differentially expressed lncNATs between purple and orange

carrots. Interestingly, we found two of these lncNATs (*asDcMYB6* and *asDcMYB7*)

transcribed in opposite direction to *DcMYB6* and *DcMYB7*, respectively. Moreover,

asDcMYB6 and *asDcMYB7* exhibited concordant expression patterns with their

corresponding sense transcripts opening the possibility that non-coding RNA antisense

transcription is a new player in the regulation of carrot anthocyanin biosynthesis,

putatively affecting the expression of *DcMYB7* (and/or *DcMYB6*). This regulation

maybe linked to the previously described genetic factors[26].

Antisense transcripts, particularly lncNATs, present in many genomes of diverse

kingdoms, showed either positively or negatively correlated expression with their

corresponding sense transcripts. In many cases, lncNATs regulate the expression of

their sense transcripts in a negative or positive way, by means of different

transcriptional or post-transcriptional mechanisms. In particular cases, upregulation of

sense gene expression may be explained by the participation of a lncNAT in the

inhibition of other factors at translational level, such as efficient translation initiation or

elongation[57–59].

In plants, both repression and activation roles have been assigned to some lncNATs in

response to environmental conditions. While *COOLAIR* and *COLDAIR* negatively

regulates *FLC* in vernalization responses[39,60], and *SVALKA* controls *CBF1* expression to

consequently regulate freezing tolerance[38], the expression of another member of the

FLC family (*MAF4*) is activated by the lncNAT MAS to fine-tune flowering time[37].

Anthocyanins are known to participate in abiotic stress responses and adaptation to

environmental variations[3,4,61], so the evolutionary role of the newly identified antisense

transcrpts *asDcMYB7* and *asDcMYB6* may be linked to the activation of anthocyanin

biosynthesis through *DcMYB7* and *DcMYB6*. Hence, our work hints to new antisense

regulations potentially involved in the variable expression of anthocyanin genes among

carrot ecotypes.

METHODS

Sample preparation and plant material

Total RNA was obtained independently from three biological replicates of phloem and

xylem root samples of two *Daucus carota* L genotypes: 'Nightbird', a purple root

hybrid (purple phloem and xylem) and 'Musica', a non-anthocyanin pigmentated root

hybrid. Plants were germinated from seeds and roots were collected after 12 weeks.

Frozen samples were grinded using liquid nitrogen and RNA was extracted using TRI

Reagent® (Sigma-Aldrich) and purified using SV Total RNA Isolation System

(Promega). RNA samples were quantified, and purity measured using a

spectrophotometer (AmpliQuant AQ-07). RNA integrity and potential genomic DNA

contaminations were checked through agarose gel electrophoresis.

Libraries construction and RNA sequencing

Twelve samples (two genotypes x two tissues x three biological replicates) were sent to

the Macrogen sequencing service (Seoul, Korea). Once in destiny they were checked for

total RNA integrity using a Bioanalyzer RNA Nano 6000 chip. All the samples

qualified to proceed with the library construction having an RNA Integrity Number

(RIN) ≥7. NGS transcriptomic libraries were constructed using a TruSeq Stranded

mRNA LT Sample Prep Kit (Illumina). To verify the size of PCR enriched fragments,

the template size distribution was checked on an Agilent Technologies 2100

Bioanalyzer using a DNA 1000 chip. The sequencing of libraries was performed as

paired-end 101 bp reads on an Illumina HiSeq 2500 platform. The quality of the raw

reads in the FastQ files was checked through FastQC[62] and were then trimmed for

sequencing adaptor and low quality sequences using Trimmomatic[63] using

'ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 LEADING:21 TRAILING:21 MINLEN:30'

as parameters. For removing reads corresponding to remaining ribosomal RNA,

trimmed reads were mapped to the rRNA reference using SortMeRNA[64] using '--ref

silva-bac-16s-id90.fasta --ref silva-bac-23s-id98.fasta --ref silva-euk-18s-id95.fasta --ref

silva-euk-28s-id98.fasta --paired_in --fastx --log -e 1e-07 -a 4 -v' as parameters.

New transcripts assembling and lncRNA identification

Clean filtered reads were aligned on the *Daucus carota* genome[42] using the STAR

aligner[65] using '--alignIntronMin 20 --alignIntronMax 20000 --outSAMtype BAM

SortedByCoordinate --outReadsUnmapped Fastx' as parameters. Subsequently, the

aligned reads were assembled by means of StringTie[66] and new transcripts were

extracted and annotated using the GffCompare[67] program (GffCompare classes "u", "x",

to adjust). Only new transcripts whose length was greater than 200nt were kept. The

classification of the newly predicted transcript was performed according the following

steps: i) the transcripts were classified as coding if their predicted open reading frame

(ORF) was greater than 120 aa or if they were predicted as coding by CPC2[68] calculator

and classify as structural RNA in case of homology with structural RNA (tRNA, rRNA,

snRNA or snoRNA) or as non-coding in case of homology with known structured non-

coding RNA (miRNA precursors, lncRNA). The transcript without any homology in

Rfam and classified as non-coding par CPC2 were classified as non-coding.

Differential expression analysis

We performed a strand-specific read counting of coding and non-coding gene using on

the carrot official annotation and the newly predicted genes of this study for each of the

12 aligned BAM files by means of the featureCount[70] software included in the Rsubread

package[71]. The resulted count data was used for differential expression analysis with

DEseq2[72]. Differentially expressed genes were declared as having a Bonferroni's

adjusted p-value < 0.01. Reads corresponding to the strand specific expression of

mRNAs and their lncNATs were visualized with the Integrative Genomics Viewer

(IGV) software[73]. Additional Venn diagrams were performed with Venny v2.1[74].

REFERENCES

1.   Herrmann, K. M. & Weaver, L. M. The Shikimate Pathway. *Annu. Rev. Plant*

Physiol. Plant Mol. Biol. **50**, 473–503 (1999).

2.   Harborne, J. B. & Williams, C. A. Advances in flavonoid research since 1992.

Phytochemistry **55**, 481–504 (2000).

3.   Koes, R. E., Quattrocchio, F. & Mol, J. N. M. M. The flavonoid biosynthetic

pathway in plants: Function and evolution. *BioEssays* **16**, 123–132 (1994).

4.   Shirley, B. W. Flavonoid biosynthesis: 'new' functions for an 'old' pathway.

Trends Plant Sci. **1**, 377–382 (1996).

5.   Hatier, J.-H. B. & Gould, K. S. Anthocyanin Function in Vegetative Organs. in

Anthocyanins 1–19 (Springer New York, 2008). doi:10.1007/978-0-387-77335-

3_1

6.   Gould, K. S. & Lister, C. Flavonoid function in plants. *Flavonoids Chem.*

Biochem. Appl. 397–441 (2006).

7.   Lila, M. A. Anthocyanins and human health: An in vitro investigative approach.

Journal of Biomedicine and Biotechnology **2004**, 306–313 (2004).

8.   Iorizzo, M. *et al.* Genetic structure and domestication of carrot (Daucus carota

subsp. sativus) (Apiaceae). *Am. J. Bot.* **100**, 930–938 (2013).

9.   Simon, P. W. Domestication, Historical Development, and Modern Breeding of

Carrot. in *Plant Breeding Reviews* 157–190 (John Wiley & Sons, Inc., 2010).

doi:10.1002/9780470650172.ch5

10.   Arscott, S. A. & Tanumihardjo, S. A. Carrots of Many Colors Provide Basic

Nutrition and Bioavailable Phytochemicals Acting as a Functional Food. *Compr.*

Rev. Food Sci. Food Saf. **9**, 223–239 (2010).

11.   Simon, P. W., Pollak, L. M., Clevidence, B. A., Holden, J. M. & Haytowitz, D.

B. Plant Breeding for Human Nutritional Quality. in *Plant Breeding Reviews* **31**,

325–392 (John Wiley & Sons, Inc., 2009).

12.   Montilla, E. C., Arzaba, M. R., Hillebrand, S. & Winterhalter, P. Anthocyanin

composition of black carrot (Daucus carota ssp. sativus var. atrorubens Alef.)

Cultivars antonina, beta sweet, deep purple, and purple haze. *J. Agric. Food*

Chem. **59**, 3385–3390 (2011).

13.   Kammerer, D., Carle, R. & Schieber, A. Quantification of anthocyanins in black

carrot extracts (Daucus carota ssp. sativus var. atrorubens Alef.) and evaluation

of their color properties. *Eur. Food Res. Technol.* **219**, 479–486 (2004).

14.   Kammerer, D., Carle, R. & Schieber, A. Detection of peonidin and pelargonidin

glycosides in black carrots (Daucus carota ssp.sativus var.atrorubens Alef.) by

high-performance   liquid   chromatography/electrospray   ionization   mass

spectrometry. *Rapid Commun. Mass Spectrom.* **17**, 2407–2412 (2003).

15.   Mazza, G. & Miniati, E. (Enrico). *Anthocyanins in fruits, vegetables, and grains*.

(CRC Press, 1993).

16.   Xu, Z. S., Feng, K., Que, F., Wang, F. & Xiong, A. S. A MYB transcription

factor, DcMYB6, is involved in regulating anthocyanin biosynthesis in purple

carrot taproots. *Sci. Rep.* **7**, 1–9 (2017).

17. Cavagnaro, P. F. *et al.* A gene-derived SNP-based high resolution linkage map of

carrot including the location of QTL conditioning root and leaf anthocyanin

pigmentation. *BMC Genomics* **15**, 1118 (2014).

18. Jaakola, L. *et al.* Expression of genes involved in anthocyanin biosynthesis in

relation to anthocyanin, proanthocyanidin, and flavonol levels during bilberry

fruit development. *Plant Physiol* **130**, 729–39 (2002).

19. Koes, R., Verweij, W. & Quattrocchio, F. Flavonoids: a colorful model for the

regulation and evolution of biochemical pathways. *Trends Plant Sci* **10**, 236–242

(2005).

20. Winkel-Shirley, B. Flavonoid biosynthesis. A colorful model for genetics,

biochemistry, cell biology, and biotechnology. *Plant Physiol* **126**, 485–493

(2001).

21. Yildiz, M. *et al.* Expression and mapping of anthocyanin biosynthesis genes in

carrot. *Theor. Appl. Genet.* **126**, 1–14 (2013).

22. Xu, Z. S. *et al.* Transcript profiling of structural genes involved in cyanidin-based

anthocyanin biosynthesis between purple and non-purple carrot (Daucus carota

L.) cultivars reveals distinct patterns. *BMC Plant Biol.* **14**, (2014).

23. Bannoud, F. *et al.* Dissecting the genetic control of root and leaf tissue-specific

anthocyanin pigmentation in carrot (Daucus carota L.). *Theor. Appl. Genet.* **132**,

2485–2507 (2019).

24. Kodama, M. *et al.* Identification of transcription factor genes involved in

anthocyanin biosynthesis in carrot (Daucus carota L.) using RNA-Seq. *BMC*

Genomics **19**, 811 (2018).

25.   Iorizzo, M. *et al.* A Cluster of MYB Transcription Factors Regulates
443   Anthocyanin Biosynthesis in Carrot (Daucus carota L.) Root and Petiole. *Front.*
444   *Plant Sci.* **9**, 1927 (2018).

445   26.   Xu, Z.-S., Yang, Q.-Q., Feng, K. & Xiong, A.-S. Changing Carrot Color:
446   Insertions in DcMYB7 Alter the Regulation of Anthocyanin Biosynthesis and
447   Modification. *Plant Physiol.* pp.00523.2019 (2019). doi:10.1104/pp.19.00523

448   27.   Ariel, F., Romero-Barrios, N., Jégu, T., Benhamed, M. & Crespi, M. Battles and
449   hijacks: noncoding transcription in plants. *Trends Plant Sci.* **20**, 362–371 (2015).

450   28.   Forestan, C. *et al.* Stress-induced and epigenetic-mediated maize transcriptome
451   regulation study by means of transcriptome reannotation and differential
452   expression analysis. *Sci. Rep.* **6**, 1–20 (2016).

453   29.   Forestan, C. *et al.* Epigenetic signatures of stress adaptation and flowering
454   regulation in response to extended drought and recovery in Zea mays. *Plant Cell*
455   *Environ.* **43**, 55–75 (2020).

456   30.   Mercer, T. R. & Mattick, J. S. Structure and function of long noncoding RNAs in
457   epigenetic regulation. *Nature Structural and Molecular Biology* **20**, 300–307
458   (2013).

459   31.   Wu, H. J., Wang, Z. M., Wang, M. & Wang, X. J. Widespread long noncoding
460   RNAs as endogenous target mimics for microRNAs in plants. *Plant Physiol.* **161**,
461   1875–1884 (2013).

462   32.   Zhang, G. *et al.* Transcriptomic and functional analyses unveil the role of long
463   non-coding RNAs in anthocyanin biosynthesis during sea buckthorn fruit
464   ripening. *DNA Res.* **25**, 465–476 (2018).

465   33.   Yang, T. *et al.* Systematic identification of long noncoding <scp>RNA</scp> s
466   expressed during light-induced anthocyanin accumulation in apple fruit. *Plant J.*

100, 572–590 (2019).

34. Li, Z. & Rana, T. M. Molecular mechanisms of RNA-triggered gene silencing machineries. *Acc. Chem. Res.* **45**, 1122–1131 (2012).

35. Rinn, J. L. & Chang, H. Y. Genome Regulation by Long Noncoding RNAs. *Annu. Rev. Biochem.* **81**, 145–166 (2012).

36. Bonasio, R. & Shiekhattar, R. Regulation of Transcription by Long Noncoding RNAs. *Annu. Rev. Genet.* **48**, 433–455 (2014).

37. Zhao, X. *et al.* Global identification of Arabidopsis lncRNAs reveals the regulation of MAF4 by a natural antisense RNA. *Nat. Commun.* **9**, 1–12 (2018).

38. Kindgren, P., Ard, R., Ivanov, M. & Marquardt, S. Transcriptional read-through of the long non-coding RNA SVALKA governs plant cold acclimation. *Nat. Commun.* **9**, (2018).

39. Heo, J. B. & Sung, S. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science (80-.).* **331**, 76–79 (2011).

40. Thieffry, A. *et al.* Characterization of Arabidopsis thaliana promoter Bidirectionality and Antisense RNAs by Depletion of Nuclear RNA Decay Pathways. *Plant Cell* tpc.00815.2019 (2020). doi:10.1105/tpc.19.00815

41. Ewing, B., Hillier, L. D., Wendl, M. C. & Green, P. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* **8**, 175–185 (1998).

42. Iorizzo, M. *et al.* A high-quality carrot genome assembly provides new insights into carotenoid accumulation and asterid genome evolution. *Nat. Genet.* **48**, 657–666 (2016).

43. Ramírez-Sánchez, O., Pérez-Rodríguez, P., Delaye, L. & Tiessen, A. Plant Proteins Are Smaller Because They Are Encoded by Fewer Exons than Animal

Proteins. *Genomics, Proteomics Bioinforma.* **14**, 357–370 (2016).

44.   Ayabe, S. I. & Akashi, T. Cytochrome P450s in flavonoid metabolism.

Phytochemistry Reviews **5**, 271–282 (2006).

45.   Kovinich, N. *et al.* Arabidopsis MATE45 antagonizes local abscisic acid

signaling to mediate development and abiotic stress responses. *Plant Direct* **2**,

e00087 (2018).

46.   Francisco, R. M. *et al.* ABCC1, an ATP Binding Cassette Protein from Grape

Berry, Transports Anthocyanidin 3-O-Glucosides. *Plant Cell Online* (2013).

doi:10.1105/tpc.112.102152

47.   Xu, Z., Yang, Q., Feng, K., Yu, X. & Xiong, A. DcMYB113, a root-specific

R2R3-MYB, conditions anthocyanin biosynthesis and modification in carrot.

Plant Biotechnol. J. pbi.13325 (2020). doi:10.1111/pbi.13325

48.   Mol, J., Grotewold, E. & Koes, R. How genes paint flowers and seeds. *Trends*

Plant Sci **3**, 212–217 (1998).

49.   Winkel-Shirley, B. Biosynthesis of flavonoids and effects of stress. *Curr Opin*

Plant Biol **5**, 218–223 (2002).

50.   Henig, R. M. *The monk in the garden: the lost and found genius of Gregor*

Mendel, the father of genetics. (Houghton Mifflin Harcourt, 2017).

51.   Hartl, D. L. & Orel, V. What did Gregor Mendel think he discovered? *Genetics*

131, 245 (1992).

52.   Petrussa, E. *et al.* Plant Flavonoids—Biosynthesis, Transport and Involvement in

Stress Responses. *Int. J. Mol. Sci.* **14**, 14950–14973 (2013).

53.   Simon, P. W. Inheritance and expression of purple and yellow storage root color

in carrot. *J. Hered.* **87**, 63–66 (1996).

54.   Kobayashi, S., Ishimaru, M., Ding, C. K., Yakushiji, H. & Goto, N. Comparison

of UDP-glucose:flavonoid 3-O-glucosyltransferase (UFGT) gene sequences

between white grapes (Vitis vinifera) and their sports with red skin. *Plant Sci*

160, 543–550 (2001).

55.   Kobayashi, S., Ishimaru, M., Hiraoka, K. & Honda, C. Myb-related genes of the

Kyoho grape (Vitis labruscana) regulate anthocyanin biosynthesis. *Planta* **215**,

924–933 (2002).

56.   Kobayashi, S., Goto-Yamamoto, N. & Hirochika, H. Retrotransposon-induced

mutations in grape skin color. *Science (80-.).* **304**, 982 (2004).

57.   Pelechano, V. & Steinmetz, L. M. Gene regulation by antisense transcription.

Nature Reviews Genetics **14**, 880–893 (2013).

58.   Wight, M. & Werner, A. The functions of natural antisense transcripts. *Essays*

Biochem. **54**, 91–101 (2013).

59.   Faghihi, M. A. & Wahlestedt, C. Regulatory roles of natural antisense transcripts.

Nature Reviews Molecular Cell Biology **10**, 637–643 (2009).

60.   Swiezewski, S., Liu, F., Magusin, A. & Dean, C. Cold-induced silencing by long

antisense transcripts of an Arabidopsis Polycomb target. *Nature* **462**, 799–802

(2009).

61.   Rienth, M. *et al.* Day and night heat stress trigger different transcriptomic

responses in green and ripening grapevine (vitis vinifera) fruit. *BMC Plant Biol.*

14, 108 (2014).

62.   Andrews, S. FastQC: a quality control tool for high throughput sequence data.

(2010).   Available   at:

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

63.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

64. Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217

(2012).

65. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,

15–21 (2013).

66. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome

from RNA-seq reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

67. GffCompare. A Program for processing GTF/GFF files. Available at:

https://ccb.jhu.edu/software/stringtie/gffcompare.shtml.

68. Kang, Y.-J. *et al.* CPC2: a fast and accurate coding potential calculator based on

sequence intrinsic features. *Nucleic Acids Res.* **45**, W12–W16 (2017).

69. Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology

searches. *Bioinformatics* **29**, 2933–2935 (2013).

70. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**,

923–930 (2014).

71. Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster,

cheaper and better for alignment and quantification of RNA sequencing reads.

Nucleic Acids Res. **47**, e47–e47 (2019).

72. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

73. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics

Viewer (IGV): high-performance genomics data visualization and exploration.

Brief. Bioinform. **14**, 178–192 (2012).

74. Oliveros, J. C. Venny. *Venny. An interactive tool for comparing lists with Venn's*

567   *diagrams.* (2015).

570   **ACKNOWLEDGEMENTS**

571   This work was supported by Agencia Nacional de Promoción Científica y Tecnológica

572   (ANPCyT): PICT2015-0822 & PICT-2016-3134; LabEx Sciences des Plantes de Saclay

573   (SPS). Travel suport to Diego Lijavetzky was funded by Programa de Movilidad para

574   Docentes de la UNCuyo, INRA and Universite Paris Sud. We thank to Dr. Pablo

575   Cavagnaro for providing and growing the plant materials.

577   **AUTHOR CONTRIBUTIONS**

578   C.C. performed the samples collection, laboratory work for library preparation,

579   participated in the draft of the manuscript and figures preparation; T.B. designed the

580   bioinformatic pipeline related to raw data processing, new transcripts identification and

581   differential expression analysis. M.C provided the main insight on the design of the

582   transcriptomic experiment. D.L. wrote the final manuscript, designed and coordinated

583   the experiments. All authors carefully read and helped to improve the final content of

584   the manuscript.

587   **ADDITIONAL INFORMATION**

588     **Data access**: Sequence files generated during this study have been deposited into the

589     NCBI BioProject database accession PRJNA668894.

590     **Competing Interests:** The authors declare no competing interests.

592   **FIGURE LEGENDS**

Figure 1. Characteristics of carrot transcripts. (A) Distribution of coding, noncoding

and structural sequences between the known and newly annotated transcripts. (B)

Transcript length distributions for the total coding, noncoding and structural RNAs. (C)

Number of exons per transcript for the total coding and noncoding RNAs. (D)

Proportional distribution of the total coding, noncoding and structural RNAs along each

chromosome. (E) Violin plot of the expression levels of carrot total coding and

noncoding RNAs. The y-axis represents the average log2(TPM) values. T-test p-value <

0.01 is considered to be significantly different.

Figure 2. Expression of carrot coding and noncoding RNAs. (A) PCA analysis of the

global gene expression of the 12 evaluated libraries (three replicates for each color-

phenotype and tissue type combination). (B) Differentially expressed genes (up- and

down-regulated) between purple and orange carrots (Bonferroni's adjusted p-value <

0.01) distributed by coding and noncoding transcripts.

Figure 3. Strand specific expression of R2R3–MYB TFs and their lncNATs. Coverage

data for the sense (green) and antisense (red) strands corresponding to

DcMYB7/asDcMYB7 (A) and *DcMYB6/as DcMYB6* (B), respectively. Tracks

correspond to four carrot libraries: two phloem samples Purple_F1 and Orange F1; and

two xylem samples Purple_X1 and Orange_X1. Data Range of each track was set to

allow an even visualization of the mRNA and lncRNA transcripts by enlarging the last

ones (20x).

Supplementary Figure S1. Picture of the purple and orange hybrids used for the RNA-

seq experiment and scheme of the performed dissection between xylem and phloem

tissues.

Supplementary Figure S2. Tissue specific differential expression of the 26 'MBW'

TFs identified in the experiment. a) Genes differentially expressed between purple and

orange carrots both in xylem and phloem tissues; b) genes differentially expressed

between purple and orange carrots just in xylem; c) gene differentially expressed

between purple and orange carrots just in phloem; d) genes differentially expressed

between purple and orange carrots detected after the join analysis of phloem and xylem

samples.

Supplementary Table S1. Summary of NGS and quality control data regarding the 12

sequenced libraries.

Supplementary Table S2. Genome annotation of the newly identified transcripts.

Supplementary Table S3. Known and newly annotated carrot genes classified as

coding, noncoding and structural transcripts.

Supplementary Table S4. Overall differentially expressed genes (DEGs) list, including

statistical tests, *cis*-located sequences, gene lengths and gene products. The 21 identified

lncNAT/coding transcript pairs are sorted on top of the list.

Supplementary Table S5. Pearson and Spearman correlation coefficients between the

expression levels of the 19 identified lncNAT/coding transcript pairs across the 12

analyzed libraries.

Supplementary File S1. FASTA sequences of the newly annotated transcripts.

A

B