Single-cell RNA-seq analysis of human coronary arteries using an enhanced workflow

reveals SMC transitions and candidate drug targets

Wei Feng Ma[1,2], Chani J. Hodonsky[2], Adam W. Turner[2], Doris Wong[2,3], Yipei Song[2,4], Nelson B.

Barrientos[2,5], Clint L. Miller[2,3,5]*

Affiliations

[1]Medical Scientist Training Program, University of Virginia, Charlottesville, VA 22908, USA.

[2]Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA.

[3]Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA

22908, USA.

[4]Department of Computer Engineering, University of Virginia, Charlottesville, VA 22908, USA.

[5]Department of Public Health Sciences, University of Virginia, Charlottesville, VA 22908, USA.

*corresponding author

Correspondence:

Clint L. Miller, PhD. (clintm@virginia.edu)

University of Virginia

Center for Public Health Genomics

PO Box 800717

Charlottesville, VA 22908 USA

Abstract

Recent advances in single-cell RNA sequencing (scRNA-seq) methods have enabled high-resolution profiling and quantification of cellular expression and transcriptional states. Here we incorporate automated cell labeling, pseudotemporal ordering, ligand-receptor evaluation, and drug-gene interaction analysis into an enhanced and reproducible scRNA-seq analysis workflow. We applied this analysis method to a recently published human coronary artery scRNA dataset and revealed distinct derivations of chondrocyte-like and fibroblast-like cells from smooth muscle cells (SMCs). We highlighted several key ligand-receptor interactions within the atherosclerotic environment through functional expression profiling and revealed several attractive avenues for future pharmacological repurposing. This publicly available workflow will also allow for more systematic and user-friendly analysis of scRNA datasets in other disease systems.

Keywords

Atherosclerosis, single-cell RNA-seq, analysis, pseudotemporal, drug-target, workflow

Abbreviations

C7- complement component C7

CAD- coronary artery disease

CH- chondrocytes

CMP- common myeloid progenitor cells

DCN- decorin

DGIdb- drug-gene interaction database

EC- endothelial cells

EGFR- epidermal growth factor receptor

FB- fibroblasts

FBLN1- fibulin 1

GMP- granulocyte-monocyte progenitor cells.

Mø- macrophages

MYH11- myosin heavy chain 11

SC- stem cells

sc/snATAC-seq - single cell/single nucleus assay for transposase-accessible chromatin

sequencing

sc/snRNA-seq- single cell/single nucleus RNA sequencing

SMC- smooth muscle cells

UMAP- uniform manifold approximation and projection

Background

Atherosclerosis is a complex process involving chronic inflammation and hardening of the vessel wall and represents one of the major causes of coronary artery disease (CAD), peripheral artery disease, and stroke (Basatemur et al., 2019). Rupture of an unstable atherosclerotic lesion can lead to the formation of a thrombus, causing complete or partial occlusion of a coronary artery (Argraves et al., 2009). The contribution of smooth muscle cells (SMCs) to both lesion stability and progression has recently been established by numerous groups, but the exact mechanisms in which SMCs modulate the atherosclerotic microenvironment and whether pharmacological agents can be used to selectively counter SMC-related deleterious mechanisms are still under investigation (Bennett et al., 2016; Pan et al., 2020; Wirka et al., 2019).

The recent advances in single-cell RNA-sequencing (scRNA-seq) have allowed for ultra-fine gene expression profiling of many diseases at the cellular level, including coronary artery disease (Wirka et al., 2019). As sequencing costs continue to decline, there has also been a consistent growth in scRNA datasets, analysis tools and applications. Currently, a major challenge with scRNA-seq analysis is the inherent bias introduced during manual cell labeling, in which cells are grouped by cluster and their identities called collectively based on their overall differential gene expression profiles (Aran et al., 2019). Another draw-back inherent to commonly used scRNA-seq protocols is their destructive nature to the cells, making time-series analyses of the same samples impossible. Instead, these studies must rely on time-points from separate libraries to monitor processes such as clonal expansion and cell differentiation (DiRenzo et al., 2017; Wang et al., 2020).

Recently, new approaches have been developed to compensate for both of these shortcomings,

namely automatic cell labeling and pseudotemporal analysis. Tools such as 'SingleR' and

'Garnett' have been used to assign unbiased identities to individual cells using reference-based

and machine learning algorithms, respectively (Aran et al., 2019; Pliner et al., 2019). On the

other hand, tools such as 'Monocle3' and 'scVelo' align and project cells onto a pseudotemporal

space where each cell becomes a snapshot within the single-cell time continuum (Bergen et al.,

2019; Cao et al., 2019). In essence, the single scRNA-seq dataset is transformed into a time

series (Bergen et al., 2019; Cao et al., 2019; Manno et al., 2018). Although the pseudotemporal

scale does not reflect the actual time scale, it is a reliable approximation to characterize cell fate

and differentiation events, e.g., during organogenesis, disease states, or in response to SARS-

CoV-2 infection (Cao et al., 2019; Chua et al., 2020).

In this study, we present the application of an enhanced, scalable, and user-friendly scRNA-seq

analysis workflow on an existing human coronary artery scRNA-seq dataset. We performed

unbiased automatic cell identification at the single-cell level, pseudotemporal analysis, ligand-

receptor expression profiling, and drug repurposing analysis. Our results demonstrate potential

new mechanisms by which SMCs contribute to the atherosclerotic phenotype and signaling

within the lesion microenvironment. More importantly, we reveal attractive candidates for future

pre-clinical drug interventional studies. This reproducible analysis workflow can also be easily

modified and extended to incorporate different tissue data sources and single-cell modalities

such as snATAC-seq (Smith and Sheffield, 2020).

Results and Discussion

Unbiased automatic cell labeling reveals abundant cells with chondrocyte and fibroblast

characteristics.

Recently, automatic cell identifications tools have been introduced to compensate the

shortcomings of manual, cluster-based cell labeling (Aran et al., 2019). For example, 'SingleR'

and 'Garnett' uses reference-based and machine learning algorithms, respectively, to call

individual cell identities (Aran et al., 2019; Pliner et al., 2019). Using 'SingleR,' which uses

known purified cell expression data as references, we found that endothelial cells (ECs) make

up the highest proportion of cells in this dataset (16.21%, Fig. 1), followed by smooth muscle

cells (SMCs, 13.8%) and stem cells (SCs, 14.06%), where the latter could be so-called

"atherosclerotic stem cells" or normal stromal stem cells but cannot be distinguished until

specific expression profile references are developed in the future (Wang et al., 2020).

Consistent with recent scRNA-seq studies in atherosclerotic models, we identified abundant

fibroblast (FB) and chondrocyte-like (CH) cells, as well as cells with an osteoblast-like (OS)

expression profile (Fig. 1) (Pan et al., 2020). In the UMAP clusters reflecting single-cell

identities, there was a substantial presence of SMC and FB cells in the OS and SC cluster.

Such heterogeneity in cell clusters would have been overlooked in manual cluster-based cell

labeling.

Pseudotemporal ordering identifies distinct chondrocyte and fibroblast-like cell

differentiation states from smooth muscle cells.

To evaluate putative cell fate decisions or differentiation events (e.g., SMC phenotypic transition

states), we performed pseudotemporal analysis and ultra-fine clustering using 'Monocle3', a

method previously applied to normal and diseased states, e.g. embryo organogenesis and

response to COVID19 infection, respectively (Cao et al., 2019; Chua et al., 2020). We found

evidence of SMCs giving rise to CH and FB-like cells (Fig. 2A). This corroborates earlier

findings showing that SMCs may transition or de-differentiate into 'fibromyocytes'—SMCs that

have undergone a phenotypic modulation within the atherosclerotic lesions (Wang et al., 2020;

Wirka et al., 2019). Genes associated with healthy SMC phenotypes, such as *MYH11* (a

canonical marker of SMC), *IGFBP2* (associated with decreased visceral fat), and *PPP1R14A*

(which enhances smooth muscle contractions), are decreased by approximately 50-75% along

the SMC trajectory as these cells become more FB-like (Table 1, Fig. 2, $p < 0.1E{-}297$) (Bennett

et al., 2016; Carter et al., 2018). Similar results were found by another group using mouse

lineage-traced models where MYH11 expression was decreased in SMC-derived modulated

"intermediate cell states" (Pan et al., 2020). More importantly, specific inflammatory markers

and proteins associated with thrombotic events during CAD, including *C7* and *FBLN1*, are

increased along the same trajectory (Argraves et al., 2009; Carter, 2012)(Fig. 2B). Together,

and in corroboration of recent studies, our pseudotemporal analysis demonstrates that SMCs

could be a source of FB and CH-like cells, with the former associated with an intermediate

atherosclerotic cell phenotype, and the latter expresses genes associated with a more

advanced atherosclerotic phenotype (Pan et al., 2020; Wirka et al., 2019). This is further

supported by the recent study where blocking of SMC-derived intermediate cells coincides with

less severe atherosclerotic lesions (Pan et al., 2020). Precisely how these cells might influence

the overall stability of the atherosclerotic lesion and patient survival requires additional

longitudinal studies using genetic models (Bennett et al., 2016; Pan et al., 2020; Wang et al.,

2020).

Comprehensive ligand-receptor analysis shows complex intercellular communications in

the human coronary micro-environment and reveals potential drug targets.

To examine the potential cross-talk between different cell types using scRNA-seq data, we

compared the ligand and receptor expression profiles of each cell type with experimentally-

validated interactions using 'scTalk' (Farbehi et al., 2019). We found that there is an intricate

network of signaling pathways connecting different cell types; some cell types, such as OS,

have stronger and more frequent outgoing signals, whereas other cell types such as

Macrophages (Mø) have fewer and weaker incoming and outgoing signals (Fig. 3A). SMCs,

OSs, and neurons also exhibit a high degree of autocrine signaling profiles (Fig. 3A).

Specifically, SMCs are shown to have the highest number of outgoing signals and are among

those with the least number of incoming signal weights (Fig. 3B). This suggests that SMCs play

an important role in regulating the coronary microenvironment by transducing signals to

neighboring cells in the lesion.

Specifically, we identified three significant ligand-receptor interactions between SMCs and FBs,

FBLN1-ITGB1, APOD-LEPR, and DCN-EGFR (Fig. 4). We searched for potentially druggable

targets to interrupt SMC-FB communication by performing an integrative analysis of the

identified ligand-receptor interactions with known drug-gene interactions using the DGIdb 3.0

database (Cotto et al., 2017). Interestingly, anti-EGFR (epidermal growth factor receptor)-based

cancer treatments such as erlotinib, cetuximab, and gefitinib were shown as potential key

mediators of signaling pathways between SMC and FB via DCN (decorin) and EGFR (Fig. 4). It

has been shown that decorin overexpression increases SMC aggregation and SMC-induced

calcification at the atherosclerotic plaque (Fischer et al., 2004). Although the overlap between

CAD and cancer has been previously noted, the efficacy and mechanisms of chemotherapy,

such as erlotinib, in the pathogenesis and stability of CAD or their adjuvant use in cancer

patients to treat CAD continues to hold promise for future translational studies (Camaré et al.,

2017; DiRenzo et al., 2017; Tapia-Vieyra et al., 2017).

Conclusions

Our findings show that an enhanced, reproducible pipeline for scRNA-seq analysis improve on

current standard scRNA-seq bioinformatics protocols. We provide new insights into intricate

cellular differentiation and communication pathways while providing actionable and testable

targets for future studies (Fig. 5). In our combined analysis, we found that SMCs give rise to

substantial proportions of CH and FB, with the latter associated with worse prognostic markers

(Argraves et al., 2009; Carter, 2012; Carter et al., 2018; Pan et al., 2020). Several FDA-

approved drugs (e.g., erlotinib, cetuximab, and gefitinib) were shown as potential effectors of

SMCs' signaling to FB, and may be used to treat CAD in cancer patients to simplify or augment

drug regiments (Camaré et al., 2017). This is consistent with recent reports showing beneficial

effects of the acute promyelocytic drug all-trans-retinoic acid (ATRA) in atherosclerosis mouse

models (Pan et al., 2020).

Although the utilization of this workflow can compensate for many of the shortcomings of current

scRNA-seq analyses, we are still unable to perform cell-lineage tracing that reflects actual

timescale without additional molecular experiments. However, leveraging mitochondrial DNA

variants in snATAC-seq data has enabled lineage tracing analysis in human cells (Lareau et al.,

2020; Xu et al., 2019). Likewise, these analyses can ultimately be extended to integrate spatial

data and other multi-modal data (Stuart et al., 2019). In the future as spatial transcriptomics and

snATAC-seq data become more widely available, this workflow can be modified to discover

signaling pathways or differentiation events at specific tissue locations and time, allowing for

more disease-relevant drug-gene interaction analyses (Fig. 5). Nonetheless, this pipeline can be

applied immediately to datasets from other tissues/diseases to generate informative directions

for follow-up studies, and is more user-friendly and reproducible compared to standard scRNA

analyses.

Methods

Data retrieval and pre-processing

Human coronary artery scRNA data read count matrix was retrieved from the Gene Expression

Omnibus (GEO) using #GSE131780 and loaded into R 4.0, and was preprocessed using

standard parameters of the R packages 'Seurat' v.3, and 'Monocle3' as required (Satija et al.,

2015; Team, 2020; Trapnell et al., 2014; Wirka et al., 2019). Uniform manifold approximation

projections (UMAP) clusters from 'Seurat' were imported into 'Monocle3' before pseudotemporal

analysis.

Automatic cell Identification and pseudotemporal ordering

scRNA read matrices were read into SingleR as previously described for cell labeling (Aran et

al., 2019). Briefly, SingleR compares each cell's gene expression profile with known human

primary cell atlas data and gives the most likely cell identity independently. SingleR first corrects

for batch effects, then calculates the expression correlation scores for each test cell to each cell

type in the reference, and the cell identity is called based on reference cell type exhibiting the

highest correlation. Then, pseudotemporal analyses were performed as previously described in

the analysis of embryo organogenesis (Cao et al., 2019; Trapnell et al., 2014). Briefly, the

UMAP clusters were passed into Monocle3 and then the 'learn_graph()' and 'order_cells()'

functions. The SMCs and related clusters were then subsetted for detailed subclustering and

analysis. For each cluster, Moran's I statistics were calculated, which identify genes that are

differentially expressed along their trajectories. Detailed codes to reproduce the figures in this

publication can be found at the Miller Lab Github (see Availability of data and materials).

Ligand-receptor cell communication analysis

We analyzed candidate ligand-receptor interactions to infer cell communication using the R

package 'scTalk', as previously described in the analysis of glial cells (Farbehi et al., 2019).

Briefly, this method is based on permutation testing of random networks, where ligand-receptor

interactions are derived from experimentally derived interactions from the STRING database.

We exported statistically significant differentially expressed genes from 'Seurat' using the

'FindMarkers()' function and imported the preprocessed data into 'scTalk'. Then, overall edges

of the cellular communication network were calculated using the 'GenerateNetworkPaths()'

function, which reflects the overall ligand-receptor interaction strength between each cell type.

Then, the cell types of interest were specified and treeplots were generated using the

'NetworkTreePlot()' function.

Gene-drug interaction analysis

The above identified ligand and receptor interaction pairs were fed into the Drug-Gene

Interaction database (DGIdb 3.0) to reveal candidate drug-gene interactions (Cotto et al., 2017).

Briefly, ligands and receptors that were deemed significant from 'scTalk' were evaluated using

the 'queryDGIdb()' function of the 'rDGIdb' R package (Cotto et al., 2017). We included all top

FDA-approved drugs produced with verified inhibitory or antagonistic activities. Figure 4 and 5

were modified using BioRender for clarity.

Declarations

No conflicts of interest to disclose.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data are publicly available on GEO, accession number GSE131778, as previously described

(Wirka et al., 2019). All codes and analysis pipelines can be viewed at github.com/MillerLab-

CPHG/Ma_2020 and repurposed for any scRNA datasets.

Competing interests

The authors declare no competing interests.

Funding

Funding support was provided by grants from the National Institutes of Health (NIH):

R00HL125912 (CLM), R01HL148239 (CLM) and T32HL007284 (CJH and DW); American Heart

Association (AHA): POST35120545 (AWT) and Leducq Foundation Transatlantic Network of

Excellence (PlaqOmics) (CLM and AWT).

Authors' contributions

WFM designed and performed the statistical analysis. CJH, AWT, DW and YS refined the

methodology and edited the manuscript. CLM conceived and refined the project and edited the

manuscript.

Acknowledgements

Not applicable.

Works cited

Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, Chak S, Naikawadi RP, Wolters PJ, Abate AR,
Butte AJ, Bhattacharya M. 2019. Reference-based analysis of lung single-cell sequencing

reveals a transitional profibrotic macrophage. *Nat Immunol* **20**:163–172. doi:10.1038/s41590-
018-0276-y

Argraves WS, Tanaka A, Smith EP, Twal WO, Argraves KM, Fan D, Haudenschild CC. 2009.
Fibulin-1 and fibrinogen in human atherosclerotic lesions. *Histochem Cell Biol* **132**:559–565.
doi:10.1007/s00418-009-0628-7

Basatemur GL, Jørgensen HF, Clarke MCH, Bennett MR, Mallat Z. 2019. Vascular smooth
muscle cells in atherosclerosis. *Nat Rev Cardiol* **16**:727–744. doi:10.1038/s41569-019-0227-
9

Bennett MR, Sinha S, Owens GK. 2016. Vascular Smooth Muscle Cells in Atherosclerosis. *Circ*
Res **118**:692-- 702. doi:10.1161/circresaha.115.306361

Bergen V, Lange M, Peidli S, Wolf FA, Theis FJ. 2019. Generalizing RNA velocity to transient
cell states through dynamical modeling. *Biorxiv* 820936. doi:10.1101/820936

Camaré C, Pucelle M, Nègre-Salvayre A, Salvayre R. 2017. Angiogenesis in the atherosclerotic
plaque. *Redox Biol* **12**:18-- 34. doi:10.1016/j.redox.2017.01.007

Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S, Christiansen L,
Steemers FJ, Trapnell C, Shendure J. 2019. The single-cell transcriptional landscape of
mammalian organogenesis. *Nature* **566**:496–502. doi:10.1038/s41586-019-0969-x

Carter AM. 2012. Complement Activation: An Emerging Player in the Pathogenesis of
Cardiovascular Disease. *Sci* **2012**:1–14. doi:10.6064/2012/402783

Carter S, Lemieux I, Li Z, Alméras N, Tremblay A, Bergeron J, Poirier P, Després J-P, Picard F.
2018. Changes in IGFBP-2 levels following a one-year lifestyle modification program are
independently related to improvements in plasma apo B and LDL apo B levels.
Atherosclerosis **281**:89–97. doi:10.1016/j.atherosclerosis.2018.12.016

Chua RL, Lukassen S, Trump S, Hennig BP, Wendisch D, Pott F, Debnath O, Thürmann L,
Kurth F, Völker MT, Kazmierski J, Timmermann B, Twardziok S, Schneider S, Machleidt F,
Müller-Redetzky H, Maier M, Krannich A, Schmidt S, Balzer F, Liebig J, Loske J, Suttorp N,
Eils J, Ishaque N, Liebert UG, Kalle C von, Hocke A, Witzenrath M, Goffinet C, Drosten C,
Laudi S, Lehmann I, Conrad C, Sander L-E, Eils R. 2020. COVID-19 severity correlates with
airway epithelium–immune cell interactions identified by single-cell analysis. *Nat Biotechnol*
1–10. doi:10.1038/s41587-020-0602-4

Cotto KC, Wagner AH, Feng Y-Y, Kiwala S, Coffman AC, Spies G, Wollam A, Spies NC, Griffith
OL, Griffith M. 2017. DGIdb 3.0: a redesign and expansion of the drug–gene interaction
database. *Nucleic Acids Res* **46**:D1068–D1073. doi:10.1093/nar/gkx1143

DiRenzo D, Owens GK, Leeper NJ. 2017. "Attack of the Clones." *Circ Res* **120**:624--626.
doi:10.1161/circresaha.116.310091

Farbehi N, Patrick R, Dorison A, Xaymardan M, Janbandhu V, Wystub-Lis K, Ho JW, Nordon
RE, Harvey RP. 2019. Single-cell expression profiling reveals dynamic flux of cardiac
stromal, vascular and immune cells in health and injury. *Elife* **8**:e43882.
doi:10.7554/elife.43882

Fischer JW, Steitz SA, Johnson PY, Burke A, Kolodgie F, Virmani R, Giachelli C, Wight TN.
2004. Decorin Promotes Aortic Smooth Muscle Cell Calcification and Colocalizes to Calcified
Regions in Human Atherosclerotic Lesions. *Arteriosclerosis Thrombosis Vasc Biology*
24:2391–2396. doi:10.1161/01.atv.0000147029.63303.28

Lareau CA, Ludwig LS, Muus C, Gohil SH, Zhao T, Chiang Z, Pelka K, Verboon JM, Luo W,
Christian E, Rosebrock D, Getz G, Boland GM, Chen F, Buenrostro JD, Hacohen N, Wu CJ,
Aryee MJ, Regev A, Sankaran VG. 2020. Massively parallel single-cell mitochondrial DNA
genotyping and chromatin profiling. *Nat Biotechnol* 1–11. doi:10.1038/s41587-020-0645-6

Manno GL, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, Lidschreiber K, Kastriti
ME, Lönnerberg P, Furlan A, Fan J, Borm LE, Liu Z, Bruggen D van, Guo J, He X, Barker R,
Sundström E, Castelo-Branco G, Cramer P, Adameyko I, Linnarsson S, Kharchenko PV.
2018. RNA velocity of single cells. *Nature* **560**:494--498. doi:10.1038/s41586-018-0414-6

Pan H, Xue C, Auerbach BJ, Fan J, Bashore AC, Cui J, Yang DY, Trignano SB, Liu W, Shi J,
Ihuegbu CO, Bush EC, Worley J, Vlahos L, Laise P, Solomon RA, Connolly ES, Califano A,
Sims PA, Zhang H, Li M, Reilly MP. 2020. Single-Cell Genomics Reveals a Novel Cell State
During Smooth Muscle Cell Phenotypic Switching and Potential Therapeutic Targets for
Atherosclerosis in Mouse and Human. *Circulation*. doi:10.1161/circulationaha.120.048378

Pliner HA, Shendure J, Trapnell C. 2019. Supervised classification enables rapid annotation of
cell atlases. *Nat Methods* **16**:983–986. doi:10.1038/s41592-019-0535-3

Satija R, Farrell JA, Gennert D, Schier AF, Regev A. 2015. Spatial reconstruction of single-cell
gene expression data. *Nat Biotechnol* **33**:495-- 502. doi:10.1038/nbt.3192

Smith JP, Sheffield NC. 2020. Analytical Approaches for ATAC-seq Data Analysis. *Curr Protoc*
Hum Genetics **106**. doi:10.1002/cphg.101

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M,
Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. *Cell* **177**:1888-
1902.e21. doi:10.1016/j.cell.2019.05.031

Tapia-Vieyra JV, Delgado-Coello B, Mas-Oliva J. 2017. Atherosclerosis and Cancer; A
Resemblance with Far-reaching Implications. *Arch Med Res* **48**:12–26.
doi:10.1016/j.arcmed.2017.03.005

Team RC. 2020. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing.

Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ,
Mikkelsen TS, Rinn JL. 2014. The dynamics and regulators of cell fate decisions are

revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* **32**:381--386.
doi:10.1038/nbt.2859

Wang Y, Nanda V, Direnzo D, Ye J, Xiao S, Kojima Y, Howe KL, Jarr K-U, Flores AM, Tsantilas
P, Tsao N, Rao A, Newman AAC, Eberhard AV, Priest JR, Ruusalepp A, Pasterkamp G,
Maegdefessel L, Miller CL, Lind L, Koplev S, Björkegren JLM, Owens GK, Ingelsson E,
Weissman IL, Leeper NJ. 2020. Clonally expanding smooth muscle cells promote
atherosclerosis by escaping efferocytosis and activating the complement cascade. *P Natl*
Acad Sci Usa **117**:15818–15826. doi:10.1073/pnas.2006348117

Wirka RC, Wagh D, Paik DT, Pjanic M, Nguyen T, Miller CL, Kundu R, Nagao M, Coller J,
Koyano TK, Fong R, Woo YJ, Liu B, Montgomery SB, Wu JC, Zhu K, Chang R, Alamprese
M, Tallquist MD, Kim JB, Quertermous T. 2019. Atheroprotective roles of smooth muscle cell
phenotypic modulation and the TCF21 disease gene as revealed by single-cell analysis. *Nat*
Med **25**:1280--1289. doi:10.1038/s41591-019-0512-5

Xu J, Nuno K, Litzenburger UM, Qi Y, Corces MR, Majeti R, Chang HY. 2019. Single-cell
lineage tracing by endogenous mutations enriched in transposase accessible mitochondrial
DNA. *Elife* **8**:e45105. doi:10.7554/elife.45105

Figures and Figure Legends

Figure 1

Fig. 1. Unbiased automatic cell labeling of human coronary scRNA data using 'SingleR'

reveals abundant cells with chondrocyte (CH) and fibroblast (FB) gene expression

patterns. (A) UMAP clustering of 9798 cells derived from human coronary artery explants. **(B)**

Population breakdown by percentage. SMC: smooth muscle cells, EC: endothelial cells, CH:

chondrocytes, FB: fibroblasts, Mø: macrophages, SC: stem cells, CMP: common myeloid

progenitor cells, GMP: granulocyte-monocyte progenitor cells.

Figure 2

Fig. 2. Pseudotemporal and ultra-fine clustering reveals derivation of CH and FB-like

cells from SMCs. (A) Left, RNA trajectory (line) shows path two direct paths from the SMC

starting nodes (grey circles). **Middle,** ultra-fine clustering shows the logical transition stages

from SMC to CH and FB. **Right,** pseudotemporal analysis confirms that the cell and clusters

existing along a logical single-cell continuum. **(B)** Selected genes that were shown to vary over

pseudotime by Moran's I test were visualized.

Figure 3

Fig. 3. Comprehensive ligand-receptor analysis reveals intricate intracellular

communications in the coronary environment. (A) circle plot representation of the inferred

intercellular communications within the coronary artery environment. **(B)** Incoming and outgoing

signal summed weights by cell type.

Figure 4

Fig. 4. Fine ligand-receptor analysis coupled with automatic DGIdb analysis reveals

potential drug therapies could interrupt SMC to CH and FB communication. 'scTalk' shows

that SMCs interacts with FB through three pairs of experimentally verified interactions. Of these

three, *DCN-EGFR* signaling is the most druggable as revealed by DGIdb 3.0.

Figure 5

Fig. 5. Roadmap of enhanced scRNA-seq analysis process. Instead of cluster-based

grouping, our pipeline uses automatic cell labeling, coupled with pseudotime trajectory, cellular

network interaction and drug targeting, and provides a reproducible process for scRNA

datasets. From this roadmap, it is easy to add additional analysis tools as each cell has its own

identity and can be re-projected into different spaces such as t-SNE or 3D spaces.

Tables

Table 1.

	p_value	morans_test_statistic	morans_I
MYH11	<1E-297	169.0285919	0.66505236
IGFBP2	<1E-297	129.2299117	0.50844307
PPP1R14A	<1E-297	169.4172149	0.66663148
FBLN1	<1E-297	154.3399713	0.60730135
C7	<1E-297	156.2248703	0.61472417
LUM	<1E-297	170.4635813	0.67075391
RAMP1	<1E-297	161.486949	0.63531256
TCF21	<1E-297	75.75779399	0.29798686
SERPINF1	<1E-297	157.9946249	0.62168635
FN1	<1E-297	125.4354275	0.49354242
CNN1	<1E-297	170.9288702	0.67251583
TNFRSF11B	<1E-297	97.66515083	0.38400709

418 **Table 1. Selected Moran's I statistics for genes listed in Figure 2.** Moran's I statistics was

419 used to identify spatial correlations within the single-cell trajectory. +1 indicates that nearby cells

420 are perfectly similar, 0 indicates no similarity or pattern, and -1 indicates total dissimilarity (Cao

421 et al., 2019)