Single-cell RNA-seq analysis of human coronary arteries using an enhanced workflow

reveals SMC transitions and candidate drug targets

Wei Feng Ma[1,2], Chani J. Hodonsky[2], Adam W. Turner[2], Doris Wong[2,3], Yipei Song[2,4], Nelson B.

Barrientos[2,5], Jose Verdezoto Mosquera[2,3], Clint L. Miller[2,3,5]*

Affiliations

[1]Medical Scientist Training Program, University of Virginia, Charlottesville, VA 22908, USA.

[2]Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA.

[3]Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA

22908, USA.

[4]Department of Computer Engineering, University of Virginia, Charlottesville, VA 22908, USA.

[5]Department of Public Health Sciences, University of Virginia, Charlottesville, VA 22908, USA.

*corresponding author

Correspondence:

Clint L. Miller, PhD. (clintm@virginia.edu)

University of Virginia

Center for Public Health Genomics

PO Box 800717

Charlottesville, VA 22908 USA

Abstract

Background and Aims

The atherosclerotic plaque microenvironment is highly complex, and selective agents that

modulate plaque stability or other plaque phenotypes are not yet available. We sought to

investigate the human atherosclerotic cellular environment using scRNA-seq to uncover

potential therapeutic approaches. We aimed to make our workflow user-friendly, reproducible,

and applicable to other disease-specific scRNA-seq datasets.

Methods

Here we incorporate automated cell labeling, pseudotemporal ordering, ligand-receptor

evaluation, and drug-gene interaction analysis into an enhanced and reproducible scRNA-seq

analysis workflow. Notably, we also developed an R Shiny based interactive web application to

enable further exploration and analysis of the scRNA dataset.

Results

We applied this analysis workflow to a human coronary artery scRNA dataset and revealed

distinct derivations of chondrocyte-like and fibroblast-like cells from smooth muscle cells

(SMCs), and show the key changes in gene expression along their de-differentiation path. We

highlighted several key ligand-receptor interactions within the atherosclerotic environment

through functional expression profiling and revealed several attractive avenues for future

pharmacological repurposing in precision medicine. Further, our interactive web application,

PlaqView (www.plaqview.com), allows other researchers to easily explore this dataset and

benchmark applicable scRNA-seq analysis tools without prior coding knowledge.

Conclusions

These results suggest novel effects of chemotherapeutics on the atherosclerotic cellular

environment and provide future avenues of studies in precision medicine. This publicly available

workflow will also allow for more systematic and user-friendly analysis of scRNA datasets in

other disease and developmental systems. *PlaqView* allows for rapid visualization and analysis

of atherosclerosis scRNA-seq datasets without the need of prior coding experience. Future

releases of *PlaqView* will feature additional larger scRNA-seq and scATAC-seq atherosclerosis-

related datasets, thus providing a critical resource for the field by promoting data harmonization

and biological interpretation.

Keywords

Atherosclerosis, scRNA, analysis, pseudotemporal, drug-target, pipeline, web application

Abbreviations

C7- complement component C7

CAD- coronary artery disease

CH- chondrocytes

CMP- common myeloid progenitor cells

DCN- decorin

DGIdb- drug-gene interaction database

EC- endothelial cells

EGFR- epidermal growth factor receptor

FB- fibroblasts

FBLN1- fibulin 1

GMP- granulocyte-monocyte progenitor cells.

Mø- macrophages

MYH11- myosin heavy chain 11

SC- stem cells

sc/snATAC-seq - single cell/single nucleus assay for transposase-accessible chromatin

sequencing

sc/snRNA-seq- single cell/single nucleus RNA sequencing

TI- trajectory inference

SMC- smooth muscle cells

UMAP- uniform manifold approximation and projection

Background

Atherosclerosis is a complex process involving chronic inflammation and hardening of the

vessel wall and represents one of the major causes of coronary artery disease (CAD),

peripheral artery disease, and stroke [1]. Rupture of an unstable atherosclerotic lesion can lead

to the formation of a thrombus, causing complete or partial occlusion of a coronary artery [2].

The contribution of smooth muscle cells (SMCs) to both lesion stability and progression has

recently been established by numerous groups. However, the exact mechanisms by which

SMCs modulate the atherosclerotic microenvironment and whether pharmacological agents can

be used to selectively counter SMC-related deleterious mechanisms are still under investigation

[3–5]

Recent advances in single-cell RNA-sequencing (scRNA-seq) have enabled ultra-fine gene

expression profiling of many diseases at the cellular level, including atherosclerotic coronary

artery disease [5]. As sequencing costs continue to decline, there has also been a consistent

growth in scRNA datasets, data analysis tools and applications [6]. Currently, a major challenge

with scRNA-seq analysis is the inherent bias introduced during manual cell labeling, in which

cells are grouped by clusters and their identities called collectively based on their overall

differential gene expression profiles [7]. Another draw-back inherent to commonly used scRNA-

seq protocols is their destructive nature to the cells, making time-series analyses of the same

cells impossible. Instead, these studies must rely on time-points from separate libraries to

monitor processes such as clonal expansion and cell differentiation [8,9].

Recently, new approaches have been developed to compensate for both of these shortcomings,

namely automatic cell labeling, pseudotemporal analysis, and trajectory inference. Tools such

as 'SingleR', 'scCATCH', and 'Garnett' have been used to assign unbiased identities to

individual cells using reference-based and machine learning algorithms, respectively [7,10,11].

Moreover, tools such as 'Monocle3' and 'scVelo' align and project cells onto a pseudotemporal

space where each cell becomes a snapshot within the single-cell time continuum [12,13]. In

essence, the single scRNA-seq dataset is transformed into a time series [12–14]. Although the

pseudotemporal scale does not reflect the actual time scale, it is a reliable approximation to

characterize cell fate and differentiation events, e.g., during organogenesis, disease states, or in

response to SARS-CoV-2 infections [13,15]

In this study, we present the application of an enhanced, scalable, and user-friendly scRNA-seq

analysis workflow on an existing human coronary artery scRNA-seq dataset. We performed

unbiased automatic cell identification at the single-cell level, pseudotemporal analysis, ligand-

receptor expression profiling, and drug repurposing analysis. Our results demonstrate potential

new mechanisms by which SMCs contribute to the atherosclerotic phenotype and signaling

within the lesion microenvironment. More importantly, we revealed attractive candidate avenues

for future pharmacological interventional studies. We also developed an interactive web

application to allow other users to explore this dataset. This reproducible analysis pipeline and

application can also be easily modified to incorporate different tissue data sources and single-

cell modalities such as scATAC-seq [16] or CITE-seq [17], and could serve as a template to

analyze and visualize single-cell datasets in other disease models.

Results and Discussion

Unbiased automatic cell labeling reveals abundant cells with chondrocyte and fibroblast

characteristics. Recently, automatic cell identification tools have been introduced to

compensate for the shortcomings of manual, cluster-based cell labeling [7]. For example,

'SingleR' and 'Garnett' use reference-based and machine learning algorithms, respectively, to

call individual cell identities [7,10,18]. Using 'SingleR,' which uses known purified cell

expression data as references, we found that endothelial cells (ECs) make up the highest

proportion of cells in this dataset (16.21%, Figure 1A-B), followed by smooth muscle cells

(SMCs, 13.8%) and stem cells (SCs, 14.06%), where the latter could be so-called

"atherosclerotic stem cells" or normal stromal stem cells but cannot be distinguished until

specific expression profile references are developed in the future [8]. Consistent with recent

scRNA-seq studies in atherosclerotic models, we identified abundant fibroblast (FB) and

chondrocyte-like (CH) cells, as well as cells with an osteoblast-like (OS) expression profile

(Figure 1B) [4]. In the UMAP clusters reflecting single-cell identities, there was a substantial

presence of SMC and FB cells in the OS and SC cluster. Such heterogeneity in cell clusters

would have been overlooked in manual cluster-based cell labeling. We also applied another

reference-based cell calling tool 'scCATCH,' and found that it underperforms relative to

'SingleR' and fails to provide consistent cell type assignment when provided with similar tissue

priors (Supplementary Table 1).

Pseudotemporal ordering identifies distinct chondrocyte and fibroblast-like cell

differentiation states from smooth muscle cells. To evaluate putative cell fate decisions or

differentiation events (e.g., SMC phenotypic transition states), we performed pseudotemporal

analysis and ultra-fine clustering using 'Monocle3', a method previously applied to normal and

diseased states, e.g., embryo organogenesis and response to coronavirus infection,

respectively [13,15]. We compared over 60 other trajectory inference (TI) methods including

Slingshot [19], PAGA [20], and SCORPIUS using the 'Dynverse' package [21]. However, we

found that most TI algorithms were unsuitable for complex tissue environments such as the

atherosclerotic plaque due to their inability to distinguish disconnected topologies

(Supplementary Table 2). In particular, we found evidence of SMCs giving rise to both

chondrocyte (CH) and fibroblast (FB)-like cells (Figure 1C). This corroborates earlier findings

showing that SMCs may transition or de-differentiate into 'fibromyocytes'—SMCs that have

undergone a phenotypic modulation to an extracellular matrix producing cell type within the

atherosclerotic lesions [5,8]. Genes associated with healthy SMC phenotypes, such as *MYH11*

(a canonical marker of SMC), *IGFBP2* (associated with decreased visceral fat), and *PPP1R14A*

(which enhances smooth muscle contraction), are decreased by approximately 50-75% along

the SMC trajectory as these cells become more FB-like (Table 1, Figure 1D, p < 0.1E-297)

[3,22]. Similar results were found by another group using mouse lineage-traced models where

MYH11 expression was decreased in SMC-derived modulated "intermediate cell states" [4].

More importantly, specific inflammatory markers and proteins associated with thrombotic events

during CAD, including complement proteins *C7* and *C3, FBLN1*, and *CXCL12* are increased

along the same trajectory (Figure 1D) [2,23]. Recent evidence suggests that CAD-associated

CXCL12 secreted from endothelial cells may promote atherosclerosis[24]. Our results point to a

potentially new source of *CXCL12* that could be targeted to inhibit SMC-to-FB dedifferentiation.

Together, and in corroboration of recent studies, our pseudotemporal analysis demonstrates

that SMCs could be a source of both FB and CH-like cells, associated with intermediate and

advanced atherosclerotic phenotypes, respectively [4,5]. This is further supported by a recent

study, in which blocking of SMC-derived intermediate cells coincides with less severe

atherosclerotic lesions [4]. Precisely how these cells might influence the overall stability of

atherosclerotic lesions and clinical outcomes requires additional longitudinal studies using

genetic models and deep phenotyping of human tissues [3,4,8].

Comprehensive ligand-receptor analysis shows complex intercellular communications in

the human coronary micro-environment and reveals potential drug targets. To examine

the potential cross-talk between different cell types using scRNA-seq data, we compared the

ligand and receptor expression profiles of each cell type with experimentally-validated

interactions using 'scTalk' [25]. We found that there is an intricate network of signaling

pathways connecting different cell types; some cell types, such as OS, have stronger and more

frequent outgoing signals, whereas other cell types such as Macrophages (Mø) have fewer and

weaker incoming and outgoing signals (Figure 2A). SMCs, OSs, and neurons also exhibit a high

degree of autocrine signaling profiles (Figure 2A). Specifically, SMCs are shown to have the

highest number of outgoing signals and are among those with the least number of incoming

signal weights (Figure 2B). This suggests that SMCs play an important role in regulating the

coronary microenvironment by transducing signals to neighboring cells in the lesion.

Specifically, ten signaling pathways were identified between SMC, FB, CH, and OS

(Supplemental Figure 1). These pathways involve C3 complement, fibulin-1 *(FBLN1)*,

apolipoprotein D *(APOD)*, decorin *(DCN)*, and matrix metalloprotease 2 (*MMP2*, Figure 2C,

Supplemental Figure 1). We searched for potentially druggable targets to interrupt SMC-FB

communication by performing an integrative analysis of the identified ligand-receptor

interactions and the druggable genome using several drug-gene interaction databases,

including DGIdb 3.0 and Pharos [26]. We found that SMCs signals to FBs through the

complement protein *C3* and syndecan-2 *(SDC2)* via *MMP2,* and these two pathways can be

disrupted by drugs such as compstatin and tanomastat, respectively (Figure 2C). Multiple

studies have linked *C3* and the complement system to atherosclerotic lesion maturation in

mouse models, and a recent case study showed that the C3 targeted inhibitor, compstatin

(AMY-101), may prevent cardiovascular complications in patients with severe COVID-19

pneumonia [27–30]. Our results provide a potential mechanistic explanation by which SMCs can

modulate the inflammatory environment and plaque formation. Further, a recent study

demonstrated that *microRNA-9* repression of *SDC2* impedes atherosclerosis formation [31],

while MMP2 alteration also contributes to atherosclerosis in mouse models [32]. Here, we show

that SMCs signal to FBs within the atherosclerotic environment via *SDC2-MMP2*, and reveal

additional upstream candidate drug therapies that may influence atherosclerosis progression.

Interestingly, anti-*EGFR* (epidermal growth factor receptor)-based cancer treatments such as

erlotinib, cetuximab, and gefitinib were identified as potential key mediators of signaling

pathways between SMCs and FBs via decorin (*DCN*) and *EGFR* (Figure 2C). It has been shown

that *DCN* overexpression increases SMC aggregation and SMC-induced calcification within

atherosclerotic plaque [33]. Although the overlap between CAD and cancer etiology has been

previously noted, the long-term efficacy and cardiovascular impact of chemotherapy drugs, such

as erlotinib, requires further translational studies to investigate their potential use in cancer

patients to treat CAD [34–36].

Furthermore, integrative analysis of gene expression variation along the SMC-to-FB RNA

trajectory revealed similar results; as shown earlier in Figure 1D, the expression of complement

genes such as *C3* and *C7,* and chemokine *CXCL12* are increased as SMCs become more FB-

like. Although CXCL12 derived from endothelial cells have been recognized to promote CAD in

mouse models [24], here we provide a potentially new source of CXCL12 using human data and

found several pharmacological agents such as tinzaparin, an FDA-approved anticoagulant, to

investigate in future interventional studies. Together, this combination of independent analyses

of a scRNA-seq dataset reaffirms the druggable potential of these target genes.

PlaqView is a user-friendly web application to explore atherosclerosis-related datasets.

To enable other researchers to explore the transcriptomic landscape of the atherosclerotic

environment, we developed a web interface called _PlaqView_ (www.plaqview.com, Figure 3A).

This interactive, R Shiny-based tool allows for multiple gene queries and comparisons of gene

expression, cell-labeling methods, RNA-trajectory tools, integrative drug-gene analysis, and

outputs high quality graphs and detailed tables. To our knowledge, there are no publicly

available tools to visualize atherosclerosis-related single-cell datasets without prior coding

knowledge. Further, _PlaqView_ is under active development and will be releasing new datasets

coincidently with future atherosclerosis-related publications. As the database in _PlaqView_

expands along with the growing number of single-cell datasets, we anticipate that it will become

an essential tool for the atherosclerosis research field.

Limitations

Despite the advances presented in this workflow, we are still working to improve several

limitations. For instance, the default reference in 'SingleR' cannot identify more recently

discovered cellular phenotypes such as fibromyocytes [5,7,37], which may require a

combination of manual and automated labeling methods for accurate identification. Factors such

as the intrinsic heterogeneity of the tissue sample, disease stage, and tissue processing

artifacts are difficult to isolate computationally and could influence automated labeling methods.

Nonetheless, interactive viewers and tools such as _PlaqView_ which incorporate multiple

methods and visualizations in one location, could help users separate out the technical and

biological variation from various single-cell datasets. Additionally, the modular nature of

PlaqView will also allow for future improvement of labeling methods as more precise reference

datasets are made available. Lastly, the true efficacy of proposed drugs cannot be verified

without extensive pre-clinical testing and clinical trials. Nonetheless, these findings may catalyze

future investigative efforts to develop more targeted therapies.

Conclusions

Our findings show that an enhanced, reproducible pipeline for scRNA-seq analysis has the

potential to improve upon current standard scRNA-seq bioinformatics protocols. For instance,

we provide new insights into intricate vascular cell differentiation and communication pathways

while providing actionable and testable targets for future experimental studies (Figure 3B). In

our combined analysis, we found that SMCs give rise to substantial proportions of CH and FB,

with the latter associated with worse prognostic markers [2,4,22,23]. SMCs signal to FBs via a

series of pathways involving C3 complement and MMP2, whose expression coincidentally

increases along the SMC-to-FB trajectory. We revealed possible therapeutic avenues that may

disrupt these cell communication pathways and alter the atherosclerotic pathology.

Furthermore, several FDA-approved drugs (e.g., erlotinib, cetuximab, and gefitinib) were shown

as potential effectors of SMC signaling to FB, and may be used to treat CAD in cancer patients

to simplify or augment drug regiments [34]. This is consistent with recent reports showing

beneficial effects of the acute promyelocytic drug all-trans-retinoic acid (ATRA) in

atherosclerosis mouse models [4]. Further investment in scRNA-seq may also help resolve the

balance of anti-tumor efficacy and atheroprotection for immune checkpoint inhibitors as well as

immunomodulators at the interface of cardio-immuno-oncology [38].

Although the utilization of this workflow can compensate for many of the shortcomings of current

scRNA-seq analyses, we are still unable to perform cell-lineage tracing that reflects actual

timescales without additional gene engineering experiments *in vivo* [39]. However, leveraging

mitochondrial DNA variants in snATAC-seq data has enabled lineage tracing analysis in human

cells [40,41]. Likewise, these analyses can ultimately be extended to integrate spatial omics and

other multi-modal data [42]. In the future as spatial transcriptomics, scATAC-seq, and/or CITE-

seq data become more widely available, this workflow can be modified to discover signaling

pathways or differentiation events at specific tissue locations and timepoints, allowing for more

disease-relevant drug-gene interaction analyses (Figure 3B). Nonetheless, this pipeline can be

applied immediately to datasets from other tissues/diseases to generate informative directions

for follow-up studies, and is more user-friendly and reproducible compared to standard scRNA

analyses. Lastly, building web applications that democratize the access and analysis of single-

cell data will promote collaboration and innovation across disciplines [43]. As *PlaqView*

incorporates additional relevant single-cell datasets, we anticipate that this application will

become an indispensable resource for the community.

Methods

Data retrieval and pre-processing

Human coronary artery scRNA data read count matrix was retrieved from the Gene Expression

Omnibus (GEO) using #GSE131780 and loaded into R 4.0, and was preprocessed using

standard parameters of the R packages 'Seurat' v.3, and 'Monocle3' as required [5,44–46].

Uniform manifold approximation projections (UMAP) clusters from 'Seurat' were imported into

'Monocle3' before pseudotemporal analysis.

Automatic cell Identification and pseudotemporal ordering

scRNA read matrices were read into SingleR as previously described for cell labeling [7].

SingleR compares each cell's gene expression profile with known human primary cell atlas data

and gives the most likely cell identity independently. SingleR first corrects for batch effects, then

calculates the expression correlation scores for each test cell to each cell type in the reference,

and the cell identity is called based on reference cell type exhibiting the highest correlation.

Then, pseudotemporal analyses were performed as previously described in the analysis of

embryo organogenesis [13,45]. Briefly, the UMAP clusters were passed into Monocle3 and then

the 'learn_graph()' and 'order_cells()' functions. The SMCs and related clusters were then

subsetted for detailed subclustering and analysis. For each cluster, Moran's I statistics were

calculated, which identify genes that are differentially expressed along their trajectories.

Additionally, we explored 60+ other trajectory inference methods using the 'Dynverse' package,

where we simulated mock dataset of 5000 cells with known disconnected graph topologies [21].

Detailed codes to reproduce the figures in this publication can be found at the Miller Lab Github

(see availability of data and materials).

Ligand-receptor cell communication analysis

We analyzed candidate ligand-receptor interactions to infer cell communication using the R

package 'scTalk', as previously described in the analysis of glial cells [25]. This method is

based on permutation testing of random networks, where ligand-receptor interactions are

derived from experimentally derived interactions from the STRING database. We exported

statistically significant differentially expressed genes from 'Seurat' using the 'FindMarkers()'

function and imported the preprocessed data into 'scTalk.' Then, overall edges of the cellular

communication network were calculated using the 'GenerateNetworkPaths()' function, which

reflects the overall ligand-receptor interaction strength between each cell type. Then, the cell

types of interest were specified and treeplots were generated using the 'NetworkTreePlot()'

function.

Gene-drug interaction analysis

The above identified ligand and receptor interaction pairs were fed into the Drug-Gene

Interaction database (DGIdb 3.0) to reveal candidate drug-gene interactions [26]. Ligands and

receptors that were deemed significant from 'scTalk' were evaluated using the 'queryDGIdb()'

function of the 'rDGIdb' R package [26]. Additionally, we queried CTD2, OMIM, ClinVar, Pharos,

GnomAD, and the ExAC databases using the docker-based tool 'DrugThatGene' [47]. We

included all top FDA-approved drugs produced with verified inhibitory or antagonistic activities,

as well as drugs that may influence or influenced by changes in the receptor. Figures 2C, 2D,

and 3B were modified using BioRender for clarity.

Development of *PlaqView*

PlaqView is written in R and Shiny, and is hosted on the web at www.plaqview.com using

dedicated shinyapps.io servers, and can be run locally through RStudio. The raw data was first

processed as previously described, but packaged into .rds objects. These objects were then

written into a Shiny script to allow for interactive display. This application is open-sourced and

its code and data is available at github.com/MillerLab-CPHG/PlaqView. Further, *PlaqView* is

actively recruiting available datasets on the github page.

Declarations

No conflicts of interest to disclose.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data are publicly available on GEO, accession number GSE131778, as previously described

[5]. All codes and analysis pipelines can be viewed at github.com/MillerLab-CPHG/Ma_2020

and repurposed for other scRNA datasets.

Competing interests

The authors declare no competing interests.

Funding

Funding support was provided by grants from the National Institutes of Health (NIH):

R00HL125912 (CLM), R01HL148239 (CLM) and T32HL007284 (CJH and DW); American Heart

Association (AHA): POST35120545 (AWT) and Leducq Foundation Transatlantic Network of

Excellence (PlaqOmics) (CLM and AWT).

Authors' contributions

WFM designed and performed the statistical analysis. CJH, AWT, DW and YS refined the

methodology and edited the manuscript. JVM helped with scripting. CLM conceived and refined

the project and edited the manuscript.

Acknowledgements

Works cited

[1] G.L. Basatemur, H.F. Jørgensen, M.C.H. Clarke, M.R. Bennett, Z. Mallat, Vascular smooth
muscle cells in atherosclerosis, Nat Rev Cardiol. 16 (2019) 727–744.
https://doi.org/10.1038/s41569-019-0227-9.

[2] W.S. Argraves, A. Tanaka, E.P. Smith, W.O. Twal, K.M. Argraves, D. Fan, C.C.
Haudenschild, Fibulin-1 and fibrinogen in human atherosclerotic lesions, Histochem Cell Biol.
132 (2009) 559–565. https://doi.org/10.1007/s00418-009-0628-7.

[3] M.R. Bennett, S. Sinha, G.K. Owens, Vascular Smooth Muscle Cells in Atherosclerosis., Circ
Res. 118 (2016) 692-- 702. https://doi.org/10.1161/circresaha.115.306361.

[4] H. Pan, C. Xue, B.J. Auerbach, J. Fan, A.C. Bashore, J. Cui, D.Y. Yang, S.B. Trignano, W.
Liu, J. Shi, C.O. Ihuegbu, E.C. Bush, J. Worley, L. Vlahos, P. Laise, R.A. Solomon, E.S.
Connolly, A. Califano, P.A. Sims, H. Zhang, M. Li, M.P. Reilly, Single-Cell Genomics Reveals a
Novel Cell State During Smooth Muscle Cell Phenotypic Switching and Potential Therapeutic
Targets for Atherosclerosis in Mouse and Human, Circulation. (2020).
https://doi.org/10.1161/circulationaha.120.048378.

[5] R.C. Wirka, D. Wagh, D.T. Paik, M. Pjanic, T. Nguyen, C.L. Miller, R. Kundu, M. Nagao, J.
Coller, T.K. Koyano, R. Fong, Y.J. Woo, B. Liu, S.B. Montgomery, J.C. Wu, K. Zhu, R. Chang,
M. Alamprese, M.D. Tallquist, J.B. Kim, T. Quertermous, Atheroprotective roles of smooth
muscle cell phenotypic modulation and the TCF21 disease gene as revealed by single-cell
analysis, Nat Med. 25 (2019) 1280--1289. https://doi.org/10.1038/s41591-019-0512-5.

[6] C. Yifan, Y. Fan, P. Jun, Visualization of cardiovascular development, physiology and
disease at the single-cell level: Opportunities and future challenges, J Mol Cell Cardiol. 142
(2020) 80–92. https://doi.org/10.1016/j.yjmcc.2020.03.005.

[7] D. Aran, A.P. Looney, L. Liu, E. Wu, V. Fong, A. Hsu, S. Chak, R.P. Naikawadi, P.J. Wolters,
A.R. Abate, A.J. Butte, M. Bhattacharya, Reference-based analysis of lung single-cell

sequencing reveals a transitional profibrotic macrophage, Nat Immunol. 20 (2019) 163–172.
https://doi.org/10.1038/s41590-018-0276-y.

[8] Y. Wang, V. Nanda, D. Direnzo, J. Ye, S. Xiao, Y. Kojima, K.L. Howe, K.-U. Jarr, A.M.
Flores, P. Tsantilas, N. Tsao, A. Rao, A.A.C. Newman, A.V. Eberhard, J.R. Priest, A.
Ruusalepp, G. Pasterkamp, L. Maegdefessel, C.L. Miller, L. Lind, S. Koplev, J.L.M. Björkegren,
G.K. Owens, E. Ingelsson, I.L. Weissman, N.J. Leeper, Clonally expanding smooth muscle cells
promote atherosclerosis by escaping efferocytosis and activating the complement cascade., P
Natl Acad Sci Usa. 117 (2020) 15818–15826. https://doi.org/10.1073/pnas.2006348117.

[9] D. DiRenzo, G.K. Owens, N.J. Leeper, "Attack of the Clones," Circ Res. 120 (2017) 624--
626. https://doi.org/10.1161/circresaha.116.310091.

[10] H.A. Pliner, J. Shendure, C. Trapnell, Supervised classification enables rapid annotation of
cell atlases, Nat Methods. 16 (2019) 983–986. https://doi.org/10.1038/s41592-019-0535-3.

[11] X. Shao, J. Liao, X. Lu, R. Xue, N. Ai, X. Fan, scCATCH: automatic annotation on cell-types
of clusters from single-cell RNA-seq data, Iscience. 23 (2020) 100882.
https://doi.org/10.1016/j.isci.2020.100882.

[12] V. Bergen, M. Lange, S. Peidli, F.A. Wolf, F.J. Theis, Generalizing RNA velocity to transient
cell states through dynamical modeling, Biorxiv. (2019) 820936. https://doi.org/10.1101/820936.

[13] J. Cao, M. Spielmann, X. Qiu, X. Huang, D.M. Ibrahim, A.J. Hill, F. Zhang, S. Mundlos, L.
Christiansen, F.J. Steemers, C. Trapnell, J. Shendure, The single-cell transcriptional landscape
of mammalian organogenesis, Nature. 566 (2019) 496–502. https://doi.org/10.1038/s41586-
019-0969-x.

[14] G.L. Manno, R. Soldatov, A. Zeisel, E. Braun, H. Hochgerner, V. Petukhov, K. Lidschreiber,
M.E. Kastriti, P. Lönnerberg, A. Furlan, J. Fan, L.E. Borm, Z. Liu, D. van Bruggen, J. Guo, X.
He, R. Barker, E. Sundström, G. Castelo-Branco, P. Cramer, I. Adameyko, S. Linnarsson, P.V.
Kharchenko, RNA velocity of single cells, Nature. 560 (2018) 494--498.
https://doi.org/10.1038/s41586-018-0414-6.

[15] R.L. Chua, S. Lukassen, S. Trump, B.P. Hennig, D. Wendisch, F. Pott, O. Debnath, L.
Thürmann, F. Kurth, M.T. Völker, J. Kazmierski, B. Timmermann, S. Twardziok, S. Schneider,
F. Machleidt, H. Müller-Redetzky, M. Maier, A. Krannich, S. Schmidt, F. Balzer, J. Liebig, J.
Loske, N. Suttorp, J. Eils, N. Ishaque, U.G. Liebert, C. von Kalle, A. Hocke, M. Witzenrath, C.
Goffinet, C. Drosten, S. Laudi, I. Lehmann, C. Conrad, L.-E. Sander, R. Eils, COVID-19 severity
correlates with airway epithelium–immune cell interactions identified by single-cell analysis, Nat
Biotechnol. (2020) 1–10. https://doi.org/10.1038/s41587-020-0602-4.

[16] J.P. Smith, N.C. Sheffield, Analytical Approaches for ATAC-seq Data Analysis, Curr Protoc
Hum Genetics. 106 (2020). https://doi.org/10.1002/cphg.101.

[17] M. Stoeckius, C. Hafemeister, W. Stephenson, B. Houck-Loomis, P.K. Chattopadhyay, H.
Swerdlow, R. Satija, P. Smibert, Simultaneous epitope and transcriptome measurement in
single cells, Nat Methods. 14 (2017) 865–868. https://doi.org/10.1038/nmeth.4380.

[18] M. Slyper, C.B.M. Porter, O. Ashenberg, J. Waldman, E. Drokhlyansky, I. Wakiro, C.
Smillie, G. Smith-Rosario, J. Wu, D. Dionne, S. Vigneau, J. Jané-Valbuena, T.L. Tickle, S.
Napolitano, M.-J. Su, A.G. Patel, A. Karlstrom, S. Gritsch, M. Nomura, A. Waghray, S.H. Gohil,
A.M. Tsankov, L. Jerby-Arnon, O. Cohen, J. Klughammer, Y. Rosen, J. Gould, L. Nguyen, M.
Hofree, P.J. Tramontozzi, B. Li, C.J. Wu, B. Izar, R. Haq, F.S. Hodi, C.H. Yoon, A.N. Hata, S.J.
Baker, M.L. Suvà, R. Bueno, E.H. Stover, M.R. Clay, M.A. Dyer, N.B. Collins, U.A. Matulonis, N.
Wagle, B.E. Johnson, A. Rotem, O. Rozenblatt-Rosen, A. Regev, A single-cell and single-
nucleus RNA-Seq toolbox for fresh and frozen human tumors, Nat Med. 26 (2020) 792--802.
https://doi.org/10.1038/s41591-020-0844-1.

[19] K. Street, D. Risso, R.B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, S. Dudoit,
Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics, Bmc Genomics.
19 (2018) 477. https://doi.org/10.1186/s12864-018-4772-0.

[20] F.A. Wolf, F.K. Hamey, M. Plass, J. Solana, J.S. Dahlin, B. Göttgens, N. Rajewsky, L.
Simon, F.J. Theis, PAGA: graph abstraction reconciles clustering with trajectory inference
through a topology preserving map of single cells, Genome Biol. 20 (2019) 59.
https://doi.org/10.1186/s13059-019-1663-x.

[21] W. Saelens, R. Cannoodt, H. Todorov, Y. Saeys, A comparison of single-cell trajectory
inference methods, Nat Biotechnol. 37 (2019) 547–554. https://doi.org/10.1038/s41587-019-
0071-9.

[22] S. Carter, I. Lemieux, Z. Li, N. Alméras, A. Tremblay, J. Bergeron, P. Poirier, J.-P. Després,
F. Picard, Changes in IGFBP-2 levels following a one-year lifestyle modification program are
independently related to improvements in plasma apo B and LDL apo B levels, Atherosclerosis.
281 (2018) 89–97. https://doi.org/10.1016/j.atherosclerosis.2018.12.016.

[23] A.M. Carter, Complement Activation: An Emerging Player in the Pathogenesis of
Cardiovascular Disease, Sci. 2012 (2012) 1–14. https://doi.org/10.6064/2012/402783.

[24] Y. Döring, E.P.C. van der Vorst, J. Duchene, Y. Jansen, S. Gencer, K. Bidzhekov, D. Atzler,
D. Santovito, D.J. Rader, D. Saleheen, C. Weber, CXCL12 Derived From Endothelial Cells
Promotes Atherosclerosis to Drive Coronary Artery Disease, Circulation. 139 (2019) 1338–
1340. https://doi.org/10.1161/circulationaha.118.037953.

[25] N. Farbehi, R. Patrick, A. Dorison, M. Xaymardan, V. Janbandhu, K. Wystub-Lis, J.W. Ho,
R.E. Nordon, R.P. Harvey, Single-cell expression profiling reveals dynamic flux of cardiac
stromal, vascular and immune cells in health and injury, Elife. 8 (2019) e43882.
https://doi.org/10.7554/elife.43882.

[26] K.C. Cotto, A.H. Wagner, Y.-Y. Feng, S. Kiwala, A.C. Coffman, G. Spies, A. Wollam, N.C.
Spies, O.L. Griffith, M. Griffith, DGIdb 3.0: a redesign and expansion of the drug–gene
interaction database, Nucleic Acids Res. 46 (2017) D1068–D1073.
https://doi.org/10.1093/nar/gkx1143.

[27] C. Buono, C.E. Come, J.L. Witztum, G.F. Maguire, P.W. Connelly, M. Carroll, A.H.
Lichtman, Influence of C3 Deficiency on Atherosclerosis, Circulation. 105 (2002) 3025–3031.
https://doi.org/10.1161/01.cir.0000019584.04929.83.

[28] S. Mastaglio, A. Ruggeri, A.M. Risitano, P. Angelillo, D. Yancopoulou, D.C. Mastellos, M.
Huber-Lang, S. Piemontese, A. Assanelli, C. Garlanda, J.D. Lambris, F. Ciceri, The first case of
COVID-19 treated with the complement C3 inhibitor AMY-101, Clin Immunol. 215 (2020)
108450. https://doi.org/10.1016/j.clim.2020.108450.

[29] A. Java, A.J. Apicelli, M.K. Liszewski, A. Coler-Reilly, J.P. Atkinson, A.H.J. Kim, H.S.
Kulkarni, The complement system in COVID-19: friend and foe?, Jci Insight. 5 (2020).
https://doi.org/10.1172/jci.insight.140711.

[30] D. Pellegrini, R. Kawakami, G. Guagliumi, A. Sakamoto, K. Kawai, A. Gianatti, A. Nasr, R.
Kutys, L. Guo, A. Cornelissen, L. Faggi, M. Mori, Y. Sato, I. Pescetelli, M. Brivio, M. Romero, R.
Virmani, A.V. Finn, Microthrombi As A Major Cause of Cardiac Injury in COVID-19: A Pathologic
Study, Circulation. (2021). https://doi.org/10.1161/circulationaha.120.051828.

[31] R. Zhang, B. Song, X. Hong, Z. Shen, L. Sui, S. Wang, microRNA-9 Inhibits Vulnerable
Plaque Formation and Vascular Remodeling via Suppression of the SDC2-Dependent FAK/ERK
Signaling Pathway in Mice With Atherosclerosis, Front Physiol. 11 (2020) 804.
https://doi.org/10.3389/fphys.2020.00804.

[32] M. Kuzuya, K. Nakamura, T. Sasaki, X.W. Cheng, S. Itohara, A. Iguchi, Effect of MMP-2
Deficiency on Atherosclerotic Lesion Formation in ApoE-Deficient Mice, Arteriosclerosis
Thrombosis Vasc Biology. 26 (2006) 1120–1125.
https://doi.org/10.1161/01.atv.0000218496.60097.e0.

[33] J.W. Fischer, S.A. Steitz, P.Y. Johnson, A. Burke, F. Kolodgie, R. Virmani, C. Giachelli,
T.N. Wight, Decorin Promotes Aortic Smooth Muscle Cell Calcification and Colocalizes to
Calcified Regions in Human Atherosclerotic Lesions, Arteriosclerosis Thrombosis Vasc Biology.
24 (2004) 2391–2396. https://doi.org/10.1161/01.atv.0000147029.63303.28.

[34] C. Camaré, M. Pucelle, A. Nègre-Salvayre, R. Salvayre, Angiogenesis in the atherosclerotic
plaque., Redox Biol. 12 (2017) 18-- 34. https://doi.org/10.1016/j.redox.2017.01.007.

[35] J.V. Tapia-Vieyra, B. Delgado-Coello, J. Mas-Oliva, Atherosclerosis and Cancer; A
Resemblance with Far-reaching Implications, Arch Med Res. 48 (2017) 12–26.
https://doi.org/10.1016/j.arcmed.2017.03.005.

[36] P. Libby, G.K. Hansson, Taming Immune and Inflammatory Responses to Treat
Atherosclerosis *, J Am Coll Cardiol. 71 (2018) 173–176.
https://doi.org/10.1016/j.jacc.2017.10.081.

[37] N.A. Mabbott, J.K. Baillie, H. Brown, T.C. Freeman, D.A. Hume, An expression atlas of
human primary cells: inference of gene function from coexpression networks, Bmc Genomics.
14 (2013) 632. https://doi.org/10.1186/1471-2164-14-632.

[38] V.G. Zaha, W.C. Meijers, J. Moslehi, Cardio-Immuno-Oncology, Circulation. 141 (2020) 87–
89. https://doi.org/10.1161/circulationaha.119.042276.

[39] M. Liu, D. Gomez, Smooth Muscle Cell Phenotypic Diversity: At the Crossroads of Lineage
Tracing and Single-Cell Transcriptomics, Arteriosclerosis Thrombosis Vasc Biology. 39 (2019)
1715–1723. https://doi.org/10.1161/atvbaha.119.312131.

[40] C.A. Lareau, L.S. Ludwig, C. Muus, S.H. Gohil, T. Zhao, Z. Chiang, K. Pelka, J.M. Verboon,
W. Luo, E. Christian, D. Rosebrock, G. Getz, G.M. Boland, F. Chen, J.D. Buenrostro, N.
Hacohen, C.J. Wu, M.J. Aryee, A. Regev, V.G. Sankaran, Massively parallel single-cell
mitochondrial DNA genotyping and chromatin profiling, Nat Biotechnol. (2020) 1–11.
https://doi.org/10.1038/s41587-020-0645-6.

[41] J. Xu, K. Nuno, U.M. Litzenburger, Y. Qi, M.R. Corces, R. Majeti, H.Y. Chang, Single-cell
lineage tracing by endogenous mutations enriched in transposase accessible mitochondrial
DNA, Elife. 8 (2019) e45105. https://doi.org/10.7554/elife.45105.

[42] T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W.M. Mauck, Y. Hao, M.
Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of Single-Cell Data, Cell. 177
(2019) 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031.

[43] J. Gao, B.A. Aksoy, U. Dogrusoz, G. Dresdner, B. Gross, S.O. Sumer, Y. Sun, A.
Jacobsen, R. Sinha, E. Larsson, E. Cerami, C. Sander, N. Schultz, Integrative Analysis of
Complex Cancer Genomics and Clinical Profiles Using the cBioPortal, Sci Signal. 6 (2013) pl1–
pl1. https://doi.org/10.1126/scisignal.2004088.

[44] R. Satija, J.A. Farrell, D. Gennert, A.F. Schier, A. Regev, Spatial reconstruction of single-
cell gene expression data., Nat Biotechnol. 33 (2015) 495-- 502.
https://doi.org/10.1038/nbt.3192.

[45] C. Trapnell, D. Cacchiarelli, J. Grimsby, P. Pokharel, S. Li, M. Morse, N.J. Lennon, K.J.
Livak, T.S. Mikkelsen, J.L. Rinn, The dynamics and regulators of cell fate decisions are revealed
by pseudotemporal ordering of single cells, Nat Biotechnol. 32 (2014) 381--386.
https://doi.org/10.1038/nbt.2859.

[46] R.C. Team, R: A Language and Environment for Statistical Computing, R Foundation for
Statistical Computing, Vienna, Austria, 2020. https://www.R-project.org.

[47] M.C. Canver, D.E. Bauer, T. Maeda, L. Pinello, DrugThatGene: integrative analysis to
streamline the identification of druggable genes, pathways and protein complexes from CRISPR
screens, Bioinformatics. 35 (2018) 1981–1984. https://doi.org/10.1093/bioinformatics/bty913.

Figure Legends

Figure 1. Unbiased automatic cell labeling of human coronary scRNA data using 'SingleR' reveals abundant cells with chondrocyte (CH) and fibroblast (FB) gene expression patterns, and pseudotemporal and ultra-fine clustering reveals derivation of CH and FB-like cells from SMCs. **(A)** UMAP clustering of 9798 cells derived from human coronary artery explants with labels based on singleR annotation (Top) and cluster-based manual annotation based on Wirka et al., 2019 (Bottom). **(B)** population breakdown by percentage. SMC: smooth muscle cells, EC: endothelial cells, CH: chondrocytes, FB: fibroblasts, Mø: macrophages, SC: stem cells, CMP: common myeloid progenitor cells, GMP: granulocyte-monocyte progenitor cells. **(C) Left,** RNA trajectory (line) shows path two direct paths from the SMC starting nodes (grey circles). **Middle,** ultra-fine clustering shows the logical transition stages from SMC to CH and FB. **Right,** pseudotemporal analysis confirms that the cell and clusters existing along a logical single-cell continuum. **(D)** selected genes that were shown to vary over pseudotime by Moran's I test were visualized.

Figure 2. Comprehensive ligand-receptor analysis coupled to drug-gene interaction analysis reveals potential drug therapies to interrupt SMC to CH and FB communication and SMC trajectory (A) circle plot representation of the inferred intercellular communications within the coronary artery environment. **(B)** incoming and outgoing signal summed weights by cell type. **(C)** 'scTalk' shows that SMCs interacts with FB through three pairs of experimentally verified interactions. Of these three, complement and *DCN-EGFR* signaling is the most druggable as revealed by DGIdb 3.0. **(D)** druggable genome analysis revealed agents that can target genes such as complement signaling (C3), C-X-C Motif Chemokine Ligand 12 (CXCL12), and matrix metalloprotease (MMP2), which increases along the SMC-to-FB phenotype trajectory.

Figure 3. (A) screenshot of the PlaqView web application. **(B)** a roadmap of enhanced scRNA-

seq analysis process. Instead of cluster-based grouping, our pipeline uses automatic cell

labeling, coupled with pseudotime trajectory, cellular network interaction and drug targeting, and

provides a reproducible process for scRNA datasets. From this roadmap, it is easy to add

additional analysis tools and modify workflow as more tools and datasets become available.

Tables and Table Legends

Table 1.

	p_value	morans_test_statistic	morans_I
MYH11	<1E-297	169.0285919	0.66505236
IGFBP2	<1E-297	129.2299117	0.50844307
PPP1R14A	<1E-297	169.4172149	0.66663148
CNN1	<1E-297	170.9288702	0.67251583
TNFRSF11B	<1E-297	97.66515083	0.38400709
C7	<1E-297	156.2248703	0.61472417
C3	<1E-297	103.4495	0.7252684
SERPINF1	<1E-297	157.9946249	0.62168635
FBLN1	<1E-297	154.3399713	0.60730135
CXCL12	<1E-298	48.92096	0.3428102
MMP2	<1E-299	62.57231	0.4385507
FN1	<1E-297	125.4354275	0.49354242

579 **Table 1. Selected Moran's I statistics for genes listed in Figure 1D.** Moran's I statistics was

580 used to identify spatial correlations within the single-cell trajectory. +1 indicates that nearby cells

581 are perfectly similar, 0 indicates no similarity or pattern, and -1 indicates total dissimilarity [13].

A

B