Title: AKAP350 regulates cytoskeleton remodeling and LFA-1 organization during NK cytolytic response

Authors: Alejandro P. Pariani[1], Evangelina Almada[1], Florencia Hidalgo[1], Carla Borini-Etichetti[1], Rodrigo Vena[2], Cristián Favre[1], James R. Goldenring[3], M. Cecilia Larocca[1]*

Affiliation: [1]Instituto de Fisiología Experimental, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario (UNR), (2000) Rosario, Argentina.

[2] Instituto de Biología Molecular y Celular de Rosario, CONICET, Facultad de Ciencias Bioquímicas y Farmacéuticas, UNR, Rosario, Argentina.

[3] Epithelial Biology Center and Department of Cell & Developmental Biology, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

Corresponding Author's Information: M. Cecilia Larocca, Instituto de Fisiología Experimental, Facultad de Ciencias Bioquímicas y Farmacéuticas. Suipacha 570, 2000 Rosario, Argentina. Tel: 54-341-4305799. E-mail: larocca@ifise-conicet.go

Abstract

Natural killer (NK) cell cytotoxicity requires extensive cytoskeleton remodeling and

receptors reorganization at the site of NK/target cell interaction (immune synapse, IS).

How these processes are integrated remains unclear. We found that AKAP350, a scaffold

protein that regulates microtubule dynamics, was required for NK cytolytic activity.

Reduction of AKAP350 expression inhibited actin and LFA-1 reorganization and

centrosome translocation to the IS. AKAP350KD cells showed decreased microtubule

nucleation at the Golgi apparatus and reduced localization at the centrosome and at the IS

of CIP4, an AKAP350 interacting protein that regulates centrosome translocation to the

IS. AKAP350 delocalization from the Golgi or pharmacological disruption of

microtubule dynamics or Golgi function impaired LFA-1 clustering at the IS, preserving

actin remodeling. These data support a role for AKAP350 in the integration of actin and

microtubule remodeling during NK activation and reveal for the first time the

participation of the Golgi apparatus in NK-IS maturation.

Introduction

Natural killer cells (NK) are the cytolytic effectors of the innate immune system, which

constitute the first line of immunological defense and are particularly relevant for the

elimination of viral infected and neoplastic cells. NK cytotoxic effector function relies on

the formation of a specialized junction with its target cell, generally referred to as immune

synapse (IS). In the case of NK lytic IS, the persistent outside-in signaling generated by

the interaction of NK activating receptors with their ligands in a susceptible target cell,

in a context of absence of inhibitory signals, triggers a multistep, highly regulated process

that culminates with the directed secretion of lytic molecules contained in specialized

lysosomes (lytic granules) (Mace et al., 2014). An early event in this process is actin

cytoskeleton remodeling into an F-actin-rich supramolecular activation cluster (SMAC)

at the IS, and the subsequent clustering of cell-surface receptors at this site. IS maturation

requires actin organization as a compact mesh at the SMAC periphery and a less dense

net of branching filaments at the IS center (Brown et al., 2011), where lytic degranulation

takes place (Rak et al., 2011). Immediately after actin reorganization at the IS, the

centrosome, usually regarded as the main microtubule organizing center (MTOC) in NK

cells, translocates to the IS, enabling lytic granules translocation and directed secretion

of lytic granules contents into the target cell (Mace et al., 2014). Therefore, the

development of NK lytic response requires an extensive, synchronized actin and

microtubule remodeling. Even though some proteins that participate in actin and

microtubule dynamics during the NK lytic IS, such as WASP (Orange et al., 2002),

CDC42 (Carlin et al., 2011) and the CDC42 interacting protein 4 (CIP4) (Banerjee et al.,

2007) have been identified, the mechanisms that coordinate actin and microtubule

remodeling remain poorly understood.

The A-kinase anchoring protein 350 (AKAP350), also known as AKAP450, CG-NAP or

AKAP9, is a centrosome and Golgi apparatus-scaffold protein (Schmidt et al, 1999) that

arises as a putative candidate to regulate cytoskeleton remodeling during the development

of the NK lytic response. AKAP350 has a prominent role in microtubule nucleation

(Takahashi et al., 2002, Rivero et al., 2009, Kolobova et al., 2019). AKAP350 also

participates in mitotic spindle orientation (Almada et al., 2017) and in centrosome

translocation towards the leading edge in migratory cells (Tonucci et al., 2015).

AKAP350 participation in microtubule remodeling is related to its capacity to scaffold

specific proteins at the centrosome and at the Golgi apparatus, including the members of

the γ-TURC CGP2/3 (Takahashi et al 2002), the casein kinase δ (Sillibourne et al., 2002),

CIP4 (Larocca et al., 2004), the component of the dynein/dynactin complex p150 glued

(Kim et al., 2007) and the plus end binding protein 1 (EB1) (Almada el al., 2017). The

involvement of AKAP350 in microtubule reorganization in response to outside-in

signaling in immune cells was first revealed by studies in T cells (El Din El Homasany et

al., 2005). Those studies demonstrated that the interference with AKAP350 function at

the centrosome leads to inhibition of lymphocyte function-associated antigen-1 (LFA-1

or αLβ2 integrin; CD11a/CD18) induced-cell migration. Later studies indicated that,

during antigen recognition by T-cells, AKAP350 participates in LFA-1 conformational

activation and centrosome translocation towards the IS (Robles Valero et al., 2010).

Regarding the mechanisms involved, they have not been elucidated.

In the present study we analyzed AKAP350 participation in the development of NK

cytotoxic effector function and characterized its role in actin and microtubule

cytoskeleton remodeling, and in LFA-1 reorganization at the lytic IS. We further

characterized microtubule nucleation in activated NK cells and evaluated the role of Golgi

associated AKAP350 in NK-IS maturation.

Results

1. AKAP350 localization at the Golgi apparatus in NK/YTS cells is up-regulated during

cell activation

To determine if AKAP350 could have a role in NK cytotoxic response, we first evaluated

its expression in YTS immortalized NK cells. Western blot analysis of YTS cell-extracts

indicates that AKAP350 is expressed in YTS cells. Immunofluorescence analysis of

AKAP350 staining in resting YTS cells showed that AKAP350 colocalized with γ-tubulin

and GM-130, what is indicative of AKAP350 localization at the centrosome and at the

Golgi apparatus (Figure 1A, B first row). KT86 cells are derived from K562

erythroleukemia cells and constitute susceptible targets for YTS cells. The analysis of

YTS:KT86 conjugates indicates that AKAP350 localization at the centrosome was

reduced (-50%), whereas its localization at the Golgi apparatus was increased (+44%) in

activated NK cells (Figure 1A, B, second row and bars). These results indicate that

activation of YTS cells induces a partial translocation of AKAP350 from the centrosome

to the Golgi apparatus.

2. Decrease of endogenous AKAP350 expression impairs NK cytolytic activity

To analyze the impact of AKAP350 expression in the development of NK cytolytic

effector function, we generated YTS cells with decreased AKAP350 expression

(AKAP350KD) using a lentiviral short hairpin RNA expression system targeting two

different AKAP350 mRNA sequences, as we have previously described (Tonucci et al.,

2015). Western blot analysis of AKAP350 protein levels showed that the expression of

either shRNA1 or shRNA4 efficiently decreased AKAP350 levels to less than 10% of

Figure 1. AKAP350 localization in YTS cells. Isolated YTS cells or YTS:KT86 conjugates were stained and analyzed by confocal microscopy (see Materials and methods). Merge images show YTS cells (upper row) or YTS:KT86 conjugates (lower row) staining for AKAP350 (red) and γ-tubulin (A) or GM130 (B) (green). Asterisk denotes YTS cells in YTS:KT86 conjugates. Bars represent the mean fraction of AKAP350 fluorescence present at centrosomes (A) or at the Golgi apparatus (B) in isolated or conjugated YTS cells, expressed as percentage of total AKAP350-fluorescence, representative of three independent experiments. At least 20 cells were analyzed for each experiment. Error bars represent SEM. *p < 0.05. Scale bars, 5 μm.

control levels (Figure 2A). To assess AKAP350 participation in NK adhesion to target

cells, control and AKAP350KD YTS cells were labeled with TrackerTM Deep Red and

KT86 cells were labeled with CFSE. Cells were incubated at different NK:target cell

ratios, for different periods, and the formation of cell conjugates positive for both

fluorophores was analyzed by FACS (Supplementary Figure 1). Our results showed that

the decrease in AKAP350 expression did not elicit any evident effect in the initial

adherence between NK and their susceptible target cells. In order to analyze NK cytolytic

activity, control and AKAP350KD YTS cells generated by expression of shRNA1 or

shRNA4 were incubated with CFSE-labeled KT86 cells at different NK:target cell ratios

and target cell lysis was analyzed by FACS analysis of propidium iodide stained cells.

Our results showed that the decrease in AKAP350 levels by expression of either shRNA1

or shRNA4 impaired NK cytolytic activity (Figure 2B,C).

3. AKAP350 participates in centrosome translocation towards the lytic IS

Our previous studies demonstrated that AKAP350 regulates centrosome translocation

towards the leading edge in migratory cells, a process with some homology to the

development of the IS. Considering that centrosome translocation to the IS is a critical

step in NK cytolytic response, centrosome localization in control and AKAP350KD YTS

cells conjugated with KT86 target cells was evaluated by confocal microscopy. To

discriminate NK from target cells, KT86 cells were preloaded with CFSE. γ-tubulin

staining was used to identify the centrosome, and the IS was defined as the region of

YTS/KT86 contact delimited in the DIC channel. Centrosome translocation was impaired

in YTS cells with decreased expression of AKAP350, as denoted by the greater distance

between the MTOC and the centroid of the IS region in AKAP350KD cells ((Figure 3A).

In non cytolytic T cells, AKAP350 participates in LFA-1 activation and the subsequent

translocation of the centrosome to the IS (Robles Valero et al, 2010). Considering that

Figure 2. Reduction of AKAP350 expression levels diminishes YTS cells' cytotoxicity. Two different specific shRNAs (shRNA1 and shRNA4) were used to generate YTS cells with decreased AKAP350 expression (AKAP350KD1 and AKAP350KD2, respectively), as described in Materials and methods. A) Western blot analysis of AKAP350 expression in control, AKAP350KD1 and AKAP350KD2 YTS cells. CIP4 was used as loading control. B,C) KT86 cells were stained with CFSE and mixed with YTS cells at 2:1 (B) or 10:1 (C) effector: target ratios. After the incubation period, cells were stained with PI and analyzed by flow cytometry. YTS cytotoxic activity was estimated as the fraction of total CFSE positive events that were also positive for PI. Bars chart represents the mean percentage of double positive events for control, AKAP350KD1 and AKAP350KD2 cells, representative of three independent experiments. Error bars represent SEM. *p < 0.05.

LFA-1 activation is sufficient to induce MTOC translocation in NK cells, we analyzed

AKAP350 role in MTOC translocation upon specific activation of LFA-1 or CD28, an

additional NK receptor that induces MTOC translocation as well (Chen et al., 2006). YTS

cells activated on slides coated with ICAM-1 or with an activating anti-CD28 antibody

were stained and analyzed by confocal microscopy (Figure 3B,C). In both conditions,

AKAP350KD cells showed an impairment in MTOC translocation towards the activating

surface. Therefore, AKAP350 participates in MTOC translocation by regulating

processes that are common to the activation of both LFA-1 and CD28 receptors.

4. AKAP350 regulates CIP4 localization in NK cells

CIP4 is a CDC42 effector protein involved in regulation of membrane curvature and actin

remodeling. In activated NK cells, CIP4 localizes at the MTOC and at the IS and regulates

MTOC translocation towards the IS (Banerjee et al., 2007). Considering that, in epithelial

cells, AKAP350 interacts with CIP4 and regulates its subcellular localization (Larocca et

al 2004, Tonucci et al 2015), we evaluated CIP4 localization at the centrosome in control

and AKAP350KD YTS cells. Immunofluorescence analysis by confocal microscopy

showed that CIP4 localization at the MTOC was significantly decreased in AKAP350KD

YTS cells (Figure 4A). In order to validate those results, subcellular fractions enriched in

centrosomes were prepared by differential centrifugation on a Ficoll cushion, and CIP4

levels at those fractions were analyzed by western blot. Estimates considering total

protein recovery and CIP4 band density at total cell lysates and at centrosomal fractions

indicated that 3.0 % of total CIP4 localized at the centrosome in control cells, whereas

CIP4 yield at centrosomal fraction was reduced to 1.5 % in AKAP350KD cells (Figure

4B), thus ratifying that AKAP350 participates in CIP4 recruitment to the centrosome in

YTS cells. In order to analyze the relevance of AKAP350 in CIP4 localization during the

Figure 3. AKAP350 is required for centrosome polarization during YTS cells activation. A) Control and AKAP350KD YTS cells were incubated with CFSE labeled KT86 cells at a 2:1 ratio. Cell conjugates were fixed, stained and analyzed by confocal microscopy. Merge images show staining for AKAP350 (red), γ-tubulin (cyan) and DAPI (blue), and CFSE labeled target cells (green). White arrows indicate the position of γ-tubulin-labeled centrosomes in YTS cells. Bars represent the mean distance from the centrosome to the IS, representative of five independent experiments. B,C) Control and AKAP350KD YTS cells were activated using coverslips coated with the LFA-1 activating ligand ICAM-1 (B) or an activating anti-CD28 antibody (C). Activated cells were fixed, stained and analyzed by confocal microscopy. Merge images show γ-tubulin (green) and DAPI (blue) staining. White arrow indicates the position of γ-tubulin-defined centrosomes. White dashed lines denote the cell surface in contact with the coverslip. Bars represent the mean distance from the centrosomes to the cell-coverslip contact surface, representative of three independent experiments. At least 30 conjugates (A) or cells (B,C) were analyzed for each experiment. Error bars represent SEM. Scale bars, 5 μm. *p<0.05.

Figure 4. Reduction of AKAP350 expression levels alters CIP4 localization in YTS cells. A) Merge images show staining for CIP4 (red) and γ-tubulin (green) in control and AKAP350KD YTS cells. White arrows mark γ-tubulin defined centrosomes. Bars represent the mean fraction of CIP4 associated fluorescence located at the centrosome, expressed as percentage of total cell fluorescence in the CIP4 channel, representative of three independent experiments. At least 30 conjugates were analyzed for each experiment. Error bars represent SEM. B) Centrosomes enriched subcellular fractions were prepared by differential centrifugation of control or AKAP350KD cell lysates (see Materials and Methods). Total cells lysates and centrosomal fractions were analyzed for CIP4 expression by western blot. Bars represent the mean fraction of CIP4 recovered at centrosome-fractions, expressed as percentage of total CIP4 recovery of three independent experiments. C) CIP4 localization during establishment of the IS was analyzed in control and AKAP350KD YTS:KT86 conjugates (2:1 ratio). Merge images show conjugates staining for CIP4 (red) and γ-tubulin (blue). White arrows indicate the position of γ-tubulin-defined centrosomes. Bars represent the mean fraction of CIP4 fluorescence located at the centrosome or at the IS, expressed as percentage of total cell fluorescence for the CIP4 channel, representative of three independent experiments. At least 30 conjugates were analyzed for each experiment. Error bars represent SEM. D) Control or AKAP350KD cells were incubated in plates coated with the activating anti-CD28 antibody and ICAM-1. Fractions of stabilized microtubules were obtained with non-ionic detergent extraction (see Material and methods). CIP4 and α-tubulin recovery in total cell lysates and microtubule-fractions were analyzed by western blot. Bars represent the mean fraction of CIP4 or α-tubulin recovered at microtubule-fractions, expressed as percentage of total yield in cell lysates, and the mean fraction of CIP4 recovered at the microtubule fraction related to α-tubulin recovery for three independent experiments. Scale bars, 5 μm. *p < 0.05

development of the lytic IS, CIP4 localization in YTS:KT86 conjugates was analyzed by

confocal microscopy. Similarly to our results in resting cells, the analysis of YTS:KT86

conjugates showed that CIP4 localization at the centrosome was decreased in

AKAP350KD cells (Figure 4C). Furthermore, CIP4 localization at the IS was also

impaired in those cells (Figure 4C). By means of its amino-terminal domain CIP4 can

interact with the microtubules, therefore, we further analyzed CIP4 in situ interaction with

microtubules. Control and AKAP350KD cells were activated with ICAM-1 and extracts

of stabilized microtubules and associated proteins were prepared. Western blot analysis

of those extracts indicated that a significant fraction of CIP4 was associated with

microtubules in control cells and that this fraction was significantly reduced in

AKAP350KD cells (Figure 4D). It is noteworthy that α-tubulin recovery in those

microtubule extracts was also diminished in AKAP350KD cells. In order to evaluate if

this decrease in α-tubulin recovery in microtubule extracts was associated with decreased

microtubule stability, we analyzed the relative levels of acetylated α-tubulin in NK cell

lysates by western blot. Similarly to what was previously described in motile T cells (Ong

et al., 2018), the decrease in AKAP350 expression was associated with a decrease in the

levels of acetylated α-tubulin in activated YTS cells, thus indicating decreased

microtubule stability in activated AKAP350KD NK cells (Supplementary Figure 2A).

CIP4 yield in AKAP350KD microtubule extracts relativized to α-tubulin yield was also

reduced (Figure 4D), suggesting that AKAP350 participates in CIP4 recruitment to the

microtubule cytoskeleton, independently of its role in the regulation of microtubule

stability.

5. The decrease in AKAP350 expression impairs actin remodeling in activated NK cells

We further analyzed if the decrease in AKAP350 expression had an impact on actin

remodeling during NK cell activation. Analysis of YTS:KT86 conjugates by confocal

microscopy indicated decreased actin accumulation at the IS in AKAP350KD cells

(Figure 5A). Analysis of YTS:KT86 conjugates morphology indicated that the surface of

interaction between both cells was also diminished in AKAP350KD cells, which was

verified by a reduction in the IS area to 50% of control levels (p<0.01). The decrease in

the area of effector:target cell interaction is usually associated with decreased integrin-

clustering. Consistently, LFA-1 reorganization at the IS was impaired in AKAP350KD

cells (Figure 5A). Similarly, inhibition of actin and LFA-1 reorganization was also

observed in AKAP350KD cells when they were specifically activated with ICAM-1

(Figure 5B) or with the activating anti-CD28 antibody (Figure 5C). The analysis of actin

distribution within the IS indicate that control YTS cells showed the characteristic actin

pattern of a compact ring in the periphery with less dense actin organization at the center

of the IS, whereas AKAP350KD cells showed a more homogeneous actin organization

(Figure 5D). LFA-1 clustering at the periphery of the IS was also impaired in

AKAP350KD cells (Figure 5D). Those results indicate that AKAP350 participate in

regulatory events that enable actin remodeling and LFA-1 clustering at the site of NK-

target cell interaction.

6. Microtubule cytoskeleton integrity is necessary for LFA-1 organization at the IS

In certain conditions, microtubule targeting drugs inhibits NK cytotoxic response (Fong

et al., 2019). Nevertheless, microtubule participation in the maturation of NK lytic IS has

not been thoroughly evaluated. We analyzed actin remodeling and LFA-1 clustering at

the IS in conditions where microtubule integrity was compromised, in the presence of

microtubule-targeting drugs at levels that inhibit NK cytotoxicity. We found that neither

cold-induced microtubule depolymerization in the presence of Nocodazole nor

suppression of microtubule dynamic instability with Taxol have a significant effect on

Figure 5. Reduction of AKAP350 expression levels results in defective actin and LFA-1 organization at the IS. A) Control and AKAP350KD YTS cells were incubated with CFSE labeled KT86 cells at a 2:1 ratio. Merge images show YTS:KT86 conjugates staining for actin (red), LFA-1 (cyan) and DAPI (blue). Individual actin and LFA-1 channels are shown in pseudocolor. Bars represent the mean fraction of actin or LFA-1 fluorescence located at the IS, expressed as percentage of total cell fluorescence for each channel, for three independent experiments. At least 30 conjugates were analyzed for each experiment. Error bars represent SEM. B, C) Control and AKAP350KD YTS cells were activated using coverslips coated with ICAM-1 (B) or with the activating anti-CD28 antibody (C).

Merge images show actin (red), LFA-1 (green) and DAPI (blue) staining. Individual actin and LFA-1channels are shown in pseudocolor. White dashed lines indicate the cell surface in contact with the coverslip. Bars represent the mean fraction of actin or LFA-1 fluorescence located at the cell surface in contact with the ICAM-1 (B) or anti-CD28 (C) coated coverslip, expressed as percentage of total cell fluorescence for each channel representative of three independent experiments. At least 30 conjugates were analyzed for each experiment. Error bars represent SEM. D) Control and AKAP350KD YTS cells were incubated with CFSE labeled KT86 cells. Images show the X-Z projection of the IS region corresponding to control and AKAP50KD YTS cells. Merge images show LFA-1 (green) and actin (red) staining. Individual LFA-1 and actin channels are shown in pseudocolor. Dashed lines denote the radius of the IS where LFA-1 and actin intensity of fluorescence were quantified. Charts represent the profile of intensity of fluorescence quantified for each channel, representative of at least ten IS analyzed in three independent experiments. Scale bars, 5 μm. *p<0.05.

total actin accumulation at the IS (Figure 6). Regarding LFA-1 distribution at the IS, both

treatments elicited similar effects, inhibiting LFA-1 accumulation (Figure 6A) and

organization (Figure 6B) at the IS.

7. AKAP350 participates in microtubule nucleation at the Golgi apparatus in activated

NK cells

Previous studies demonstrate that AKAP350 can modulate microtubule nucleation at the

Golgi apparatus and at the centrosomes in different systems (Takahashi et al, 2002;

Rivero et al, 2009; Ori-McKenney et al, 2012; Maia et al, 2013; Tonucci et al, 2018). To

our knowledge, microtubule nucleation at the Golgi apparatus in NK cells has not been

analyzed so far. Therefore, we analyzed microtubule nucleation at the centrosome and at

the Golgi apparatus in activated YTS cells, using ice recovery assays as previously

described (Grimaldi et al, 2013). In conditions of YTS cell activation with ICAM-1 and

anti-CD28 receptor antibody, control cells showed Golgi and centrosome nucleation of

microtubules (Figure 7, first row). AKAP350KD cells, in contrast, showed almost

exclusive centrosome nucleation of microtubules (Figure 7, second row). Quantitative

analysis of newly nucleated microtubules indicated a significant reduction in Golgi

associated microtubules in AKAP350KD cells (Figure 7, upper bars), whereas no

differences were observed for centrosome-associated microtubules (Figure 7, lower bars).

Similar results were obtained at 3 min (Figure 7) or 7 min (not shown) of recovery.

8. Golgi apparatus integrity is essential for the IS maturation in NK cells

As delineated above, during YTS cell activation AKAP350 translocates to the Golgi

apparatus (Figure 1) where it is essential for microtubule nucleation (Figure 7). In order

to analyze if AKAP350 participation in NK IS maturation could be associated with its

function at the Golgi apparatus, we analyzed the IS development in NK cells subjected to

Figure 6. Microtubule dynamics is required for LFA-1 organization at the IS. Control YTS cells or YTS cells treated with nocodazole or taxol (see Materials and methods) were mixed with KT86 cells at a 2:1 ratio. Images show x,y planes (A) or x,z projections (B) of the IS region corresponding to control and nocodazole or taxol treated YTS cells. Merge images show cell conjugates staining for actin (red) and LFA-1 (green). LFA-1 channel is also shown in pseudocolor. A) Bars represent the mean fraction of LFA-1 fluorescence located at the IS, expressed as percentage of total cell fluorescence, representative of three independent experiments. At least 30 conjugates were analyzed for each experiment. B) Dashed lines denote the radius of the IS where LFA-1 intensity of fluorescence was quantified. Charts represent the profile of intensity of fluorescence quantified for the LFA-1 channel, representative of at least ten IS analyzed in three independent experiments. Scale bars, 5 µm. Error bars represent SEM. *p<0.05.

Figure 7. AKAP350 is essential for microtubule nucleation at the Golgi apparatus (GA) in activated YTS cells. Control or AKAP350KD YTS cells were activated using ICAM-1 and anti-CD28 antibody for 30 minutes and then subjected to ice recovery assays. Merge images show staining for GM130, as Golgi marker, in red and for α-tubulin in green. Bar charts represent the mean number of microtubules derived from Golgi apparatus or from the centrosome and the mean fraction of α-tubulin fluorescence associated to each organelle, expressed as percentage of total cell fluorescence for the α-tubulin channel, representative of three independent experiments. At least 10 cells were analyzed for each experiment. Error bars represent SEM. Scale bars, 5 μm. *p<0.05

Brefeldin A (BFA) treatment. BFA inhibits the activation of ADP-ribosylation factors,

leading to the collapse of the Golgi and its redistribution into the ER (Chardin and

McCormick, 1999). Previous studies showed that BFA treatment induced relocalization

of AKAP450 from the Golgi apparatus to ER exit sites and decreased MT nucleation at

the Golgi apparatus (Rivero et al. 2009). Analysis of LFA-1 distribution at the IS showed

that BFA-treatment inhibited LFA-1 organization at the IS (Figure 8A,B). Interestingly,

actin organization was preserved in BFA treated cells (data not shown). To specifically

examine the relevance of AKAP350 localization at the Golgi in the maturation of NK

lytic IS, we prepared YTS cells with stable expression of AKAP350(1-1229)

(AKAP350GBD) domain, which acts as a dominant negative construct that displaces the

protein from the Golgi apparatus, inhibiting microtubule nucleation at this organelle

(Hurtado et al 2011). The analysis of AKAP350 colocalization with GM130 confirmed

that AKAP350 localization at the Golgi apparatus was significantly decreased in

AKAP350GBD cells (Figure 8C). Relative levels of acetylated α-tubulin were decreased

in AKAP350GBD cells (Supplementary figure 2B), suggesting that the decreased α-

tubulin acetylation observed in AKAP350KD cells was associated with the decreased

expression of AKAP350 at the Golgi apparatus. Similarly to what was observed in BFA

treated cells, LFA-1 reorganization was compromised in AKAP350GBD cells (Figure

8D,E), whereas actin organization at the IS was preserved. Overall, this set of data

showed, for the first time, that the Golgi apparatus is directly involved in LFA-1

organization at the IS in NK cells and, considering the evidence presented in Figures 6

and 7, gave strong evidence supporting a role for Golgi derived microtubules in this

effect.

Figure 8. AKAP350 localization at the Golgi apparatus is required for IS maturation. A, B) YTS cells were treated with brefeldin A (BFA) (see Material and methods). A) Control and BFA treated YTS cells were incubated with KT86 cells. Merge images show staining for LFA-1 (green) and GM130 (gray). LFA-1 channel is also shown in pseudocolor. Bars represent the mean fraction of LFA-1 fluorescence located at the IS, expressed as percentage of total LFA-1 cell fluorescence, representative of three independent experiments. B) Images show the x,z projection of the IS region. Dashed lines denote the radius of the IS where LFA-1 intensity of fluorescence was quantified. Charts represent the profile of intensity of fluorescence, representative of at least 10 IS. C-E) YTS cells with stable expression of AKAP350 Golgi binding domain (AKAP350GBD) cells were prepared as described in Materials and methods. C) Merge images show staining for AKAP350 (green) and GM130 (red). Bars represent the mean Pearson´s correlation coefficient, used to estimate AKAP350/GM130 colocalization, for 10 different cells. D,E) Control and AKAP350GBD YTS cells were incubated with KT86 cells. D) Merge images show YTS:KT86 conjugates staining for actin (red) and LFA-1 (green). Individual actin and LFA-1 channels are shown in pseudocolor. Bars represent the mean fraction of LFA-1 fluorescence located at the IS, expressed as percentage of total cell fluorescence. More than 30 conjugates from 3 different experiments were analyzed. E) Images show the x,z projection of the IS region corresponding to control and AKAP350GBD YTS cells. Dashed lines denote the radius of the IS where LFA-1 intensity of fluorescence was quantified. Charts represent the profile of intensity of fluorescence, representative of at least ten IS. Scale bars, 5 μm. Error bars represent SEM. *p<0.05.

Discussion

NK cytotoxic effector function depends on the extensive remodeling of actin and microtubule cytoskeleton, which has been extensively characterized (reviewed in Lagrue et al, 2013). Nevertheless, the mechanisms coordinating cytoskeleton reorganization are not completely understood. In the present study, we provide evidence that AKAP350 acts as a scaffold protein that integrates events that are central for both microtubule and actin organization, therefore regulating the development of NK lytic IS. Furthermore, our results indicate that the Golgi apparatus acts as a MTOC in activated NK cells and reveals a novel role for this organelle in the maturation of the lytic IS.

In the present study, we found that normal levels of AKAP350 expression are required for MTOC translocation to the IS during the development of NK lytic response. Previous studies by Banerjee and col (2007) demonstrated that CIP4 localizes to the centrosome and to the IS in activated NK cells and that it has a central role in centrosome translocation to the IS. CIP4 is an F-BAR protein initially identified by its ability to interact with the activated form of CDC42 (Aspenström 1997). In the present work, using biochemical and immunofluorescence microscopy-assays, we found that AKAP350 is essential for CIP4 localization at the centrosome both in resting and in activated YTS cells. These results give central cues on how CIP4, a protein essentially associated with phospholipid membranes, interacts with the only non-membranous organelle in NK cells and suggest that, homologously to what was described in migratory cells, AKAP350/CIP4 protein complex could act in concert to regulate centrosome translocation towards the IS. Our data further showed that AKAP350 conditions CIP4 localization at the IS. Besides its F-BAR domains, CIP4 also contains a Src Homology 3 (SH3) domain at the carboxyl terminus that mediates its interaction with the proline-rich domains of WASP (Tian et al, 2000). WASP is an activator of the Arp2/3 complex that promotes the formation of

branched actin filaments, being positively regulated in this function by CDC42 (Leibfried

et al., 2008). In NK cells, WASP colocalizes with actin at the IS, playing a central role in

the development of the cytotoxic response (Orange et al., 2002). Interestingly, CDC42

activity increases at the IS very early during NK cell activation (Carlin et al, 2011). In

this scenario, CIP4 could act as an adaptor that enables WASP activation by CDC42 at

NK lytic IS. Moreover, by means of its SH3 domain, CIP4 also interacts with formins

Dia and DAAM1 (Tsujita et al., 2006; Aspenström et al., 2006). Formins generate linear

actin filaments which, in T cells, constitute the central region of the IS and are essential

for integrin-clustering and IS maturation (Murugesan et al., 2016). In NK cells, Dia

localizes centrally at the IS and participates in targeting microtubules to this site, where

lytic degranulation occurs (Butler and Cooper, 2009). Therefore, CIP4 localization at the

IS could be central for the maturation of the F-actin-rich SMAC at the IS and microtubule

interaction with this site. In this regard, the decrease in CIP4 expression does not affect

total actin accumulation at the IS, indicating that CIP4 does not participate in initial actin

nucleation, but it is involved in later events that are central for the maturation of the IS

(Banerjee et al., 2007). How AKAP350, which specifically localizes to the Golgi

apparatus and the centrosome, conditions CIP4 localization at the IS is not clear. In this

regard, there is evidence supporting two putative mechanisms: 1. AKAP350 regulates

CIP4 T225 phosphorylation by PKA, which controls its localization in invadopodia and

CIP4 interaction with CDC42 (Tonucci et al., 2019); similarly, AKAP350 could integrate

regulatory events that condition CIP4 localization at the IS. 2. CIP4 recovery in

microtubule extracts was decreased in AKAP350KD cells. As we discuss below,

microtubule dynamics is altered in AKAP350KD cells. Therefore, AKAP350 could

regulate CIP4 localization at the IS by enabling cellular events that condition CIP4

trafficking towards the IS.

Microtubules can regulate actin remodeling in dissimilar scenarios where polarized actin

organization is required, such as integrin mediated phagocytosis (Lewkowicz et al.,

2008), directional cell migration (Waterman-Storer & Salmon, 1999) and epithelial apical

pole-differentiation (Mattaloni et al, 2012; Tonucci et al, 2018). Microtubule´s role in

actin-organization at the IS was not clear. Microtubule targeting agents have paradoxical

effects on NK cells cytotoxicity, what seems to be related to the concentration and the

type of microtubule targeting agent evaluated (Katz et al., 1982; Fong et al., 2019). We

found that levels of nocodazole that actually inhibits NK cytotoxicity (Katz et al., 1982),

impair LFA-1 organization at the IS, but does not elicit any evident effect on actin

organization. Orange et al have previously demonstrated that neither cell-surface receptor

nor F-actin accumulation at the activating NK IS were disrupted with colchicine (Orange

et al., 2003). Colchicine and nocodazole have similar mechanisms of action: both drugs

bind to the β tubulin subunit blocking α/β heterodimers polymerization (Florian and

Mitchison 2016). Nevertheless, in our study we additionally induced microtubule

depolymerization by ice incubation, which leads to a more robust disruption of

microtubule integrity. Our results also showed that micromolar levels of the microtubule

stabilizing agent Taxol, which impedes microtubule growing and shortening events

(Derry et al., 1995) and inhibit NK cytotoxicity (Chuang et al., 19941), did not affect

actin accumulation or organization, but inhibited LFA-1 reorganization at the IS. At

micromolar levels, taxol inhibits the association of microtubule plus end binding proteins

such as EB1 (Morrison et al., 1998) and CLIP170 (Perez et al., 1999). Therefore,

microtubule dynamic instability and/or the presence of specific microtubule plus end

binding proteins could be required for LFA-1 reorganization at the IS. In this regard, in

T cells, the α-tubulin deacetylase HDAC6, translocates to the IS where it regulates α-

tubulin deacetylation and LFA-1 organization (Serrador et al., 2004). HDAC6 increases

microtubule dynamics instability, which correlates with its interaction with EB1

(Zilberman et al., 2009) and with CLIP170 (Li et al., 2014). Therefore, it would be

interesting to investigate if HDAC6 and/or specific microtubule plus end binding proteins

are involved in LFA-1 reorganization at the lytic IS in NK cells.

Our results demonstrate that, upon NK cell interaction with susceptible targets, AKAP350

partially translocates from the centrosome to the Golgi apparatus. Therefore, we analyzed

if there was a specific role of AKAP350 at the Golgi apparatus. Even though it is well

established that the Golgi apparatus translocates to the IS during NK lytic response

(Kupfer et al., 1983), to our knowledge, the role of this organelle in the development of

NK lytic IS has not been examined so far. We found that AKAP350 delocalization from

the Golgi apparatus inhibits LFA-1 reorganization at the IS, underscoring the relevance

of AKAP350 location at this organelle for the development of NK lytic IS. Furthermore,

interference with the Golgi apparatus function with BFA, inhibited LFA-1 reorganization

at the IS as well, which reassures that the Golgi apparatus has regulatory effects in the

maturation of the IS. The presence of AKAP350 at the Golgi apparatus has been mainly

associated with the role of this organelle in microtubule nucleation (Rivero et al, 2009,

Ori-McKenney et al, 2012; Maia et al, 2013, Tonucci et al, 2018). Microtubule nucleation

in lymphocytes has been typically attributed to the centrosome, which, in fact, is

commonly denoted as MTOC. We found that LFA-1/CD28 activated NK cells nucleate

microtubules both at the centrosome and at the Golgi apparatus. In those conditions, the

decrease in AKAP350 expression specifically inhibits microtubule nucleation at the

Golgi. Those results are in line with studies that showed that the decrease in AKAP350

expression inhibits microtubule nucleation at the Golgi, without having a significant

impact on centrosomal nucleation of microtubules (Larocca et al, 2006; Rivero et al,

2009; Tonucci et al, 2018). Golgi-derived microtubules are particularly stable, which

correlates with their enrichment in post-translationally modified tubulin (reviewed in Sanders et al., 2015). In fact, the study that first characterized the Golgi apparatus role as a MTOC showed that Golgi-nucleated microtubules become rapidly acetylated (Chabin-Brion et al., 2001). A recent study in motile T cells showed that AKAP350 participates in microtubule nucleation at the centrosome and at extra-centrosomal nucleation centers (Ong et al., 2018). Although this study did not identify microtubule nucleation at the Golgi, it showed that the decrease in AKAP350 expression led to decreased levels of acetylated α-tubulin. Consistently, we found that either the decrease in AKAP350 expression or its delocalization from the Golgi apparatus elicited a decrease in acetylated α-tubulin levels. Altogether, our results demonstrate that the Golgi apparatus participates in cytolytic IS maturation in NK cells, and strongly suggest that Golgi derived microtubules are involved in this effect. How Golgi-derived microtubules can affect NK cytolytic IS maturation deserves further investigation. In this regard, different studies indicate that those non-radial, highly stable tracks support the directionality of Golgi associated polarized trafficking, (reviewed in Sanders and Kaverina, 2015). In connection with that point, studies performed in T cells show that the Golgi apparatus regulates the two way vesicle trafficking to the IS of signaling proteins that are essential for IS maturation (Carpier et al., 2018). Our future studies will address how this particular set of microtubules modulate Golgi derived trafficking and the relevance of this trafficking for the maturation of the lytic IS in NK cells.

Materials and methods

NK and target cell lines

The immortalized NK YTS cells [Drexler, and Matsuo. 2000] and KT86 cells, derived from the MHC class I-negative K562 erythroleukemia cell line stably expressing CD86

[Parry et al, 2003], used as susceptible target cells, were obtained from Dr. Jordan Orange lab (Texas Children's Hospital, Houston, TX, USA). Cells were maintained in RPMI medium 1640 supplemented with 10% FBS, 1% L-glutamine, 1% non-essentials amino acids and 1% streptomycin/ampicillin mixture at 37° C and 5% CO_2 atmosphere.

In order to prepare YTS cells with reduced expression of AKAP350 (AKAP350KD), we proceeded as we have described previously [Tonucci et al., 2015]. Constructs were made by annealing and ligating oligonucleotides targeting two specific AKAP350 sequences (shRNA1, 5′-CCCAGCTCACTGCTAATTT-3′; shRNA4, 5′-GCAAGAACTAGAACGAGAA-3′) or a scrambled control into the AgeI and EcoRI cloning sites of pLKO.1-puro vector (details at http://www.addgene.org). These constructs were sequenced and used to co-transfect human embryonic kidney 293 FT cells with Virapower lentiviral packaging mix (Invitrogen, Carlsbad, CA). The next day, transfection complexes were removed, and cells were allowed to produce virus for 24 h. Media containing viruses were collected and used to directly transduce YTS cells overnight. The cells were then allowed to recover for 24 h and subjected to puromycin selection (2 μg/ml) for 2 weeks. Silencing was confirmed by western blotting and immunofluorescence analyses.

In order to delocalize AKAP350 from the Golgi apparatus, the nucleotide sequence codifying for AKAP350(1-1229) (AKAP350GBD) was cloned into pInducer20 vector (details at http://www.addgene.org). The lentiviral particles were constructed as described above. Transduced YTS cells were subjected to G418 selection (1 μg/ml) for two weeks. AKAP350GBD expression was induced with Doxycycline (2 μg/ml) 24 h before the experiments.

CFSE-based Cytotoxic assay

In order to determine the lytic capacity of YTS cells, we proceed as described [Jedema et

al. 2004]. KT86 cells were washed with PBS, resuspended at 2×10^6 cells/mL and labeled

with 300 nM CFSE (BD Horizont) for 15 min at 37°C. The reaction was stopped by the

addition of an equal volume of fetal bovine serum (FCS), followed by a 2-min incubation

at room temperature. CFSE-labeled target cells were washed twice and resuspended in

RPMI complete medium. The cell concentration was adjusted to 5×10^5 cells/mL, and

100 µL/tube were plated in Eppendorf tubes. YTS cells were added at different

effector:target ratios. Tubes were incubated in a humidified atmosphere of 5% CO2 and

37°C for 3 hours. For dead cells staining, propidium iodide (1 µg/mL) was added, and

samples were mixed properly and directly analyzed by flow cytometry using a BD

FACSAria II flow cytometer. 10000 events were acquired for each sample. Double

positive events for CFSE and PI were considered dead target cells. The percentage of

specific lysis was calculated as follows: specific lysis (%) = [(Double positive cells -

spontaneous dead cells) / CFSE positive cells] x 100.

FACS-based conjugation assay

YTS cells were washed with PBS, resuspended to 1×10^6 cells/ml and labeled with 1.5

mM Cell TrackerTM Deep Red (Invitrogen) for 20 min at 37° C. The reaction was

stopped by the addition of an equal volume of fetal bovine serum, followed by a 5 min

incubation at room temperature. After two washes with PBS, the YTS-labeled cells were

resuspended in RPMI complete medium. Target cells were labeled with 300 nM CFSE

(BD Horizon) as described above. Cells were combined at an effector to target ratio of

2:1 or 10:1 and incubated at 37º C for different periods, as indicated. The samples were

gently vortexed for 3 s and immediately fixed with 2% paraformaldehyde (PFA). Samples

were run in triplicates and 50000 events were counted for each replicate. The frequency

of double positive events was determined within the CFSE positive population using

FlowJo software. The following gating strategy was used: First, cells were gated to

exclude debris. Compensation adjustments were made on this population using single-

positive cells stained for either Cell TrackerTM Deep Red or CFSE. Gates were set to

differentiate between the double positives from the single positives and double negatives.

Immunofluorescence

KT86 were labeled with 300 nM CFSE (BD Horizon) as described above. Conjugates

between YTS cells and KT86 at a 2:1 ratio were established in suspension for 15 min at

37 ° C and adhered to poly-lysine-coated glass slides (Polyprep; Sigma-Aldrich) for

additional 15 min. In experiments evaluating NK cell activation by means of specific

activating receptors, poly– l -lysine – coated glass slides were coated with anti-CD28, or

ICAM-1 by overnight incubation in PBS. Slides were washed, and NK cells were

incubated on the slide for 30 min at 37 ° C. Non adherent cells were washed and cells

adhering to the slide were fixed with 4% PFA at room temperature or methanol at -20°

C. Fixed cells were permeabilized and blocked with 0.3%Triton X-100/1% bovine serum

albuminin/PBS, pH 7.4, for 10 min. Then, they were incubated for 2 h with mouse

monoclonal antibody anti CIP4 (BD transduction-612556), anti-LFA-1 (Biolegend

301202) or anti AKAP350 (Schmidt et al, 2001) and rabbit monoclonal antibody anti γ-

tubulin (Sigma-T5192) . The coverslips were washed, incubated for 1 h with the

secondary antibodies conjugated to Alexa 488, Alexa 560 or Alexa 633 or with

phalloidin-Alexa 568 (Molecular probes-A34055, 1:200) for actin staining and with 4′,6-

diamidino-2-phenylindole (DAPI) and mounted with ProLong (Invitrogen). Fluorescence

was detected using LSM880 confocal with an ObserverZ1 inverted microscope. Serial

optical 0.2 μm thick sections were collected in the z-axis. Z-stacks were built, and

projections were obtained using ImageJ tools. In preparing the figures, adjustment in

brightness and contrast were equally applied to the entire images using Adobe Photoshop

software to improve visualization of fluorescence.

Image analysis

Centrosome polarization during IS formation. YTS:KT86 conjugates were stained with

anti γ-tubulin and DAPI, for centrosome and nucleus identification. The IS was defined

as the cell to cell contact region identified in the images obtained by differential

interference microscopy (DIC). The cell perimeter of the YTS cell forming the IS was

drawn using the same images. The centrosome region was delimited by setting a threshold

on the γ-tubulin channel to define a mask, which was used to automatically outline the

centrosomal pixels. The IS, centrosome, and cell centroid were determined using the

appropriate ImageJ tool and the distances from the cell or the centrosome centroid to the

IS centroid were calculated. Alternatively, for analysis of centrosome translocation during

specific LFA-1 or CD28 activation, the surface of YTS interaction with the ICAM-1 or

anti-CD28 coated slides was considered as the IS site, and the distance was measured at

the *(y,z)* plane where the centrosome was best visualized. At least 30 cells from three

independent experiments were analyzed for each condition.

For protein accumulation at the IS, YTS cells conjugated with KT86 cells were analyzed.

We used DIC images for IS and cell selections, as described above. We measured

fluorescence intensity at YTS IS and at the YTS cell (T). The percentage of protein

accumulation was calculated as the ratio IS*100/T.

Centrosomal localization of CIP4 or AKAP350 was determined in images obtained by

confocal microscopy by setting a threshold on the γ-tubulin channel as described above,

and a threshold on CIP4 or AKAP350 channels to define a mask to automatically outline

total voxels for each protein staining. Average and total intensity of fluorescence in CIP4

or AKAP350 channels was measured in each region of interest, and the percentage of

centrosomal levels for each protein calculated.

Golgi levels of AKAP350 were determined in images obtained by confocal microscopy

by setting a threshold on GM130 channel to define a mask, which was used to

automatically outline the Golgi voxels. Total intensity of fluorescence in the AKAP350

channel was measured in each region of interest. In addition, total AKAP350 levels were

determined in a region delimited by a mask defined using the AKAP350 channel, and the

percentage of AKAP350 fluorescence at the Golgi apparatus calculated. AKAP350

colocalization with GM130 in AKAP350GBD cells was determined by estimating the

Pearson´s coefficient using the Image J colocalization tools.

Immunoblotting

Cells were harvested at 400 g for 5 min at room temperature and washed with cold PBS.

Pelleted cells were lysed in ice-cold lysis buffer (50 mM Tris-HCl [pH 7.5], 100 mM

NaCl, 15 mM EDTA and 1% Triton X-100, supplemented with protease inhibitors) and

subjected to two freeze–thaw cycles. Lysates were centrifuged at 1000 g for 5 min at 4°

C, and the clear supernatants were conserved. Total protein concentration was measured

according to Lowry et al. (1951). Samples were heated for 10 min at 90 °C in sample

buffer (20 mM Tris-HCl, pH 8.5, 1% SDS, 400 μM DTT, 10% glycerol). Samples

containing equal amounts of proteins were subjected to SDS polyacrylamide gel

electrophoresis (6–10% gradient to assess AKAP350 levels or 10% otherwise). The

proteins in the 6% gel were transferred to nitrocellulose membranes (Amersham

Pharmacia Biotech), whereas the 10% gel proteins were transferred to polyvinyl

difluoride membranes (Perkin Elmer Life Sciences). Blots were blocked with 5% nonfat

dry milk in PBS, 0.3% Tween-20 (PBS-Tween). Nitrocellulose blots were then probed

with the monoclonal mouse anti-AKAP350 antibody (1:500), and polyvinyl difluoride

membranes were probed with mouse monoclonal antibodies anti-α-tubulin (Sigma-

T9026, 1:5000), anti-γ-tubulin (Sigma-T6557, 1:2000) or anti-CIP4 (BD transduction-

612556, 1:500). The blots were washed and incubated with the horseradish peroxidase-

conjugated corresponding secondary antibodies, and bands were detected by enhanced

chemiluminescence (Pierce, Thermo Scientific). Autoradiographs were obtained by

exposing the blots to Kodak XAR film. The bands were quantitated by densitometry using

the NIH Image J program.

Preparation of centrosome and microtubule enriched subcellular fractions.

Centrosome-enriched fractions were prepared by centrifugation on a Fycoll cushion,

using a method based on Blomberg-Wirschell and Doxsey [1998]. Briefly, cells were

treated with Nocodazole (1.65 μM) and Cytochalasin (390 ηM) for 30 min, and washed

with PBS. Cells were resuspended and gently sonicated in a lysis buffer containing 1Mm

TRIS-HCl [pH 8], 0.5% β-Mercaptoethanol, 0.5% Triton X-100 and protease inhibitors.

Components were added to cell lysates to achieve a final concentration of 10 mM PIPES,

1 mM EDTA. In order to eliminate unbroken cells and nuclear fractions, cell extracts

were centrifuged at 1,500 g for 15 min. The supernatant was filtered through a nylon mesh

and loaded on the top of a 20% (w/v) Ficoll cushion in buffer 10 mM PIPES/1 mM EDTA

with 0.1% Triton X-100 and tubes were spun at 25,000 g in a swinging bucket rotor.

Centrosomes were taken from the top of the Ficoll cushion using a Pasteur pipet, and

CIP4 levels at this fraction were analysed by western blot.

Cellular extracts enriched in stable microtubules were prepared based on a non-ionic

detergent extraction protocol described by Brown et al (1992). A total of 4×10^6 YTS cells

were harvested and added on plastic wells coated with ICAM-1 and CD28 for 30 minutes

at 37°C. Cells were rinsed once with PBS and once with PHEM (60 mM Pipes, 25 mM

Hepes, 10 mM EGTA, 2 mM MgC12, pH 6.9) and then extracted for 5 min in PHEM

containing 0.1% Tritón X-100, 10 µM taxol and a mixture of protease inhibitors. Then,

the cells were washed with PHEM without Tritón and scrapped with a 100 µL of sample

buffer. The protein-contained sample buffer was collected into eppendorf tubes and

incubated for 10 minutes at 90°C . The proteins associated with microtubules were

analyzed by western blot.

YTS cells treatments

In order to assess the role of microtubules in the IS formation, YTS cells were harvested

and 2.10^6 cells were treated for 1 hour with nocodazole (17 µM) on ice or for 1 hour or

with taxol (10 µM) at room temperature and then allowed to form conjugates with KT86

cells at a 2:1 ratio for 15 min at 37° C. In order to evaluate the role of the Golgi apparatus

in IS formation, YTS cells were treated with Brefeldin A (5 µg/ml) for 30 min at 37 °C.

After Brefeldin A treatment, YTS cells were allowed to form conjugates with KT86 cells

at a 2:1 ratio for 15 min at 37 °C. YTS cells without treatment were used as control for

each experiment. Conjugates were analyzed by confocal microscopy as described above.

Ice recovery assay

A total of $2x10^5$ YTS cells were placed on poly-l-lysine-coated coverslips coated with

ICAM-1 and CD28 and incubated for 30 minutes at 37°C. Then, coverslips were

incubated on ice 50 min and allowed to recover at room temperature. For analysis of

Golgi and centrosome nucleation of microtubules, cells were recovered for 3 or 7 minutes

and immediately treated 45 s with extraction buffer (60 mM PIPES, 25 mM HEPES, 10

mM EGTA, 2 mM MgCl2, 0.1% Tritón X-100, pH 6.9, supplement with 0.25 nM

nocodazole and 0.25 nM taxol). Cells were then fixed with methanol and stained with

antibody anti GM130 (Abcam-EP892Y) and antibody anti α-tubulin (Sigma-T9026).

Acknowledgments

We thank Dr. Andrés Zucchetti for his critical discussion of the manuscript. This work

was supported by Grants PUE 0089 from CONICET and PICT2015-2755 and PICT2016-

4758 from ANPCyT. The funders had no role in study design, data collection and

analysis, decision to publish, or paper preparation.

Competing interests

There are not competing interests to declare.

References

Aspenström PA. Cdc42 target protein with homology to the non-kinase domain of FER

has a potential role in regulating the actin cytoskeleton. *Curr Biol*. 1997; 7: 479-87. DOI:

https://doi.org/10.1016/S0960-9822(06)00219-3

Aspenström P, Richnau N, Johansson AS. The diaphanous-related formin DAAM1

collaborates with the Rho GTPases RhoA and Cdc42, CIP4 and Src in regulating cell

morphogenesis and actin dynamics. Exp. Cell Res. 2006; 312:2180-2194. DOI:

https://doi.org/10.1016/j.yexcr.2006.03.013

Banerjee PP, Pandey R, Zheng R, Suhoski MM, Monaco-Shawver L, Orange JS. Cdc42-

interacting protein-4 functionally links actin and microtubule networks at the cytolytic

NK cell immunological synapse. J Exp Med. 2007; 204:2305-20. DOI:

10.1084/jem.20061893

Blomberg-Wirschell, M. & Doxsey, S. J. Rapid isolation of centrosomes. *Methods*

Enzymol. **298**, 228–38 (1998). DOI: 10.1016/s0076-6879(98)98022-3

Brown ACN, Oddos S, Dobbie IM, Alakoskela J-M, Parton RM, et al. (2011)

Remodelling of Cortical Actin Where Lytic Granules Dock at Natural Killer Cell Immune

Synapses Revealed by Super-Resolution Microscopy. PLoS Biol 9(9): e1001152. DOI:

10.1371/journal.pbio.1001152

Brown A, Slaughter T, Black MM. Newly Assembled Microtubules Are Concentrated in

the Proximal and Distal Regions of Growing Axons. J Cell Biol. 1992; 119: 867-882.

DOI: 10.1083/jcb.119.4.867.

Butler B, Cooper JA. Distinct roles for the actin nucleators Arp2/3 and hDia1 during NK-

mediated cytotoxicity. Curr Biol 2009; 19: 1886–1896. DOI: 10.1016/j.cub.2009.10.029.

Carlin LM, Evans R, Milewicz H, Fernandes L, Matthews DR, Perani M, Levitt J,

Keppler MD, Monypenny J, Coolen T, Barber PR, Vojnovic B, Suhling K, Fraternali F,

Ameer-Beg S, Parker PJ, Thomas NS, Ng T. A targeted siRNA screen identifies

regulators of Cdc42 activity at the natural killer cell immunological synapse Sci Signal.

2011; 4:ra81. DOI: 10.1126/scisignal.2001729

Carpier JM et al. Rab6-dependent retrograde traffic of LAT controls immune synapse

formation and T cell activation. J. Exp. Med. (2018) 215, 1245–1265. DOI:

10.1084/jem.20162042

Chabin-Brion K, Marceiller J, Perez F, Settegrana C, Drechou A, Durand G, Poüs C . The

Golgi complex is a microtubule-organizing organelle. Mol Biol Cell. 2001; 12:2047-60.

DOI: 10.1091/mbc.12.7.2047

Chardin P, McCormick F. Brefeldin A: The Advantage of Being Uncompetitive. Cell.

1999; 97:153-155. DOI: 10.1016/s0092-8674(00)80724-2

Chen X, Allan DS, Krzewski K, Ge B, Kopcow H, Strominger JL. CD28-stimulated

ERK2 phosphorylation is required for polarization of the microtubule organizing center

and granules in YTS NK cells. Proc. Natl. Acad. Sci. USA. 2006; 103:10346–10351.

DOI: 10.1073/pnas.0604236103

Chuang, L.T.; Lotzová, E.; Heath, J.; Cook, K.R.; Munkarah, A.; Morris, M.; Wharton,

J.T. Alteration of lymphocyte microtubule assembly, cytotoxicity, and activation by the

anticancer drug taxol. Cancer Res. 1994, 54, 1286–1291.

Derry WB, Wilson L, Jordan MA. Substoichiometric Binding of Taxol Suppresses

Microtubule Dynamics. Biochemistry 1995,34, 2203-2211. DOI: 10.1021/bi00007a014

Drexler, H. G., Y. Matsuo. 2000. Malignant hematopoietic cell lines: in vitro models for

the study of natural killer cell leukemia-lymphoma. Leukemia 14: 777-782. DOI:

10.1038/sj.leu.2401778

El Din El Homasany BS, Volkov Y, Takahashi M, Ono Y, Keryer G, Delouvée A, Looby

E, Long A, Kelleher D. 2005. The scaffolding protein CG-NAP/AKAP450 is a critical

integrating component of the LFA-1-induced signaling complex in migratory T cells. J

Immunol. 175:7811-8. DOI: 10.4049/jimmunol.175.12.7811

Florian S, Mitchison TJ. Anti-Microtubule Drugs. Methods Mol Biol. 2016;1413:403-21.

doi: 10.1007/978-1-4939-3542-0_25. DOI: 10.1007/978-1-4939-3542-0_25

Fong A, Durkin A, and Lee H. Int. The Potential of Combining Tubulin-Targeting

Anticancer Therapeutics and Immune Therapy. J. Mol. Sci. 2019, 20, 586;

doi:10.3390/ijms20030586. DOI: 10.3390/ijms20030586

Grimaldi AD, Fomicheva M, Kaverina I. Ice recovery assay for detection of Golgi-

derived microtubules. Methods Cell Biol. 2013; 118:401-15. DOI: 10.1016/B978-0-12-

417164-0.00024-0

Hurtado L, Caballero C, Gavilan MP, Cardenas J, Bornens M, Rios RM. Disconnecting

the Golgi ribbon from the centrosome prevents directional cell migration and ciliogenesis.

J Cell Biol. 2011 May 30;193(5):917-33. DOI: 10.1083/jcb.201011014

Jedema, Inge, et al. New CFSE-based assay to determine susceptibility to lysis by cytotoxic T cells of leukemic precursor cells within a heterogeneous target cell population. *Blood* 103.7 (2004): 2677-2682. DOI: 10.1182/blood-2003-06-2070

Katz P, Zaytoun AM and Lee J H Jr. Mechanisms of human cell-mediated cytotoxicity. III. Dependence of natural killing on microtubule and microfilament integrity. J Immunol 1982, 129: 2816-2825.

Kupfer A, Dennert G, Singer SJ. Polarization of the Golgi apparatus and the microtubule-organizing center within cloned natural killer cells bound to their targets. Proc Natl Acad Sci USA 1983; 80: 7224–7228. DOI: 10.1073/pnas.80.23.7224

Kim HS, Takahashi M, Matsuo K, Ono Y. Recruitment of CG-NAP to the Golgi apparatus through interaction with dynein-dynactin complex. Genes Cells. 2007; 12:421-34. DOI: 10.1111/j.1365-2443.2007.01055.x

Kolobova E, Roland JT, Lapierre LA, Williams JA, Mason TA, Goldenring JR. The C-terminal region of A-kinase anchor protein 350 (AKAP350A) enables formation of microtubule-nucleation centers and interacts with pericentriolar proteins. J Biol Chem. 2017; 292: 20394–20409. doi: 10.1074/jbc.M117.806018. DOI: 10.1074/jbc.M117.806018

Lagrue K, Carisey A, Oszmiana A, Kennedy PR, Williamson DJ, Cartwright A, Barthen C, Davis DM.. The central role of the cytoskeleton in mechanisms and functions of the NK cell immune synapse. Immunological Reviews 2013. Vol. 256: 203–221. DOI: 10.1111/imr.12107

Larocca MC, Jin M, Goldenring JR. AKAP350 modulates microtubule dynamics. Eur J Cell Biol. 2006; 85:611-9. DOI: 10.1016/j.ejcb.2005.10.008

Larocca MC, Shanks RA, Tian L, Nelson DL, Stewart DM, Goldenring JR. AKAP350

interaction with cdc42 interacting protein 4 at the Golgi apparatus. Mol Biol Cell. 2004;

15:2771-81. DOI:: 10.1091/mbc.E03-10-0757

Leibfried A, Fricke R, Morgan MJ, Bogdan S, Bellaiche Y. Drosophila Cip4 and WASp

define a branch of the Cdc42-Par6-aPKC pathway regulating E-cadherin endocytosis.

Curr Biol. 2008; 18:1639-48. DOI: 10.1016/j.cub.2008.09.063

Li D, Sun X, Zhang L, Yan B, Xie S, Liu R, Liu M, Zhou J. Histone deacetylase 6 and

cytoplasmic linker protein 170 function together to regulate the motility of pancreatic

cancer cells. Protein Cell. 2014; 5: 214–223. DOI: 10.1007/s13238-013-0010-3

Maia A. R., Zhu X., Miller P., Gu G., Maiato H., Kaverina I. (2013). Modulation of Golgi-

associated microtubule nucleation throughout the cell cycle. Cytoskeleton 70, 32–43.

10.1002/cm.21079. DOI: 10.1002/cm.21079

Mace EM, Dongre P, Hsu H-T, Sinha P, James AM, Mann SS, Forbes LR, Watkin LB,

Orange JS. Cell biological steps and checkpoints in accessing NK cell cytotoxicity.

Immunol Cell Biol. 2014; 92: 245–255. DOI: 10.1038/icb.2013.96

Mattaloni SM, Kolobova E, Favre C, Marinelli RA, Goldenring JR, Larocca MC.

AKAP350 Is involved in the development of apical "canalicular" structures in hepatic

cells HepG2. J Cell Physiol. 2012; 227:160-71. doi: 10.1002/jcp.22713. DOI:

10.1002/jcp.22713

Morrison MM, Wardleworth BN, Askham JM, Markham AF, Meredith DM. EB1, a

Protein Which Interacts With the APC Tumour Suppressor, Is Associated With the

Microtubule Cytoskeleton Throughout the Cell Cycle. Oncogene 1998;17(:3471-7. doi:

10.1038/sj.onc.1202247. DOI: 10.1038/sj.onc.1202247

Murugesan S, Hong J, Yi J, Li D, Beach JR, Shao L, Meinhardt J, Madison G, Wu X,

Betzig E, Hammer JA. Formin-generated actomyosin arcs propel T cell receptor

microcluster movement at the immune synapse. J Cell Biol. 2016; 215:383-399. DOI:

10.1083/jcb.201603080

Ong ST, Chalasani MLS, Fazil MHUT, Prasannan P, Kizhakeyil A, Wright GD, Kelleher

D, Verma NK. Centrosome- and Golgi-Localized Protein Kinase N-Associated Protein

Serves As a Docking Platform for Protein Kinase A Signaling and Microtubule

Nucleation in Migrating T-Cells. Front Immunol. 2018; 9:397. DOI:

10.3389/fimmu.2018.00397

Orange JS, et al. Wiskott-Aldrich syndrome protein is required for NK cell cytotoxicity

and colocalizes with actin to NK cell-activating immunologic synapses. Proc Natl Acad

Sci USA 2002;99:11351–11356. DOI: 10.1073/pnas.162376099

Orange JS, Harris KE, Andzelm MM, Valter MM, Geha RS, Strominger JL. The Mature

Activating Natural Killer Cell Immunologic Synapse Is Formed in Distinct Stages. Proc

Natl Acad Sci USA. 2003; 100:14151-6. DOI: 10.1073/pnas.1835830100.

Ori-McKenney, K. M., Jan, L. Y., & Jan, Y.-N. (2012). Golgi outposts shape dendrite

morphology by functioning as sites of acentrosomal microtubule nucleation in neurons.

Neuron, 76, 921– 930. DOI: 10.1016/j.neuron.2012.10.008

Parry, Richard V., et al. CD28 and inducible costimulatory protein Src homology 2

binding domains show distinct regulation of phosphatidylinositol 3-kinase, Bcl-xL, and

IL-2 expression in primary human CD4 T lymphocytes. The Journal of immunology,

2003, vol. 171, no 1, p. 166-174. DOI: 10.4049/jimmunol.171.1.166

Perez, F., G.S. Diamantopoulos, R. Stalder, and T.E. Kreis. 1999. CLIP-170 highlights

growing microtubule ends in vivo. Cell. 96:517–527. DOI: 10.1016/s0092-

8674(00)80656-x

Rak GD, Mace EM, Banerjee PP, Svitkina T, Orange JS. 2011. Natural killer cell lytic

granule secretion occurs through a pervasive actin network at the immune synapse. PLoS

Biol 9:e1001151. DOI: 10.1371/journal.pbio.1001151.

Rivero S, Cardenas J, Bornens M, Rios RM. Microtubule nucleation at the cis-side of the

Golgi apparatus requires AKAP450 and GM130. EMBO J. 2009, 281016-28. DOI:

10.1038/emboj.2009.47

Roberts-Galbraith RH & Gould KL. Setting the F-BAR: functions and regulation of the

F-BAR protein family. *Cell Cycle* 2010; 9:4091-4097. DOI: 10.4161/cc.9.20.13587

Robles-Valero J, Martín-Cófreces NB, Lamana A, Macdonald S, Volkov Y, Sánchez-

Madrid F. (2010) Integrin and CD3/TCR activation are regulated by the scaffold protein

AKAP450. Blood. 115:4174-84. DOI: 10.1182/blood-2009-12-256222

Sanders AA, Kaverina I. Nucleation and dynamics of Golgi-derived microtubules. Front

Neurosci. 2015;9:431. doi: 10.3389/fnins.2015.00431. DOI: 10.3389/fnins.2015.00431

Schmidt PH, Dransfield DT, Claudio JO, Hawley RG, Trotter KW, Milgram SL,et al.

AKAP350, a multiply spliced protein kinase A-anchoring protein associated with

centrosomes. J Biol Chem. 1999; 274:3055-66. DOI: 10.1074/jbc.274.5.3055

Serrador JM, Cabrero JR, Sancho D, Mittelbrunn, M, Urzainqui A, Sánchez-Madrid F,

2004 HDAC6 Deacetylase Activity Links the Tubulin Cytoskeleton with Immune

Synapse Organization. Immunity, 20, 417–428. DOI: 10.1016/s1074-7613(04)00078-0

Sillibourne JE, Milne DM, Takahashi M, Ono Y, Meek DW. Centrosomal anchoring of

the protein kinase CK1delta mediated by attachment to the large, coiled-coil scaffolding

protein CG-NAP/AKAP450. J Mol Biol. (2002) 322:785–97. DOI: 10.1016/S0022-
2836(02)00857-4.

Takahashi M, Yamagiwa A, Nishimura T, Mukai H, Ono Y. Centrosomal Proteins CG-
NAP and Kendrin Provide Microtubule Nucleation Sites by Anchoring γ-Tubulin Ring
Complex. Mol Biol Cell. 2002; 13: 3235–3245. DOI: 10.1091/mbc.E02-02-0112

Tian L, Nelson DL, Stewart, DM. Cdc42-interacting protein 4 mediates binding of the
Wiskott-Aldrich syndrome protein to microtubules. *J. Biol. Chem.* 2000; 275:7854-7861.
DOI: 10.1074/jbc.275.11.7854

Tonucci FM, Almada E, Borini-Etichetti C, Pariani A, Hidalgo F, Rico MJ, Girardini J,
Favre C, Goldenring JR, Menacho-Marquez M, Larocca MC. Identification of a CIP4
PKA phosphorylation site involved in the regulation of cancer cell invasiveness and
metastasis. Cancer Lett. 2019; 461:65-77. DOI: 10.1016/j.canlet.2019.07.006.

Tonucci FM, Ferretti A, Almada E, Cribb P, Vena R, Hidalgo F, Favre C, Tyska MJ,
Kaverina I, Larocca MC. Microtubules regulate brush border formation. J Cell Physiol.
2018; 233:1468-1480. DOI: 10.1002/jcp.26033.

Tonucci FM, Hidalgo F, Ferretti A, Almada E, Favre C, Goldenring JR, Kaverina I,
Kierbel A, Larocca MC. Centrosomal AKAP350 and CIP4 act in concert to define the
polarized localization of the centrosome and Golgi in migratory cells. J Cell Sci. 2015;
128:3277-89. DOI: 10.1242/jcs.170878.

Tsujita K, Suetsugu S, Sasaki N, Furutani M, Oikawa T, Takenawa T. Coordination
between the actin cytoskeleton and membrane deformation by a novel membrane
tubulation domain of PCH proteins is involved in endocytosis. J. Cell Biol. 2006;
172:269-279. DOI: 10.1083/jcb.200508091

Zilberman, Y., Ballestrem, C., Carramusa, L., Mazitschek, R., Khochbin, S., and

Bershadsky, A. Regulation of microtubule dynamics by inhibition of the tubulin

deacetylase HDAC6. J Cell Sci 2009; 122, 3531–3541. DOI: 10.1242/jcs.046813

Supplementary figure 1. Reduction of AKAP350 expression does not affect YTS:KT86 conjugate formation. YTS control and AKAP350KD cells were stained with cell tracker Deep red and mixed with KT86 cells stained with CFSE. Conjugate formation was analyzed by flow cytometry. The conjugates were determined as CFSE/cell tracker Deep red double positive events. Effector and target cells were mixed at a 2:1 (A) or 10:1 (B) ratio and incubated at 37 °C and 5 % CO_2 for the indicated periods (A) or for 30 min (B). Bars represent the mean percentage of conjugates formation from three independent experiments. Error bars represent SEM. *$p < 0.05$.

Supplementary figure 2. AKAP350 expression and localization at the Golgi apparatus is required for proper microtubule acetylation. Acetylated α-tubulin and total α-tubulin levels in control, AKAP350KD (A) and AKAPGBD (B) cell lysates were analyzed by WB. The bars chart represents the mean density of the band corresponding to acetylated α-tubulin relative to α-tubulin for three independent experiments. Error bars represent SEM. *p< 0.05.