Genetic diversity and population structure of razor clam *Sinonovacula constricta*

in Ariake Bay, Japan, revealed using RAD-Seq SNP markers.

Ryo Orita[1,*], Yukio Nagano[2], Yoshio Kawamura[1], Kei Kimura[1], and Genta Kobayashi[1]

[1]*Faculty of Agriculture, Saga University, 1 Honjo-machi, Saga, 840-8502, Japan*

[2]*Analytical Research Center for Experimental Sciences, Saga University, 1 Honjo-machi,*

Saga, 840-8502, Japan

*Corresponding author: Ryo Orita, E-mail: ss7427@cc.saga-u.ac.jp

Abstract

The razor clam *Sinonovacula constricta* is a commercially important bivalve in Japan.

The current distribution of this species in Japan is limited to Ariake Bay, where the

fishery stock is declining. It is necessary to understand the genetic population structure of

this species in order to restore the fishery stock while preserving the genetic diversity of

the clam. Here, we report for the first time the genetic population structure of *S. constricta*

in Ariake Bay, Japan. Paired-end restriction site-associated DNA sequencing (RAD-Seq)

analyzed samples of *S. constricta* collected from seven mudflats located along Ariake

Bay. Two different genetic populations exist in Ariake Bay, one inhabiting wild habitats

and the other inhabiting the transplanted area of artificial seedlings. Our results suggest

that genetic differentiation occurred between these two populations (F_{st} value = 0.052),

and a high level of genetic differentiation is maintained between the two groups. In the

future, monitoring the interbreeding status of the two genetically distinct populations and

the genetic differentiation within each population is important for conserving the genetic

diversity of *S. constricta* in Japan.

Introduction

The technique of artificially producing seedlings and releasing them into fishing grounds

is an effective method of aquaculture enhancement. This technique has been

implemented with a variety of fishery species (e.g., finfish [1, 2], flounder [3], shellfish

[4-7]). In artificial seed production, large numbers of seedlings are often produced by a

few broodstocks and released into natural seawater, which reduces the genetic diversity

of wild populations in the field. Therefore, when seedlings are produced and released into

natural seawater, there is a need to understand the genetic structure within the wild

population in order to preserve genetic diversity.

The razor clam *Sinonovacula constricta* (Lamarck, 1818) is an important

economic species in East Asian coastal waters [8-10] and is widely farmed in China [10].

This species was once distributed from central to western Japan [11-12] but is now

restricted to the mudflats around Ariake Bay on the west coast of Kyushu in western

Japan. In Japan, razor clams are not farmed, and the catch depends on the amount of

resources in natural seawaters. According to a report by Ministry of the Environment,

Japan [13], the annual catch of *S. constricta* in Ariake Bay often recorded more than

1000 tons per year from the 1900s to the early the 1920s, but it decreased to less than 500

tons in most years from the latter half of the 1920s to the latter half of the 1980s, and no

catches have been made since 1994 because of the marked decline in the abundance of

this species. In order to restore the *S. constricta* resources, the Saga Prefectural Ariake

Fisheries Research and Development Center has developed seed production techniques

[14, 15], and since 1988, artificial seedlings of *S. constricta* have been released into the

transplanting areas established in natural seawaters [16] and have been conserved in

transplanting areas as populations to produce subsequent generations. However, the

genetic population structure of *S. constricta* within Ariake Bay has not been examined.

In recent years, technological innovations have made the genome-wide analysis of

non-model organisms more widely available, and these techniques are useful for

examining the genetic population structure of non-model aquatic organisms [17-20]. In

the present study, to clarify the genetic population structure of *S. constricta* in Ariake Bay,

we conducted field sampling surveys at 11 mudflats located along the bay in 2019 (Fig. 1)

and used restriction-site-associated DNA sequencing (RAD-Seq) to analyze population

genetics and phylogenetics. We propose guidelines concerning the optimal seed

production and seed release methods for the conservation of the genetic diversity of *S.*

constricta in Japan.

Results

Sampling data.   We collected 28 razor clams from seven of the 11 mudflats and found

no razor clams on the remaining four mudflats (Table 1, Fig. 1). The mean CPUE of the

seven mudflats where individuals could be collected was 3.2 ($\pm$ 3.0 SD) ind. person^{-1}

hour^{-1}, and the range of CPUE at each site was 0.3 to 8.5 ind. person^{-1} hour^{-1}. The shell

length of the collected individuals was 39.7 to 94.6 mm, implying that they contained

specimens from 1 to 5 years old. The mean shell length of individuals collected from the

transplanted area was 86.0 ($\pm$ 4.5 SD) mm for the 3-year-old seedlings produced in 2016

(i.e., sample ID: 2016.1-2016.7) and 92.9 (± 1.1 SD) mm for the 5-year-old seedlings

produced in 2014 (i.e., sample ID: 2014.1-2014.5).

RAD-Seq data.   We used 30 individuals for RAD-Seq analysis, including 16

individuals collected from wild habitats, 12 individuals collected from the transplanted

area of artificial seedlings, and 2 individuals stored at the Saga Prefectural Ariake

Fisheries Research and Development Center (Table 1). In all, 401,109,332 raw reads

were generated from the HiSeq X run, and 249,040,001 reads were retained after

discarding reads with low quality and adapters. The final dataset included 137,246 loci,

with an average of 679.38 bp (SE ± 0.18) of genotyped sites per locus.

Population genetics structure.   The genetic relationships among sampling sites were

clearly divided into individuals collected from wild habitat and those from the

transplanted area of artificial seedlings along the axis of PC1 (Fig. 2a). The PC2 axis

represents the differences within each habitat type. In the PCA plot, the coordinates of the

transplanted samples were more widely scattered than those of the wild samples were,

showing more diverse genetic relationships among transplanted samples. When

evaluated by shell length (Fig. 2b), genetic diversity was higher in the 75–90 mm size

group than that in the younger age group with smaller shells. The results of admixture

analysis also showed a significant genetic difference between the individuals collected

from the wild versus those collected from the transplanted areas of artificial seedlings

(Fig. 3). The F_{st} value between populations consisting of individuals collected from the

wild and the transplanted areas was 0.052. Owing to the calculation of the pairwise F_{st}

values for each individual within the two populations, there was no significant difference

in the mean values between these populations (Fig. 4). The pairwise F_{st} values within the

population were more varied in the transplanted area (from 0.44 to 0.64) than those in the

wild habitat (from 0.53 to 0.56). There were no significant differences in shell

morphology between these two genetically different populations (Supplementary Fig.

S1).

Phylogenetic tree analysis.   ML tree and SVDquartets analysis of RAD-Seq data

showed that the phylogenies were highly consistent with each other (Fig. 5a and 5b). The

obtained phylogenetic trees were clearly divided into individuals collected from the wild

habitat and those collected from the transplanted area of artificial seedlings, with the

exception of one sample (i.e., sample ID: 2016.7).

Discussion

Genetic population structure of *S. constricta* in the Ariake Bay, Japan.   All our

analyses of the RAD-Seq data showed that individuals collected from wild habitats were

genetically different from those collected from sites where artificial seedlings were

transplanted. Only one sample collected from the area transplanted in 2016 (sample ID:

2016.7) showed different results from other seedlings collected in the same area, and

belonged to a population consisting of individuals collected from the wild habitat (Fig. 3,

Fig. 5a and 5b). Since this individual had a distinctly larger shell length than other

3-year-old seedlings in the same area, and its size was the same as that of 5-year-old

seedlings (Table 1), it was likely a wild individual inhabiting the transplanted area prior

to the 2016 transplantation. Therefore, we consider that there were two genetically

distinct populations in Ariake Bay in 2019.

The reason for the differences between these two populations is the different

parental origins of the seedlings. The parental origins can be divided into the Nanaura

mudflat and the Kashima mudflat for a number of reasons. Artificial seedlings of *S.*

constricta have been released by the Saga Prefectural Ariake Fisheries Research and

Development Center into transplanting areas established in several mudflats along the

coast of Ariake Bay. The seedlings collected from the transplanted area in this study

(Takori_b and Takori_c) originated from parents at the Nanaura mudflat (hereinafter

referred to as Nanaura lineage population), since the parents were collected from the

Nanaura mudflat before 2010 and released into the transplanting areas of the Takori

mudflat. In contrast, individuals collected from the wild habitat of Kashima, Nanaura,

and Takori mudflats are likely to have originated from parents of Kashima mudflat

(hereinafter referred to as the Kashima lineage population). In 2017 and 2018, the

seedlings produced by parents collected from Kashima mudflat were released into new

transplanting areas at Takori and Nanaura mudflats due to declining populations in wild

and transplanted areas in these mudflats. Therefore, the 1- to 2-year old individuals

collected in wild habitats in 2019 were likely the offspring of the transplanted Kashima

seedlings in 2017 and 2018.

Artificial transplantation of aquatic organisms has been reported to alter the

geographic genetic structure of wild populations [21-24], including that of *S. constricta*

(in this case, along the Chinese coast) [25]. Anthropogenic transplantation likely strongly

influenced the distribution of the genetic population structure of *S. constricta* in Japan.

Comparison of the genetic distribution of species with a life history similar to that of *S.*

constricta but those that are not exploited for fisheries can enhance our understanding of

anthropogenic impacts on the genetic distribution of *S. constricta* in the Ariake Bay.

Genetic differentiation among and within populations.   For artificial seed

production and release of seedlings, it is necessary to understand the distribution of

populations with different genetic structures in natural seawater. Therefore, it is

necessary to select broodstocks and release areas according to the degree of genetic

differentiation among and within populations. F_{st} values are widely used as a measure of

the degree of genetic differentiation among and within populations [26-28]. According to

Niu et al. [25], the F_{st} value among populations of S. constricta at 10 sites along the coast

of mainland China is 0.044. The F_{st} value between the two populations in the present

study was 0.052, suggesting that a similar level of genetic differentiation occurred

between the two populations in Ariake Bay, Japan.

For these two populations, the genetic differentiation within each population was

maintained at a high level, and its variation was greater within the population consisting

of individuals collected from the transplanted area (Fig. 4). There are three possible

reasons why the variation in the degree of genetic differentiation within populations

differed between the two populations. The first is that the degree of variation in genetic

differentiation in these populations originally may have been different. Second, the

methods of seedling production may lead to greater variation of genetic differentiation

within populations in the transplanted area. In seedling production, broodstocks were

randomly selected with closely related individuals collected from the same transplanted

area or distantly related individuals collected from different transplanted areas. After

stimulating the mature individuals to spawn, 20 to 30 individuals were placed in the same

tank to produce fertilized eggs. Thus, the degree of genetic differentiation may have been

more varied in the transplanted populations than in the wild habitat population. Third, the

sample size may have been too small to accurately estimate the population. However, in

the current situation in which the number of wild populations is small, it is difficult to

verify the validity of a larger sample size.

Guidelines concerning the best seed production and seed release methods.   The

current abundance of *S. constricta* is estimated to be very low based on the CPUE results

of this study (Table 1). These results suggest that artificial seed production and release of

seedlings could be an effective strategy for the recovery of *S. constricta* abundance in

Ariake Bay, Japan. The present study suggested genetic differentiation between the two

populations of *S. constricta* in Ariake Bay, and a high level of genetic differentiation

within each population was maintained. Therefore, maintaining the two populations

without interbreeding seems to be an appropriate conservation strategy. As of 2019, the

Kashima lineage population is widely distributed in wild habitats, whereas the Nanaura

lineage population is limited to the Takori mudflats. In order to avoid hybridization with

the Kashima lineage population, the release of seedlings of the Nanaura lineage

population should continue to be restricted to the mudflats within the current cove, with a

transplanting area.

To produce seedlings to slow down the decline in genetic diversity within a

population, mathematical models have evaluated five strategies: I) avoidance of line

breeding, II) selective use of wild-born individuals, III) line breeding, IV) avoidance of

artificial breeding of males, and V) selective use of wild-born males [29]. Among them,

strategy II has been considered the most effective [29, 30]. The best way to produce

seedlings of the Kashima lineage population would be to follow strategy II; all parental

individuals would be randomly collected from wild habitats and previously produced

seedlings would not be used as parents. In contrast, the Nanaura lineage population has

already been maintained by a similar approach to strategy I; all parental individuals will

be randomly collected from transplanted areas without excluding released seedlings. This

approach gradually reduces the genetic diversity within the population. Indeed, lower

pairwise F_{st} values were detected in the transplanted population (Fig. 4). Therefore, in the

Nanaura lineage population, countermeasures should be taken to increase the number of

parents used for seed production or to avoid using the produced seedlings as parents of

the next generation.

In conclusion, we detected two genetically distinct populations of *S. constricta* in

the Ariake Bay, Japan. Additionally, our results suggests that genetic differentiation

within each population is maintained at a high level. In the future, monitoring the

interbreeding status of the two genetically distinct populations and the genetic

differentiation within each population is important to enhance the stock of *S. constricta* in

Japan by producing and releasing seedlings, while preserving genetic diversity. As there

was no significant difference in the morphology of the shells of these genetically distinct

205 populations (Supplementary Fig. S1), DNA markers may be effective for discriminating

206 between the two populations.

208 **Methods**

209 **Sampling and genomic DNA extraction.** Sampling of the razor clam *S. constricta*

210 was conducted at 11 mudflats located along Ariake Bay (Fig. 1). Information on the

211 sampling day and habitat types at each site is shown in Table 1. Habitats are divided into

212 two types: transplanted area of artificial seedlings, and wild habitat. As mentioned above,

213 artificial seedlings of *S. constricta* have been released into the transplanting areas

214 established in natural seawaters since 1988. Therefore, artificially produced individuals

215 live within the transplanted area, and the wild habitat would be populated by natural

216 individuals and offspring of seedlings from the transplanted areas. The ages of

217 individuals in the transplanted areas were estimated by calculating backwards from the

218 year of release. At each site, razor clams were explored by hand and their abundance was

219 assessed using the catch per unit effort index (CPUE), calculated as the number of clam

individuals collected divided by the number of explorers and sampling time (i.e., ind.

person^{-1} hour^{-1}). The samples were stored at -80°C until genomic DNA extraction. In

addition to the collected individuals, the oldest seedlings (produced in 2005, $n = 2$) stored

at the Saga Prefectural Ariake Fisheries Research and Development Center, were

included in the RAD-Seq analysis for comparison with seedlings produced in the past.

Genomic DNA was extracted from finely chopped foot tissue of the specimens using

DNAs-ici!-F (RIZO, Tsukuba, Japan), and the concentration of the DNA was assessed

using a Qubit 3.0 Fluorometer (Thermo Fisher Scientific).

RAD-Seq analysis.   The preparation for *the Eco*RI RAD-Seq library has been

described previously [28], based on the established method [31]. The library was

sequenced with 150 bp paired-end reads in one lane of an Illumina HiSeq X (Illumina,

San Diego, CA, USA). The RAD-Seq data were analyzed using the Stacks program

(version 2.52) [32]. The data were quality filtered using the process_radtags program in

the Stacks package (-r -c -q, - adapter_1, adapter_2, and adapter_mm2). Filtered

sequences were registered in the DRA (DRA accession number: DRA010922). The data

were aligned with the reference genome of *S. constricta* [33] (NCBI/DDBJ BioProject

number: PRJNA508451) using BWA-MEM (version 0.7.17) [34]. Aligned data were

analyzed using the ref_map.pl script of the Stacks package.

Population genetics analysis.   Principal component analysis (PCA) was conducted

based on a variant call format (VCF) file using the SNPrelate program (v.1.18.1) [35].

The VCF file was created by the populations program of the Stacks package

(--rm-pcr-duplicates, -R 0.7, --write-single-snp, --min-maf 0.05) [36]. The admixture

analysis was conducted based on the PLINK file using ADMIXTURE (v. 1.3.0) [37]. The

PLINK file was created with the same Stacks populations program that created the VCF

file [38]. The genetic diversity of razor clams among and within populations was assessed

using AMOVA F_{st} values obtained from 'populations.fst_files', which were calculated

by the Stacks populations program. To visualize the distribution of all data points,

pairwise F_{st} values between individuals in each population are shown in boxplot with

beeswarm using R (package: 'beeswarm') [39].

Phylogenetic tree analysis.   Phylogenetic tree analysis was performed using two

different approaches: the maximum likelihood (ML) method and SVDquartets analysis.

Multiple alignments within the cluster were obtained using the Stacks package

populations program mentioned earlier. The phylogenetic tree based on the ML method

was constructed using the RAxML program (version 8.2.12) [40] (-f = a, -x = 12,345, -p =

12,345, -N (bootstrap value) = 1,000, and -m = GTRGAMMA). The SVDquartets

analysis based on the coalescent model was computed as implemented in PAUP ver. 4.0a

167 using multiple alignments [41, 42]. The number of bootstrap analyses replicates was

1,000. In each analysis, the oldest samples (i.e., sample ID, 2005.1; 2005.2), were used as

the root.

References

1. Fushimi, H. Production of juvenile marine finfish for stock enhancement in Japan.

Aquaculture **200,** 33–53. https://doi.org/10.1016/S0044-8486(01)00693-7 (2001).

2. Masuda, R. & Tsukamoto, K. Stock enhancement in Japan: review and perspective.

Bull. Mar. Sci. **62,** 337–358.

https://www.ingentaconnect.com/content/umrsmas/bullmar/1998/00000062/000000

02/art00005 (1998).

3. Sekino, M., Saitoh, K., Yamada, T., Hara, M. & Yamashita, Y. Genetic tagging of

released Japanese flounder (*Paralichthys olivaceus*) based on polymorphic DNA

markers. *Aquaculture* **244**, 49–61. https://doi.org/10.1016/j.aquaculture.2004.11.006

(2005).

4. Arnold, W. S. Bivalve enhancement and restoration strategies in Florida, USA.

Hydrobiologia **465,** 7–19. https://doi.org/10.1023/A:1014596909319 (2001).

5. Castell, L. L., Naviti, W. & Nguyen, F. Detectability of cryptic juvenile *Trochus*

niloticus Linnaeus in stock enhancement experiments. *Aquaculture* **144,** 91–101.

https://doi.org/10.1016/S0044-8486(96)01320-8 (1996).

6. McCormick, T. B., Herbinson, K., Mill, T. S. & Altick, J. A review of abalone seeding,

possible significance and a new seeding device. *Bull. Mar. Sci.* **55,** 680–693.

https://www.ingentaconnect.com/content/umrsmas/bullmar/1994/00000055/f002000

2/art00035 (1994).

7. Zohar, Y. *et al.* The Chesapeake Bay blue crab (*Callinectes sapidus*): A

multidisciplinary approach to responsible stock replenishment. *Rev. Fish. Sci.* **16,** 24–

34. https://doi.org/10.1080/10641260701681623 (2008).

8. Funge-Smith, S., Briggs, M. & Miao, W. Regional Overview of Fisheries and

Aquaculture in Asia and the Pacific 2012. *RAP Publication (FAO)* (2012).

9. Mao, Y., Lin, F., Fang, J., Fang, J., Li, J. & Du, M. Chapter 4. Bivalve production in

China. In *Goods and Services of Marine Bivalves.* (eds Smaal, A. C. *et al.*) 51–72

(Springer, Cham, 2019).

10. Ran, Z. *et al.* Fatty acid and sterol changes in razor clam *Sinonovacula constricta*

(Lamarck 1818) reared at different salinities. *Aquaculture* **473,** 493–500.

https://doi.org/10.1016/j.aquaculture.2017.03.017 (2017).

11. Suzuki, T., Inoue, K. & Ozawa, T. Environmental degradation in Ise and Mikawa

Bays after 1960's as viewed from intertidal molluscan community. *Boll. Nagoya Univ.*

Museum **22,** 31–64. http://doi.org/10.18999/bulnum.022.04 (2006).

12. Nakamura, T., Nishida, I., Takada, H., Okuno, M., Minami, M., & Oda, H. Marine

reservoir effect deduced from 14C dates on marine shells and terrestrial remains at

archeological sites in Japan. *Nucl. Instrum. Methods Phys. Res., B* **259**, 453–459.

https://doi.org/10.1016/j.nimb.2007.01.186 (2007).

13. Ministry of the Environment, Japan. *Ariake Sea and Yatsushiro Sea Comprehensive*

Survey Evaluation Committee Report.

https://www.env.go.jp/council/20ari-yatsu/rep061221/all.pdf (2006).

14. Ito, S., Eguchi, T. & Kawahara, I. Rearing experiment on planktonic larvae of the

JackKnife clam, *Sinonovacula constricta*. *Boll. Saga Prefect. Ariake Fish. Res. Dev.*

Cent. **20,** 49–53. https://agriknowledge.affrc.go.jp/RN/2010813185.pdf (2001).

15. Ohkuma, H., Ymagachi, T., Kawahara, I., Eguchi, T. & Ito, S. A study on the

development of techniques for mass production of seeds of jackknife clam,

Sinonovacula constricta. Boll. Saga Prefect. Ariake Fish. Res. Dev. Cent. **22,** 47–54.

https://agriknowledge.affrc.go.jp/RN/2030813210.pdf (2004).

16. Tsukuda, M. *et al.* Variation in the Distribution of the JackKnife Clam, *Sinonovacula*

constricta, on the Muddy Tidal flat of Ariake Sound off Saga Prefecture. *Boll. Saga*

Prefect. Ariake Fish. Res. Dev. Cent. **28,** 47–49.

https://agriknowledge.affrc.go.jp/RN/2010925685.pdf (2017).

17. Holman, L. E., Garcia de la serrana, D., Onoufriou, A., Hillestad, B. & Johnston, I. A.

A workflow used to design low density SNP panels for parentage assignment and

traceability in aquaculture species and its validation in Atlantic salmon. *Aquaculture*

476, 59–64. https://doi.org/10.1016/j.aquaculture.2017.04.001 (2017).

18. Li, Y. H. & Wang, H. P. Advances of genotyping-by-sequencing in fisheries and

aquaculture. *Rev. Fish Biol. Fish.* **27,** 535–559.

https://doi.org/10.1007/s11160-017-9473-2 (2017).

19. Robledo, D., Palaiokostas, C., Bargelloni, L., Martínez, P. & Houston, R.

Applications of genotyping by sequencing in aquaculture breeding and genetics. *Rev.*

Aquac **10,** 670–682. https://doi.org/10.1111/raq.12193 (2018).

20. You, X., Shan, X. & Shi, Q. Research advances in the genomics and applications for

molecular breeding of aquaculture animals. *Aquaculture* **526,** 735357.

https://doi.org/10.1016/j.aquaculture.2020.735357 (2020).

21. Arthington, A. H. Ecological and genetic impacts of introduced and translocated

freshwater fishes in Australia. *Can. J. Fish. Aquat. Sci.* **48,** 33–43.

https://doi.org/10.1139/f91-302 (1991).

22. Habtemariam, B. T., Arias, A., García-Vázquez, E. & Borrell, Y. J. Impacts of

supplementation aquaculture on the genetic diversity of wild *Ruditapes decussatus*

from northern Spain. *Aquacult. Environ. Interact.* **6,** 241–254.

https://doi.org/10.3354/aei00128 (2015).

23. Williams, S. L. & Orth, R. J. Genetic diversity and structure of natural and

transplanted eelgrass populations in the Chesapeake and Chincoteague Bays.

Estuaries **21,** 118–128. https://doi.org/10.2307/1352551 (1998).

24. Yamakawa, A. Y. & Imai, H. PCR-RFLP typing reveals a new invasion of Taiwanese

Meretrix (Bivalvia: Veneridae) to Japan. *Aquat. Invasions* **8,** 407–415.

http://doi.org/10.3391/ai.2013.8.4.04 (2013).

25. Niu, D. H., Feng, B. B., Liu, D. B., Zhong, Y. M., Shen, H. D. & Li, J. L. Significant

genetic differentiation among ten populations of the razor clam *Sinonovacula*

constricta along the coast of china revealed by a microsatellite analysis. *Zool. Stud.*

51, 406–414. http://zoolstud.sinica.edu.tw/Journals/51.3/406.pdf. (2012).

26. DeFaveri, J., Shikano, T., Ghani, N. I. A. & Merilä, J. Contrasting population

structures in two sympatric fishes in the Baltic Sea basin. *Mar. Biol.* **159,** 1659–1672.

https://doi.org/10.1007/s00227-012-1951-4 (2012).

27. Jeffery, N. W. *et al.* RAD sequencing reveals genomewide divergence between

independent invasions of the European green crab (*Carcinus maenas*) in the

Northwest Atlantic. *Ecol. Evol.* **7,** 2513–2524. https://doi.org/10.1002/ece3.2872

(2017).

28. Kato, D. *et al.* Evaluation of the population structure and phylogeography of the

Japanese Genji firefly, *Luciola cruciata*, at the nuclear DNA level using RAD-Seq

analysis. *Sci. Rep.* **10,** 1533. https://doi.org/10.1038/s41598-020-58324-9 (2020).

29. Yokota, M., Harada, Y. & Iizuka, M. Genetic drift in a hatchery and the maintenance

of genetic diversity in hatchery-wild systems. *Fish Sci.* **69,** 101–109.

https://doi.org/10.1046/j.1444-2906.2003.00593.x (2003).

30. Harada, Y., Yokota, M., & Iizuka, M. Genetic risk of domestication in artificial fish

stocking and its possible reduction. *Popul. Ecol.* **40**, 311–324.

https://doi.org/10.1007/BF02763463 (1998).

31. Etter, P. D., Bassham, S., Hohenlohe, P. A., Johnson, E. A. & Cresko, W. A. SNP

discovery and genotyping for evolutionary genetics using RAD sequencing, in:

Molecular methods for evolutionary genetics. *Methods Mol Biol. Humana Press* **772,**

157–178. https://doi.org/10.1007/978-1-61779-228-1_9 (2011).

32. Rochette, N.C. & Catchen, J.M. Deriving genotypes from RAD-seq short-read data

using Stacks. *Nat. Protoc.* **12,** 2640–2659. https://doi.org/10.1038/nprot.2017.123

(2017).

33. Ran, Z. *et al.* Chromosome-level genome assembly of the razor clam *Sinonovacula*

constricta (Lamarck, 1818). *Mol. Ecol. Resour.* **19,** 1647–1658.

https://doi.org/10.1111/1755-0998.13086 (2019).

34. Li, H. Aligning sequence reads, clone sequences and assembly contigs with

BWA-MEM. arXiv preprint arXiv:1303.3997. https://arxiv.org/abs/1303.3997

(2013).

35. Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C. & Weir, B. S. A

high-performance computing toolset for relatedness and principal component

analysis of SNP data. *Bioinformatics* **28,** 3326–3328.

https://doi.org/10.1093/bioinformatics/bts606 (2012).

36. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27,** 2156–

2158. http://doi.org/10.1093/bioinformatics/btr330 (2011).

37. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry

in unrelated individuals. *Genome Res.* **19,** 1655–1664.

https://pubmed.ncbi.nlm.nih.gov/19648217/ (2009).

38. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and

population-based   linkage   analyses. *Am. J. Hum. Genet.* **81,** 559–575.

https://doi.org/10.1086/519795 (2007).

39. Eklund, A. R package beeswarm: the bee swarm plot, an alternative to stripchart.

https://cran.r-project.org/web/packages/beeswarm/beeswarm.pdf (2016).

40. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of   large   phylogenies.   *Bioinformatics*   **30,**   1312–1313.

https://doi.org/10.1093/bioinformatics/btu033 (2014).

41. Chifman, J. & Kubatko, L. Quartet inference from SNP data under the coalescent

model. *Bioinformatics* **30,** 3317–3324. https://doi.org/10.1093/bioinformatics/btu530

(2014).

42. Swofford, D. L. PAUP*. Phylogenetic analysis using parsimony (*and other

methods). Version 4. (Sinauer Associates, 2002).

Acknowledgments

The authors are grateful to the Saga Prefectural Ariake Fisheries Research and

Development Center for supporting the field survey and providing samples and

information on seedling production. This work was supported by the "Projects for

sophistication of production and utilization technology supporting local agriculture and

marine industry" from Saga University.

404   Author contributions

405   R.O. designed and wrote the original manuscript draft. R.O. and Y.N. performed the

406   RAD-Seq analysis and bioinformatic analysis. All authors reviewed drafts of the

407   manuscript and approved its final version.

409   Competing interests

410   The authors declare no competing interests.

413 Table 1. Sampling information for *Sinonovacula constricta* in the Ariake Bay, Japan

Sampling site	Sampling day	Habitat type	CPUE* (ind. person^{-1} h^{-1})	Sample ID	Shell length (mm)
	Nov. 11, 2019	Wild	0.3	Kashima_a.1	39.7
Kashima_b	Oct. 11, 2019	Wild	1.3	Kashima_b.1	46.6
				Kashima_b.2	45.2
				Kashima_b.3	54.7
				Kashima_b.4	54.0
Kashima_c	Nov. 19, 2019	Wild	4.0	Kashima_c.1	44.7
				Kashima_c.2	51.4
				Kashima_c.3	48.1
				Kashima_c.4	67.0
	Oct. 23, 2019	Wild	N/A[†]	Kashima_c.5	70.2
Kashima_d	Nov. 11, 2019	Wild	0.0	–	–
Nanaura_a	Nov. 13, 2019	Wild	0.7	Nanaura_a.1	47.0
				Nanaura_a.2	53.9
Tara_a	Nov.13, 2019	Wild	0.0	–	–
Takori_a	Nov. 15, 2019	Wild	1.8	Takori_a.1	75.0
				Takori_a.2	55.7
				Takori_a.3	81.7
				Takori_a.4	94.6
Takori_b	Nov. 15, 2019	Transplanted area[‡]	5.5	2014.1	92.8
				2014.2	94.8
				2014.3	92.1
				2014.4	91.9
				2014.5	92.9
Takori_c	Nov. 15, 2019	Transplanted area[‡]	8.5	2016.1	88.4
				2016.2	87.0
				2016.3	83.7
				2016.4	82.6
				2016.5	86.7
				2016.6	79.7
				2016.7	93.6
Higashiyoka_a	Nov. 12, 2019	Wild	0.0	–	–
Higashiyoka_b	Oct. 18, 2019	Wild	0.0	–	–
N/A[†]	Dec. 10, 2006	Transplanted area[‡]	N/A[†]	2005.1	66.7
				2005.2	74.7

*: Catch per unit effort (individual person^{-1} hour^{-1})

[†]: Not available

[‡]: Transplanted area of artificial seedlings

414 **Figures**

418 **Figure 1.** Sampling sites. The dotted line shows the area explored.

Figure 2. Biplots of principal component analysis. (a) Showing by sampling areas. (b)

Showing by shell sizes.

Figure 3. Admixture analysis of *S. constricta* in the Ariake Bay, Japan.

440 **Figure 4.** Pairwise F_{st} values between individuals in each population.

Figure 5. Phylogenetic trees of *S. constricta* inferred from RAD-Seq dataset (a)

Maximum likelihood phylogeny estimated using RAxML. (b) Species tree inferred using

SVDquartets. Numbers at the nodes indicate bootstrap values (1000 replicates).

455 **Supplementary material**

Fig. S1 Comparison of shell morphology. WH: individuals in wild habitat (n=17), TA: individuals in transplanted area (n=11). NS[1]: non-significant difference between the comparison groups using the Student's t test. NS[2]: non-significant difference between the comparison groups using the Wilcoxon rank sum test. Bar indicate the SD.