Genetic diversity and population structure of razor clam *Sinonovacula constricta*

in Ariake Bay, Japan, revealed using RAD-Seq SNP markers.

Ryo Orita[a,*], Yukio Nagano[b], Yoshio Kawamura[a], Kei Kimura[a], and Genta Kobayashi[a]

[a]*Faculty of Agriculture, Saga University, 1 Honjo-machi, Saga, 840-8502, Japan*

[b]*Analytical Research Center for Experimental Sciences, Saga University, 1*

Honjo-machi, Saga, 840-8502, Japan

*Corresponding author: Ryo Orita, E-mail: ss7427@cc.saga-u.ac.jp, Tel: +81-952-28-

8763

Abstract:

The razor clam *Sinonovacula constricta* is a commercially important bivalve in Japan.

The current distribution of this species in Japan is limited to Ariake Bay, where the

fishery stock is declining. It is necessary to understand the genetic population structure

of this species in order to restore the fishery stock while preserving the genetic

diversity of the clam. Here, we report for the first time the genetic population structure

of *S. constricta* in Ariake Bay, Japan. Paired-end restriction site-associated DNA

sequencing (RAD-Seq) analyzed samples of *S. constricta* collected from seven

mudflats located along Ariake Bay. Two different genetic populations exist in Ariake

Bay, one inhabiting wild habitats and the other inhabiting the transplanted area of

artificial seedlings. Our results suggest that genetic differentiation occurred between

these two populations (F_{st} value = 0.052), and a high level of genetic differentiation is

maintained between the two groups. In the future, the two genetically distinct

populations need to be available as fishery resources, while taking into account their

conservation and hybridization status.

1. Introduction

The technique of artificially producing seedlings and releasing them into fishing grounds is an effective method of aquaculture enhancement. This technique has been implemented with a variety of fishery species (e.g., finfish, Fushimi, 2001; Masuda and Tsukamoto, 1998; flounder, Sekino et al., 2005; shellfish, Arnold, 2001; Castell et al., 1996; McCormick et al., 1994; Zohar et al., 2008). In artificial seed production, large numbers of seedlings are often produced by a few broodstocks and released into natural seawater, which reduces the genetic diversity of wild populations in the field. Therefore, when seedlings are produced and released into natural seawater, there is a need to understand the genetic structure within the wild population in order to preserve genetic diversity.

The razor clam *Sinonovacula constricta* (Lamarck, 1818) is an important economic species in East Asian coastal waters (Funge-Smith et al., 2012; Mao et al., 2019; Ran et al., 2017). Shell records suggest that *S. constricta* was once distributed in tidal flats in western Japan (Suzuki et al., 2006), but its current distribution is limited to

the mudflats around Ariake Bay located on the west coast of Kyusu, Japan. According

to a report by Ministry of the Environment, Japan (2006), the annual catch of *S.*

constricta in Ariake Bay often recorded more than 1000 tons per year from the 1900s to

the early the 1920s, but it decreased to less than 500 tons in most years from the latter

half of the 1920s to the latter half of the 1980s; no catches have been made since 1994.

In order to restore the *S. constricta* resources, the Saga Prefectural Ariake Fisheries

Research and Development Center has developed seed production techniques (Ito et al.,

2001; Ohkuma et al., 2004), and since 1988, artificial seedlings of *S. constricta* have

been released into the transplanting areas established in natural seawaters (Tsukuda et

al., 2017). However, the genetic population structure of *S. constricta* within Ariake Bay

has not been examined.

In recent years, technological innovations have made the genome-wide analysis

of non-model organisms more widely available, and these techniques are useful for

examining the genetic population structure of non-model aquatic organisms (Holman et

al., 2017; Li and Wang, 2017; Robledo et al., 2018; You et al., 2020). In the present

study, to clarify the genetic population structure of *S. constricta* in Ariake Bay, we

conducted field sampling surveys at 11 mudflats located along the bay in 2019 and used

restriction-site-associated DNA sequencing (RAD-Seq) to analyze population genetics

and phylogenetics. We propose guidelines concerning the optimal seed production and

seed release methods for the conservation of the genetic diversity of *S. constricta* in

Japan.

2. Materials and methods

2.1. Sampling and genomic DNA extraction

Sampling of the razor clam *S. constricta* was conducted at 11 mudflats located

along Ariake Bay (Fig. 1). Information on the sampling day and habitat types at each

site is shown in Table 1. Habitats are divided into two types: transplanted area of

artificial seedlings, and wild habitat. As mentioned above, artificial seedlings of *S.*

constricta have been released into the transplanting areas established in natural

seawaters since 1988. Therefore, artificially produced individuals live within the

transplanted area, and the wild habitat would be populated by natural individuals and

offspring of seedlings from the transplanted areas. At each site, razor clams were

explored by hand and their abundance was assessed using the catch per unit effort index

(CPUE), calculated as the number of clam individuals collected divided by the number

of explorers and sampling time (i.e., ind. person^{-1} hour^{-1}). The samples were stored at

-80°C until genomic DNA extraction. In addition to the collected individuals, the oldest

seedlings (produced in 2005, n = 2) stored at the Saga Prefectural Ariake Fisheries

Research and Development Center, were included in the RAD-Seq analysis for

comparison with seedlings produced in the past. Genomic DNA was extracted from

finely chopped foot tissue of the specimens using DNAs-ici!-F (RIZO, Tsukuba, Japan),

and the concentration of the DNA was assessed using a Qubit 3.0 Fluorometer (Thermo

Fisher Scientific).

2.2. RDA-Seq analysis

The preparation for *the Eco*RI RAD-Seq library has been described previously

(Kato et al., 2020), based on the established method (Etter et al., 2011). The library was

sequenced with 150 bp paired-end reads in one lane of an Illumina HiSeq X (Illumina,

San Diego, CA, USA). The RAD-Seq data were analyzed using the Stacks program

(version 2.52) (Rochette and Catchen, 2017). The data were quality filtered using the

process_radtags program in the Stacks package (-r -c -q, - adapter_1, adapter_2, and

adapter_mm2). Filtered sequences were registered in the DRA (DRA accession number:

DRA010922). The data were aligned with the reference genome of *S. constricta* (Ran et

al., 2019; NCBI/DDBJ BioProject number: PRJNA508451) using BWA-MEM (version

0.7.17) (Li, 2013). Aligned data were analyzed using the ref_map.pl script of the Stacks

package.

2.3. Population genetics analysis

Principal component analysis (PCA) was conducted based on a variant call

format (VCF) file using the SNPrelate program (v.1.18.1) (Zheng et al., 2012). The

VCF file was created by the populations program of the Stacks package

(--rm-pcr-duplicates, -R 0.7, --write-single-snp, --min-maf 0.05) (Danecek et al., 2011).

The admixture analysis was conducted based on the PLINK file using ADMIXTURE (v.

1.3.0) (Alexander et al., 2009). The PLINK file was created with the same Stacks

populations program that created the VCF file (Purcell et al., 2007). The genetic

diversity of razor clams among and within populations was assessed using AMOVA F_{st}

values obtained from 'populations.fst_files', which were calculated by the Stacks

populations program. To visualize the distribution of all data points, pairwise F_{st} values

between individuals in each population are shown in boxplot with beeswarm using R

(package: 'beeswarm', Eklund, 2016).

2.4. Phylogenetic tree analysis

Phylogenetic tree analysis was performed using two different approaches: the

maximum likelihood (ML) method and SVDquartets analysis. Multiple alignments

within the cluster were obtained using the Stacks package populations program

mentioned earlier. The phylogenetic tree based on the ML method was constructed

using the RAxML program (version 8.2.12) (Stamatakis, 2014) (-f = a, -x = 12 345, -p =

12 345, -N (bootstrap value) = 1000, and -m = GTRGAMMA). The SVDquartets

analysis based on the coalescent model was computed as implemented in PAUP ver.

4.0a 167 using multiple alignments (Chifman and Kubatko, 2014; Swofford, 2002). The

number of bootstrap analyses replicates was 1000. In each analysis, the oldest samples

(i.e., sample ID, 2005.1; 2005.2), were used as the root.

3. Results

3.1. Sampling data

We collected 28 razor clams from seven of the 11 mudflats and found no razor

clams on the remaining four mudflats (Table 1). The mean CPUE of the seven mudflats

where individuals could be collected was 3.2 ($\pm$ 3.0 SD) ind. person^{-1} hour^{-1}, and the

range of CPUE at each site was 0.3 to 8.5 ind. person^{-1} h $^{-1}$. The shell length of the

collected individuals was 39.7 to 94.6 mm, implying that they contained specimens

from 1 to 5 years old. The mean shell length of individuals collected from the

transplanted area was 86.0 (± 4.5 SD) mm for the 3-year-old seedlings produced in

2016 (i.e., sample ID: 2016.1-2016.7) and 92.9 (± 1.1 SD) mm for the 5-year-old

seedlings produced in 2014 (i.e., sample ID: 2014.1-2014.5).

3.2. RAD-Seq data

We used 30 individuals for RAD-Seq analysis, including 16 individuals

collected from wild habitats, 12 individuals collected from the transplanted area of

artificial seedlings, and 2 individuals stored at the Saga Prefectural Ariake Fisheries

Research and Development Center (Table 1). In all, 401 109 332 raw reads were

generated from the HiSeq X run, and 249 040 001 reads were retained after discarding

reads with low quality and adapters. The final dataset included 137 246 loci, with an

average of 679.38 bp (SE ± 0.18) of genotyped sites per locus.

3.3. Population genetics structure

The genetic relationships among sampling sites were clearly divided into

individuals collected from wild habitat and those from the transplanted area of artificial

seedlings along the axis of PC1 (Fig. 2 (a)). The PC2 axis represents the differences

within each habitat type. In the PCA plot, the coordinates of the transplanted samples

were more widely scattered than those of the wild samples were, showing more diverse

genetic relationships among transplanted samples. There were no distinct genetic

relationships among generations, and the generation with a shell length of 75 to 90 mm

showed the most genetic diversity (Fig. 2(b)). The results of admixture analysis also

showed a significant genetic difference between the individuals collected from the wild

versus those collected from the transplanted areas of artificial seedlings (Fig. 3). The F_{st}

value between populations consisting of individuals collected from the wild and the

transplanted areas was 0.052. Owing to the calculation of the pairwise F_{st} values for

each individual within the two populations, there was no significant difference in the

mean values between these populations (Fig. 4). The pairwise F_{st} values within the

population were more varied in the transplanted area (from 0.44 to 0.64) than in the

wild habitat (from 0.53 to 0.56).

3.4. Phylogenetic tree analysis

ML tree and SVDquartets analysis of RAD-Seq data showed that the

phylogenies were highly consistent with each other (Fig. 5(a) and (b)). The obtained

phylogenetic trees were clearly divided into individuals collected from the wild habitat

and those collected from the transplanted area of artificial seedlings, with the exception

of one sample (i.e., sample ID: 2016.7).

4. Discussion

4.1. Genetic population structure of S. constricta in the Ariake Bay, Japan

All our analyses of the RAD-Seq data showed that individuals collected from

wild habitats were genetically different from those collected from sites where artificial

seedlings were transplanted. Only one sample collected from the area transplanted in

2016 (sample ID: 2016.7) showed different results from other seedlings collected in the

same area, and belonged to a population consisting of individuals collected from the

wild habitat (Fig. 1(a), Fig. 3, 4). Since this individual had a distinctly larger shell

length than other 3-year-old seedlings in the same area, and its size was the same as that

of 5-year-old seedlings (Table 1), it was likely a wild individual inhabiting the

transplanted area prior to the 2016 transplantation. Therefore, we consider that there

were two genetically distinct populations in Ariake Bay in 2019.

The reason for the differences between these two populations is the different

parental origins of the seedlings. The parental origins can be divided into the Nanaura

mudflat and the Kashima mudflat for a number of reasons. Artificial seedlings of *S.*

constricta have been released by the Saga Prefectural Ariake Fisheries Research and

Development Center into transplanting areas established in several mudflats along the

coast of Ariake Bay. The seedlings collected from the transplanted area in this study

(Takori_b and Takori_c) originated from parents at the Nanaura mudflat (hereinafter

referred to as Nanaura lineage population), since the parents were collected from the

Nanaura mudflat before 2010 and released into the transplanting areas of the Takori

mudflat. In contrast, individuals collected from the wild habitat of Kashima, Nanaura,

and Takori mudflats are likely to have originated from parents of Kashima mudflat

(hereinafter referred to as the Kashima lineage population). In 2017 and 2018, the

seedlings produced by parents collected from Kashima mudflat were released into new

transplanting areas at Takori and Nanaura mudflats due to declining populations in wild

and transplanted areas in these mudflats. Therefore, the 1- to 2-year old individuals

collected in wild habitats in 2019 were likely the offspring of the transplanted Kashima

seedlings in 2017 and 2018.

Artificial transplantation of aquatic organisms has been reported to alter the

geographic genetic structure of wild populations (Arthington, 1991; Habtemariam et al.,

2015; Williams and Orth, 1998; Yamakawa and Imai, 2013), including that of *S.*

constricta, in this case along the Chinese coast (Niu et al., 2012). Anthropogenic

transplantation likely strongly influenced the distribution of the genetic population

structure of *S. constricta* in Japan. From the dataset of this study, it is unclear why the

origins of the parents differed between the Kashima mudflat and the Nanaura mudflat,

which are in close proximity to each other. In future analyses, comparisons with

ancestral samples from the coast of the continent of China may lead to further

understanding of the phylogenetic relationship between these two populations.

4.2. Genetic differentiation among and within populations

For artificial seed production and release of seedlings, it is necessary to

understand the distribution of populations with different genetic structures in natural

seawater. Therefore, it is necessary to select broodstocks and release areas according to

the degree of genetic differentiation among and within populations. F_{st} values are

widely used as a measure of the degree of genetic differentiation among and within

populations (DeFaveri et al., 2012; Jeffery et al., 2017; Kato et al., 2020). According to

Niu et al. (2012), the F_{st} value among populations of *S. constricta* at 10 sites along the

coast of mainland China is 0.044. The F_{st} value between the two populations in the

present study was 0.052, suggesting that a similar level of genetic differentiation

occurred between the two populations in Ariake Bay, Japan.

For these two populations, the genetic differentiation within each population

was maintained at a high level, and its variation was greater within the population

consisting of individuals collected from the transplanted area (Fig. 4). There are two

possible reasons why the variation in the degree of genetic differentiation within

populations differed between the two populations. The first is that the degree of

variation in genetic differentiation in these populations originally may have been

different. Second, the methods of seedling production may lead to greater variation of

genetic differentiation within populations in the transplanted area. In seedling

production, broodstocks were randomly selected with closely related individuals

collected from the same transplanted area or distantly related individuals collected from

different transplanted areas. After stimulating the mature individuals to spawn, 20 to 30

individuals were placed in the same tank to produce fertilized eggs. Thus, the degree of

genetic differentiation may have been more varied in the transplanted populations than

in the wild habitat population.

4.3. Guidelines concerning the best seed production and seed release methods

The current abundance of *S. constricta* is estimated to be very low based on the

CPUE results of this study (Table 1). These results suggest that artificial seed

production and release of seedlings could be an effective strategy for the recovery of *S.*

constricta abundance in Ariake Bay, Japan. The present study suggested genetic

differentiation between the two populations of *S. constricta* in Ariake Bay, and a high

level of genetic differentiation within each population was maintained. Therefore,

maintaining the two populations without interbreeding seems to be an appropriate

conservation strategy. As of 2019, the Kashima lineage population is widely distributed

in wild habitats, whereas the Nanaura lineage population is limited to the Takori

mudflats. In order to avoid hybridization with the Kashima lineage population, the

release of seedlings of the Nanaura lineage population should continue to be restricted

to the mudflats within the current cove, with a transplanting area.

To produce seedlings to slow down the decline in genetic diversity within a

population, mathematical models have evaluated five strategies: I) avoidance of line

breeding, II) selective use of wild-born individuals, III) line breeding, IV) avoidance of

artificial breeding of males, and V) selective use of wild-born males (Yokota et al.,

2003). Among them, strategy II has been considered the most effective. The best way to

produce seedlings of the Kashima lineage population would be to follow strategy II; all

parental individuals would be randomly collected from wild habitats and previously

produced seedlings would not be used as parents. In contrast, the Nanaura lineage

population has already been maintained by a similar approach to strategy I; all parental

individuals will be randomly collected from transplanted areas without excluding

released seedlings. This approach gradually reduces the genetic diversity within the

population. Indeed, lower pairwise F_{st} values were detected in the transplanted

population (Fig. 4). Therefore, in the Nanaura lineage population, countermeasures

should be taken to increase the number of parents used for seed production or to avoid

using the produced seedlings as parents of the next generation.

RAD-Seq analysis is an inexpensive way to identify populations, but it requires

a lot of manipulation and time. Several DNA markers, such as SSRs, SNPs, and InDels,

have been developed for simple and inexpensive identification of populations, and these

techniques are useful for monitoring the genetic hybridization of each population in a

short period of time (Liu and Cordes, 2004; Maqsood and Ahmad, 2017). For example,

PCR and agarose gel electrophoresis alone can detect short indels (Noda et al., 2020).

Monitoring the interbreeding status of the two genetically distinct populations and the

genetic differentiation within each population is important to enhance the stock of *S.*

constricta in Japan by producing and releasing seedlings, while preserving genetic

diversity.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or

personal relationships that could have appeared to influence the work reported in this

paper.

Acknowledgments

The authors are grateful to the Saga Prefectural Ariake Fisheries Research and

Development Center for supporting the field survey and providing samples and

information on seedling production. This work was supported by the "Projects for

sophistication of production and utilization technology supporting local agriculture and

marine industry" from Saga University.

References

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of

ancestry in unrelated individuals. Genome Res. 19, 1655–1664. Available online:

https://pubmed.ncbi.nlm.nih.gov/19648217/.

Arnold, W.S., 2001. Bivalve enhancement and restoration strategies in Florida, USA.

Hydrobiologia 465, 7–19. https://doi.org/10.1023/A:1014596909319.

Arthington, A.H., 1991. Ecological and genetic impacts of introduced and translocated

freshwater fishes in Australia. Can. J. Fish. Aquat. Sci. 48, 33–43.

https://doi.org/10.1139/f91-302.

Castell, L.L., Naviti, W., Nguyen, F., 1996. Detectability of cryptic juvenile *Trochus*

niloticus Linnaeus in stock enhancement experiments. Aquaculture 144, 91–101.

https://doi.org/10.1016/S0044-8486(96)01320-8.

Chifman, J., Kubatko, L., 2014. Quartet inference from SNP data under the coalescent

model. Bioinformatics 30, 3317–3324.

https://doi.org/10.1093/bioinformatics/btu530.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A.,

Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R.,

1000 Genomes Project Analysis Group, 2011. The variantcall format and VCFtools.

Bioinformatics 27, 2156–2158. http://doi.org/10.1093/bioinformatics/btr330.

308 DeFaveri, J., Shikano, T., Ghani, N.I.A., Merilä, J., 2012. Contrasting population

309 structures in two sympatric fishes in the Baltic Sea basin. Mar. Biol. 159, 1659–

310 1672. https://doi.org/10.1007/s00227-012-1951-4.

311 Eklund, A., 2016. R package beeswarm: the bee swarm plot, an alternative to stripchart.

312 Available online: https://cran.r-project.org/web/packages/beeswarm/beeswarm.pdf.

313 Etter, P.D., Bassham, S., Hohenlohe, P.A., Johnson, E.A., Cresko, W.A., 2011. SNP

314 discovery and genotyping for evolutionary genetics using RAD sequencing, in:

315 Molecular methods for evolutionary genetics. Methods Mol Biol. Humana Press

316 772, 157–178. https://doi.org/10.1007/978-1-61779-228-1_9.

317 Funge-Smith, S., Briggs, M., Miao, W., 2012. Regional Overview of Fisheries and

318 Aquaculture in Asia and the Pacific 2012. RAP Publication (FAO).

319 Fushimi, H., 2001. Production of juvenile marine finfish for stock enhancement in

320 Japan. Aquaculture 200, 33–53. https://doi.org/10.1016/S0044-8486(01)00693-7.

321 Habtemariam, B.T., Arias, A., García-Vázquez, E., Borrell, Y.J., 2015. Impacts of

322 supplementation aquaculture on the genetic diversity of wild *Ruditapes decussatus*

323 from northern Spain. Aquacult. Environ. Interact. 6, 241–254.

324 https://doi.org/10.3354/aei00128.

325 Holman, L.E., Garcia de la serrana, D., Onoufriou, A., Hillestad, B., Johnston, I.A.,

326 2017. A workflow used to design low density SNP panels for parentage assignment

327 and traceability in aquaculture species and its validation in Atlantic salmon.

328 Aquaculture 476, 59–64. https://doi.org/10.1016/j.aquaculture.2017.04.001.

329 Ito, S., Eguchi, T., Kawahara, I., 2001. Rearing experiment on planktonic larvae of the

330 JackKnife clam, *Sinonovacula constricta*. Boll. Saga Prefect. Ariake Fish. Res. Dev.

331 Cent. 20, 49–53. Available online:

332 https://agriknowledge.affrc.go.jp/RN/2010813185.pdf.

333 Jeffery, N.W., DiBacco, C., Van Wyngaarden, M., Hamilton, L. C., Stanley, R.R.,

334 Bernier, R., FitzGerald, J., Matheson, K., McKenzie, C.H., Ravindran, P.N., Beiko,

335 R. Bradbury, I.R., 2017. RAD sequencing reveals genomewide divergence between

336 independent invasions of the European green crab (*Carcinus maenas*) in the

337 Northwest Atlantic. Ecol. Evol. 7, 2513–2524. https://doi.org/10.1002/ece3.2872.

Kato, D., Suzuki, H., Tsuruta, A., Maeda, J., Hayashi, Y., Arima, K., Ito, Y., Nagano,

Y., 2020. Evaluation of the population structure and phylogeography of the

Japanese Genji firefly, *Luciola cruciata*, at the nuclear DNA level using RAD-Seq

analysis. Sci. Rep. 10, 1533. https://doi.org/10.1038/s41598-020-58324-9.

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with

BWA-MEM.   arXiv   preprint   arXiv:1303.3997.   Available   online:

https://arxiv.org/abs/1303.3997.

Li, Y.H., Wang, H.P., 2017. Advances of genotyping-by-sequencing in fisheries and

aquaculture.   Rev.   Fish   Biol.   Fish.   27,   535–559.

https://doi.org/10.1007/s11160-017-9473-2.

Liu, Z.J., Cordes, J.F., 2004. DNA marker technologies and their applications in

aquaculture genetics. Aquaculture 238, 1–37.

https://doi.org/10.1016/j.aquaculture.2004.05.027.

Mao, Y., Lin, F., Fang, J., Fang, J., Li, J., Du, M., 2019. Chapter 4. Bivalve production

in China, in: Goods and Services of Marine Bivalves. Springer, Cham, pp. 51–72.

Maqsood, H.M., Ahmad, S.M., 2017. Advances in molecular markers and their

applications in aquaculture and fisheries. Genet. Aquat. Org. 1, 27–41. Available

online: http://www.genaqua.org/uploads/pdf_1.pdf.

Masuda, R., Tsukamoto, K., 1998. Stock enhancement in Japan: review and perspective.

Bull. Mar. Sci. 62, 337–358. Available online:

https://www.ingentaconnect.com/content/umrsmas/bullmar/1998/00000062/000000

02/art00005.

McCormick, T.B., Herbinson, K., Mill, T.S., Altick, J., 1994. A review of abalone

seeding, possible significance and a new seeding device. Bull. Mar. Sci. 55, 680–

693. Available online:

https://www.ingentaconnect.com/content/umrsmas/bullmar/1994/00000055/f00200

02/art00035.

Ministry of the Environment, Japan., 2006. Committee Report. Ministry of the

Environment, Bay, A., Sea, Y. Comprehensive Survey Evaluation Committee, 52

pp. Available online: https://www.env.go.jp/council/20ari-yatsu/rep061221/all.pdf.

Niu, D.H., Feng, B.B., Liu, D.B., Zhong, Y.M., Shen, H.D., Li, J.L., 2012. Significant

genetic differentiation among ten populations of the razor clam *Sinonovacula*

constricta along the coast of china revealed by a microsatellite analysis. Zool. Stud.

51, 406–414. Available online: http://zoolstud.sinica.edu.tw/Journals/51.3/406.pdf.

Noda, T., Daiou, K., Mihara, T., Nagano, Y., 2020. Development of indel markers for

the selection of *Satsuma mandarin* (*Citrus unshiu* Marc.) hybrids that can be used

for low-cost genotyping with agarose gels. Euphytica 216, 1–13.

https://doi.org/10.1007/s10681-020-02654-2.

Ohkuma, H., Ymagachi, T., Kawahara, I., Eguchi, T., Ito, S., 2004. A study on the

development of techniques for mass production of seeds of jackknife clam,

Sinonovacula constricta. Boll. Saga Prefect. Ariake Fish. Res. Dev. Cent. 22, 47–

54. Available online: https://agriknowledge.affrc.go.jp/RN/2030813210.pdf.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D.,

Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., Sham, P.C., 2007. PLINK: a

tool set for whole-genome association and population-based linkage analyses. Am.

J. Hum. Genet. 81, 559–575. https://doi.org/10.1086/519795.

Ran, Z., Chen, H., Ran, Y., Yu, S., Li, S., Xu, J., Liao, K., Yu, X., Zhong, Y., Ye, M.,

Yan, X., 2017. Fatty acid and sterol changes in razor clam *Sinonovacula constricta*

(Lamarck 1818) reared at different salinities. Aquaculture 473, 493–500.

https://doi.org/10.1016/j.aquaculture.2017.03.017.

Ran, Z., Li, Z., Yan, X., Liao, K., Kong, F., Zhang, L., Cao, J., Zhou, C., Zhu, P., He, S.,

Huang, W., Xu, J., 2019. Chromosome level genome assembly of the razor clam

Sinonovacula constricta (Lamarck, 1818). Mol. Ecol. Resour. 19, 1647–1658.

https://doi.org/10.1111/1755-0998.13086.

Robledo, D., Palaiokostas, C., Bargelloni, L., Martínez, P., Houston, R., 2018.

Applications of genotyping by sequencing in aquaculture breeding and genetics.

Rev. Aquac 10, 670–682. https://doi.org/10.1111/raq.12193.

Rochette, N.C., Catchen, J.M., 2017. Deriving genotypes from RAD-seq short-read data

using Stacks. Nat. Protoc. 12, 2640–2659. https://doi.org/10.1038/nprot.2017.123.

Sekino, M., Saitoh, K., Yamada, T., Hara, M., Yamashita, Y., 2005. Genetic tagging of

released Japanese flounder (*Paralichthys olivaceus*) based on polymorphic DNA

markers. Aquaculture 244, 49–61.

https://doi.org/10.1016/j.aquaculture.2004.11.006.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and

post-analysis of large phylogenies. Bioinformatics 30, 1312–1313.

https://doi.org/10.1093/bioinformatics/btu033.

Suzuki, T., Inoue, K., Ozawa, T., 2006. Environmental degradation in Ise and Mikawa

Bays after 1960's as viewed from intertidal molluscan community. Boll. Nagoya

Univ. Museum 22, 31–64. http://doi.org/10.18999/bulnum.022.04.

Swofford, D. L. 2002. PAUP*. Phylogenetic analysis using parsimony (*and other

methods). Version 4. Sinauer Assoc. Sunderland, Massachusetts.

Tsukuda, M., Kanzaki, H., Tsujo, K., Fukumoto, T., Umeda, T., Aramaki, H., Ito, S.,

2017. Variation in the Distribution of the JackKnife Clam, *Sinonovacula constricta*,

on the Muddy Tidal flat of Ariake Sound off Saga Prefecture. Boll. Saga Prefect.

Ariake Fish. Res. Dev. Cent. 28, 47–49. Available online:

https://agriknowledge.affrc.go.jp/RN/2010925685.pdf.

Williams, S.L., Orth, R.J., 1998. Genetic diversity and structure of natural and

transplanted eelgrass populations in the Chesapeake and Chincoteague Bays.

Estuaries 21, 118–128. https://doi.org/10.2307/1352551.

Yamakawa, A.Y., Imai, H., 2013. PCR-RFLP typing reveals a new invasion of

Taiwanese Meretrix (Bivalvia: Veneridae) to Japan. Aquat. Invas. 8, 407–415.

http://doi.org/10.3391/ai.2013.8.4.04.

Yokota, M., Harada, Y., Iizuka, M., 2003. Genetic drift in a hatchery and the

maintenance of genetic diversity in hatchery-wild systems. Sci 69, 101–109.

https://doi.org/10.1046/j.1444-2906.2003.00593.x.

You, X., Shan, X., Shi, Q., 2020. Research advances in the genomics and applications

for molecular breeding of aquaculture animals. Aquaculture 526.

https://doi.org/10.1016/j.aquaculture.2020.735357,

Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., Weir, B.S., 2012. A

high-performance computing toolset for relatedness and principal component

analysis of SNP data. Bioinformatics 28, 3326–3328.

https://doi.org/10.1093/bioinformatics/bts606.

Zohar, Y., Hines, A.H., Zmora, O., Johnson, E.G., Lipcius, R.N., Seitz, R.D., Eggleston,

D.B., Place, A.R., Schott, E.J., Stubblefield, J.D., Chung, J.S., 2008. The

Chesapeake Bay blue crab (*Callinectes sapidus*): A multidisciplinary approach to

responsible stock replenishment. Rev. Fish. Sci. 16, 24–34.

https://doi.org/10.1080/10641260701681623.

Table 1. Sampling infromation for *Sinonovacula constricta* in the Ariake Bay, Japan

Sampling site	Sampling day	Habitat type	CPUE* (ind. person^{-1} h^{-1})	Sample ID	Shell length (mm)
Kashima_a	Nov. 11, 2019	Wild	0.3	Kashima_a.1	39.7
Kashima_b	Oct. 11, 2019	Wild	1.3	Kashima_b.1	46.6
				Kashima_b.2	45.2
				Kashima_b.3	54.7
				Kashima_b.4	54.0
Kashima_c	Nov. 19, 2019	Wild	4.0	Kashima_c.1	44.7
				Kashima_c.2	51.4
				Kashima_c.3	48.1
				Kashima_c.4	67.0
	Oct. 23, 2019	Wild	N/A[†]	Kashima_c.5	70.2
Kashima_d	Nov. 11, 2019	Wild	0.0	–	–
Nanaura_a	Nov. 13, 2019	Wild	0.7	Nanaura_a.1	47.0
				Nanaura_a.2	53.9
Tara_a	Nov.13, 2019	Wild	0.0	–	–
Takori_a	Nov. 15, 2019	Wild	1.8	Takori_a.1	75.0
				Takori_a.2	55.7
				Takori_a.3	81.7
				Takori_a.4	94.6
Takori_b	Nov. 15, 2019	Tranplanted area[‡]	5.5	2014.1	92.8
				2014.2	94.8
				2014.3	92.1
				2014.4	91.9
				2014.5	92.9
Takori_c	Nov. 15, 2019	Tranplanted area[‡]	8.5	2016.1	88.4
				2016.2	87.0
				2016.3	83.7
				2016.4	82.6
				2016.5	86.7
				2016.6	79.7
				2016.7	93.6

Higashiyoka_a	Nov. 12, 2019	Wild	0.0	–	–
Higashiyoka_b	Oct. 18, 2019	Wild	0.0	–	–
N/A[†]	Dec. 10, 2006	Tranplanted area[‡]	N/A[†]	2005.1	66.7
				2005.2	74.7

*: Catch per unit effort (individual person^{-1} hour^{-1})

[†]: Not available

[‡]: Transplanted area of airtficial seedlings

438   **Figures**

441   **Fig. 1.** Sampling sites. The dotted line shows the area explored.

Fig. 2. Biplots of principal component analysis. (a) Showing by sampling areas. (b)

444   Showing by generations.

447 **Fig. 3.** Admixture analysis of *S. constricta* in the Ariake Bay, Japan.

449   **Fig. 4.** Pairwise F_{st} values between individuals in each population.

Fig. 5. Phylogenetic trees of *S. constricta* inferred from RAD-Seq dataset (a) Maximum

likelihood phylogeny estimated using RAxML. (b) Species tree inferred using

SVDquartets. Numbers at the nodes indicate bootstrap values (1000 replicates).