Genetic landscape of recessive diseases in the Vietnamese population

from large-scale clinical exome sequencing

Ngoc Hieu Tran[1], Thanh-Huong Nguyen Thi[2], Hung-Sang Tang[1,3], Le-Phuc Hoang[2], Trung-Hieu

Le Nguyen[4,9], Nhat-Thang Tran[5], Thu-Huong Nhat Trinh[6], Van Thong Nguyen[7], Bao-Han Huu

Nguyen[2], Hieu Trong Nguyen[1], Loc Phuoc Doan[1], Ngoc-Minh Phan[1,3], Kim-Huong Thi

Nguyen[1,3], Hong-Dang Luu Nguyen[1,3], Minh-Tam Thi Quach[1,3], Thanh-Phuong Thi Nguyen[1,3],

Vu Uyen Tran[1], Dinh-Vinh Tran[8], Quynh-Tho Thi Nguyen[3], Thanh-Thuy Thi Do[3], Nien Vinh

Lam[9], Phuong Cao Thi Ngoc[1], Dinh Kiet Truong[3], Hoai-Nghia Nguyen[9,*], Minh-Duy Phan[1,*],

Hoa Giang[1,3,*]

[1]Gene Solutions, Ho Chi Minh city, Vietnam

[2]Children Hospital 1, Ho Chi Minh city, Vietnam

[3]Medical Genetics Institute, Ho Chi Minh city, Vietnam

[4]Department of Neurology, Children Hospital 2, Ho Chi Minh city, Vietnam

[5]University Medical Center, Ho Chi Minh city, Vietnam

[6]Tu Du Hospital, Vietnam

[7]Hung Vuong Hospital, Vietnam

[8]Da Nang Hospital for Women and Children, Vietnam

[9]University of Medicine and Pharmacy at Ho Chi Minh city, Vietnam

*Corresponding authors.

Hoai-Nghia Nguyen (nhnghia81@gmail.com)

Minh-Duy Phan (pmduy@yahoo.com); ORCID: 0000-0002-3426-1044

Hoa Giang (gianghoa@gmail.com)

Abstract

Purpose: Accurate profiling of population-specific recessive diseases is essential for the design

of cost-effective carrier screening programs. However, minority populations and ethnic groups,

including Vietnamese, are still under-represented in existing genetic studies. Here we reported

the first comprehensive study of recessive diseases in the Vietnamese population.

Methods: Clinical exome sequencing (CES) data of 4,503 disease-associated genes obtained

from a cohort of 985 Vietnamese individuals was analyzed to identify pathogenic variants,

associated diseases and their carrier frequencies in the population.

Results: Eighty-five recessive diseases were identified in the Vietnamese population, among

which seventeen diseases had carrier frequencies of at least 1% (1 in 100 individuals). Three

diseases were especially prevalent in the Vietnamese population with carrier frequencies of 2-

12 times higher than in other East Asia or the world populations, including Beta-thalassemia (1

in 25), citrin deficiency (1 in 33) and phenylketonuria (1 in 40). Seven novel pathogenic and

three likely pathogenic variants associated with nine recessive diseases were also discovered.

Conclusions: The comprehensive profile of recessive diseases identified in this study shall

enable the design of cost-effective carrier screening programs specific to the Vietnamese

population. The newly discovered pathogenic variants may also exist in other populations at

extremely low frequencies, thus representing a valuable resource for future research. Our study

has demonstrated the advantage of population-specific genetic studies to advance the

knowledge and practice of medical genetics.

Keywords: carrier frequency; carrier screening; recessive diseases.

INTRODUCTION

As high-throughput sequencing technologies are getting more popular and affordable, carrier screening has become a routine, essential tool for preventive healthcare and offers a great resource of information to guide public health policies [1-3]. Individuals identified as carriers of pathogenic genes and associated disorders may take preventive steps to reduce the risk of having their offspring inherit these disorders, such as preimplantation genetic diagnosis of embryos and/or early prenatal genetic testing. In addition, newborn screening programs enable early diagnosis and effective treatment of affected children, which not only significantly improve the outcomes but also reduce the treatment costs and efforts.

However, there are more than a thousand of Mendelian inherited disorders that have been documented, according to the Online Mendelian Inheritance in Man (OMIM) database [4]. Most of the diseases are rare and their prevalence depends heavily on specific populations and ethnicities. Thus, accurate profiling of population- or ethnicity-specific inherited diseases is essential for the design of cost-effective and comprehensive carrier screening programs. For example, cystic fibrosis is recommended for carrier screening for individuals of Caucasian or Ashkenazi Jewish ancestry, Tay–Sachs disease for individuals of Ashkenazi Jewish ancestry, and Beta-thalassemia for individuals from Mediterranean regions [1]. The critical problem, however, is the under-representation of minority populations and ethnic groups in existing genetic studies and databases [5-7]. Gurdasani *et al.* found that nearly 78% of the participants in genetic studies had European ancestries, whereas the two major populations, Asian and African, only accounted for 11% and 2.4%, respectively [6]. Indeed, genetic research on the Vietnamese population still lags behind other Western and Asian populations, despite recent efforts of genome sequencing projects in the country [8, 9]. There has been no research to study the prevalence of inherited disorders in the Vietnamese population. Even for well-known diseases in the population such as non-syndromic hearing loss and deafness or Beta-

thalassemia, their exact carrier frequencies and associated pathogenic variants are still

unknown. Carrier screening in Vietnam is still in its infancy.

In this study, we reported the first comprehensive profile of 85 recessive diseases and their

prevalence in the Vietnamese population. The study was performed on the clinical exome

sequencing data of 4,503 genes obtained from a cohort of 985 Vietnamese individuals. We

analyzed the genetic variants obtained from these individuals and identified all pathogenic

variants, genes, associated diseases, and their carrier frequencies in the Vietnamese

population. We also compared the results to other populations and highlighted three diseases

that were found to be specific to the Vietnamese population. Finally, we identified seven novel

pathogenic and three likely pathogenic variants and discussed how they might cause severe

damages in nine associated diseases. Our study made an important step to advance the

practice of medical genetics in Vietnam by providing the first inclusive picture of recessive

diseases in the population and facilitating the development of carrier screening programs in the

country.

MATERIALS AND METHODS

Recruitment of study participants

In this study, 985 individuals were recruited from 51 hospitals and clinics across Vietnam. The

participants have approved and given written informed consent to the anonymous re-use of their

genomic data for this study. The data was de-identified and aggregated for genetic analysis of

the Vietnamese population. The study was approved by the institutional ethics committee of the

University of Medicine and Pharmacy, Ho Chi Minh city, Vietnam

Gene panel

Targeted exome sequencing with a panel of 4,503 clinically relevant genes was performed to

study inherited diseases in the Vietnamese population. The full list of genes is provided in

Supplementary Table S1.

Clinical exome sequencing

Libraries were prepared from 2 ng of DNA using the NEBNext Ultra II FS DNA library prep kit

(New England Biolabs, USA) following the manufacturer's instructions. Subsequently, libraries

were pooled prior to hybridization with the xGen Lockdown probes for 4,503 targeted genes

(Integrated DNA Technologies, USA). Exome sequencing was performed using NextSeq

500/550 High output kits v2 (150 cycles) on Illumina NextSeq 550 system (Illumina, USA) with

the coverage of 100x

Variant calling and analysis

Quality control and alignment of sequencing data to the human reference genome (build

GRCh38) was performed following an established analysis workflow with FastQC [10],

trimmomatic [11], bwa [12], samtools [13], and bedtools [14]. Variant calling was performed

using GATK 3.8, followed by standard filters of quality and sequencing coverage [15]. We also

filtered out variants with allele frequencies less than 0.1% and variants that were located outside

of the target regions of our gene panel. The final variant call set was annotated against dbSNP

(version 151, [16]) and ClinVar (version 20191231, [17]) databases, and was analyzed for their

potential consequences using VEP [18]. Principal component analysis was performed using

PLINK (version 1.9, [19]).

RESULTS

Study cohort

The cohort in our study included 985 participants who were recruited from 51 hospitals and

clinics across Vietnam. The ages and types of samples of the participants are summarized in

Table 1. The average age was 23.8 weeks gestational for fetuses, 4.4 years for children (54%

male, 46% female), and 39.5 years for adults (43% male, 57% female). Among types of

samples, most were blood (65.7%), followed by amniotic fluid (18.3%), buccal swab (12.1%),

placental (1.32%), umbilical cord (0.8%) and others (1.83%).

Summary of genetic variants in the study cohort

The aggregated variant call set from 985 individuals, denoted as G4500, consisted of 67,140

variants, including 61,327 SNPs (91.3%) and 5,813 indels (8.7%). Figures 1a-c show the

comparison of the G4500 call set to that of the KHV population (Kinh in Ho Chi Minh City,

Vietnam) from the 1000 genomes project [7] and the dbSNP database (for this comparison, we

only considered variants located within the target regions of our gene panel). We found that

27,655 variants (41.2%) of the G4500 call set had been reported earlier in the KHV call set

(Figure 1a), and their allele frequencies were consistent between the two call sets with a strong

Pearson correlation of 99.0% (Figure 1c). We also noted that the G4500 call set missed 8,634

KHV variants, and further investigation showed that most (91.7%) of those variants were rare,

appearing only in one single allele in the KHV population. The G4500 call set included 39,485

variants (58.8%) that had not been reported in the KHV call set. Among them, 30,681 (45.7%)

were found in the dbSNP database and the remaining 8,804 (13.1%) were novel. Most of the

novel variants had allele frequencies less than 5% (Figure 1b).

We also performed principal component analysis (PCA) on the G4500 population and other

East Asia populations from the 1000 Genome Project (JPT: Japanese in Tokyo, Japan; CHB:

Han Chinese in Beijing, China; CHS: Southern Han Chinese; CDX: Chinese Dai in

Xishuangbanna, China; KHV: Kinh in Ho Chi Minh City, Vietnam). Overall, Figure 1d shows that

the PCA clustering of the populations was consistent with their respective geographic locations.

The G4500 and KHV populations closely clustered together as both represented the

Vietnamese population. They were also located closer to the CDX population than to the CHS,

CHB, and JPT populations, agreeing with the respective geographical distances.

We then used Variant Effect Predictor (VEP [18]) to predict potential effects of variants in the

G4500 call set (Figure 1e). Majority of them were missense variants (45.1%), followed by

synonymous variants (29.0%), and intron or splice region variants (16.7%). Notably, 4.5% of the

variants were predicted to have high-impact consequences, including stop-gained, stop-lost,

start-lost, frameshift, splice receptor and splice donor. Those high-impact variants may lead to

protein truncation and are critical for clinical interpretation, as we shall show in the next

sections.

Carrier frequencies of genetic diseases in the Vietnamese population

We annotated the G4500 variant call set against the ClinVar database to identify pathogenic

variants, genes, associated diseases, and estimated their carrier frequencies in the Vietnamese

population. We found 21,151 variants with ClinVar annotations, and among them, 158 variants

had been reviewed as "Pathogenic" or "Likely pathogenic". These 158 variants were located on

116 genes: 84 genes were associated with autosomal recessive (AR) diseases, 18 genes with

autosomal dominant (AD) diseases, 9 genes with both AD and AR diseases, one gene with X-

linked dominant disease (XLD) and one gene with X-linked recessive disease (XLR). In this

study, we focused on 114 pathogenic variants on 85 genes that were associated with recessive

diseases (84 AR and one XLR).

Twenty-three individuals in our cohort were identified as homozygous or compound

heterozygous carriers for 5 genes associated with recessive diseases, including *GJB2* (n=12),

HFE (n=5), *VPS13B* (n=4), *CBS* (n=1), and *GBA* (n=1) (Supplementary Table S2). Since our

cohort data was obtained from pre-existing hospital records rather than a randomized study

design, we took a conservative approach and excluded these 23 individuals before calculating

the carrier frequencies of the respective genes and diseases. Overall, the carrier frequencies

were reduced by 0.1%-1% by this exclusion (Supplementary Table S3).

Figure 2 shows a summary of 114 pathogenic variants on 85 genes and associated recessive

diseases identified from our G4500 dataset. The complete details are provided in

Supplementary Table S4. As shown in Figure 2a, majority (54%) of these variants were protein-

truncating (including stop gained, frameshift, splice acceptor or donor), followed by missense

variants (41%). While most of the 85 genes only had one pathogenic variant, 20 of them

(23.5%) had at least two pathogenic variants per gene (Figure 2b), such as *GAA* (5 variants),

GJB2 and *HBB* (3 variants each), *VPS13B* (2 variants), etc (Supplementary Table S4). By

taking into account all pathogenic variants of each gene, our study provided more accurate

estimates of disease carrier frequencies than a targeted genotyping approach that only focused

on major variants [2]. The carrier frequency distribution is presented in Figure 2c. 17/85 genes

(20%) were estimated to have carrier frequencies of more than 1% (1 in 100), among which

seven diseases appeared in more than 2% (1 in 50), including three appeared in more than 5%

(1 in 20) of the Vietnamese population.

Figure 2d shows the top seven genes and associated recessive diseases with carrier

frequencies of more than 2% (1 in 50) in the Vietnamese population. Deafness, autosomal

recessive 1A associated with gene *GJB2* was the most prevalent disorder with a carrier

frequency of 17.2% (1 in 6). The prevalence of *GJB2*, in particular, the SNP rs72474224 C>T, in

the Vietnamese population and other East Asian populations, as compared to Western

populations, had been reported previously in [9]. Two other autosomal recessive diseases were

found with relatively high carrier frequencies, including hemochromatosis type 1 (*HFE*, 9.4% or

1 in 11) and Cohen syndrome (*VPS13B*, 8.1% or 1 in 12). Hemochromatosis type 1 is a

metabolic disorder that causes the body to absorb too much iron (iron overload). Cohen

syndrome is a multisystem disorder characterized by many clinical features, including

developmental delay, intellectual disability and facial dysmorphis. Both diseases are common

genetic disorders among Western populations, but we found that they appeared less frequently

in the Vietnamese population (Table 2). We also observed three other disorders that are among

the most commonly encountered diseases by local medical doctors in Vietnam, including Beta-

thalassemia (*HBB*, 4% or 1 in 25), citrin deficiency (*SLC25A13*, 3% or 1 in 33), and

phenylketonuria (*PAH*, 2.5% or 1 in 40).

Beta-thalassemia, Citrin Deficiency, and Phenylketonuria

We further compared the allele frequencies of pathogenic variants of the top seven diseases-

genes between the Vietnamese, the East Asia, and the global populations (gnomAD [20]).

Figure 2e and Table 2 show that several pathogenic variants appeared 2-12 times more

frequent in the Vietnamese population, especially for three diseases Beta-thalassemia, citrin

deficiency and phenylketonuria. Beta-thalassemia is a blood disorder that reduces the

production of hemoglobin; its major type can lead to severe or life-threatening outcomes and

requires frequent blood transfusions for red blood cell supply. The prevalence and severe

consequences of Beta-thalassemia is well-known among the Vietnamese population, yet no

research has been done to study its genetic patterns in the population. Here we found that the

allele frequency of the SNP rs33950507 C>T in gene *HBB* was 12 times higher in the

Vietnamese population than in the East Asia population (1.57% and 0.13%, respectively).

Furthermore, rs33950507 and two other pathogenic variants in gene *HBB* collectively

contributed to a carrier frequency of 4% (1 in 25) for Beta-thalassemia in the Vietnamese

population. The global carrier frequency of Beta-thalassemia had been estimated previously as

0.7% (1 in 143), i.e. 5.7 times lower than in the Vietnamese population [2].

Another two SNPs, rs192592111 C>A and rs199475650 G>T, in gene *PAH* and associated with

phenylketonuria were also found to have allele frequencies 9 times higher in the Vietnamese

population than in the East Asia population (Figure 2e, Table 2). Phenylketonuria is a metabolic

disorder that causes phenylalanine to build up in the body, and if not treated, may lead to

intellectual disability and other serious health problems. This disease is a very rare genetic

condition in the world with a carrier frequency of 0.7%, mostly observed in Southern Europe or

Hispanic, but not among the East Asia population [2]. However, we found two of its variants and

estimated that its carrier frequency was 2.5% (1 in 40) in the Vietnamese population.

Similarly, we found two SNPs rs80338720 and rs80338725 in gene *SLC25A13* that were

associated with citrin deficiency, and their respective allele frequencies were 2.4 times and 1.6

times higher in the Vietnamese population than in the East Asia population. The total carrier

frequency of citrin deficiency was estimated as 3% (1 in 33) in the Vietnamese population,

which was in line with recent results for South East Asian populations in Singapore [3]. Citrin

deficiency is a metabolic disorder that manifests in newborns as neonatal intrahepatic

cholestasis or in adulthood as recurrent hyperammonemia with neuropsychiatric symptoms in

citrullinemia type II. Without appropriate treatment, severe liver problems may develop and

require liver transplantation.

In addition to the top seven genes with carrier frequencies of more than 2% (1 in 50), ten other

genes had carrier frequencies of at least 1% (1 in 100), and the remaining 68 genes had carrier

frequencies of less than 1% in the Vietnamese population. The complete profile of pathogenic

variants, genes, recessive diseases, and their frequencies in the Vietnamese population is

provided in Supplementary Table S4. Some other examples of high carrier frequencies include

Pompe disease (*GAA*, 1.9% or 1 in 52), Zellweger syndrome (*PEX1*, 1.6% or 1 in 62), Stargardt

disease (*ABCA4*, 1.3% or 1 in 76), Krabbe disease (*GALC*, 1.3% or 1 in 76), Bestrophinopathy,

autosomal recessive (*BEST1*, 1.1% or 1 in 90), and Wilson disease (*ATP7B*, 0.9% or 1 in 110).

Identifying new pathogenic variants for the Vietnamese population

We next attempted to identify new pathogenic variants for the Vietnamese population from the

G4500 call set. We focused on the variants that were predicted by VEP to have high-impact

consequences but had not been reported in ClinVar. We identified 131 variants that may cause

protein truncation, including stop-gained, stop-lost, start-lost, frameshift, and splice receptor or

donor disruptions. Their distribution is presented in Figure 3a. We then manually reviewed these

variants according to the American College of Medical Genetics (ACMG) classification

guidelines [21] and classified seven of them as "Pathogenic" and three as "Likely pathogenic"

variants. Their details are presented in Figure 3b and Supplementary Table S5.

The seven new pathogenic variants include four stop-gained and three frameshift variants that

are rare or not present in public databases. In particular, four of them were found in gnomAD

with global allele frequencies ≤0.1% and three of them were only found in our G4500 dataset.

Their allele frequencies in the Vietnamese population were several times higher than in the East

Asia and the world populations. For instance, the SNP rs185805779 G>A had allele frequencies

of 1.52%, 0.17%, and 0.03% in the Vietnamese, the East Asia, and the global populations,

respectively (Figure 3b). This stop-gained variant in gene *GCNT2* leads to a premature

termination codon p.Trp5Ter at the beginning of the protein NP_663624.1 and disrupts this

whole protein. Similar nonsense, loss-of-function variants in gene *GCNT2* had been reported as

pathogenic and associated with the cataract 13 with adult i phenotype, an autosomal recessive

disorder of i and I antigens in blood that may lead to congenital cataract (OMIM 600429). Thus,

we classified this variant as pathogenic (evidence categories PVS1, PM2 and PM4 in ACMG

guidelines).

Notably, we identified three novel pathogenic variants that had never been reported before in

any databases. In particular, the SNP chr8:93755784 C>A in gene *TMEM67* is a stop-gained

variant that causes a premature termination codon p.Ser77Ter on the protein NP_714915.3.

Two other stop-gained, loss-of-function variants on this gene and its protein had been reported

in ClinVar as pathogenic, including ClinVar 506012 (NP_714915.3:p.Arg172Ter) and ClinVar

1376 (NP_714915.3:p.Arg208Ter). Note that the mutated amino acid of the new SNP is located

at position 77 and hence results in a shorter truncated protein than the other two mutations,

causing even more severe damages. We classified this new SNP as pathogenic for *TMEM67*-

associated Joubert syndrome (OMIM 609884).

Another novel stop-gained variant that we classified as pathogenic was the SNP chr4:78448244

A>T in gene *FRAS1*, which causes a premature termination codon p.Lys2068Ter on the protein

NP_079350.5. Note that a missense variant, rs1578330963 A>G, had been reported at the

same location in dbSNP for the Korean population [22]. Two other stop-gained, loss-of-function

variants in *FRAS1* and NP_079350.5 had been reported in ClinVar as pathogenic, including

ClinVar 197861 (NP_079350.5:p.Arg124Ter) and ClinVar 435260 (NP_079350.5:p.Gln907Ter).

Thus, we classified the new SNP as pathogenic for *FRAS1*-associated Fraser syndrome 1

(OMIM 607830). Fraser syndrome is a rare genetic disorder characterized by cryptophthalmos,

cutaneous syndactyly, and abnormalities of the genitalia and the urinary tract.

Similarly, we classified a novel deletion variant, chr6:152293724 TAGAG>T, as pathogenic for

SYNE1-associated Spinocerebellar ataxia-8. This variant causes a frameshift p.Leu5887fs on

the protein NP_149062.2, for which several loss-of-function frameshift variants had been

reported as pathogenic (ClinVar IDs 204299, 436905, 199228). Spinocerebellar ataxia-8 is a

slowly progressive neurodegenerative disorder characterized by gait ataxia and other cerebellar

signs, such as nystagmus and dysarthria (OMIM 608441).

Last but not least, we classified three new splice acceptor or donor variants as likely pathogenic

(Supplementary Table S5). These variants were predicted to disrupt mRNA splicing and result

in an absent or disrupted protein product. They were not found or appeared at less than 0.01%

frequency in gnomAD. Similar splice acceptor or donor variants on the same genes had been

reported as pathogenic or likely pathogenic. For instance, the SNP rs1183832067 A>C is a

splice donor in gene *RFX5*, and we found that its corresponding splice acceptor rs748270285

G>A for the same exon 6 of transcript NM_001025603.2 had been reported as pathogenic for

Bare lymphocyte syndrome, type II, complementation group c (ClinVar 7646). Since more data

is needed to establish the pathogenicity, we classified the three splice acceptor or donor

variants in Supplementary Table S5 as likely pathogenic (evidence categories PVS1 and PM2 in

ACMG guidelines).

DISCUSSION

In this paper, we analyzed the clinical exome sequencing data of 4,503 genes obtained from a

cohort of 985 individuals to study recessive diseases in the Vietnamese population. We

identified a comprehensive variant call set named G4500 that includes 61,327 SNPs and 5,813

indels. We showed that the G4500 variant call set accurately represented the genetic

characteristics of the Vietnamese population and also demonstrated how they are related to

other East Asia populations.

Most importantly, our work is the first study that provided a comprehensive picture of 85 most

common recessive diseases and their prevalence in the Vietnamese population. Among them,

seven diseases had carrier frequencies of more than 2% (1 in 50) and ten diseases had carrier

frequencies of at least 1% (1 in 100). For each disease, we provided complete details of its

pathogenic variants, gene, and carrier frequency in the Vietnamese population as compared to

other populations. For instance, *GJB2*-associated deafness autosomal recessive was the most

prevalent disorder with a carrier frequency of 17.2% and consisted of three pathogenic variants.

Notably, we found three diseases that were specific to the Vietnamese population with carrier

frequencies of several times higher than in other East Asia or the world populations, including

Beta-thalassemia (*HBB*, 4% or 1 in 25), citrin deficiency (*SLC25A13*, 3% or 1 in 33), and

phenylketonuria (PAH, 2.5% or 1 in 40).

We also discovered seven new pathogenic and three new likely pathogenic variants that had

not been reported in ClinVar. These new variants were associated with nine autosomal

recessive diseases in autoimmune, hematology, ophthalmology, and neurology. Notably, two

new pathogenic variants revealed much higher carrier frequencies of *TMEM67*-associated

Joubert syndrome and *GCNT2*-associated cataract 13 with adult i phenotype in the Vietnamese

population (2.64% and 3.04%, respectively) than previously estimated. Some of these variants

and diseases might also appear in other populations at extremely low frequencies, e.g. *GCNT2*-

associated cataract 13 with adult i phenotype and *RP1L1*-associated retinitis pigmentosa, thus

representing a great resource for further studies. We also discussed how these new variants

were related to previously reported pathogenic variants on the corresponding genes and

proteins.

One limitation of this study was that our cohort was sampled from pre-existing hospital records

rather than a randomized study design. To remove potential bias in our estimation of allele and

carrier frequencies due to this type of sampling, we took a conservative approach by

considering only recessive diseases and excluding 23 individuals identified as homozygous or

compound heterozygous carriers for 5 genes. Thus, our estimated carrier frequencies for these

5 genes and diseases may be considered as lower bounds. A more properly designed study

with sufficiently large dataset could offer a more accurate representative of the Vietnamese

population.

In conclusion, our study has significantly improved the knowledgebase and the practice of

medical genetics in Vietnam in many aspects. Our findings offer a great resource to inform local

public health policies to understand and better align with the specific landscape of genetic

diseases in the Vietnamese population. Carrier or newborn genetic screening programs can be

re-designed for cost-effectiveness and comprehensiveness. The results also help clarify and

expand existing knowledge of popular inherited diseases in the local population by providing the

extra dimension of molecular genetic information. By demonstrating the underlying fundamental role of genetics in inherited diseases, our work also contributes to the development of genetics education, genetics counseling, and genetics screening among the local population. The identification of three inherited diseases specific to the Vietnamese population affirms the necessity of population-specific genetic studies and that larger and more comprehensive population genetic studies dedicated to the Vietnamese population are highly desired.

Ethics approval and consent to participate

The study was approved by the institutional ethics committee of the University of Medicine and Pharmacy, Ho Chi Minh city, Vietnam. The study has followed the guidelines set by the University of Medicine and Pharmacy, Ho Chi Minh city, Vietnam, in handling human genetic data of the participants. The participants have approved and given written informed consent to the anonymous re-use of their genomic data for this study.

Consent for publication

All authors have read and approved the manuscript for publication.

Availability of data and materials

The G4500 variant call set is available upon reasonable request to the corresponding authors, subject to our policy of data privacy.

Competing interests

This study was funded by Gene Solutions, Vietnam. The funder did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

NHT, HST, HTN, LPD, NMP, KHTN, HDLN, MTTQ, TPTN, VUT, PTCN, HG and MDP are

current employees of Gene Solutions, Vietnam. The other authors declare no competing

interests.

Authors' contributions

THNT, HST, LPH, THLN, NTT, THNT, VTN, BHHN, NMP. KHTN, HDLN, MTTQ, TPTN, DVT,

QTTN recruited patients and performed clinical analysis.

HNN, TTTD, NVL, VUT, PCTN, DKT, HTN, LPD, designed experiments and analyzed data.

NHT, HG, MDP designed the experiments, analyzed the data and wrote the manuscript.

HNN supervised the project.

References

1. Antonarakis, S.E. Carrier screening for recessive disorders. *Nat. Rev. Genet.* **20**, 549-561

(2019).

2. Lazarin, G.A. *et al.* An empirical estimate of carrier frequencies for 400+ causal Mendelian

variants: results from an ethnically diverse clinical sample of 23,453 individuals. *Genet.*

Med. **15**, 178-186 (2013).

3. Bylstra, Y. *et al.* Population genomics in South East Asia captures unexpectedly high

carrier frequency for treatable inherited disorders. *Genet. Med.* **21**, 207-212 (2019).

4. Hamosh, A. *et al.* Online Mendelian Inheritance in Man (OMIM), a knowledgebase of

human genes and genetic disorders. *Nucleic Acids Res.* **33**, D514-D517 (2005).

5. Editorial. Diversity matters. *Nat. Rev. Genet.* **20**, 495 (2019).

6. Gurdasani, D., Barroso, I., Zeggini, E. & Sandhu, M.S. Genomics of disease risk in globally

diverse populations. *Nat. Rev. Genet.* **20**, 520-535 (2019).

7. The 1000 Genomes Project Consortium. A global reference for human genetic variation.

Nature **526**, 68-74 (2015).

8. Le, V.S. *et al.* A Vietnamese human genetic variation database. *Hum. Mutat.* **40**, 1664-

1675 (2019).

9. Tran, N.H. *et al.* Genetic profiling of Vietnamese population from large-scale genomic

analysis of non-invasive prenatal testing data. bioRxiv 868588; doi:

https://doi.org/10.1101/868588 (2020).

10. FastQC: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

11. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

12. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997v2 [q-bio.GN].

13. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-

2079 (2009).

14. Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841-842 (2010).

15. Van der Auwera, G.A. *et al.* From FastQ data to high confidence variant calls: the Genome

Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1-11.10.33

(2013).

16. Sherry, S.T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* **29**,

308-311 (2001).

17. Landrum, M.J. *et al.* ClinVar: public archive of relationships among sequence variation and

human phenotype. *Nucleic Acids Res.* **42**, D980-D985 (2014).

18. McLaren, W. The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).

19. Chang, C.C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015).

20. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**,

285-291 (2016).

21. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a

joint consensus recommendation of the American College of Medical Genetics and

Genomics and the Association for Molecular Pathology. *Genet. Med.* **17**, 405-424 (2015).

22. Jeon, S. *et al.* Korean Genome Project: 1094 Korean personal genomes with clinical

information. *Sci. Adv.* **6**, eaaz7835 (2020).

Figure 1. Summary of the G4500 variant call set. (a) Comparison of G4500, KHV, and dbSNP. (b) Allele frequency distribution of the G4500 call set. (c) Comparison of allele frequency between G4500 and KHV. (d) Principal component analysis of the G4500 call set and other East Asia populations (JPT: Japanese in Tokyo, Japan; CHB: Han Chinese in Beijing, China; CHS: Southern Han Chinese; CDX: Chinese Dai in Xishuangbanna, China; KHV: Kinh in Ho Chi Minh City, Vietnam). (e) Distribution of variant consequences of the G4500 call set (high-impact: stop-gained, stop-lost, start-lost, frameshift, splice receptor, and splice donor).

Figure 2. Summary of pathogenic variants, genes, and associated recessive diseases identified from the G4500 dataset. (a) Distribution of coding consequences of pathogenic variants. (b) Distribution of pathogenic variants per gene. (c) Distribution of carrier frequencies of pathogenic genes. (d) Top seven diseases-genes with carrier frequencies of more than 2%. (e) Allele frequencies of pathogenic variants of the top seven diseases-genes in the Vietnamese G4500 population (x-axis) and how they are compared to the frequencies in the East Asia population (y-axis). Some genes may have multiple variants, e.g. *GJB2* has three variants. Red points indicate variants with allele frequencies different by more than two folds between the two populations (i.e. $\log_2$ fold change is less than -1 or greater than 1). The details of these variants are provided in Table 2.

(a)

(b)

variant ID	rs ID	G4500 AF	gnomAD EAS AF	gnomAD AF	gene	consequence	amino acid change	OMIM	disease
in KHV, dbSNP, and G4500									
chr6_10528925_G_A	rs185805779	1.52%	0.17%	0.03%	GCNT2	stop gained	NP_663624.1:p.Trp5Ter	600429	Cataract 13 with adult i phenotype
in dbSNP and G4500 only									
chr8_10607568_A_C	rs777475406	0.61%	0.13%	0.01%	RP1L1	stop gained	NP_849188.4:p.Leu2177Ter	608581	Retinitis pigmentosa 88
chr8_10610076_GTT_G	rs1491506199	0.61%	0.07%	0.10%	RP1L1	frameshift	NP_849188.4:p.Glu1340fs	608581	Retinitis pigmentosa 88
chr10_123041350_C_CT	rs755014798	0.66%	n.a.	n.a.	ACADSB	frameshift	NP_001600.1:p.Val219fs	600301	2-methylbutyrylglycinuria
in G4500 only									
chr8_93755784_C_A	n.a.	1.22%	n.a.	n.a.	TMEM67	stop gained	NP_714915.3:p.Ser77Ter	609884	Joubert syndrome 6
chr6_152293724_TAGAG_T	n.a.	0.51%	n.a.	n.a.	SYNE1	frameshift	NP_149062.2:p.Leu5887fs	608441	Spinocerebellar ataxia, autosomal recessive 8
chr4_78448244_A_T	n.a.	0.61%	n.a.	n.a.	FRAS1	stop gained	NP_079350.5:p.Lys2068Ter	607830	Fraser syndrome 1

Figure 3. (a) Distribution of variants identified from the G4500 dataset that had high-impact consequences but had not been reported in the ClinVar database. (b) Seven new pathogenic variants that we selected from (a), reviewed, and classified as "pathogenic" according to the ACMG guidelines. (AF: allele frequency; EAS: East Asia; ACMG: American College of Medical Genetics).

Table 1. Summary of participants (n=985)

Types of ages	Types of samples						OVERALL TOTAL
	Blood	Amniotic fluid	Placental	Umbilical cord	Buccal swab	Others	
Fetus	0	180	13	8	0	0	201 (20.4%)
Child (age<18)	212	0	0	0	90	5	307 (31.2%)
Adult (age≥18)	435	0	0	0	29	13	477 (48.4%)
OVERALL TOTAL	647 (65.7%)	180 (18.3%)	13 (1.3%)	8 (0.8%)	119 (12.1%)	18 (1.8%)	

Table 2. Pathogenic variants of the seven most prevalent diseases-genes in the Vietnamese population.
(AF: allele frequency; CF: carrier frequency; EAS: East Asia)

variant ID	rs ID	G4500 AF	gnomAD EAS AF	gnomAD AF	gene	ClinVar ID	ClinVar disease
chr13_20189473_C_T	rs72474224	8.17%	8.54%	0.35%	GJB2	17023	Deafness, autosomal recessive 1A
chr13_20189346_AG_A	rs80338943	0.31%	0.93%	0.02%	GJB2	17014	Deafness, autosomal recessive 1A
chr13_20189281_CAT_C	rs111033204	0.15%	0.13%	0.00%	GJB2	44736	Deafness, autosomal recessive 1A
chr6_26090951_C_G	rs1799945	4.69%	2.65%	10.13%	HFE	10	Hemochromatosis type 1
chr8_99832368_G_T	rs386834119	3.72%	6.32%	4.50%	VPS13B	56699	Cohen syndrome
chr8_99832367_A_T	rs74485751	2.24%	7.57%	7.08%	VPS13B	555020	Cohen syndrome
chr11_5226943_C_T	rs33950507	1.57%	0.13%	0.03%	HBB	15161	Beta Thalassemia
chr11_5226970_T_A	rs33986703	0.25%	0.08%	0.01%	HBB	15401	Beta Thalassemia
chr11_5226762_CAAAG_C	rs80356821	0.15%	0.22%	0.01%	HBB	15417	Beta Thalassemia
chr7_96189371_TCATA_T	rs80338720	1.47%	0.61%	0.01%	SLC25A13	225472	Citrin Deficiency
chr7_96121928_G_GCCCG GGCAGCCACCTGTAATCTC	rs80338725	0.10%	0.06%	0.00%	SLC25A13	6003	Citrin Deficiency
chr12_102855326_C_A	rs192592111	1.17%	0.13%	0.00%	PAH	664621	Phenylketonuria
chr12_102846924_G_T	rs199475650	0.10%	0.01%	0.00%	PAH	102904	Phenylketonuria
chr2_31529325_C_T	rs9332964	1.17%	0.73%	0.02%	SRD5A2	3351	5-alpha reductase deficiency