Revealing tissue-specific metabolic crosstalk after a myocardial infarction

Muhammad Arif[1, #], Martina Klevstig[2, #], Rui Benfeitas[3], Stephen Doran[4], Hasan Turkez[5], Mathias Uhlén[1], Maryam Clausen[6], Johannes Wikström[7], Damla Etal[6], Cheng Zhang[1], Malin Levin[2], Adil Mardinoglu[1, 4, *], Jan Boren[2, *]

[1]Science for Life Laboratory, KTH - Royal Institute of Technology, Stockholm, SE-17121, Sweden.

[2]Department of Molecular and Clinical Medicine, University of Gothenburg, The Wallenberg Laboratory, Sahlgrenska University Hospital, Gothenburg, Sweden.

[3]National Bioinformatics Infrastructure Sweden (NBIS), Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, S-10691 Stockholm, Sweden

[4]Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral & Craniofacial Sciences, King's College London, London, SE1 9RT, United Kingdom.

[5]Department of Medical Biology, Faculty of Medicine, Atatürk University, Erzurum, 25240, Turkey.

[6]Translational Genomics, BioPharmaceuticals R&D, Discovery Sciences, AstraZeneca, Gothenburg, Sweden.

[7]Bioscience Cardiovascular, Research and Early Development, Cardiovascular, Renal and Metabolism (CVRM), BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden

[#]Contributed equally.

*Corresponding authors: Adil Mardinoglu and Jan Boren

Email: adilm@scilifelab.se; Jan.Boren@wlab.gu.se

Abstract/Summary

Myocardial infarction (MI) promotes a range of systemic effects, many of which are

unknown. Here, we investigated the alterations associated with MI progression in heart and

other metabolically active tissues (liver, skeletal muscle, and adipose) in a mouse model of

MI (induced by ligating the left ascending coronary artery) and sham-operated mice. We

performed a genome-wide transcriptomic analysis on tissue samples obtained 6- and 24-hours

post MI or sham operation. By generating tissue-specific biological networks, we observed:

(1) dysregulation in multiple biological processes (including immune system, mitochondrial

dysfunction, fatty-acid beta-oxidation, and RNA and protein processing) across multiple

tissues post MI; and (2) tissue-specific dysregulation in biological processes in liver and heart

post MI. Finally, we validated our findings in independent MI cohorts, using both bulk and

single-cell transcriptomic data. Overall, our integrative analysis highlighted both common and

specific biological responses to MI across a range of metabolically active tissues.

Keywords

Systems biology; network analysis; whole-body modelling; cardiovascular disease;

Myocardial infarction; multi-tissue; metabolically active tissues; liver; adipose; muscle

Introduction

Cardiovascular disease (CVD) is the leading cause of death worldwide, accounting for more than 17 million deaths globally in 2016 (WHO, 2019). Myocardial infarction (MI) is one of the most common causes of CVD-related death, and is the result of severe coronary artery disease that develops from tapered arteries or chronic blockage of the arteries caused by accumulation of cholesterol or plaque (atherosclerosis). MI has been linked to multiple behavioral risk factors (including unhealthy diet, physical inactivity, excessive use of alcohol, and tobacco consumption) that can lead to significant alterations in metabolism that are responsible for hypertension, obesity, diabetes, and hyperlipidemia. These abnormalities are known as the high-risk factors of MI and CVDs in general.

Systems biology has been used in many studies to reveal the underlying molecular mechanisms of complex human diseases and to answer important biological questions related to the progression, diagnosis and treatment of the diseases. The use of systems biology has aided the discovery of new therapeutic approaches in multiple diseases (Mardinoglu et al., 2017b; Mardinoglu and Nielsen, 2015; Nielsen, 2017) by identifying novel therapeutic agents and repositioning of existing drugs (Turanli et al., 2019). Systems biology has also been employed in the identification of novel biomarkers, characterization of patients and stratification of heterogenous cancer patients (Benfeitas et al., 2019; Bidkhori et al., 2018; Lee et al., 2016). Specifically, integrated networks (INs) (Lee et al., 2016) and co-expression networks (CNs) (Lee et al., 2017) have been proven to be robust methods for revealing the key driver of metabolic abnormalities, discovering new therapy strategies, as well as gaining systematic understanding of diseases (Bakhtiarizadeh et al., 2018; Mukund and Subramaniam, 2017).

Previously, multiple studies in individual tissues have been performed and provided new insights into the underlying mechanisms of diseases (Das et al., 2019; Ounzain et al., 2014; Pedrotty et al., 2012; Williams et al., 2018). However, the crosstalk between different tissues and their dysregulation has not been examined in MI and other CVD-related complications (Priest and Tontonoz, 2019). Here we performed an integrated analysis of heart and other metabolically active tissues (liver, skeletal muscle and adipose tissue) using a mouse model of MI. We used several systems biology approaches to obtain a systematic picture of the

83   metabolic alterations that occur after an MI (**Figure 1A**), and validated our findings in human

84   and mouse datasets.

Results

Differential expression analysis shows a pronounced effect on gene expression 24 h post MI

To study global biological alterations and systemic whole-body effects associated with MI, we obtained heart, liver, skeletal muscle, and white adipose tissue from mice 6 h and 24 h after either an MI (induced by ligating the left ascending coronary artery) or a sham operation (as control). We generated transcriptomics data and identified differentially expressed genes (DEGs) 6 and 24 h post MI and sham operation in all tissues, with the most significant differences occurring after 24 h **(Table S1, Figure 1B)**. Principal component analysis (PCA) showed a close clustering between the control (for both time points) and MI (6 h and 24 h separately) samples for heart tissue but clustering by extraction time points (6 h and 24 h clusters) for the other tissues (**Figure S1**). We present the transcriptional changes associated with MI in **Table S1** and the DEGs (FDR < 5%) using an UpSet plot (Lex et al., 2014) in **Figure 1C**.

All tissues showed a more pronounced effect in terms of the number of DEGs 24 h post MI (**Figure 1C**). As expected, the most affected tissue was the heart (393 DEGs at 6 h, 3318 DEGs at 24 h, and 318 DEGs were the same at both time points). By contrast, 136, 641 and 374 genes were significantly changed in liver, skeletal muscle and adipose tissues 24 h post MI compared to control, respectively. More than 33% of the DEGs that significantly changed in the other tissues also changed in the heart (**Figure 1C**). Interestingly, more than 97% of the shared DEGs between heart and skeletal muscle changed in the same direction, with corresponding numbers of 88% and 64% in adipose and liver, respectively.

Functional analysis reveals widespread mitochondrial, fatty acid, immune, and protein and RNA-related alterations post MI with liver shows contrasting trend

We performed gene-set enrichment analysis (GSEA) with KEGG pathways (**Table S2, Figure 1D**) and gene ontology (GO) biological processes (BPs) (**Table S3, Figure 2A**) to identify altered biological functions and pathways 24 h after an MI. Mitochondrial functions (specifically, mitochondrial translation, respiratory chain and oxidative phosphorylation) were significantly downregulated in the heart, muscle and adipose tissues but not in the liver. Processes related to oxidative stress were upregulated in the heart and skeletal muscle. Fatty acid beta-oxidation was downregulated in the heart and adipose but upregulated in the liver.

Processes and pathways related to immune systems were significantly upregulated in the heart
and skeletal muscle but significantly downregulated in liver. Processes associated with protein
and RNA processing, ribosome biogenesis and protein targeting endoplasmic reticulum were
upregulated in all tissues except liver whereas protein processing in endoplasmic reticulum
and RNA transport pathways were upregulated in all tissues.
We also observed that liver was showing opposite trends compared to the other tissues in
other important functions, such as fatty acid metabolism and immune response. By checking
regulation at the gene level, we observed that only 16 DEGs in liver showed opposite
regulation compared to the other tissues whereas 97 out of the 136 DEGs in liver were not
DEGs in any other tissues (**Table S4**). Therefore, the differences we observed in liver were
mainly due to different DEGs rather than opposite regulation compared to other tissues.
129

Tissue-specific altered biological functions point to specificity of metabolic and signaling
responses to MI

The functional analysis also indicated that several metabolic pathways (including cholesterol,
ascorbate and aldarate, linoleic acid, and sphingolipid metabolism pathways) and signaling
pathways (including GnRH, FoxO, cAMP and prolactin signaling pathways) were
significantly upregulated in heart 6 h after an MI (**Table S2**, **Figure S2A**). We also observed
significant down regulation of tryptophan metabolism and upregulation of glycosaminoglycan
biosynthesis in heart 24 h after an MI (**Table S2**, **Figure S2A**). Processes related to retinol
metabolism were upregulated in heart at both timepoints. Pathways that were previously
associated with cardiac hypertrophy and cardiac remodeling (e.g. JAK-STAT, MAPK,
estrogen, and TNF signaling pathways, and ECM-receptor interaction) were significantly
upregulated in heart 6 and 24 h after an MI (**Figure S1B**).
142

Our analysis also indicated significant metabolic differences in adipose tissue 24 h after an MI
(**Figure S2B**). Fructose and mannose metabolism, glyoxylate and dicarboxylate metabolism,
glycolysis/gluconeogenesis, and pentose phosphate pathways, glycine, serine and threonine
metabolism and pyrimidine metabolism, as well as endocrine systems (e.g. insulin signaling
pathway and regulation of lipolysis in adipocytes) were downregulated in adipose tissue.
We observed that the PPAR signaling pathway was upregulated whereas glutathione was
downregulated in liver 24 h post-infarction (**Figure S2B**). We found that sphingolipid

metabolism and immune-related pathways were upregulated in skeletal muscle 24 h post-

infarction (**Figure S2B**).

Reporter metabolite analyses show significant alterations in fatty acid, amino acid, retinol,

and estrogen metabolism post MI

To predict the effect of the transcriptional changes on metabolism, we performed reporter

metabolite analyses (**Table S5**) using the gene-to-metabolites mapping from the Mouse

Metabolic Reaction database (Mardinoglu et al., 2015); results in each tissue 24 h after MI are

shown in **Figure 2B**. In agreement with our analyses above, reporter metabolites related to

oxidative phosphorylation, such as *ubiquinol*, *ubiquinone*, *NADH* and *NAD+*, were

downregulated in all tissues except liver. Moreover, *linolenoyl-CoA, acetyl CoA,* and several

other fatty acyl-CoA-related metabolites were downregulated in heart and adipose tissue but

upregulated in liver. We also found that several *5-S-glutathionyl* metabolite forms, known to

be related to phenylalanine, tyrosine and tryptophan biosynthesis, were downregulated in

heart, liver and skeletal muscle. The same pattern of downregulation was also observed for

metabolites related to estrogen metabolism, specifically metabolites related to oestrone and its

glutathione conjugate derivative. Moreover, *12-keto-LTB4* and *12-oxo-c-LTB3*, related to

leukotriene metabolism, and *hepoxilin A3,* an arachidonic acid, were also found to be

downregulated in heart, liver, and skeletal muscle.

The liver showed the highest alteration in reporter metabolites, which is attributed to its role

as one of the most metabolically active tissues. We found that several reporter metabolites

related to retinol metabolism, namely *retinal*, *retinol*, *retinoate*, and *all-trans-18-*

hydroxyretinoic acid, were significantly downregulated only in liver tissue. Retinol

metabolism has been previously associated with MI (Lima et al., 2018; Palace et al., 1999).

Network analyses unveil universal and tissue-specific clusters and mechanisms post MI

The use of co-expression network (CN) analyses can assist in elucidating the functional

relationships between genes in a specific cell and tissue (Lee et al., 2017). Here, we

performed CN analysis to reveal the functional relationship between the DEGs by generating

tissue-specific CNs and selected highly connected genes (the top 5% positively correlated

genes that fulfilled FDR < 0.05) (**Table 1**). To better define the structure of the networks, we

used the Leiden clustering algorithm (Traag et al., 2019) by maximizing the modularity scores

(**Figure 3A-D**) and selected the clusters that include more than 30 genes. Next, we

superimposed DEGs 24 h post-infarction onto the network (**Table S1**) and identified the

components of the clusters that were affected by an MI. We also used functional analysis with

GO BP and KEGG pathways to understand the specific functions associated with each cluster

by using the Enrichr algorithm (FDR < 0.05) (Chen et al., 2013; Kuleshov et al., 2016). We

summarized the GO BP terms with Revigo (**Table S6**) (Supek et al., 2011) and checked the

average clustering coefficient to define the centrality of each cluster (**Table S6**) (Lee et al.,

2017). Among the clusters, we identified the key clusters as those with the highest average

clustering coefficient, allowing us to identify sets of genes whose time-dependent coordinated

changes showed the strongest relationships.

Interestingly, key clusters contained genes with similar functionalities including RNA

processing, transports, and RNA metabolic processes in all tissue-specific CNs (**Table S6**). In

addition, we found that the majority of the DEGs associated with those clusters were

significantly upregulated. These observations strengthen the findings of the functional

analysis above (**Figure 2A**) and further highlight how embryonically distinct tissues display

similar functional responses to MI, with the most highly connected groups of genes preserved

between different tissues (**Table S6**, **Figure 3E**).

Community detection reveals tissue-specific clusters post MI

We investigated the tissue specificity of each cluster by performing enrichment analysis with

data from the Mouse Gene Atlas (Su et al., 2004), which involved counting the number of

tissue-specific genes.

The heart network showed the highest number of tissue-specific genes in cluster Heart-3 (302

genes). Based on DEG analysis, we found that 522 genes were downregulated and 192 genes

were upregulated in the cluster. The enriched GO BP terms in the cluster were mitochondrial

transport, protein processing and respiratory chain, cardiac muscle cell action potential,

response to muscle stretch, and heart contraction (**Figure 3F**). We observed that the results of

the KEGG pathway enrichment analysis were consistent with those obtained from GO BP

analysis (**Table S6**).

In the liver network, cluster Liver-2 showed the highest tissue specificity (479 genes). In this

cluster, we found that 15 genes were significantly downregulated and 17 genes were

significantly upregulated. Based on GO BP enrichment analysis, the genes in this cluster were

associated with cholesterol metabolism and homeostasis, lipid transport, glutathione

metabolism, lipoprotein metabolism, and glucose 6-phosphate metabolism (**Table S6**). KEGG

enrichment analysis also showed that the genes in the cluster were related to retinol,

carbohydrate, lipid and amino-acid metabolism (**Table S6**).

The muscle network had two clusters with high tissue specificity: cluster Muscle-4 (276

genes) and Muscle-5 (143 genes). Muscle-4 showed association with GO BP terms such as

mitochondrial transport, protein processing and respiratory chain, response to muscle stretch,

and muscle contraction (**Table S6**). In contrast, the KEGG pathway in this cluster showed

relation to glycolysis/glucogenesis, propanoate metabolism, glyoxylate and dicarboxylate

metabolism, and several signaling pathways (e.g. oxytocin, glucagon, cGMP-PKG and HIF-1)

(**Table S6**). Muscle-5 was enriched in GO BP terms associated with protein

dephosphorylation, muscle contraction and intracellular protein transport (**Table S6**). We also

found that insulin, MAPK and Wnt signaling pathways were associated to Muscle-5 from the

KEGG enrichment analysis (**Table S6**).

The adipose tissue network showed tissue specificity in cluster Adipose-2 (33 genes), which

is associated with GO BP processes including mRNA processing, regulation of mitotic cell

cycle phase, ribosome biogenesis, and viral processes (**Table S6**). We observed that the

results of the KEGG pathway enrichment analysis were consistent with those obtained from

GO BP analysis, with additional associations with multiple signaling and regulatory pathways

(**Table S6**).

Tissue-specific clusters show important tissue-specific changes post MI

To understand the specific behavior of each tissue, we further studied the tissue-specific

clusters in the CNs (**Figure 4A**). Heart specific cluster, Heart-3, was driven by several central

genes including *Pln*, *Pde4b*, and *Atp2a2* (related to regulation of cardiac muscle contraction)

and Pdha*1* and *Vdac1* (related to mitochondrial functions). These genes were also found to be

significantly differentially expressed in heart 24 hours post MI (**Table S1**). Genes in the

heart-specific cluster were related to multiple other processes/pathways, e.g. oxytocin

signaling pathway, and several metabolic pathways (glycogen, inositol phosphate and purine)

(**Table S6**).

Mitochondrial dysfunction in the heart leads to disturbance of energy (ATP) production

(Kiyuna et al., 2018; Palaniyandi et al., 2010) and, in the presence of oxygen, to

accumulation of reactive oxygen species (ROS), which can cause oxidative stress. *Vdac1*, a

key gene for regulation of mitochondria function and one of the central genes in the heart-

specific cluster (see above), is significantly downregulated in MI (Camara et al., 2017). *Vdac1*

is located in the outer mitochondrial membrane and is involved directly in cardioprotection

(Schwertz et al., 2007) within the cGMP/PKG pathway (**Figure S3A**). In the same pathway,

we also observed down-regulation of the reporter metabolite hydrogen peroxide (**Table S5**), a

ROS that is related to cardioprotection (Schwertz et al., 2007; Yada et al., 2006). We also

observed downregulation of *Pdha1*, which is known to have a substantial role in both the

HIF-1 signaling pathway and the pyruvate metabolism pathway that converts pyruvate to

acetyl-CoA in the mitochondria (**Figure S3B**). Acetyl-CoA is used in the TCA cycle to

produce NADH and FADH2, which are both needed for ATP production and were

downregulated in our reporter metabolite analysis of the heart. Our findings are thus

consistent with dysfunctional mitochondria and ATP production in the heart in response to an

MI. *Pdha1* has been also been linked to the heart sensitivity during to ischemic stress, where

its deficiency can compromise AMP-activated protein kinase activation (Sun et al., 2016).

In skeletal muscle and adipose tissue, we found that central genes in their respective tissue-

specific clusters related to fatty acid metabolism and lipid metabolism were significantly

altered (**Table S6**, **Figure 5**). In liver-specific cluster, we found that their central genes were

related to fatty-acid beta oxidation (*Cyp4a31*, *Cyp4a32*) and glutathione metabolism (*Gstm3*)

(**Table S6**, **Figure 5A**). Alterations of fatty acid beta-oxidation and glutathione metabolism

have previously been reported in non-alcoholic fatty liver disease, a known risk factor of

CVD (Alexander et al., 2019; Mardinoglu et al., 2017a). Moreover, in liver, we also found

that retinol metabolism was uniquely related to genes in the liver-specific cluster, mainly

driven by four significantly differentially expressed central genes of the clusters, i.e.

Cyp26a1, *Cyp4a31*, *Cyp4a32*, and *Hsd17b6* (**Table S6**). A previous study showed that

mortality from CVD in older individuals was accompanied by impaired liver ability to store

retinol (Lima et al., 2018).

Multi-tissue modeling reveals key metabolic pathways affected post MI

To investigate the metabolic responses to MI in and across tissues in the mice, we constructed

a multi-tissue genome-scale metabolic model. The model consisted of five tissue-specific

genome scale metabolic models, namely heart, liver, skeletal muscle, adipose, and small

intestine. The small intestine model (for which we do not have transcriptomic data) was added

to include ingestion and conversion of dietary nutrients into chylomicrons, which are directly

secreted into blood and transport lipids to other tissues (Mardinoglu et al., 2015). The final

mouse multi-tissue model included 19,859 reactions, 13,284 metabolites, 7,116 genes and 41

compartments. We predicted the metabolic fluxes in mice 24 h after an MI or sham operation

by integrating the dietary input, tissue-specific resting energy expenditure and transcriptomics

data.

The modeling showed that oxygen uptake, carbon dioxide production and the oxidative

phosphorylation pathway in heart, adipose and skeletal muscle were decreased in MI mice, in

agreement with the downregulation of oxidative phosphorylation we observed in these tissues

(**Table S7**). By contrast, liver showed slightly increased oxygen uptake, which might due to

the slightly (not statistically significant) upregulated oxidative phosphorylation (**Table S7**).

These findings indicate that the changes in oxygen and carbon dioxide fluxes and the

oxidative phosphorylation pathway could serve as a positive control for predicting the

changes due to MI in the fluxes.

Next, we investigated the tissue-specific metabolic flux changes in the same model (**Table**

S7). We found that the pentose phosphate pathway was upregulated in heart 24 hours post MI,

consistent with upregulated glucose metabolism after an MI. Elevated glycolysis could allow

the heart to rapidly generate energy under stress conditions, and the enhanced pentose

phosphate pathway could increase the NADPH level, which could help maintain the level of

reduced glutathione in heart (Tran and Wang, 2019). We also found that adipose tissue

secreted more ketone bodies, including acetoacetate and butyrate, into plasma; the plasma

level of ketone bodies has been reported as a stress marker in acute MI (Miyamoto et al.,

1999). Notably, relatively small metabolic changes were found in liver and skeletal muscle,

which is probably due to the small number of transcriptomic changes in metabolic pathways

in these tissues.

Single-cell analysis highlights cell-type patterns among central genes in a heart-specific

cluster

We then questioned whether and how our gene set from both central and heart-specific

clusters identified by bulk RNA-seq data would perform at the single cell level. We examined

the DEGs from most central and heart-specific cluster genes in an independent single-cell

dataset of heart failure in human (Wang et al., 2020). We observed that the majority of the

heart-specific genes clustered together in both cardiomyocyte- (CM) and non-CM-enriched

data during heart failure, with higher expression levels in CM-enriched data (**Figure S4**). On

the other hand, the genes from the most central cluster showed no particular clustering

pattern. We also analyzed the clustered tissue-specific genes, and found three distinct groups:

group 1, which consisted of genes related to the TCA cycle, glycolysis and HIF-1 signaling

pathways; group 2, which consisted of genes with higher expression in all cell types and were

related to heart-specific functions and propanoate and pyruvate metabolism; and group 3,

which consisted of genes related to cholesterol and lipid metabolism and cGMP-PKG

signaling pathways.

Although these gene sets were identified through analysis of bulk RNA-seq data, a clustering

analysis through UMAP (McInnes et al., 2018) highlighted clear distinctions between CM-

and non-CM-enriched data at the single-cell level (**Figure 6A**). We found that CM-enriched

cells (consisting of ~43% smooth muscle cells and ~36% fibroblasts) (**Figure 6B, top left**)

were dominated by the mean expression of genes from the heart-specific cluster (**Figure 6C**),

with very high expression of genes in groups 1 and 2 (**Figure 6D-E**). By contrast, non-CM

cells showed a large heterogeneity in terms of cell type (**Figure 6B, right**). Furthermore, this

cluster showed low mean expression of gene groups 1 and 3, and lower mean expression of

gene group 2 than did CM-enriched cells (**Figure 6D-F**). Approximately 22% of the non-CM

cells were dominated by genes from the most central cluster (**Figure 6C**), whereas 76% of

those cells were endothelial cells. Finally, we observed that the central cluster (**Figure 6C**

orange) tended to display overall lower expression of gene group 2 (**Figure 6E bottom**

right). Altogether, these observations indicate that heart-specific and central clusters tend to

display distinct patterns of gene expression and distribution among cell subtypes, both in CM

and non-CM-enriched cells.

Validating our findings with publicly available datasets

We validated our observations in heart tissue in two independent cohorts of bulk RNA-seq

data from mouse heart (**Table S8**). We filtered both validation cohorts to get and analyzed

only 24 hours post-MI data. We found that there were 2169 DEGs in heart 24 h after

infarction from our data were validated in at least one of the independent cohort (959 in both)

(**Figure 6G**). We also found that 109 out of the 123 most connected genes in our heart-

specific cluster were also significantly differentially expressed in at least one of the

independent cohorts (81 in both). By performing functional analysis of the validation cohorts,

we found that ~61% of GO BP and 84% of KEGG pathways identified in our analysis of the

heart were also present in at least one of the validation cohorts 24 h after infarction (**Figure

6H-I**). In both cohorts, we observed downregulation of mitochondrial functions and fatty acid

metabolism processes. We also observed upregulation of processes and pathways related to

retinol metabolism and inflammatory response in both validation cohorts.

Identification of driver genes in MI

We observed that *Flnc*, *Lgals3*, *Prkaca* and *Pprc1* showed important response to MI. These

genes were 4 of 16 genes that were DEGs in at least three tissues and validated in both

validation cohorts (**Table S9**). *Flnc*, *Lgals3* and *Pprc1* were upregulated in heart, skeletal

muscle, and adipose, whereas *Prkaca* was downregulated in these three tissues. We further

retrieved their neighbors at each tissue specific CNs, showed their regulations from

differential expression results, and performed functional analysis in **Table S9**.

Flnc, which encodes filamin-C, was part of heart and skeletal muscle-specific CN cluster and

also part of gene group 2 at the single-cell level (**Figure S4**). Its neighbor genes were found to

be significantly (FDR < 0.05) associated to several functions, including TCA cycle, pyruvate

metabolism, glycolysis pathway, and involved in mitochondrial functions. Specifically, they

were related to heart-specific processes in heart, VEGF signaling pathway in muscle,

carbohydrate metabolism in adipose, and to MAPK signaling pathway and muscle contraction

in heart and muscle.

Lgals3 (encodes galectin-3) and *Prkaca* were among the most central genes in central clusters

(**Table S6**). The neighbors of *Lgals3* were significantly related to cell cycle and protein

digestion and absorption pathway in all tissues, and to RNA and mRNA related-processes in

muscle and adipose tissue. The neighbors of *Prkaca* were related to insulin signaling pathway

in heart and adipose, and several mitochondrial functions in adipose. *Pprc1* was part of most

central clusters in heart and adipose tissue CN, and its neighbors were related to ribosomal

RNA processing and ribosome biogenesis.

The median expression of these genes at the single-cell level is shown in **Table S9, Figure**

S5. We found that all genes were expressed in all cell types in CM and non-CM enriched

cells, except *Pprc1*. In CM cells, *Flnc* showed highest median expression in macrophages,

whilst *Prkaca* showed highest median expression in fibroblasts. *Lgals3* had similar median

expression in all cell types. More heterogeneity was seen in non-CM cells: *Flnc* showed high

median expression in macrophages and fibroblasts, whilst *Lgals3* were observed to have high

median expression in fibroblasts.

Discussion

CVD has a complex etiology and is responsible for a range of systemic effects, hindering our understanding of its consequences on different tissues. Here, we took advantage of the technological advances in high-throughput RNA-seq and applied integrative network analyses to comprehensively explore the underlying biological effects of MI. Specifically, we generated RNA-seq data from heart, liver, skeletal muscle and adipose tissue obtained from mice 6 and 24 h after an MI or sham operation. We used transcriptomics data analyses (differential expression, functional analysis, and reporter metabolites analysis) to determine the systemic effects of the MI across multiple tissues. Moreover, we performed CN analyses to pinpoint important key and tissue-specific clusters in each tissue, and identified the key genes in each cluster. Finally, we used a whole-body modelling approach to identify the crosstalk between tissues and reveal the global metabolic alterations, before finally validating our findings with publicly available independent MI cohorts.

Based on our analyses, we observed downregulation of heart-specific functions and upregulation of lipid metabolism and inflammatory response in heart, muscle, and adipose tissue after an MI (**Figure 4B**). Liver showed a distinct response with respect to the other three tissues, including downregulation of inflammatory response. We observed that fatty acid metabolism was downregulated in heart and adipose tissue, whereas fatty acid beta-oxidation was upregulated and glutathione metabolism was downregulated in liver. We also observed upregulation of oxidative stress in heart and skeletal muscle. We also observed downregulation of mitochondrial functions in heart, muscle, and adipose tissue. Furthermore, we found upregulation of retinol metabolism in heart and downregulation of retinol metabolites in liver (**Figure 4B**).

We hypothesized that downregulation of fatty acid metabolism from adipose tissue was due to exchange of fatty acids with other tissues (liver and muscle) (**Figure 4B**). We also observed the flow of retinol from liver to heart during MI, consistent with previous reports (Palace et al., 1999). These MI-associated alterations lead to dysfunctional mitochondria and decreased energy production, especially in heart and skeletal muscle.

We further examined our results with single cell RNA-seq data. Although our gene set was extracted from bulk RNA-seq data analysis, it still had a strong presence at the single cell

RNA level, in both CM and NCM cells. We also validated our results with publicly available

MI datasets generated in separate independent studies. The validation results strengthened our

findings on the altered functions/pathways and the important heart-specific genes after an MI.

Importantly, our analyses of gene clusters highlighted multiple key genes in the response to

MI in different tissues. Specifically, we observed that *Flnc*, *Prkaca, Lgals3*, and *Pprc1*

showed important responses in heart, skeletal muscle, and adipose tissue. Moreover, *Flnc*,

Lgals3, and *Prkaca* showed strong presence at the single cell level. *Flnc* is involved in actin

cytoskeleton organization in heart and skeletal muscle, and previous studies have shown that

this gene has critical role in CVD (Hall et al., 2020; Zhou et al., 2020). Similarly, *Prkaca,* an

important metabolic gene, has also been shown to play an important function during CVD

(Bers, 2008; Diviani et al., 2011; Turnham and Scott, 2016). *Lgals3*, related to acute

inflammation response, has been studied intensively in recent years as a key gene in CVD,

and as a potential CVD therapy target (Suthahar et al., 2018; Zhong et al., 2019). Lastly,

Pprc1, as important regulator of mitochondrial biogenesis, has not been explored for its direct

relationship with CVD; however, mitochondrial biogenesis appears to be an important

response to CVD (Piantadosi and Suliman, 2012; Ren et al., 2010; Siasos et al., 2018).

In summary, we systematically unveiled the deregulation of biological processes and

pathways that resulted from MI in heart, liver, muscle, and adipose tissue by integrating

transcriptomic data and the use of biological networks. We also identified the key clusters and

central genes using generated tissue-specific CNs. In this study, we demonstrated a strategy to

utilize multi-tissue transcriptomic data to identify alteration of biological processes and

pathways to systemically explore the effect of a disease.

Limitation of the Study

We recognized several limitations to be noted on this research. First, only transcriptomic data

was analyzed in this research, hence the sensitivity might be limited especially for short

timepoint, e.g. 6 hours after MI. Second, we focused our analysis in this research only on

protein-coding genes. Third, to explore more about the shift in metabolism due to MI, longer

timepoints needs to be explored. This opens new opportunities for future research, including

analyzing the non-protein-coding gene signatures and longer timepoints.

Author Contribution

MK performed the animal experiments, MA performed the computational analysis and analyzed the clinical data together with RB, SD, HT, MU, MC, JW, DE, CZ, AM, and JB coordinated the generation of the clinical data. MA, MK, AM and JB wrote the paper and all authors were involved in editing the paper.

Acknowledgements

This work was financially supported by the Knut and Alice Wallenberg Foundation, Swedish Research Foundation and Swedish Heart-Lung Foundation.

Conflict of Interest

JW, MC, DE are employees at AstraZeneca. The other authors declare no conflict of interest.

Figure Legends

Figure 1 *(A) Overview of this study (B) Number of differentially expressed genes for each*

tissue at each time point. Effect of MI shown to be more pronounced after 24 h. (C) UpSet

plot to show intersection between differentially expressed genes (FDR < 5%) in different

tissues. The plot showed that each tissue has its specific set of genes that were affected by MI.

(D) KEGG pathway analysis (FDR < 0.05 in at least 3 tissues) for 24 hours post MI

compared to its control for each tissue. We observed that 141 (5 upregulated) and 125 (14

upregulated) pathways are significantly altered in heart 6 and 24 h after infarction,

respectively. For other tissues, we found that 24 (9 upregulated), 61 (54 upregulated) and 48

(15 upregulated) pathways are altered in liver, muscle, and adipose, respectively.

Figure 2 *(A) Functional analysis with GO (FDR < 0.05% in at least 3 tissues) revealed that*

944 (919 upregulated) and 1019 (970 upregulation) BPs are significantly altered in heart 6

and 24 h after infarction, respectively. The results also showed 38 (16 upregulated), 376 (357

upregulated) and 193 (116 upregulated) BPs are significantly altered 24 h after infarction in

liver, muscle and adipose, respectively. Most tissues show significant alterations in multiple

biological processes, including mitochondrial functions, RNA processes, cell adhesion,

ribosome and immune systems. The results of this analysis showed alterations concordant

with those observed for KEGG pathways. (B) Reporter metabolites analysis shows significant

alternation in important metabolites. Our analysis revealed that 169, 324, 118 and 51

reporter metabolites are significantly altered in heart, liver, skeletal muscle and adipose

tissues, respectively, at 24 h post-infarction (Table S4)

Figure 3 *Network analyses. (A) Heart co-expression network clusters with superimposed*

DEGs 24 h post-infarction (Blue = down-regulated, Red = up-regulated) marked with the

cluster numbers. The edges between the clusters were aggregation of the inter-cluster edges

(B) Liver. (C) Muscle. (D) Adipose. (E) Intersection of the most central clusters in all tissues

shows that the central architecture of the network was conserved in all tissues. We found 4

sub-clusters within the network intersection. Top 10 most connected genes are marked in

black. (F) Enriched GO BP in heart-specific cluster generated by Revigo.

Figure 4 *(A) Similarity of functions in the most central cluster and specific functions of each*

tissue-specific cluster. (B) Functional analysis for each tissue and hypothesized flow of

metabolites

Figure 5 (A) Significantly differentially expressed central genes of each tissue-specific cluster

to fatty acid metabolism, as one of the most affected metabolic process. (B) Lipid metabolism.

Red = upregulated, blue = downregulated.

Figure 6 (A-C) UMAP clustering of the single cell RNA-seq data with our gene set based on

the CM enrichment, cell types, and network gene cluster type (D-F) colored based on the

mean expression of the gene groups (G) DEGs intersection of our data and validation cohort

(H) & (I) Intersection of functional analysis results (GO BP and KEGG Pathways) of our

data and validation cohort

Table

Table 1 Properties of the co-expression network

Tissue	# of Genes	# of Edges	# of Clusters	Modularity Scores
Heart	8793	1570898	7	0.540179085
Liver	7760	1103589	6	0.577273459
Muscle	8834	1660603	7	0.521239124
Adipose	10790	2636378	8	0.495469439

STAR Methods

Induction of MI

10-week-old male C57Bl/6N mice were fasted for 4 h before induction of myocardial infarction. The mice were then anesthetized with isoflurane, orally intubated, and connected to a small-animal ventilator (SAR-830, Geneq, Montreal, Canada) distributing a mixture of oxygen, air and 2–3% isoflurane. ECG electrodes were placed on the extremities, and cardiac rhythm was monitored during surgery. An incision was made between the 4th and 5th ribs to reveal the upper part of the anterior left ventricle (LV) wall and the lower part of the left atrium. Myocardial infarction was induced by ligating the left anterior descending (LAD) coronary artery immediately after the bifurcation of the left coronary artery 1. The efficacy of the procedure was immediately verified by characteristic ECG changes, and akinesis of the LV anterior wall. After verification of the infarction, the lungs were hyperinflated, positive end-expiratory pressure was applied, and the chest was closed. Sham mice were handled identically, but no ligation of the LAD coronary artery was performed (and thus, no ischemia was induced in these mice). The mice received an intraperitoneal injection of 0.1 ml buprenorphine to relieve postoperative pain and were allowed to recover spontaneously after stopping isoflurane administration. Mice were killed with an overdose of isoflurane 6 h or 24 h after occlusion or sham operation. We collected the left ventricle (the whole left ventricle containing mainly infarcted tissue) of the heart, whereas white adipose tissue (WAT) was collected from the abdomen and musculus soleus was taken as the muscle tissue. Mouse hearts and biopsies from the liver, muscle and WAT were snap-frozen in liquid nitrogen and stored at -80°C until analysis. All mice studies were approved by the local animal ethics committee and conform to the guidelines from Directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes.

Echocardiography in mice

Echocardiographic examination, using VisualSonics VEVO 2100 system (VisualSonics Inc, Ontario, Canada), which includes an integrated rail system for consistent positioning of the ultrasound probe was performed 6 and 24 h after an MI to determine the size of the MI. We calculated infarct size based on wall motion score index (WMSI) 24 h after myocardial infarction by a 16-segments model on 3 short axis images, as 0 for normal, ½ for reduced wall thickening and excursion in a segment and 1 for no wall thickening and excursion in a segment. WMSI was calculated as the sum of scores divided by the total number of segments.

Hair removal gel was applied to isofluorane-anesthetized (1.2%) mice chest to minimize

resistance to ultrasonic beam transmission. The mice were then placed on a heating pad and

extremities were connected to an ECG. A 55 MHz linear transducer (MS550D) was used for

imaging. An optimal parasternal long axis (LAX) cine loop of >1000 frames/s was acquired

using the ECG-gated kilohertz visualization technique. Parasternal short axis cine-loops were

acquired at 1, 3, and 5 mm below the mitral annulus. Infarct size was calculated based on wall

motion score index 6 and 24 hours after myocardial infarction by a 16-segments model on

LAX and 3 short axis images view, as 0 for normal, ½ for reduced wall thickening and

excursion in a segment and 1 for no wall thickening and excursion in a segment. The data

were evaluated using VevoStrain™ software system (VisualSonics Inc, Ontario, Canada).

RNA extraction and sequencing

Total RNA was isolated from homogenized heart tissue using RNeasy Fibrous Tissue Mini

Kit (Qiagen). cDNA was synthesized with the high-capacity cDNA Reverse Transcription Kit

(Applied Biosystems) and random primers. mRNA expression of genes of interest was

analyzed with TaqMan real-time PCR in a ViiA™ 7 system (Applied Biosystems). RNA

sequencing library were prepared with Illumina RNA-Seq with Poly-A selections.

Subsequently, the libraries were sequenced on NovaSeq6000 (NovaSeq Control Software

1.6.0/RNA v3.4.4) with a 2x51 setup using 'NovaSeqXp' workflow in 'S1' mode flow cell.

The Bcl was converted to FastQ by bcl2fastq_v2.19.1.403 from CASAVA software suite

(Sanger/phred33/Illumina 1.8+ quality scale).

RNA-sequencing data analysis

The raw RNA-sequencing results were processed using Kallisto (Bray et al., 2016) with index

file generated from the Ensembl mouse reference genome (Release-96) (Zerbino et al., 2017).

The output from Kallisto, both estimated count and TPM (Trancript per kilobase million),

were subsequently mapped to gene using the mapping file retrieved from Ensembl BioMart

website, by filtering only protein coding genes and transcripts. Genes with mean expression

less than 1 TPM in each condition were filtered. For data exploration, we used PCA from

sklearn package (Pedregosa et al., 2011) in Python 3.7 and used TPM values as the input.

Subsequently, we performed differential gene expression analysis using DESeq2 package in

R. We utilized the capabilities from DESeq2 to normalize the rounded estimated count data

and to correct for confounding factors (such as time). To define a gene as differentially

expressed (DEGs), a gene has to fulfill a criterion of FDR < 5%. The results of differential

expression analysis were then used for functional analysis.

We checked the tissue specificity of the DEGs in each tissue with the data from Mouse Gene

Atlas (Su et al., 2004). For all the tissue-specific genes, we also checked their human-

homolog genes in the human secretome database (Uhlén et al., 2019).

Functional analysis

We performed functional analysis using the R package PIANO (Väremo et al., 2013). As the

input, we used the fold changes and p-values from the DESeq2, and also GO BP and KEGG

pathways gene-set collections from Enrichr (Chen et al., 2013; Kuleshov et al., 2016), and

metabolites from Mouse Metabolic Reaction database (Mardinoglu et al., 2015). To define a

process or pathway as significant, we used a cut off of FDR < 5% for the distinct direction of

PIANO (both up and down).

Co-expression network generation

We generated the co-expression network by generating gene-gene Spearman correlation ranks

within a tissue type, using *spearmanr* function from SciPy (Jones et al., 2001) in Python 3.7.

Using the same environment, we performed multiple hypothesis testing using Benjamini-

Hochberg method from *statsmodels* (Perktold et al., 2017). Correlation data were filtered with

criterion of adjusted p-value < 5%.

The top 5% of filtered correlation results were then loaded into iGraph module (Csardi and

Nepusz, 2006) in Python 3.7 as an unweighted network. To find the subnetworks, we

employed the Leiden clustering algorithm (Traag et al., 2019) with *ModularityVertexPartition*

method. Each cluster was analyzed by using Enrichr (Chen et al., 2013; Kuleshov et al., 2016)

to get the enriched GO BP and KEGG pathways. Criterion FDR < 0.05 were used to find the

significantly enriched terms. Clusters with less than 30 genes were discarded, to be able to get

significant functional analysis results. Since GO BP was relatively sparse, we used Revigo

(Supek et al., 2011) to summarize the GO BP into a higher level. Revigo was further

employed to build a GO BP network. Clustering coefficient was calculated based on the

average local clustering coefficient function within iGraph.

Multi-tissue metabolic modeling

We combined tissue-specific models (of heart, liver, muscle, adipose and small intestine) constructed previously (Mardinoglu et al., 2015) in a multi-tissue model by adding an additional compartment representing the plasma, which allows the exchange of metabolites among different tissues. Blocked reactions that could not carry fluxes (and the unused metabolites and genes linked to these reactions) were removed from the models. In addition, the dietary input reactions and constraints were added to the small intestine model to simulate the food intake (**Table S7**). Specifically, we assumed that the mice weighed 30 g and consumed 4.5 g chow diet per day (15 g/100 g body weight) based on a previous study (Kummitha et al., 2014). We also calculated the tissue-specific resting energy expenditures and set them as mandatory metabolic constraints based on previous studies and resting energy expenditure for other tissues was incorporated by including a mandatory glucose secretion flux out from the system with the lower bound calculated based on ATP (**Table S7**) (Kummitha et al., 2014).

To simulate the metabolic flux distribution in the sham-operated mice, we set the lipid droplet accumulation reaction in adipose tissue (m3_Adipose_LD_pool) as the objective function so that the energy additional to the resting energy expenditure will be stored as fat; we used parsimonious FBA to calculate the flux distribution. To simulate the flux distribution after an MI, we used the previously developed method Relative Metabolic Difference ver. 2 (RMetD2) (Zhang et al., 2019) to integrate transcriptomic data. In brief, RMetD2 calculates the expected fluxes of reactions based on their reference fluxes and fold changes of gene expression, and searches for a flux distribution that is close to the expected fluxes while subject to the model constraints.

Validation of the results

We validated our findings by performing similar steps of RNA sequencing and functional analysis for the publicly available mouse MI datasets GSE104187 and GSE52313 (Ounzain et al., 2014; Williams et al., 2018). Single-cell RNA-seq human data for heart failure was retrieved from GSE109816 (Wang et al., 2020). Hierarchical clustering of the single-cell data was performed with *clustermap* function from *seaborn* module in Python.

Data and code availability

All raw RNA-sequencing data generated from this study can be accessed through accession

number GSE153485. Codes used during the analysis are available on

https://github.com/sysmedicine/ArifEtAll_2020_MultiTissueMI

Supplementary Tables

Table S1 *Differential Expression Analysis Results*

Table S2 *KEGG Pathways*

Table S3 *Gene Ontology Biological Processes*

Table S4 *DEG comparison between Liver and other tissues*

Table S5 *Reporter Metabolite Analysis*

Table S6 *Enrichment Analyses of Clusters, Clusters properties*

Table S7 *Food Intake, Energy Expenditure, and Flux Balance Analysis (FBA) of Whole-Body*
Modeling

Table S8 *Validation Result (Differential Expression and Functional Analysis)*

Table S9 *Detailed Information of 16 Key Genes that are DEGs in at least 3 tissues,*
Neighbors and Functional Analysis Results of The Neighbors of 4 key genes, and mean
expression of 4 key genes in each cell-type in the scRNA data.

References

Alexander, M., Loomis, A.K., van der Lei, J., Duarte-Salles, T., Prieto-Alhambra, D., Ansell,
D., Pasqua, A., Lapi, F., Rijnbeek, P., and Mosseveld, M. (2019). Non-alcoholic fatty liver
disease and risk of incident acute myocardial infarction and stroke: findings from matched
cohort study of 18 million European adults. bmj *367*, l5367.

Bakhtiarizadeh, M.R., Hosseinpour, B., Shahhoseini, M., Korte, A., and Gifani, P. (2018).
Weighted Gene Co-expression Network Analysis of Endometriosis and Identification of
Functional Modules Associated With Its Main Hallmarks. Frontiers in Genetics *9*.

Benfeitas, R., Bidkhori, G., Mukhopadhyay, B., Klevstig, M., Arif, M., Zhang, C., Lee, S.,
Cinar, R., Nielsen, J., and Uhlen, M. (2019). Characterization of heterogeneous redox
responses in hepatocellular carcinoma patients using network analysis. EBioMedicine *40*,
471-487.

Bers, D.M. (2008). Calcium cycling and signaling in cardiac myocytes. Annu Rev Physiol *70*,
23-49.

Bidkhori, G., Benfeitas, R., Klevstig, M., Zhang, C., Nielsen, J., Uhlen, M., Boren, J., and
Mardinoglu, A. (2018). Metabolic network-based stratification of hepatocellular carcinoma
reveals three distinct tumor subtypes. Proceedings of the National Academy of Sciences *115*,
E11874-E11883.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nature biotechnology *34*, 525.

Camara, A.K., Zhou, Y., Wen, P.-C., Tajkhorshid, E., and Kwok, W.-M. (2017).
Mitochondrial VDAC1: a key gatekeeper as potential therapeutic target. Frontiers in
physiology *8*, 460.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and
Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. BMC Bioinformatics *14*, 128.

Csardi, G., and Nepusz, T. (2006). The igraph software package for complex network
research. InterJournal, Complex Systems *1695*, 1-9.

Das, S., Frisk, C., Eriksson, M.J., Walentinsson, A., Corbascio, M., Hage, C., Kumar, C.,
Asp, M., Lundeberg, J., and Maret, E. (2019). Transcriptomics of cardiac biopsies reveals
differences in patients with or without diagnostic parameters for heart failure with preserved
ejection fraction. Scientific reports *9*, 3179.

Diviani, D., Dodge-Kafka, K.L., Li, J., and Kapiloff, M.S. (2011). A-kinase anchoring
proteins: scaffolding proteins in the heart. American Journal of Physiology-Heart and
Circulatory Physiology *301*, H1742-H1753.

Hall, C.L., Gurha, P., Sabater-Molina, M., Asimaki, A., Futema, M., Lovering, R.C., Suárez,
M.P., Aguilera, B., Molina, P., and Zorio, E. (2020). RNA sequencing-based transcriptome
profiling of cardiac tissue implicates novel putative disease mechanisms in FLNC-associated
arrhythmogenic cardiomyopathy. International journal of cardiology *302*, 124-130.

Jones, E., Oliphant, T., and Peterson, P. (2001). SciPy: Open source scientific tools for
Python.

Kiyuna, L.A., e Albuquerque, R.P., Chen, C.-H., Mochly-Rosen, D., and Ferreira, J.C.B.
(2018). Targeting mitochondrial dysfunction and oxidative stress in heart failure: challenges
and opportunities. Free Radical Biology and Medicine *129*, 155-168.

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev,
S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A.*, et al.* (2016). Enrichr: a comprehensive gene
set enrichment analysis web server 2016 update. Nucleic Acids Research *44*, W90-W97.

Kummitha, C.M., Kalhan, S.C., Saidel, G.M., and Lai, N. (2014). Relating tissue/organ
energy expenditure to metabolic fluxes in mouse and human: experimental data integrated
with mathematical modeling. Physiological reports *2*.

Lee, S., Zhang, C., Kilicarslan, M., Piening, B.D., Bjornson, E., Hallström, B.M., Groen,
A.K., Ferrannini, E., Laakso, M., and Snyder, M. (2016). Integrated network analysis reveals
an association between plasma mannose levels and insulin resistance. Cell metabolism *24*,
172-184.

Lee, S., Zhang, C., Liu, Z., Klevstig, M., Mukhopadhyay, B., Bergentall, M., Cinar, R.,
Ståhlman, M., Sikanic, N., and Park, J.K. (2017). Network analyses identify liver☐specific
targets for treating liver diseases. Molecular systems biology *13*.

Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R., and Pfister, H. (2014). UpSet:
visualization of intersecting sets. IEEE transactions on visualization and computer graphics
20, 1983-1992.

Lima, I., Peres, W., Cruz, S., and Ramalho, A. (2018). Association of Ischemic
Cardiovascular Disease with Inadequacy of Liver Store of Retinol in Elderly Individuals.
Oxidative medicine and cellular longevity *2018*.

Mardinoglu, A., Bjornson, E., Zhang, C., Klevstig, M., Söderlund, S., Ståhlman, M., Adiels,
M., Hakkarainen, A., Lundbom, N., and Kilicarslan, M. (2017a). Personal model☐assisted
identification of NAD+ and glutathione metabolism as intervention target in NAFLD.
Molecular systems biology *13*.

Mardinoglu, A., Boren, J., Smith, U., Uhlen, M., and Nielsen, J. (2017b). The employment of
systems biology in gastroenterology and hepatology. Nat Rev Gastroenterol Hepatol.

Mardinoglu, A., and Nielsen, J. (2015). New paradigms for metabolic modeling of human
cells. Current Opinion in Biotechnology *34*, 91-97.

Mardinoglu, A., Shoaie, S., Bergentall, M., Ghaffari, P., Zhang, C., Larsson, E., Bäckhed, F.,
and Nielsen, J. (2015). The gut microbiota modulates host amino acid and glutathione
metabolism in mice. Molecular Systems Biology *11*, 834.

McInnes, L., Healy, J., and Melville, J. (2018). Umap: Uniform manifold approximation and
projection for dimension reduction. arXiv preprint arXiv:180203426.

Miyamoto, T., Katayama, Y., and Harano, Y. (1999). Blood ketone body as a stress marker in
acute myocardial infarction. Nihon Kyukyu Igakukai Zasshi *10*, 621-622.

Mukund, K., and Subramaniam, S. (2017). Co-expression network approach reveals
functional similarities among diseases affecting human skeletal muscle. Frontiers in
physiology *8*, 980.

Nielsen, J. (2017). Systems biology of metabolism: a driver for developing personalized and
precision medicine. Cell metabolism *25*, 572-579.

Ounzain, S., Micheletti, R., Beckmann, T., Schroen, B., Alexanian, M., Pezzuto, I., Crippa,
S., Nemir, M., Sarre, A., and Johnson, R. (2014). Genome-wide profiling of the cardiac
transcriptome after myocardial infarction identifies novel heart-specific long non-coding
RNAs. European heart journal *36*, 353-368.

Palace, V.P., Hill, M.F., Khaper, N., and Singal, P.K. (1999). Metabolism of vitamin A in the
heart increases after a myocardial infarction. Free Radical Biology and Medicine *26*, 1501-
1507.

Palaniyandi, S.S., Qi, X., Yogalingam, G., Ferreira, J.C.B., and Mochly-Rosen, D. (2010).
Regulation of mitochondrial processes: a target for heart failure. Drug Discovery Today:
Disease Mechanisms *7*, e95-e102.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., and Dubourg, V. (2011). Scikit-learn: Machine learning in
Python. Journal of machine learning research *12*, 2825-2830.

Pedrotty, D.M., Morley, M.P., and Cappola, T.P. (2012). Transcriptomic biomarkers of
cardiovascular disease. Progress in cardiovascular diseases *55*, 64-69.

Perktold, J., Seabold, S., and Taylor, J. (2017). Statsmodels: Statistics in python (Oct).

Piantadosi, C.A., and Suliman, H.B. (2012). Transcriptional control of mitochondrial
biogenesis and its interface with inflammatory processes. Biochimica et Biophysica Acta
(BBA)-General Subjects *1820*, 532-541.

Priest, C., and Tontonoz, P. (2019). Inter-organ cross-talk in metabolic syndrome. Nature
Metabolism.

Ren, J., Pulakat, L., Whaley-Connell, A., and Sowers, J.R. (2010). Mitochondrial biogenesis
in the metabolic syndrome and cardiovascular disease. Journal of molecular medicine *88*,
993-1001.

Schwertz, H., Carter, J.M., Abdudureheman, M., Russ, M., Buerke, U., Schlitt, A., Müller-
Werdan, U., Prondzinsky, R., Werdan, K., and Buerke, M. (2007). Myocardial
ischemia/reperfusion causes VDAC phosphorylation which is reduced by cardioprotection
with a p38 MAP kinase inhibitor. PROTEOMICS *7*, 4579-4588.

Siasos, G., Tsigkou, V., Kosmopoulos, M., Theodosiadis, D., Simantiris, S., Tagkou, N.M.,
Tsimpiktsioglou, A., Stampouloglou, P.K., Oikonomou, E., and Mourouzis, K. (2018).
Mitochondria and cardiovascular diseases—from pathophysiology to treatment. Annals of
translational medicine *6*.

Su, A.I., Wiltshire, T., Batalov, S., Lapp, H., Ching, K.A., Block, D., Zhang, J., Soden, R.,
Hayakawa, M., Kreiman, G.*, et al.* (2004). A gene atlas of the mouse and human protein-
encoding transcriptomes. Proceedings of the National Academy of Sciences of the United
States of America *101*, 6062-6067.

Sun, W., Quan, N., Wang, L., Yang, H., Chu, D., Liu, Q., Zhao, X., Leng, J., and Li, J.
(2016). Cardiac-specific deletion of the Pdha1 gene sensitizes heart to toxicological actions of
ischemic stress. Toxicological Sciences *151*, 193-203.

Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). REVIGO summarizes and
visualizes long lists of gene ontology terms. PloS one *6*, e21800.

Suthahar, N., Meijers, W.C., Silljé, H.H., Ho, J.E., Liu, F.-T., and de Boer, R.A. (2018).
Galectin-3 activation and inhibition in heart failure and cardiovascular disease: an update.
Theranostics *8*, 593.

Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing
well-connected communities. Scientific reports *9*.

Tran, D.H., and Wang, Z.V. (2019). Glucose Metabolism in Cardiac Hypertrophy and Heart
Failure. Journal of the American Heart Association *8*, e012673.

Turanli, B., Zhang, C., Kim, W., Benfeitas, R., Uhlen, M., Arga, K.Y., and Mardinoglu, A.
(2019). Discovery of therapeutic agents for prostate cancer using genome-scale metabolic
modeling and drug repositioning. EBioMedicine *42*, 386-396.

Turnham, R.E., and Scott, J.D. (2016). Protein kinase A catalytic subunit isoform PRKACA;
History, function and physiology. Gene *577*, 101-108.

Uhlén, M., Karlsson, M.J., Hober, A., Svensson, A.-S., Scheffel, J., Kotol, D., Zhong, W.,
Tebani, A., Strandberg, L., Edfors, F.*, et al.* (2019). The human secretome. Science Signaling
12, eaaz0274.

Väremo, L., Nielsen, J., and Nookaew, I. (2013). Enriching the gene set analysis of genome-
wide data by incorporating directionality of gene expression and combining statistical
hypotheses and methods. Nucleic Acids Research *41*, 4378-4391.

Wang, L., Yu, P., Zhou, B., Song, J., Li, Z., Zhang, M., Guo, G., Wang, Y., Chen, X., and
Han, L. (2020). Single-cell reconstruction of the adult human heart during heart failure and
recovery reveals the cellular landscape underlying cardiac function. Nature Cell Biology *22*,
108-119.

WHO (2019). Cardiovascular diseases (CVDs) Fact sheets.

Williams, A.L., Khadka, V., Tang, M., Avelar, A., Schunke, K.J., Menor, M., and Shohet,
R.V. (2018). HIF1 mediates a switch in pyruvate kinase isoforms after myocardial infarction.
Physiological genomics *50*, 479-494.

Yada, T., Shimokawa, H., Hiramatsu, O., Haruna, Y., Morita, Y., Kashihara, N., Shinozaki,
Y., Mori, H., Goto, M., and Ogasawara, Y. (2006). Cardioprotective role of endogenous
hydrogen peroxide during ischemia-reperfusion injury in canine coronary microcirculation in
vivo. American Journal of Physiology-Heart and Circulatory Physiology *291*, H1138-H1146.

Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M R., Barrell, D., Bhai, J., Billis, K.,
Cummins, C., Gall, A., Girón, C.G.*, et al.* (2017). Ensembl 2018. Nucleic Acids Research *46*,
D754-D761.

Zhang, C., Lee, S., Bidkhori, G., Benfeitas, R., Lovric, A., Chen, S., Uhlen, M., Nielsen, J.,
and Mardinoglu, A. (2019). RMetD2: a tool for integration of relative transcriptomics data
into Genome-scale metabolic models. BioRxiv, 663096.

Zhong, X., Qian, X., Chen, G., and Song, X. (2019). The role of galectin 3 in heart failure
and cardiovascular disease. Clinical and Experimental Pharmacology and Physiology *46*, 197-
203.

Zhou, Y., Chen, Z.e., Zhang, L., Zhu, M., Tan, C., Zhou, X., Evans, S.M., Fang, X., Feng, W.,
and Chen, J. (2020). Loss of Filamin C Is Catastrophic for Heart Function. Circulation *141*,
869-871.

A Heart
3 2 1 4 5 6 0

B Liver
0 5 2 4 1 3

C Muscle
4 1 6 5 2 0 3

D Adipose
1 4 6 2 5 3 7 0

E

Reg. of transcription from RNAP-II
mRNA Processing

Foxj3, Tln1, Specc1, Mark3, Pum1, Atp6-v0e2, Ccdc82, Atp2a3, Abca7, Ywhah

Regulation of Mitotic Cell Cycle Phase
Ubiquitin-dependent Protein Catabolic Process

Rnps1, Lyrm1, Efcab14, Armc8, Aoc3, Smad4, 4933434-E20Rik, Fbln5, Drosha, Cct4

Amyloid Fibril formation
Histone H3-K36 Demethylation

Zfp329, Tbc1d1, Mtrex, Gpatch2, Brd1, Fzd4, Cdc42e, Rnf11, Ccdc66, Smarcal1

Nucleic Acid-templated Transcription
Vesicle-mediated Transport

Btbd2, Ptdss2, Git2, Akt2, Kansl3, Dmwd, Mark2, Fut8, Fam-219b, Trps1

F

muscle contraction
heart contraction
ATP metabolism
reg. calcium in sarcoplasmic reticulum
regulation ATPase activity
myofibril assembly
cardiac muscle cell membrane repolarization
mitochondrial ATP synthesis
ubiquinone biosynthesis
regulation of acetyl-CoA biosynthesis
cardiac muscle cell actionpotential
inner mitochondrial membrane org.
respiratory electron transport chain
actomyosin structure
mitochondrial respiratory chain
mitochondrion organization
NADH dehydrogenase complex assembly
mitochondrial respiratory chain

Figure S1 (A) PCA plots of each tissue showing data from mice 6 and 24 h after an MI or sham operation. The plot showed that heart was affected the most by the change in conditions and the rest were most affected by time shifts. (B) KEGG pathways related to cardiac problems show activation after an MI.

Figure S2 KEGG pathway analysis results for each tissue (A) Heart 6- and 24- hours post MI (B) Liver, Muscle, and Adipose tissue 24 hours post MI.

Figure S3 (A) cGMP-PKG & (B) HIF-1 signaling pathway with overlay data from differential expression and reporter metabolites analysis.

Figure S4 Expression level of central genes from most central and tissue-specific clusters and their hierarchical clustering on the single cell level.activation after an MI.

Figure S5 scRNA Expression of Flnc, Lgals3, and Prkaca.