10-28-2020

DNAJB12 and Hsp70 Facilitate the Conformation Specific Degradation of Arrested N1303K-CFTR Intermediates by ERQC-Autophagy

Lihua He, Andrew S. Kennedy, Scott Houck, Andrei Aleksandrov, Nancy L. Quinney, Deborah M. Cholon, Martina Gentzsch, Scott H. Randell, Hong Yu Ren, Douglas M. Cyr*

Department of Cell Biology and Physiology and the Cystic Fibrosis/Pulmonary Research and Treatment

Center, the University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

Running Title: Triage of CFTR intermediates between ERAD and ERQC-Autophagy

*Address correspondence to Douglas M. Cyr, Ph.D., Department of Cell Biology and Physiology Biology,

Tel. 919-843-4805; E-mail: DMCYR@med.unc.edu

Keywords: Cystic Fibrosis, autophagy, Hsp70, Hsp40, DNAJB12, endoplasmic reticulum, lysosome,

proteasome, triage, folding modulator

SUMMARY

Proteotoxic stress, intrinsic folding inefficiencies, and disease-related mutations challenge cellular quality control systems tasked with proteome biogenesis and suppressing proteotoxicity. The ER-transmembrane Hsp40 DNAJB12 and Hsp70 cooperate in protein maintenance by triaging nascent membrane proteins for folding versus degradation. N1303K is the second most common CF causing mutation in CFTR, but unlike F508del-CFTR, its biogenic and functional defects are resistant to correction by VX-809. VX-809 stimulates the accumulation of kinetically trapped N1303K-CFTR intermediates that exhibit conformational features of folded CFTR but remain associated with DNAJB12-Hsp70. Stable N1303K-CFTR intermediates are resistant to ERAD and become concentrated in ER-tubules associated with autophagy initiation sites containing WIPI1, FIP200, and LC3. Budding of autophagosomes containing N1303K-CFTR from omegasomes occurs where autolysosomes dock to WIPI1 rings. Mechanisms will be discussed for Hsp40 and Hsp70 action in the sorting of N1303K-CFTR intermediates for secretion versus degradation by ERAD or ERQC-autophagy.

INTRODUCTION

Protein triage mechanisms sort misfolded proteins between pathways for folding, degradation, and sequestration to suppress the accumulation of toxic species that disrupt cellular homeostasis and kill cells. Missense mutations that lead to misfolding of membrane proteins, such as ion channels and G protein-coupled receptors, cause diseases including cystic fibrosis (CF), retinitis pigmentosa, hypercholesterolemia, and hypogonadism (Houck and Cyr 2012). CF has been linked to over 1700 different mutations in the CFTR gene with global misfolding, premature degradation of F508del-CFTR being the most cause (Veit, Avramescu, et al. 2016). Folding modulators help overcome defects in F508del-NBD1 folding and assembly and, when used in combination with ion channel potentiators, restore CFTR function to therapeutic levels. However, it is not clear how changes in the conformation of a membrane protein elicited by a folding modulator would impact its recognition by molecular chaperones and sorting within the ER-membrane system for secretion versus retention and degradation.

A challenge to addressing questions about membrane protein triage mechanisms is that ensembles of intermediates accumulate at different stages of folding and post-translation assembly that are difficult to resolve. The use of folding modulators as tools provides an approach to increase pools of on-pathway intermediates and identify rate-limiting steps in the folding and escape from the endoplasmic reticulum (ER) protein quality control of WT and mutant CFTR. CFTR is a 1480 amino acid anion channel containing two membrane-spanning domains (MSD1 and MSD2), two nucleotide-binding domains (NBD1 and NBD2), and regulatory R-domain. Assembly of CFTR involves domain folding and interaction of its NBDs with themselves and cytosolic loops (CL) located between different pairs of transmembrane helices (TM){Cyr, 2005 #2294}. The F508del mutation is located in NBD1 so F508del-NBD1 is unstable and fails to correctly assemble with NBD2 and CLs transmembrane spans. The modulator VX-809 partially restores F508del-CFTR folding by stabilizing MSD1 and F508del-NBD1,

making downstream assembly reactions involving contact formation with CLs more efficient. However, VX-809 only restores around 5% of CFTR function in F508del-CFTR patients, and a second modulator is needed to restore lung function to a therapeutic 10% of normal {Keating, 2018 #8227} {Donaldson, 2018 #9217}. Enhancement of modulator therapy for CF will require identifying the rate-limiting steps in CFTR folding that limit the efficacy of modulators and mechanisms for triage of membrane proteins by ER quality control.

Assembly of CFTR is complicated because it exposes surfaces in the cytosol, ER-membrane, and lumen, and is facilitated by the sequential action of cytosolic DNAJA1/Hsp70, ER-lumenal calnexin, and Hsp90. Defects in CFTR assembly are also detected by Hsp70 complexed with the ER-associated DNAJB12, instead of DNAJA1, and protein quality control (PQC) E3 ubiquitin-ligase complexes (Grove, Fan et al. 2011). The majority of CFTR mutations cause global misfolding of CFTR due to domain destabilization, with intermediates being rapidly degraded by the ER-associated degradation pathway (ERAD). However, there are mutant forms of CFTR that have more subtle defects causing non-functional ion channels to fold and escape the ERQC and accumulate at the cell surface. Missense mutations in C-terminal NBD2 that arrest late steps in CFTR assembly and are resistant to F508del-CFTR modulators are also encountered (Veit, Avramescu et al. 2016). How the cell triages non-functional or partially assembled proteins that accumulate as biochemically stable intermediates is not clear. This is an essential question because such intermediates are prone to adopt toxic folds that damage membranes and kill cells.

The N1303K mutation in NBD2 is the 2nd most prevalent (2%) CFTR mutation (Cutting 2017), and the N1303K-CFTR function is not restored by modulators that restore the F508del-CFTR function. Still, little else is known about mechanisms for modulator-resistant misfolding and triage of N1303K. Herein, we report on the use of VX-809 as a tool to identify biogenic defects that must be overcome to restore the N1303K-CFTR function. These studies lead to the identification of a requirement for DNAJB12 (JB12) and

Hsp70 in the clearance of stable N1303K-CFTR intermediates from the ER via an ERQC-autophagy

pathway. ERQC-autophagy is distinct from ER-phagy because ER-tubules containing N1303K-CFTR

become clients of autophagy initiation machinery painted on adjacent tubules that convert N1303K-

tubes into phagophores containing LC3. Autophagosomes having N1303K-CFTR and LC3 then bud and

appear to fuse with lysosomes docked to the WIPI1 rings on which there were formed.   These results

provide new insights into mechanisms by which Hsp70 and DNAJB12 interact with ERAD-resistant

membrane proteins and autophagy initiation machinery during ERQC-autophagy.

RESULTS

Impact of Folding Modulators and Cl-channel Potentiators on N1303K CFTR Function

    To initiate the evaluation of the functional and folding defects caused by N1303K, we measured the

response of endogenous N1303K-CFTR expressed in native airway cells obtained from CF patients to

folding modulators (Figure 1). The effect on N1303K CFTR function of the Class I modulator VX-809 and

the Class II modulators C3 and C4 (Rosser, Grove et al. 2008, Grove, Rosser et al. 2009), which act in

synergy to restore F508del CFTR function (Ren, Grove et al. 2013), was evaluated in the absence and

presence of the potentiator VX-770. Primary airway cells with the heterozygous N1303K/W1282X

genotype were grown on semipermeable-supports at an air-liquid interface to permit the formation of a

sealed monolayer and then evaluated in USSING chambers (Fulcher, Gabriel et al. 2005). In the absence

or presence of indicated modulators, forskolin-stimulated trans-epithelial short-circuit currents for CFTR

were evaluated (Gentzsch, Ren et al. 2016, Singh, Fan et al. 2020).   W1282X CFTR mRNA is subject to

non-sense mediated decay. W1282X CFTR does not accumulate in native airway cells (Oren, Pranke et

al. 2017), so the N1303K CFTR channel function is evaluated in the absence of W1282X CFTR in primary

N1303K/W1282X cells. N1303K CFTR chloride channel activity in polarized cell monolayers was not

detected above the background (Figure 1A, top panel, black trace) and incubation of the cells with VX-

809 for 48 hrs. did not significantly increase FSK responses (Figure 1A and B, red trace, and bar graph),

nor did a mixture of modulators, including C3 and C4 (Figure 1A, green trace, and blue trace,

respectively). The acute addition of VX-770 to USSING chambers caused no significant boost in

measured currents. However, modulators appear to impact the conformation of N1303K-CFTR in native

airway cells because they increase its steady-state levels (Figure S1).

Low-temperature cell culture permits some CFTR mutants to fold, escape the ER, and traffic to the

plasma membrane (Younger, Ren et al. 2004), so N1303K-CFTR was stably expressed in BHK cells that

were grown at 27 °C. instead of 37 °C.   Single-channel measurements identified N1303K-CFTR channels

in detergent-soluble extracts made from the plasma membranes of low-temperature BHK cells (Figure

1C). N1303K-CFTR channels exhibited the same conductance but an open probability (P_o) of less than

half of WT CFTR (Figure 1C, top and middle panels). When the activity of reconstituted N1303K-CFTR

channels was measured in single-channel recordings at temperatures from 25 to 37 °C, the open

probability (P_o) of WT CFTR increased to 0.72. It remained stable during the time course of

measurements (Figure 1D, top trace). However, the P_o for N1303K CFTR decreased (Figure 1D, middle

trace), and channel activity was less stable than WT-CFTR at 37 °C. A summary of the P_o and

conductance measured at different temperatures is presented in Figure 1E. Notably, the P_o of N1303K

CFTR was stabilized by VX-770 at 25, 30, and 37 °C. These results suggest that N1303K-CFTR can

assemble into a CFTR-like ion channel when thermal energy is low. These data indicate that N1303K-

CFTR intermediates fate can be altered to suppress recognition of it by ERQC factors and restore ion

channel function.

N1303K-CFTR intermediates accumulate in a detergent-soluble and biochemically stable state

To determine the challenge presented to the ERQC-machinery by late-stage intermediates of membrane proteins that fail to complete folding, we characterized the behavior of N1303K-CFTR in ER membranes of cells cultured at 37 ^{0}C (Meacham, Lu et al. 1999, Meacham, Patterson et al. 2001, Younger, Chen et al. 2006, Ren, Grove et al. 2013, Gentzsch, Ren et al. 2016). N1303K-CFTR intermediates accumulated in an incompletely glycosylated state in the ER that is termed band B, that has faster electrophoretic mobility than the folded form, represented by band C. Band C is glycosylated further during passage through the Golgi on route to the plasma membrane (Figure 2) (Meacham, Lu et al. 1999, Meacham, Patterson et al. 2001, Dalal, Rosser et al. 2004, Cyr 2005). Levels of the B-form for N1303K were several-fold higher than levels of the B-form of F508del-CFTR (Figure 2A and Figure S2). These differences were accounted for by the observed longer half-life of N1303K-CFTR (3-hr) versus F508del-CFTR (1-hr) in cycloheximide (CHX) chase experiments (Figure 2A). A short 2-hr pre-treatment with VX-809 (3 μM) further increased the stability of N1303K-CFTR such that at the end of 3-hr CHX-chase, around 80% of B-band remained. VX-809 treatment increased the accumulation of the folded C-form of ΔF508-CFTR but only increased the half-life of its B-form to 1.5 hr. N1303K-CFTR accumulates in a conformation that is stabilized by VX-809. Yet, at physiological temperatures, this Class I modulator does not promote the assembly of N1303K-CFTR into channels that escape the ER at levels detected by western blot in the Golgi-modified C-form.

To study the intracellular behavior of N1303K-CFTR via microscopy, GFP-N1303K-CFTR, and GFP-F508del-CFTR were employed as tools in imaging studies (Figure 2-7). In controls, untagged and GFP-tagged CFTR behaved similarly (Figure 3A versus 2, 3, 4-7)), so triage of GFP-N1303K-CFTR and GFP-F508del reflects the behavior of untagged CFTR. In the characterization of CFTR intermediates the following was observed (Figure 2B-C): 1, F508del-CFTR, and N1303K-CFTR accumulate in apparently healthy tubular regions of ER in the absence of visible aggregation. 2. In live cells, GFP-F508del-CFTR and

GFP-N1303K-CFTR accumulate within dynamic ER tubules, and both diffuse at similar rates into photobleached regions. 3. The ER-localized B-form of F508del-CFTR and N1303K-CFTR do not form detergent-insoluble aggregates that partition into the pellets of centrifuged cell extracts prepared with the detergent Triton-X-100 (Figure 2C). 4. The modulator VX-809, while increasing the half-life of N1303K CFTR, did not lead it to accumulate in the detergent-insoluble fraction of cells. Also, inhibition of the proteasome (bortezomib (Bort)) or lysosomal proteases (chloroquine (CQ)) for 6-hours, which enhances the accumulation of the B-form, does not lead N1303K-CFTR to enter into detergent-insoluble aggregates. 5. N1303K-CFTR accumulates in ER tubules containing the molecular chaperone calnexin but is excluded from ER exit sites that are marked by TANGO1 and the vesicle coat protein Sec31 (Ge, Zhang et al. 2017) (Figure 2D). Overall, N1303K-CFTR is reported to accumulate in the ER in a conformation that can be stabilized by VX-809, but N1303K intermediates cannot fold, and such kinetically trapped intermediates persist within ER membranes.

N1303K-CFTR Segregates into ER Sub-domains with the Hsp40 DNAJB12 or the Autophagy Factor LC3B

We explored why N1303K-CFTR has a 3-hr half-life versus 1- hr. for F508del-CFTR by examining the respective intermediates' sensitivity to modulation of the ERAD factors JB12 and RMA1/RNF5 (Figure 3A). Accumulation of N1303K and F508del was sensitive to the elevation of JB12 or RMA1, and addition of VX-809 did not protect significant portions of intermediates from JB12 or RMA1 action. In proximity studies, APEX-JB12, which has an ER luminal APEX, labeled N1303K-CFTR in the absence and presence of VX-809 (Figure 3B). Also, the siRNA depletion of JB12 was accompanied by an increase in the steady-state levels of N1303K-CFTR. In contrast, a control depletion of an unrelated transmembrane VMP1, which regulates SERCA2B function, did not impact steady-levels of N1303K-CFTR (Figure 3C). N1303K-CFTR is similar to WT-CFTR and F508del-CFTR in that it is a client of JB12 (Grove, Fan et al. 2011). Even though VX-809 stabilizes N1303K-CFTR, VX-809 treatments do not reduce proximity labeling of N1303K-

CFTR by JB12-APEX or decrease the sensitivity of N1303K-CFTR to modulation of JB12 or RMA1 activity. N1303K-CFTR intermediates appear to become kinetically trapped, and even though VX-809 increases the accumulation of trapped intermediates, they remain clients of JB12-containing complexes.

Misfolded intermediates of membrane proteins that have long-half lives are resistant to ERAD. Those that do not form large detergent-insoluble aggregates are cleared from the ER via a selective autophagy mechanism termed ERQC-autophagy (Houck, Ren et al. 2014). To evaluate the triage of N1303K-CFTR, we explored via fluorescence microscopy the localization of its intermediates in the ER membrane system with JB12 and the autophagy initiation factor LC3B (Figure 3D). Remarkably, GFP-N1303K-CFTR was found in two different subsets of interconnected ER-tubules containing either overexpressed WT-JB12 or endogenous LC3B (Figure 3D). The inset shows JB12 (red) localized in tubules with GFP-N1303K-CFTR (green) that form a perimeter around an adjacent region of the ER enriched in GFP-N1303K-CFTR and LC3B. LC3 is conjugated to phosphatidylethanolamine within the membranes of ER-connected phagophores at initial phases of autophagy (Yla-Anttila, Vihinen et al. 2009, Matsunaga, Morita et al. 2010). ER-membranes containing N1303K-CFTR may segregate within the ER into microdomains known as omegasomes, where ER- tubules are modified and converted into phagophores (Yla-Anttila, Vihinen et al. 2009, Matsunaga, Morita et al. 2010). Also, we detected co-localized signals for GFP-N1303K-CFTR and LC3B in non-ER attached foci that resemble lysosomes.

QPD-JB12, containing a mutation in the HPD motif in the J-domain, binds clients but cannot release them because QPD-JB12 can no longer interact with Hsp70 (Grove, Fan et al. 2011, Sopha, Ren et al. 2017). QPD-JB12 co-localized with GFP-N1303K CFTR in the ER and hindered the segregation of N1303K-CFTR into ER-membranes decorated with LC3B. Hsp70 is not required for basal autophagy, and QPD-JB12 expression does not interfere with the formation of autophagic foci containing LC3B (Figure 3D). QPD-JB12's inability to interact with Hsp70 and cycle on and off N1303K-CFTR, therefore, appears to selectively inhibit the segregation of N1303K-CFTR into ER-membranes decorated with LC3B.

Endogenous JB12 was also detected via microscopy in ER microdomains with GFP-N1303K-CFTR and endogenous LC3B (Figure 3E). ER-tubules containing endogenous JB12 and GFP-N1303K-CFTR intersect with tubules decorated with endogenous LC3B that extend from out from ER-rings. Depletion of JB12 by siRNA was accompanied by an expansion of N1303k-CFTR containing ER-membranes and exclusion of N1303K-CFTR from LC3B-decorated tubular extensions of the ER.

In summary, kinetically-trapped intermediates of N1303K-CFTR are present in complexes containing JB12. The depletion of JB12 causes the expansion of ER-domains containing N1303K-CFTR and hinders N1303K-CFTR segregation into LC3B-decorated ER-tubules. JB12 depletion/inactivation does not reduce the formation of autophagic foci containing LC3B, so it appears to be required for the selective autophagic degradation of N1303K-CFTR.

N1303K-CFTR intermediates, but not F508del-CFTR, traffic from the ER to Autolysosomes

How do the N1303K-CFTR that co-localize with ER-tubular extensions and decorated with LC3B transit to autolysosomes (Figure 4)? To address this question, we first, we repeated experiments conducted with GFP-N1303K-CFTR with untagged CFTR and demonstrated that N1303K-CFTR, but not F508del-CFTR, accumulates in autophagic foci whose central core contained LC3B (Figure 4A). Three-color imaging showed that GFP-N1303K-CFTR and LC3B positive puncta also have LAMP1/2, so they are, in fact, autolysosomes (Figure 4B).

The autophagy initiation kinase (VPS34) inhibitor PIK-III blocked the entry of GFP-N1303K-CFTR into autolysosomes containing LAMP1 (Figure 4B) and did so under conditions where it abolished conversion of LC3B-I to LC3B-II (Figure S3B). LC3B II being the form that is covalently attached to the phoshatidylethanolamine headgroup of ER-lipids during initiation of phagophore formation. Likewise, siRNA of Beclin-1, treatment of cells with a ULK1 inhibitor, and overexpression of a dominant-negative form of ULK1, also blocked the transit of GFP-N1303K-CFTR from the ER to autolysosomes (Table S1).

Importantly, siRNA depletion of JB12 caused GFP-N1303K-CFTR to accumulate within biosynthetically active regions of the ER containing protein translocon associated protein Sec62 and reduced co-localization LAMP1 (Figure 4C). Sec62 and JB12 each co-localize in the ER with N1303K-CFTR, but Sec62 and JB12 are not detected with GFP-N1303K-CFTR in membranes decorated with LC3B or in autolysosomes with LAMP1. N1303K-CFTR intermediates that interact with JB12 appear separate from JB12 in a process requiring Hsp70 before entry into ER membranes containing LC3B and subsequent passage out of the ER to autolysosomes.

ER Membranes Enriched in N1303K-CFTR Protrude from WIPI1 Rings Containing Docked Autolysosomes

The route taken by N1303K-CFTR from the tubular-ER to the autolysosome was explored by visualizing its association with the omegasome factors DFCP1 and WIPI1, and LAMP1. The VPS34 kinase complex produces pools of inositol-3-phosphate head groups on membrane lipids that are bound by cytosolic DFCP1 and WIPI1, which facilitate downstream reactions in phagophore formation involving LC3 conjugation and autophagosome biogenesis. GFP-N1303K-CFTR co-localizes in tubular ER-membranes with endogenous pools of the Sec62 and mCherry-DFCP1 (Figure 5A). Similarly, GFP-N1303K-CFTR is detected with mCherry-WIPI1 in domains containing the ULK1 subunit FIP200 (Figure 5B). Interestingly, GFP-N1303K-CFTR containing ER-tubules extend out through and out from the surface of WIPI1 and FIP200 decorated ER membranes in apparent phagophores. Before exiting from the ER, it appears that N1303K-CFTR is concentrated in ER-tubules that associate with ER membranes that are decorated with FIP200, which controls ULK1 localization and thereby defines sites for focal activation of selective autophagy (Vargas, Wang et al. 2019).

Use of STED microscopy to image GFP-N1303K-CFTR, WIPI1-mCherry, and BFP-LAMP1 in live cells enabled visualization of GFP-N1303K containing tubules that emerge from WIPI1 rings to which

BFP- LAMP1 associates (Figure 4C). GFP-N1303K-CFTR is detected in ER membranes that contact WIPI1

and autolysosomes and also present are small foci of around 50 nm that contain GFP-N1303K-CFTR and

LC3B that are detached from the ER. These foci appear to be autophagosomes that have budded from

the WIPI1-ring surface at sites where autolysosomes dock.

Data presented are interpreted to suggest that N1303K-CFTR is degraded by a selective

autophagic process that requires JB12 and canonical autophagy initiation factors. N1303K-CFTR does not

aggregate or damage ER-tubules, and the autophagic intermediates detected are not being consumed

by phagophores proposed to engulf damaged ER during ER-phagy (Khaminets, Heinrich et al. 2015,

Forrester, De Leonibus et al. 2019). Instead, data are consistent with an ER-membrane containing

N1303K-CFTR associating with autophagy initiation machinery present in WIPI1 decorated ER-

microdomains. The N1303K-CFTR ER-membrane appears to be modified and converted into ER-

connected phagophores that form in association with WIPI1-rings containing docked autolysosomes.

These data suggest that ERAD-resistant degradation intermediates of membrane proteins are sorted

into ER-connected phagophores' membranes. Interpretations are consistent with results from ultra-thin

sectioning and electron microscopic studies showing connections between ER tubules and phagophores

formed in starved cells (Yla-Anttila, Vihinen et al. 2009, Le Bars, Marion et al. 2014) (Axe, Walker et al.

2008).

The Conformation of CFTR -Intermediates Controls Triage for ERAD versus ERQC-autophagy

Next, we investigated the role played by the conformation of kinetically-trapped intermediates of

CFTR in their selection for ERQC-autophagy (Figure 6). The conformation of N1303K-CFTR intermediates

was probed with a cysteine (Cys) pair cross-linking assay that monitors the formation of contacts

between NBD1 or NBD2 with different CLs identified in high-resolution models of CFTR structure (Figure 6A-C). Cys pairs were introduced at CL/NBD interfaces in Cys-less CFTR with or without N1303K, and cross-linking was conducted to evaluate the folding of CFTR to a formation that contains the indicated domain interfaces that are present in folded CFTR {He, 2013 #3203}. When a Cys pair was introduced at CL2/NBD2 interface at amino-acid residues C276C/Y1307C, cross-linking of CFTR was detected after treatment of cells with MTS disulfide bond cross-linkers of different spacer arm lengths, M3M and M8M (Figure 6C, lanes 2 and 3, marked with X). VX-809 increased signals that report on the CL2/NBD2 interface within WT-CFTR, accumulating in the low mobility C-form (Figure 6C, lanes 5 and 6). However, in the absence or presence of VX-809, no cross-linked band was detected in N1303K-CFTR that reports on the CL2/NBD2 interface (Figure 6C, lanes 7-12). Similar results were found in N1303D-CFTR (Figure S5B, lanes 1-6), indicating the introduction of a charged residue at N1303 disrupts the NBD2/CL2 interface in a manner that could not be reversed by VX-809.

The Cys pair at amino acid residues E543C/T1057C, in NBD1 and CL4, respectively, was acted upon by both MTS cross-linkers M3M and M8M to generate a lower mobility C-form of CFTR (Figure 6D, lanes 2 and 3). The introduction of N1303K into this reporter prevented the detection of the C-form of CFTR (Figure 5D, lane 7). Still, a faint cross-linked band was detected slightly above the C-band's location is seen in extracts of cells treated with M3M and M8M (Figure 6D, lanes 8 and 9, marked with an X'). VX-809 treatment increased levels of the B-form N1303K-CFTR (Figure 6D, lane 10) and the amount of band X' (Figure 6D, lanes 11 and 12). VX-809 restored the interface between CL3/NBD1 in N1303K-CFTR, as shown with the cross-linking of the E543C/T966C marked with X' 'Figure 6E).

Sub-domains of N1303K-CFTR assemble into a conformation that has features of folded WT CFTR, but folding steps involving NBD2 are defective and resistant to modulation by VX-809. The combined folding of N1303K-CFTR's-terminal domains and failure to complete late-stage domain:domain assembly reactions could cause the accumulation of kinetically-trapped intermediates.

Control of Triage Decisions by the Conformation of N1303K-CFTR

To test the model proposed for conformation-dependent triage of CFTR intermediates, we

developed a CFTR reporter that is conditionally degraded in the absence or presence of VX-809 by either

ERAD or ERQC-autophagy, respectively (Figure 7A). The disease-causing mutation P67L that is located

in the N-terminus causes global misfolding and degradation of P67L CFTR by ERAD, and VX-809

completely restores the folding and function of P67L CFTR (Ren, Grove, et al. 2013). P67L was therefore

introduced into N1303K-CFTR to create P67L-N1303K-CFTR. It was then asked: 1. What is the effect of

P67L on the triage of N1303K-CFTR (Figure 7B)? 2. Does stabilization of N-terminal domains by VX-809

cause P67L-N1303K CFTR to behave like N1303K-CFTR (Figure 7C).   P67L-N1303K-CFTR intermediates

exhibited a short half-life and were not detected in autolysosomes. Treatment with VX-809 restored

P67L-CFTR folding and accumulation in the C-form and in parallel incubations increases in the stability of

P67L-N1303K-CFTR intermediates and their accumulation in autolysosomes was now observed (Figure

7B and 7C). Globally misfolded P67L-N1303K-CFTR goes to ERAD, whereas VX-809 stabilized P67L-

N1303K-CFTR is degraded by ERQC-autophagy. At the same time, misfolded CFTR and F508-CFTR are

degraded by ERAD in the absence or presence of VX-809. These data demonstrate that the degradative

fate of a misfolded and detergent-soluble membrane protein such as CFTR is dictated by the stability of

its intermediates.

Discussion

We report that the mutation N1303K in NBD2 arrests CFTR folding at a late stage and causes kinetically trapped intermediates of it to accumulate in the ER. N1303K-CFTR intermediates are chaperoned by JB12 and Hsp70 and triaged in a conformation-specific manner for degradation by ERAD or ERQC-autophagy. Data presented suggest that globally misfolded CFTR is degraded by ERAD, whereas biochemically stable intermediates are selected for ERQC-autophagy. N1303K-CFTR intermediates have long 3-hour half-lives, and are soluble in non-ionic detergent, mobile in the ER tubular network, and contain cross-linkable domain interfaces present in folded CFTR. N1303K-CFTR intermediates are stabilized by VX-809, but stabilized N1303K-CFTR does not escape the ER and remains associated with JB12 and additional PQC factors. Disruption of early steps in CFTR folding to cause global misfolding of N1303K-CFTR via the introduction of the P67L mutation reduces the half-life of N1303K-CFTR, and P67L-N1303K-CFTR is not detected in the autophagy system. VX-809 stabilizes P67L-N1303K-CFTR intermediates causing them to be triaged to ERQC-autophagy instead of ERAD.   Data reported on N1303K-CFTR support the concept that in energized and biosynthetically active ER membranes, non-toxic intermediates of misfolded membrane proteins are selectively triaged for proteasomal or lysosomal degradation based on their conformation.

CFTR degradation intermediates are now grouped with an expanding collection of secretory and membrane proteins that misfold and accumulate in both ERAD-sensitive or ERAD-resistant conformations (Buchberger 2014, Houck, Ren, et al. 2014). ERAD requires that misfolded luminal or membrane proteins be unfolded and retrotranslocated to the cytosol, where they are translocated into the narrow opening of proteasomes' proteolytic chamber. In cases where nascent secretory proteins misfold in the ER-lumen and adopt a retrotranslocation-resistant conformation, the trapped intermediates interact with complexes containing ER-phagy receptors prior to selective ER-phagy (Fregno and Molinari 2018). We extensively evaluated the role of selective ER-phagy in the

conformation-dependent triage of N1303K-CFTR and did not identify an ER-phagy receptor or soluble autophagy receptor that was necessary for its conformation-dependent sorting and traffic to autolysosome (Table S1). Since there are a large number of LIR-containing proteins implicated in selective ER-phagy and selective autophagy, it could be that functional redundancy complicates the interpretation of siRNA and CRISPR/Cas9 loss of function experiments. However, we combined gene inactivation with imaging analysis and biochemical studies and were unable to observe physical or functional interactions between soluble or ER-membrane associated autophagy receptors with N1303K-CFTR that were suggestive of playing a role in quality control of membrane proteins.

In contrast to results reported with N1303K-CFTR, aggregates of ER-luminal alpha-1-antitrypsin and large assemblies of misfolded collagen are cleared from the ER by a selective ER-phagy process involving FAM134B and the ER exit site protein Sec31 (Forrester, De Leonibus, et al. 2019).{Omari, 2018 #9242}. N1303K-CFTR is excluded from ER-exits, and in these situations, the chaperone calnexin binds selective ER-phagy clients in the lumen and functions in a complex with FAM134B to target them for selective ER-phagy.   CFTR is a calnexin client, and calnexin facilitates assembly reactions involving C-terminal regions of CFTR, but a role for FAM134B in the degradation of N1303K-CFTR could not be demonstrated (Table S1). This might be the case because polytopic membranes expose surfaces in the cytosol that are ubiquitinated when they fail to fold correctly. N1303K-CFTR, like other misfolded forms of CFTR, is ubiquitinated, which could permit cytosolic ubiquitin chain binding autophagy receptors to usurp requirements for ER-phagy receptors. However, we were unable to demonstrate a role for p62, NBR1, FKBP8, or Tollip in the conformation-dependent sorting of misfolded membrane proteins to the autophagy initiation sites when ER-tubules contain N1303K-CFTR associate with WIPI1-rings (Table S1). Mechanisms for the conformation-dependent selection of misfolded and detergent soluble forms of CFTR for selective autophagy, therefore, appear different than those demonstrated for ERAD-resistant luminal aggregates and aggregates of cytosolic proteins.

Triage of nascent CFTR between folding and degradation is mediated by JB12, Hsp70, and ERQC-E3 ligases such as RMA1/RNF5 and CHIP (Meacham, Lu et al. 1999, Meacham, Patterson, et al. 2001, Younger, Chen et al. 2006, Grove, Fan, et al. 2011). N1303K-CFTR is present in complexes with JB12 and Hsp70, and siRNA depletion of JB12 hinders degradation of N130K-CFTR and blocks the accumulation of N1303K-CFTR in autolysosomes. The overexpression of dominant-negative QPD JB12 also prevents N1303K-CFTR from being degraded by ERAD or autophagy. QPD-JB12 can bind clients, but cannot interact with Hsp70, so QPD-JB12 can bind but fails to efficiently release clients (Grove, Fan, et al. 2011). JB12 is not required for basal autophagy, so a functional interaction of JB12 and Hsp70 appears needed for the delivery of N1303K-CFTR to the autophagy machinery of the ER. JB12 is detected in complexes with Beclin-1, and complex formation between JB12 and Beclin-1 is stimulated by ERQC-autophagy clients (Houck, Ren, et al. 2014). Therefore, stable client binding to JB12 could trigger the focal nucleation of autophagy initiation machinery (Joo, Dorsey et al. 2011, Pickles, Vigie et al. 2018) and facilitate the observed intramembrane sorting of its ERAD-resistant N1303K-CFTR to omegasomes. However, additional studies are required to determine the mechanism by which JB12 and Hsp70 assist in the sorting of stable degradation intermediates of membrane proteins to the autophagy initiation sites within the ER.

To define the pathway for N1303K-CFTR intermediates from the ER to the autolysosome, we carried out a combination of fixed and live-cell imaging at normal and super-resolution. In particular, we sought to determine if N1303K-CFTR entered autolysosomes via an ER-phagy like mechanism in which ER-subdomains containing it are consumed by phagophores or a focal mechanism involving direct entry of N1303K-CFTR into membranes for ER-derived autolysosomes (Houck, Ren, et al. 2014). To test the latter possibility that the association of ER-microdomains containing N1303K-CFTR or F508del-CFTR with initiation site machinery (WIPI1, LC3B, FIP200) that localizes within omegasomes and phagophores was

evaluated. ER membranes enriched in N1303K-CFTR and F508del-CFTR were detected on the perimeter of WIPI1 decorated ER membranes, but only N1303K-CFTR was detected in ER-associated membranes which were decorated with the phagophore marker LC3B. Likewise, N1303K-CFTR was also seen in WIPI1 enriched foci with the ULK1 subunit FIP200. In contrast, JB12 was excluded from ER-tubules that are decorated with LC3b and/or associate with WIPI1 and FIP200. It was common in live cells for LAMP1 enriched lysosomes to be detected in association with ER membranes containing N1303K-CFTR that extend from the surface of WIPI1-rings. There were also N1303K-CFTR containing foci in the 50 nm size-range that were sandwiched between the surface of WIPI1-rings and docked lysosomes. Some of such foci contained LC3b and appeared to represent autophagosomes in the act of docking with lysosomes. These imaging data are consistent with the concept of ERQC-autophagy involving the focal activation of selective autophagy in response to the accumulation of ERAD-resistant intermediates of N1303K-CFTR.

N1303K-CFTR is the second most common CF causing allele, and it was hoped that folding modulators developed to restore the F508del-CFTR function could also fix the N1303K-CFTR function. The Class I modulator VX-809 does increase the accumulation of N1303K-CFTR, but not proper folding. Treatment with VX-809 alone, or in combination with the tool compounds in the Class II modulator family, or VX-441, which is used in combination with Class I modulators clinically, does not restore N1303K-CFTR function in heterozygous or homozygous native human airways cells from CF patients {Veit, 2020 #9242}. We were, however, able to fix the folding of a small fraction of N1303K-CFTR in cells that are cultured at low temperatures. Reconstituted N1303K-CFTR channels isolated from the plasma membranes of cells cultured at low-temperature and assayed over a range of temperatures had the same conductance as WT CFTR. Rescued N1303K-CFTR channels were, however, thermodynamically unstable with a dramatically lower open probability than WT CFTR.   Notably, the CFTR channel potentiator VX-770 was found to enhance the open probability of N1303K-CFTR at temperatures between 20 and 37 $^{\circ}$C, but rescued channels are unstable.

Folding/assembly of NBD2 is a slow step in CFTR biogenesis that involves cooperative

conformational changes that result from dimerization of NBD2 with NBD1 and reorientation of

transmembrane spans and CLs (Wang, Wu et al. 2010, Kirk and Wang 2011, Veit, Xu et al. 2018). Defects

in the folding of N1303K-NBD2 could therefore block the folding progression of partially assembled

intermediates for several reasons. Class I and Class II modulators were developed to overcome defects

in F508del-NBD1 folding/assembly, and they are also able to suppress functional defects in a subset of

non-F508del CFTR mutants. Yet, rare CFTR alleles that respond to Trikaftor have common biogenic

defects that are similar to those caused by F508del (Middleton, Mall, et al. 2019). Such defects are

suppressed allosterically by stabilizing N-terminal regions of CFTR and promoting intramolecular

domain: domain assembly reactions involving F508del-NBD1 (Ren, Grove, et al. 2013, Veit, Avramescu,

et al. 2016). Clinically relevant restoration of the N1303K CFTR function might be achievable through

the development of modulators that rescue defects in NBD2 misfolding and/or assembly.

Materials and Methods

Reagents and antibodies

Reagents and antibodies were purchased from the following suppliers: HRP-conjugated goat anti-mouse IgG (31430) and goat anti-rabbit IgG (31461), Alex Fluor 488 goat anti-mouse IgG (A11029), Texas Red goat anti-mouse IgG (T-862), and goat anti-rabbit IgG (T-2767), Alexa Fluor Plus 647 goat anti-mouse IgG (A32728) and goat anti-rabbit IgG (A32733) from Thermo Fisher Scientific. Anti-beclin-1 (NB500-249), anti-FAM134B (NBP2-55248) and anti-Sec62 (NBP1-84045) from Novus Biologicals. anti-calnexin (C4731), anti-HA (H9658), anti-LC3B (L7543), anti-Myc (C3956), and anti-tubulin (T-9026) from Sigma-Aldrich. Anti-ERGIC53 (ALX-804-602) from Enzo Life Sciences. Anti-GABARAP+L1+L2 (ab109364) and anti-GM130 (ab52649) from Abcam. Anti-LAMP2 (555803) and anti-Sec31A (612351) from BD Pharmingen. Anti-p62 (H00008878-M0162) from Abnova. Anti-VMP1 (12978s) from Cell Signaling. DC Protein Assay (500-0119) and Clarity Western ECL Substrate (170-5061) from BioRad. Effectene Transfection Reagent (301427) from Qiagen. Lipofectamine RNAiMAX Reagent (100014472) from Thermo Fisher Scientific. Methanethiosulfonate (MTS) cross-linking reagents M3M (1,3-propanediyl bismethanethiosulfonate) and M8M (1,5-pentanediyl bismethanethiosulfonate) from Toronto Research Chemicals. ProLong Diamond antifade mountant without (P36970) or with DAPI (P36971) from Thermo Fisher Scientific.

Cell culture and transfection

African green monkey kidney cells (COS-7 ATCC CRL-1651), human embryonic kidney cells (HEK293, ATCC CRL-1573), and baby hamster kidney cells (BHK-21, ATCC CCL-10) were cultured in DuDulbecco'sagle minimum enriched medium (DMEM) supplemented with 10% fetal bovine serum (Sigma-Aldrich, St. Louis, MO) and 10U/ml penicillin, 10μg/ml streptomycin (GIBCO, Carlsbad, CA) at 37°C, 5% CO_2. Human lung tissue was procured under The University of North Carolina Office of Research Ethics Biomedical Institutional Review Board Approved Protocol No. 03–1396. Primary HBE cells were harvested and cultured using established procedures previously described in detail (Ref).

Plasmid transfections in cell lines were performed using Effectene Transfection Reagent following the product instruction. For knockdown experiments, all siRNAs were purchased from Thermo Fischer Scientific. For some target genes, two siRNAs to the same target gene were used to increase knockdown efficiency. COS-7 or HEK293 cells that were seeded into a 6-well plate at $3x10^5$ cells/well for 20 hr. were transfected at five pmol per well of siRNA with Lipofectamine RNAiMAX. Twenty hours post-transfection, the cells were split again to the appropriate density for transfection of plasmids using Effectene Transfection Reagent. siControl siVMP1 and sip62 sequences:

Protein sample preparation for SDS-PAGE and Western blotting

COS-7 cells or HEK293 cells were harvested from plates with citric saline (135mM KCl, 15 mM Sodium Citrate), and cell pellets were resuspended in PBST lysis solution (1% (v/v) Triton X-100 in phosphate-buffered saline) supplemented with one mM PMSF, 1x complete protease inhibitor cocktail (Roche, Basel, Switzerland). 2x Laemmli sample buffer without reducing agent was added, and the total lysates were homogenized by sonication. Protein concentrations were measured using detergent compatible DC Protein Assay Reagent, and 50 mM DTT was added before samples were resolved by SDS-PAGE and proceeded to Western blot analysis. Protein signal was detected by Clarity™ western ECL substrate and visualized by LAS680 imager (GE Life Sciences, Pittsburgh, USA). Images were analyzed by ImageJ (NIH, USA) and quantified with ImageQuant (GE Life Sciences, Pittsburgh, USA).

Co-immunoprecipitation assay

HEK293 or COS-7 cell lysates in PBST solution (1% Triton X-100 in PBS) were centrifuged at 100,000 g, at 4 °C, for 10 min to remove any insoluble fraction. Protein concentrations were measured using DC Protein Assay, and samples of an equal amount of proteins were used for immunoprecipitation analysis. To pull-down proteins interacting with CFTR, mAb596 antibodies cross-linked to Protein G agarose beads (Invitrogen) were incubated with cleared lysate for 1 hr. at 4 °C. Beads were washed 3x with PBST lysis buffer. Bound proteins were eluted from beads via incubation with 2x SDS-PAGE sample buffer without reducing agent at 37°C for 15 min. Anti-FLAG M2 beads (Sigma-Aldrich, St. Louis, USA) were used to pull-down proteins interacting with FLAG-tagged proteins. After washing, proteins bound were eluted with FLAG peptide. Eluted proteins were subjected to SDS-PAGE and Western blot analysis as described above.

Cycloheximide chase

To determine the steady-state level of different CFTR mutants, HEK293 cells transfected and grown overnight prior to incubation with 5μg/ml cycloheximide (CHX; Sigma-Aldrich, St. Louis, MO) in the presence and absence of 15μM CQ (Sigma-Aldrich, St. Louis, MO) or 10μM bortezomib (LC Laboratories, Woburn, MA). Cells were harvested at indicated time points, and protein samples were prepared for Western blot analysis as described above.

Triton X-100 solubility assay

Harvested cell pellets were lysed in PBST (1% Triton X-100 in PBS) and separated into soluble and insoluble fractions by centrifugation at 20,000xg for 30 min at 4˚C. The insoluble fraction was resuspended in a 2x sample buffer with an equal volume of soluble fraction and sonicated. Total lysates, soluble fractions, and insoluble fractions were resolved with SDS-PAGE and subjected to Western blot analysis.

Disulfide cross-linking in whole cells

Cross-linking of Cys pairs introduced in Cys-less CFTR expressed in HEK293 cells was performed essentially as previously described (Serohijos, Hegedus, et al. 2008). Briefly, HEK cells transiently transfected with Cys-less CFTR constructs with Cys pairs introduced at CL/NBD interfaces were grown in 6 well plates. The cell was harvested in citric saline, the cell pellet was washed in PBS and then resuspended in 30 µl PBS. 10 µl of cell suspension was mixed with 20 µl PBS with DMSO as vehicle control or PBS containing 300 µM MTS cross-linker to yield a final concentration of 200 µM of bifunctional cross-linkers M3M (1,3-propanediyl bismethanethiosulfonate), or M8M (1,5-Pentanediyl bismethane-thiosulfonate) (Toronto Research Chemicals). After 15 min incubation at room temperature, the cross-linking reaction was stopped with 120 µl 1.5x Laemmli sample buffer without DTT. After sonication, 30 µl of the samples were loaded on 7.5% SDS-PAGE, and anti-CFTR mAb 596 was used for Western blotting.

Immunofluorescence microscopy

COS-7 cells were used for most of the microscopy study due to their flat and spread out nature, making them ideal for resolving intracellular organelles. COS-7 cells were grown on thin coverslips (Thermo Fisher Sci. 12541A) in 6 well plates. After transfection and treatment, cells were fixed at -20 °C for 5 min with pre-cooled methanol. Cells were rehydrated by washing 3x in PBS and blocked with 3% BSA in PBS for 30 min, before immunostaining with various antibodies. Goat anti-mouse or goat anti-rabbit secondary antibodies with different fluorescent tags were used to detect proteins of interest. Coverslips were mounted onto slides using ProLong Diamond antifade mountant without (P36970) or with DAPI (P36971) (Invitrogen). Slides were allowed to dry in the dark at room temperature for at least 12 hours before imaging with an IX81 motorized inverted microscope (Olympus, Center Valley, PA) equipped with standard DAPI (blue), FITC (green), TRITC (red), Cyan (far red) filter cubes. STED images were collected

with a Leica TCS SP8 3X equipped and a FOV scanner. Acquired images were deconvoluted using

CellSens Software (San Jose, CA, USA) to remove background noise.

Live cell imaging

COS-7 cells were cultured in 35mm glass-bottom dishes (MatTek P35G-1.5-10-C). After overnight

transfection with fluorescent-tagged proteins, cells were rinsed with FluorBrite DMEM (Gibco Life Sci.)

and kept in the same medium during drug treatment and imaging. Live cell imaging was carried out

using an Olympus FV3000RS Confocal Microscope equipped with a humidified incubation chamber set

to 37 °C. Images were acquired and deconvoluted using CellSens software (Olympus).

Measurement of CFTR activity in USSING chambers

Ion transport measurements with HBEs were performed in modified Ussing chambers (Physiologic

Instruments) under voltage clamp-conditions using Acquire & Analyze (version 2.3) software (Physiologic

Instruments). Reference measurements [potential difference (P_d)] and transepithelial resistance (R_t)

were obtained for each culture. Short circuit current (I_{sc}) was measured every 20 s and recorded digitally

after bilateral equilibration at pH 7.4 for 10 min in 5 ml of pH 7.4 Krebs bicarbonate Ringers (KBR; 115

NaCl mM, 25 NaHCO$_3$ mM, 2.4 K$_2$HPO$_4$ mM, 1.2 CaCl$_2$ mM, 1.2 MgCl$_2$ mM, 0.4 KH$_2$PO$_4$ mM, and five D-

glucose mM). Chamber temperature was maintained at 36°C ± 1°C by a circulating water bath, and KBR

was bubbled with 95% O$_2$-5% CO$_2$ throughout the experiment.

Agonists and inhibitors used to modulate CFTR activity were purchased from Sigma Chemical. In

the presence of amiloride (100 µM, apical), forskolin (10 µM) was applied bilaterally to induce cAMP

activation of CFTR and VX-770 (5 µM, apical) was applied to further activate CFTR. Currents detected

were sensitive to the CFTR channel blocker CFTR$_{inh}$-172 (10 µM apical). As an internal control, UTP (100

µM) was used to activate calcium-activated chloride channels (CaCC) at the conclusion of the

experiment. Data were exported and analyzed in Microsoft Excel, and slope values were subtracted for

accuracy when necessary. Mean, SD, and SE measurements were calculated for each culture. Statistical analysis was performed by an unpaired two-tailed StStudent's-test. $P < 0.05$ was considered to indicate statistical significance. Line and Bar graphs were created using Origin 8.6 software (OriginLab).

Membrane isolation and planar bilayer based single-channel recording

BHK cells stabling expressing N1303K CFTR with or without stabilizing mutations H1402S and ΔRI/2PT/H1402S were grown to 90% confluency, and cells were moved to a 27 °C CO_2 incubator and grown with three µM VX-809 for 24 hrs. Cells were harvested and homogenized on ice in 10 mM HEPES, pH7.2, one mM EDTA containing a protease inhibitor cocktail. Centrifugation at 3000 g for 5 min removed nuclei and undisrupted cells. The supernatant was centrifuged at 100,000 g for 30 min to pellet membranes, which were then resuspended in phosphorylation buffer (10 mM Hepes, pH 7.2 containing 0.5 mM EGTA, two mM $MgCl_2$, and 250 mM sucrose).

Planar lipid bilayers were prepared by painting a 0.2 mm hole drilled in a Teflon cup with a phospholipid solution in n-decane containing a 3:1 mixture of 1-palmitoyl-2-oleoyl-sn-glycerol-3-phosphoethanolamine and 1-palmitoyl-2-oleoylsn- glycerol-3-phosphoserine (Avanti Polar Lipids). The lipid bilayer separated 1.0 ml of solution (*cis* side) from 5.0 ml of solution (*trans* side). Both chambers were magnetically stirred and thermally insulated. Heating and temperature were controlled with a Temperature Control System TC2BIP (Cell Micro Controls). CFTR ion channels were transferred into the preformed lipid bilayer by spontaneous fusion of membrane vesicles containing CFTR. To maintain uniform orientation and functional activity of CFTR channels, two mM ATP, 50 nM PKA, and membrane vesicles were added to the *cis* compartment only. All measurements were done in symmetrical salt solution (300 mM Tris/HCl; pH 7.2; 3 mM $MgCl_2$ and 1 mM EGTA) under voltage-clamp conditions, using an Axopatch 200B amplifier (Axon Instrument/Molecular Device). The membrane voltage potential of −75 mV is the difference between *cis* and *trans* (ground) compartments. The output signal was filtered

with an 8-pole Bessel low-pass filter LPBF-48DG (NPI Electronic, Tamm, Germany) with a cut-off

frequency of 50 Hz, digitized with a sampling rate of 500 Hz, and recorded with pClamp9.2software.

Origin 75 software (Origin Lab Corp., Northampton, MA, USA) was used to fit all-points histograms

(pClamp 9.2, Axon Instruments) by multi-peak Gaussians. Single-channel current was defined as the

distance between peaks on the fitting curve and used for the calculation of the single-channel

conductance. The probability of the single-channel being open (P_o) was calculated as a ratio of the area

under the peak for the open state to the total area under both peaks on the fitting curve.

Acknowledgments:

This work was supported by grants from the Cystic Fibrosis Foundation (CYR18XX0, BOUCHE19R0, GENTZS19I0) and NIH (GM056981 and P03DK065988). FRAP and STED Images were collected with the assistance of the UNC Hooker and Neuroscience Imaging Core facilities. Susan Boyle assisted with the culture of primary human bronchial epithelial cells.

Literature Cited

Axe, E. L., S. A. Walker, M. Manifava, P. Chandra, H. L. Roderick, A. Habermann, G. Griffiths, and N. T. Ktistakis (2008). "A"tophagosome formation from membrane compartments enriched in phosphatidylinositol 3-phosphate and dynamically connected to the endoplasmic reticulum." " Cell Biol **182**(4): 685-701.

Buchberger, A. (2014). "ERQC autophagy: Yet another way to die." Mol Cell **54**(1): 3-4.

Cutting, G. R. (2017). "T"eating Specific Variants Causing Cystic Fibrosis." "AMA **318**(21): 2130-2131.
Cyr, D. M. (2005). "A"rest of CFTRDeltaF508 folding." "at Struct Mol Biol **12**(1): 2-3.

Dalal, S., M. F. Rosser, D. M. Cyr and P. I. Hanson (2004). "D"stinct roles for the AAA ATPases NSF and p97 in the secretory pathway." "ol Biol Cell **15**(2): 637-648.

Forrester, A., C. De Leonibus, P. Grumati, E. Fasana, M. Piemontese, L. Staiano, I. Fregno, A. Raimondi, A. Marazza, G. Bruno, M. Iavazzo, D. Intartaglia, M. Seczynska, E. van Anken, I. Conte, M. A. De Matteis, I. Dikic, M. Molinari and C. Settembre (2019). "A selective ER-phagy exerts procollagen quality control via a Calnexin-FAM134B complex." EMBO J **38**(2).

Fregno, I. and M. Molinari (2018). "E"doplasmic reticulum turnover: ER-phagy and other flavors in selective and non-selective ER clearance." "1000Res **7**: 454.

Fulcher, M. L., S. Gabriel, K. A. Burns, J. R. Yankaskas and S. H. Randell (2005). "W"ll-differentiated human airway epithelial cell cultures." "ethods Mol Med **107**: 183-206.

Ge, L., M. Zhang, S. J. Kenny, D. Liu, M. Maeda, K. Saito, A. Mathur, K. Xu and R. Schekman (2017). "R"modeling of ER-exit sites initiates a membrane supply pathway for autophagosome biogenesis." "MBO Rep **18**(9): 1586-1603.

Gentzsch, M., H. Y. Ren, S. A. Houck, N. L. Quinney, D. M. Cholon, P. Sopha, I. G. Chaudhry, J. Das, N. V. Dokholyan, S. H. Randell and D. M. Cyr (2016). "R"storation of R117H CFTR folding and function in human airway cells through combination treatment with VX-809 and VX-770." "m J Physiol Lung Cell Mol Physiol **311**(3): L550-559.

Grove, D. E., C. Y. Fan, H. Y. Ren and D. M. Cyr (2011). "T"e endoplasmic reticulum-associated Hsp40 DNAJB12 and Hsc70 cooperate to facilitate RMA1 E3-dependent degradation of nascent CFTRDeltaF508." "ol Biol Cell **22**(3): 301-314.

Grove, D. E., M. F. Rosser, H. Y. Ren, A. P. Naren and D. M. Cyr (2009). "M"chanisms for rescue of correctable folding defects in CFTRDelta F508." "ol Biol Cell **20**(18): 4059-4069.

He, L., P. Kota, A. A. Aleksandrov, L. Cui, T. Jensen, N. V. Dokholyan and J. R. Riordan (2013). "C"rrectors of DeltaF508 CFTR restore global conformational maturation without thermally stabilizing the mutant protein." "ASEB J **27**(2): 536-545.

Houck, S. A. and D. M. Cyr (2012). "M"chanisms for quality control of misfolded transmembrane proteins." "iochim Biophys Acta **1818**(4): 1108-1114.

Houck, S. A., H. Y. Ren, V. J. Madden, J. N. Bonner, M. P. Conlin, J. A. Janovick, P. M. Conn and D. M. Cyr (2014). "Q"ality control autophagy degrades soluble ERAD-resistant conformers of the misfolded membrane protein GnRHR." "ol Cell **54**(1): 166-179.

Joo, J. H., F. C. Dorsey, A. Joshi, K. M. Hennessy-Walters, K. L. Rose, K. McCastlain, J. Zhang, R. Iyengar, C. H. Jung, D. F. Suen, M. A. Steeves, C. Y. Yang, S. M. Prater, D. H. Kim, C. B. Thompson, R. J. Youle, P. A. Ney, J. L. Cleveland and M. Kundu (2011). "H"p90-Cdc37 chaperone complex regulates Ulk1- and Atg13-mediated mitophagy." "ol Cell **43**(4): 572-585.

Khaminets, A., T. Heinrich, M. Mari, P. Grumati, A. K. Huebner, M. Akutsu, L. Liebmann, A. Stolz, S. Nietzsche, N. Koch, M. Mauthe, I. Katona, B. Qualmann, J. Weis, F. Reggiori, I. Kurth, C. A. Hubner and I. Dikic (2015). "Regulation of endoplasmic reticulum turnover by selective autophagy." Nature **522**(7556): 354-358.

Kirk, K. L. and W. Wang (2011). "A"unified view of cystic fibrosis transmembrane conductance regulator (CFTR) gating: combining the allosterism of a ligand-gated channel with the enzymatic activity of an ATP-binding cassette (ABC) transporter." " Biol Chem **286**(15): 12813-12819.

Le Bars, R., J. Marion, R. Le Borgne, B. Satiat-Jeunemaitre and M. W. Bianchi (2014). "A"G5 defines a phagophore domain connected to the endoplasmic reticulum during autophagosome formation in plants." "at Commun **5**: 4121.

Matsunaga, K., E. Morita, T. Saitoh, S. Akira, N. T. Ktistakis, T. Izumi, T. Noda and T. Yoshimori (2010). "A"tophagy requires endoplasmic reticulum targeting of the PI3-kinase complex via Atg14L." " Cell Biol **190**(4): 511-521.

Meacham, G. C., Z. Lu, S. King, E. Sorscher, A. Tousson and D. M. Cyr (1999). "T"e Hdj-2/Hsc70 chaperone pair facilitates early steps in CFTR biogenesis." "mbo J **18**(6): 1492-1505.

Meacham, G. C., C. Patterson, W. Zhang, J. M. Younger and D. M. Cyr (2001). "T"e Hsc70 co-chaperone CHIP targets immature CFTR for proteasomal degradation." "at Cell Biol **3**(1): 100-105.

Middleton, P. G., M. A. Mall, P. Drevinek, L. C. Lands, E. F. McKone, D. Polineni, B. W. Ramsey, J. L. Taylor-Cousar, E. Tullis, F. Vermeulen, G. Marigowda, C. M. McKee, S. M. Moskowitz, N. Nair, J. Savage, C. Simard, S. Tian, D. Waltz, F. Xuan, S. M. Rowe, R. Jain and V. X. S. Group (2019). "E"exacaftor-Tezacaftor-Ivacaftor for Cystic Fibrosis with a Single Phe508del Allele." " Engl J Med **381**(19): 1809-1819.

Oren, Y. S., I. M. Pranke, B. Kerem and I. Sermet-Gaudelus (2017). "T"e suppression of premature termination codons and the repair of splicing mutations in CFTR." "urr Opin Pharmacol **34**: 125-131.
Pickles, S., P. Vigie and R. J. Youle (2018). "M"tophagy and Quality Control Mechanisms in Mitochondrial Maintenance." "urr Biol **28**(4): R170-R185.

Ren, H. Y., D. E. Grove, O. De La Rosa, S. A. Houck, P. Sopha, F. Van Goor, B. J. Hoffman and D. M. Cyr (2013). "V"-809 corrects folding defects in cystic fibrosis transmembrane conductance regulator protein through action on membrane-spanning domain 1." "ol Biol Cell **24**(19): 3016-3024.

Rosser, M. F., D. E. Grove, L. Chen and D. M. Cyr (2008). "A"sembly and misassembly of cystic fibrosis transmembrane conductance regulator: folding defects caused by deletion of F508 occur before and after the calnexin-dependent association of membrane spanning domain (MSD) 1 and MSD2." "ol Biol Cell **19**(11): 4570-4579.

Serohijos, A. W., T. Hegedus, A. A. Aleksandrov, L. He, L. Cui, N. V. Dokholyan and J. R. Riordan (2008). "P"enylalanine-508 mediates a cytoplasmic-membrane domain contact in the CFTR 3D structure crucial to assembly and channel function." "roc Natl Acad Sci U S A **105**(9): 3256-3261.

Singh, A. K., Y. Fan, C. Balut, S. Alani, A. M. Manelli, A. M. Swensen, Y. Jia, T. R. Neelands, T. A. Vortherms, B. Liu, X. B. Searle, X. Wang, W. Gao, T. C. Hwang, H. Y. Ren, D. Cyr, P. R. Kym, K. Conrath and C. Tse (2020). "B"ological Characterization of F508delCFTR Protein Processing by the CFTR Corrector ABBV-2222/GLPG2222." " Pharmacol Exp Ther **372**(1): 107-118.

Sopha, P., H. Y. Ren, D. E. Grove and D. M. Cyr (2017). "E"doplasmic reticulum stress-induced degradation of DNAJB12 stimulates BOK accumulation and primes cancer cells for apoptosis." " Biol Chem **292**(28): 11792-11803.

Vargas, J. N. S., C. Wang, E. Bunker, L. Hao, D. Maric, G. Schiavo, F. Randow and R. J. Youle (2019). "S"atiotemporal Control of ULK1 Activation by NDP52 and TBK1 during Selective Autophagy." "ol Cell **74**(2): 347-362 e346.

Veit, G., R. G. Avramescu, A. N. Chiang, S. A. Houck, Z. Cai, K. W. Peters, J. S. Hong, H. B. Pollard, W. B. Guggino, W. E. Balch, W. R. Skach, G. R. Cutting, R. A. Frizzell, D. N. Sheppard, D. M. Cyr, E. J. Sorscher, J. L. Brodsky and G. L. Lukacs (2016). "F"om CFTR biology toward combinatorial pharmacotherapy: expanded classification of cystic fibrosis mutations." "ol Biol Cell **27**(3): 424-433.

Veit, G., H. Xu, E. Dreano, R. G. Avramescu, M. Bagdany, L. K. Beitel, A. Roldan, M. A. Hancock, C. Lay, W. Li, K. Morin, S. Gao, P. A. Mak, E. Ainscow, A. P. Orth, P. McNamara, A. Edelman, S. Frenkiel, E. Matouk, I. Sermet-Gaudelus, W. G. Barnes and G. L. Lukacs (2018). "S"ructure-guided combination therapy to potently improve the function of mutant CFTRs." "at Med **24**(11): 1732-1742.

Wang, W., J. Wu, K. Bernard, G. Li, G. Wang, M. O. Bevensee and K. L. Kirk (2010). "A"P-independent CFTR channel gating and allosteric modulation by phosphorylation." "roc Natl Acad Sci U S A **107**(8): 3888-3893. Yla-Anttila, P., H. Vihinen, E. Jokitalo and E. L. Eskelinen (2009). "3" tomography reveals connections between the phagophore and endoplasmic reticulum." "utophagy **5**(8): 1180-1185.

Younger, J. M., L. Chen, H. Y. Ren, M. F. Rosser, E. L. Turnbull, C. Y. Fan, C. Patterson and D. M. Cyr (2006). "S"quential quality-control checkpoints triage misfolded cystic fibrosis transmembrane conductance regulator." "ell **126**(3): 571-582.

Younger, J. M., H. Y. Ren, L. Chen, C. Y. Fan, A. Fields, C. Patterson and D. M. Cyr (2004). "A"foldable CFTR{Delta}F508 biogenic intermediate accumulates upon inhibition of the Hsc70-CHIP E3 ubiquitin ligase." " Cell Biol **167**(6): 1075-1085.

FIGURE LEGENDS

Figure 1. Function of N1303K-CFTR is partially restored by cell culture at low-temperature but is insensitive to folding modulators. (A) The effect of small molecule folding modulators and channel potentiators on CFTR activity in polarized native human bronchial epithelial (HBE) was measured in USSING chambers. Representative traces illustrate the response of CFTR expressed in HBE treated with indicated compounds and CFTR channel activators (B): Bar graphs represent the mean net response (n=4 cultures per corrector treatment) to acutely added stimuli or inhibitors in USSING chamber experiments. HBE from patients carrying N1303K/W1282X mutations were chronically treated for 48 h with 3 µM VX-809, or 5 µM C3+C4, or 0.1% DMSO, as indicated. See Figure S1A for Western blot analysis of the effect of correctors on N1303K/W1282X CFTR expression in HBE. (C) Single channel function of the WT and rescued N1303K (rN1303K) CFTR in the lipid bilayer. The ion channels were transferred into the preformed lipid bilayer by spontaneous fusion of membrane vesicles prepared from BHK cells stably expressing WT or rescued N1303K CFTR. BHK cells expressing N1303K CFTR was rescued by growing at 27 °C in the presence of 5 µM C3 and C4 for the last 24 hrs before harvesting and labeled as rN1303K on the figure. Single channels were recorded at 25 °C (C) or 37 °C (D) for WT CFTR (top panel), rescued N1303K CFTR (middle panel), and rescued N1303K CFTR in the presence of VX-770 (bottom panel). The all points histogram used to calculate single channel conductance (γ) and open probability (P_o) is shown on the left of the upper line of each group. The 5 min single channel recording used to prepare all points histogram is shown on the right of the upper line. 7 independent experiments of total 58 minutes duration, 5 independent experiments of total 35 minutes duration, and 4 independent experiments of total 17 minutes duration were used to calculate γ and P_o of WT CFTR, rN1303K and rN1303K+VX-770, respectively, at 25 °C (C). 5 independent experiments of total 42 minutes, 4 independent experiments of total 28 minutes duration, and 3 independent experiments with total 12 minutes duration were used to calculate γ and P_o of WT CFTR, rN1303K and rN1303K+VX-770, respectively, at 37 °C (D). The difference in conductance between rN1303K and WT CFTR is not significant while the difference in P_o is significant ($p<0.05$). E. Table summarizing the γ and P_o of WT and rN1303K CFTR measured at 25, 30 and 37 °C.

Figure 2. N1303K CFTR is more stable than ΔF508 CFTR and is partially degraded by autophagy
 (A). Cycloheximide (CHX) chase of N1303K and ΔF508 CFTR in the absence or presence of VX-809. HEK cells were transiently transfected with N1303K and ΔF508 CFTR. At 24 hrs. after transfection, cells were pretreated with or without 5 µM VX-809 for 2 hrs., followed by addition of 100 µM CHX and incubation for the indicated time. Cell lysates in SDS-PAGE sample buffer were subjected to 7.5% SDS-PAGE and Western blot analysis with CFTR mAb 596. Protein loading was normalized by reprobing the blots with anti-tubulin antibody. Representative blots are shown on the left, with quantification normalized to tubulin shown on the right (n=3). CHX chase of N1303K and ΔF508 CFTR after overnight VX-809 treatment gave similar results. (B). Live-Cell imaging and FRAP of EGFP-N1303K CFTR and EGFP-F508del CFTR that was transiently expressed in COS-7 cells and imaged with a Zeiss-880 microscope using a PMT detector. The swarm plot shows the diffusion coefficients for N1303K (8 trials) and ΔF508 CFTR (10 trials). (C). Fractionation of WT, F508del and N1303K CFTR into Triton X-100 soluble and insoluble fractions of cell extracts. (D) N1303K is excluded from ER exit sites and co-localizes in the tubular ER with calnexin. COS-7 cells were transiently transfected with N1303K CFTR. 18 hrs. after-transection and 6 hrs. prior fixation cells were treated with 15 µM chloroquine (CQ). Immunostaining was carried out on methanol fixed cells with anti-GFP, and with antibodies to endogenous Sec31A and calnexin (CNX)

Figure 3. N130K-CFTR is sensitive to elevation of JB12 and RMA1/RNF5 and accumulates in ER tubules with JB12 or membranes that exclude JB12 and contain LC3B. (A) Changes in steady-state levels of CFTR, F508del-CFTR, and N1303K-CFTR in the absence or presence of VX-809 that are caused by the elevation of JB12 or RMA1/RNF5. Indicated forms of CFTR were expressed in the absence or presence of VX-809 that was added to cell cultures 6-hours prior to harvest and analysis by western blot. Western blots in middle panels show levels of overexpressed JB12 (50 ng of pCDNA3-DNAJB12) or RMA1/RNF5 (100 ng of pCDNA-RMA1/RNF5) relative to endogenous pools. (B) Proximity labeling of N1303K-CFTR by JB12-APEX in the absence or presence of VX-809. JB12-APEX, which contains an APEX domain localized in the ER-lumen was co-expressed with N1303K-CFTR in the absence or presence of VX-809. Cells were incubated with hydrogen peroxide and biotin-phenol and modified adducts were then isolated from cell extracts by streptavidin-pull downs and detected by western-blot with indicated antibodies. (C) Analysis of the localization of GFP-N1303K-CFTR in the ER relative to JB12 or RMA1/RNF5. N1303K-CFTR was expressed in COS-7 cells in the absence or presence of JB12 (expressed 50 ng of pCDNA3). Cells were fixed with methanol and stained for JB12 or endogenous LC3. Immunofluorescent images were collected with an Olympus IX-81 with CellSens software and processed with ICY. (D) Overexpressed FLAG-DNAJB12 or FLAG-QPD-DNAJB12 with GFP-N1303K-CFTR in COS-7 cells. Cells were fixed in methanol at -20°C for 5 minutes, then blocked for 1 hour at ambient temperature with blocking buffer (3% bovine serum albumin in phosphate buffered solution), followed by primary and secondary antibody labeling diluted with blocking buffer. After washes with blocking buffer, coverslips were mounted using ProLong Diamond with DAPI mounting solution, and sealed immediately with nail polish. (E) siRNA (or control) for DNAJB12 and transiently transfected (effectene) with GFP-N1303K in COS-7 cells, then fixed (using same procedure as in D).

Figure 4. JB12 is Necessary for Traffic of N1303K-CFTR from the ER to Autolysosome. (A) Localization in COS-7 cells of untagged N1303K-CFTR, but not F508del-CFTR, in foci with endogenous LC3B. (B) Localization of GFP-N1303K-CFTR in autolysosomes with endogenous LAMP1 and LC3B is sensitive to PI3K-III, an inhibitor of the autophagy initiation kinase, VPS34. (C) Depletion of JB12 from COS-7 cells by siRNA reduces movement of N1303K-CFTR from the ER to LAMP1 positive foci. JB12 levels were reduced via siRNA and 18-hours later cells were transfected with GFP-N1303K-CFTR. Cells were fixed 18 hours after introduction of GFP-N1303K-CFTR and stained for endogenous LAMP1 or the ER-marker Sec62. Immunofluorescent images were collected with an Olympus IX-81 with CellSens software and processed with ICY.

Figure 5. N1303K-CFTR intermediates, but not F508del-CFTR, are detected by fluorescence microscopy with autophagy initiation machinery during sorting within the ER and trafficking to autolysosome. (A) N1303K-CFTR co-localizes in ER-tubules with the autophagy initiation factor mCherry-WIPI1 and the translocon associated Sec62. (B) ER-attached membranes containing N1303K-CFTR extend from rings of ER-membrane that is decorated with mCherry-WIPI1 and endogenous FIP200. (C) Super-resolution analysis of associations between N1303K-CFTR, RPF-WIPI1 and BFP-LC3 in live COS-7 cells. A and B display images of fixed COS-7 cells expressing GFP-N1303K-CFTR, and the indicated mCherry-tagged or endogenous marker proteins obtained with an Olympus IX-81 and CellSens software. In C images displayed were obtained with fixed cells at super-resolution with a Lecia STED Microscope.

Figure 6. N1303K CFTR folded to a conformation that contains native-like domain:domain contacts between NBD1 and ICL4, but native-like assembly of NBD2 is not detected.
(A). Model for domain structure of CFTR containing two nucleotide-binding domains (NBD1 and NBD2), two membrane-spanning domains (MSD1 and MSD2), and a regulatory region (R domain). Each MSD contains two cytoplasmic loops (CL) that form interfaces with the NBDs. F508 in NBD1, N1303 in NBD2, and residues mutated to Cys are marked. (B) The 3-dimensional structure of human CFTR (5UAK) with major domains and CLs labeled. (C-E). Cross-linking of Cys pairs introduced at different CL/NBD interfaces. HEK293 cells were transiently transfected with Cys-less CFTR or Cys-less N1303K CFTR with Cys pairs E543C/T1057C introduced at CL4/NBD1 interface (B), E543C/T966C at CL3/NBD1 interface (C) 276C/Y1307C at CL2/NBD2 interface (D), respectively. At 24 h after transfection, cells were treated with 3 μM VX-809 for 24 h at 27°C. Cells were harvested and resuspended in PBS and incubated with 200 μM M3M, M8M, or an equal amount of DMSO as vehicle control. Cell lysates in SDS-PAGE sample buffer without DTT were subjected to 7.5% SDS-PAGE and Western blot analysis with mAb 596. C, mature complex-glycosylated CFTR; B, immature core-glycosylated CFTR; X, cross-linked mature CFTR; X', cross-linked immature CFTR. Additional cross-linking experiments are presented in the supplement.

Figure 7. Introduction of N1303K in P67L CFTR renders it to be recognized by autophagy machinery after VX-809 treatment (A). Model for conformation dependent triage of misfolded CFTR intermediates (B). P67L/N1303L CFTR as a model to test conformation specific triage of CFTR degradation. COS-7 cells were transfected with various EGFP-tagged mutant CFTR. Cells were treated overnight with VX-809 before Bafilomycin A1 treatment for 6 hrs. Immunostaining was carried out with anti-LAMP2 (shown in red) and anti-LC3B (shown in blue) antibodies. (C). CHX chase of P67L and P67L/N1303K CFTR. HEK cells transfected with P67L CFTR or P67L/N1303K CFTR and CHX chase was carried out as described in Figure 2A figure legend.

	25 °C		30 °C		37 °C	
	P_o	γ (pS)	P_o	γ (pS)	P_o	γ (pS)
WT-CFTR	0.25± 0.03	10.8±0.2	0.60± 0.05	12.3±0.2	0.72±0.07	14.3±0.2
rN1303K	0.11±0.04	10.9±0.2	0.07±0.03	12.2±0.2	0.04±0.02	14.2±0.2
rN1303K+ VX-770	0.19±0.03	10.9±0.3	0.23±0.04	12.3±0.2	0.15±0.03	14.2±0.3

Figure 5

A N1303K DFCP1 CFTR Sec62 Overlay
   ΔF508

B N1303K WIPI1 CFTR FIP200 Overlay
   ΔF508

C Overlay (STED) LAMP2 N1303K LC3b Overlay (Confocal)

Figure 7

A   CFTR

P67
F508   N1303

Global misfold → ERAD

Autophagy

Late stage misfold

Green: EGPF/CFTR   Red: LAMP2   Blue: LC3B

B   N1303K   ΔF508   P67L   P67L/N1303K

Baf A1

INSET

VX-809
+ Baf A1

INSET

C   DMSO   VX-809
              0 1 2 3   0 1 2 3   CHX (hrs)

P67L
         250 —
         150 —   ◄ C
                                        ◄ B
tubulin

P67L/
N1303K
         250 —   ◄ C
         150 —   ◄ B
tubulin

● P67L   ● P67L+VX-809
● P67L/N1303K   ● P67L/N1303K+VX-809

% of Initial (B band)

CHX Chase (hr)