Ethacridine inhibits SARS-CoV-2 by inactivating viral particles in cellular models

Xiaoquan Li[1, 2#], Peter Lidsky[3#], Yinghong Xiao[3#], Chien-Ting Wu[4], Miguel Garcia-Knight[3], Junjiao Yang[1, 2], Tsuguhisa Nakayama[5], Jayakar V. Nayak[5], Peter K. Jackson[4], Raul Andino[3, *], Xiaokun Shu[1, 2, *]

Affiliations:

[1]Department of Pharmaceutical Chemistry, University of California, San Francisco, San Francisco, California, USA.

[2]Cardiovascular Research Institute, University of California, San Francisco, San Francisco, California, USA

[3]Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, California, USA.

[4]Department of Baxter Laboratory for Stem Cell Biology, Department of Microbiology & Immunology, Stanford University, California, USA
[5]Department of Otolaryngology–Head and Neck Surgery, Stanford University, Stanford, California, USA

[#]These authors contributed to the work equally.

[*]Correspondence to: Raul Andino (email: raul.andino@ucsf.edu); Xiaokun Shu (email: xiaokun.shu@ucsf.edu)

Abstract

More than a million people have now died from COVID-19, because of infection with the
SARS-CoV-2 coronavirus. Currently, the FDA has approved remdesivir, an inhibitor of
SARS-CoV-2 replication, to treat COVID-19, though very recent data from WHO
showed little if any COVID19 protective effect. Here we report that ethacridine, a safe
and potent antiseptic use in humans, effectively inhibits SARS-CoV-2, at very low
concentrations ($EC_{50} \sim 0.08$ μM). Ethacridine was identified through a high-throughput
screening of an FDA-approved drug library in living cells using a fluorescent assay.
Interestingly, the main mode of action of ethacridine is to inactivate virus particles,
preventing binding to the host cells. Thus, our work has identified a potent drug with a
distinct mode of action against SARS-CoV-2.
36

Introduction

The worldwide outbreak of the respiratory disease COVID-19 is caused by the coronavirus SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2). SARS-CoV-2 is an RNA betacoronavirus of the family *Coronaviridae*. It contains a single-stranded positive-sense RNA genome encapsulated within a membrane envelope [1-4]. Its genomic RNA is approximately 30kb with a 5′-cap structure and 3′-poly-A tail [1]. The genome of SARS-CoV-2 can be split into two main regions that contain as many as 14 open reading frames (ORFs) [5].

The first region, containing the first ORF (ORF1a/b), is about two-thirds of the genome length. After coronavirus attachment and entry into the host cell, the viral genomic RNA is released. The first region of the genomic RNA is used as a template to directly translate two polyproteins: pp1a and pp1ab. The pp1a polyprotein is translated from ORF1a. The pp1ab polyprotein comes from a -1 ribosomal frameshift between ORF1a and ORF1b. The two overlapping polyproteins are processed by a papain-like protease (PLpro) and a 3-chymotrypsin-like protease (3CLpro), which is also referred to as the main protease - Mpro. Both pp1a and pp1ab are mainly processed by Mpro, which digests the polyproteins at no less than 11 conserved sites, starting with the auto-proteolytic cleavage of this viral protease itself from pp1a and pp1ab. The functional polypeptides, 16 non-structural proteins (nsp1–16), are released from the polyproteins after extensive proteolytic processing. Nsp12 (i.e., RNA-dependent RNA polymerase (RdRp)), together with other nsps (e.g., nsp7 and nsp8), forms a multi-subunit replicase/transcriptase complex (RTC) that is associated with the virus-induced double-membrane vesicles [4,6,7]. The membrane-bound RTC synthesizes a full-length negative-strand RNA template that is used to make positive-strand viral genomic RNA.

The remaining one-third of the genomic RNA is used by the RTC to synthesize subgenomic RNAs (sgRNAs) that encode four conserved structural proteins (spike protein (S), envelope protein (E), membrane protein (M), and nucleocapsid protein (N)), and several accessory proteins. Eventually, the viral RNA-N complex and S, M, and E proteins are assembled in the ER-Golgi intermediate compartment (ERGIC) to form a mature virion that is then released via budding from the host cell. S protein is exposed on the surface of the virion and binds the virus receptor ACE2 on the host cell surface. Therefore, Mpro plays a central and critical role in the lifecycle of the coronavirus and is an attractive drug target [8-10], which also include other biological steps essential for viral replication and budding.

To identify drugs that inhibit the coronavirus, we redesigned green fluorescent protein (GFP) into an activity reporter of Mpro that becomes fluorescent only upon cleavage by the active Mpro. Using this assay, we screened a drug library in living cells and identified several drugs that inhibit Mpro activity. One highly effective drug, ethacridine, inhibits SARS-CoV-2 production by inactivating the viral particles.

RESULTS

Rational design of a fluorogenic Mpro activity reporter FlipGFP^{Mpro}

To develop an activity reporter of Mpro with a large dynamic range suitable for high-throughput screening (HTS), we applied the GFP-based protease reporter called FlipGFP

[11], which was designed by flipping one of the 11 beta-strands of a split GFP. Briefly, the
split GFP contains two parts: one part contains beta-strands 10 and 11 (i.e., GFP10 and
11), and the other contains nine other beta-strands and the central alpha helix (i.e., GFP1–
9). GFP10–11 contains the highly conserved Glu222 that is essential for catalyzing
chromophore maturation. GFP1-9 contains the three amino acids that form the
chromophore via cyclization, dehydration and oxidation [12]. GFP10-11 spontaneously
binds GFP1-9 and catalyzes the chromophore maturation, leading to a GFP. To design an
Mpro activity reporter, we "flipped" GFP10-11 using heterodimeric coiled coils (E5 and
K5) so that the flipped GFP10-11 cannot bind GFP1-9 when Mpro is inactive, and thus,
no or little fluorescence is detected (**Fig. 1a**). We incorporated an Mpro-specific cleavage
sequence AVLQ↓SGFR (↓ denotes the cleavage site) between GFP11 and K5. In this
way, when Mpro cleaves the proteolytic site, GFP11 flipped back, which allowed
GFP10-11 to bind GFP1-9, resulting in bright fluorescence (**Fig. 1a**). We named this
reporter FlipGFP^{Mpro}. To normalize the fluorescence, we added a red fluorescent protein
mCherry in the construct via a "self-cleaving" 2A peptide[13] (**Fig. 1c**).
To determine if FlipGFP^{Mpro} reports on SARS-CoV-2's Mpro activity in living
cells, we expressed the human ACE2, SARS-CoV2 receptor, in HEK293-FlipGFP^{Mpro}
cells. Next, we infected the cells with SARS-CoV-2, and at 24 hours post-infection, cells
were analyzed by immunofluorescence using antibodies directed against double-stranded
RNA (dsRNA) and by the FlipGFP^{Mpro} signal. The green fluorescence of the sensor
FlipGFP^{Mpro}, normalized by the co-expressed mCherry, was 63% greater in the infected
cells than in non-infected cells (**Fig. 1d**). Infected cells also showed dsRNA fluorescence.
In contrast, non-infected (mock) cells showed no dsRNA staining (**Fig. 1d**). These data,
thus, demonstrate that FlipGFP^{Mpro} can report on SARS-CoV-2 Mpro activity in living
cells.
Next, we established a system for screening Mpro inhibitors in living cells by
exogenously expressing Mpro in HEK293. Specifically, we co-expressed wild-type Mpro
or an inactive Mpro mutant in which the catalytic cysteine 145 is mutated to alanine. The
green fluorescence of FlipGFP^{Mpro} was barely detected in the cells expressing the inactive
Mpro/C145A mutant, whereas the red fluorescence of mCherry was observed (**Fig. 1e**,
upper panels). On the other hand, strong green fluorescence was detected in the cells
expressing Mpro with similar levels of mCherry fluorescence (**Fig. 1e**, lower panels). The
green fluorescence of FlipGFP^{Mpro}, normalized to the red fluorescence of mCherry,
revealed an ~60-fold dynamic range between inactive and active Mpro (**Fig. 1f**).
Furthermore, based on these quantified data, we calculated a *Z'-factor*[14]
118
$$Z = 1 - \frac{(3\sigma_+ + 3\sigma_-)}{|\mu_+ - \mu_-|},$$
which is ~0.8 with Mpro and its inactive mutant as positive (+) and negative (-) controls,
respectively (here σ is standard deviation, μ is mean). This suggests that the assay is
robust for HTS.
The FlipGFP^{Mpro} sensor was not responsive to the TEV protease, and the
FlipGFP-based TEV reporter (FlipGFP^{TEV}) was only activated by TEV but not by Mpro
(**Fig. 1f**). Thus, FlipGFP^{Mpro} specifically detects Mpro activity with a large dynamic
range. Therefore, our data show that we have established a robust HTS system for
screening Mpro inhibitors at a BSL2 level with 60-fold fluorescence change and a robust
z'-factor. Comparing the normalized FlipGFP^{Mpro} fluorescence in the SARS-CoV-2
infected cells (**Fig. 1d**) with that from the exogenously Mpro-expressing cells (**Fig. 1e, f**)

Fig. 1. Design and demonstration of a GFP-based activity reporter of SARS-CoV-2 main protease Mpro. (**a**) Schematic of the reporter. (**b**) Sequence of the flipped GFP10-11. (**c**) Construct of the reporter FlipGFP[Mpro]. (**d**) Fluorescence images (left) and quantitative analysis (right) of SARS-CoV-2 or mock-infected HEK293T cells that co-expressed hACE2. The images in the FlipGFP channel were brightened 30-fold compared to those in (**e**). (**e**) Fluorescence images of HEK293T cells expressing FlipGFP[Mpro] and mCherry, together with the inactive Mpro mutant C145A (upper panels) or wild type Mpro (lower panels). (**f**) Normalized FlipGFP fluorescence by mCherry. The ratio of FlipGFP/mCherry for the Mpro/C145A is normalized to 1. Data are mean $\pm$ SD (n = 5). FlipGFP[TEV] is a TEV activity reporter containing TEV cleavage sequence in FlipGFP. Scale bar: 5 μm (**d**); 10 μm (**f**).

129 suggests that the active Mpro concentration in the cytoplasm of the coronavirus-infected
130 cells is ~100-fold lower than that of the HEK293 cells exogenously expressing Mpro
131 (under a EF1α promoter).
132
HTS of drugs that inhibit Mpro activity in living cells
Next, we conducted HTS of ~1600 FDA-approved drugs (20 μM final concentration, **Fig.**
2a). The reporter construct (FlipGFP^{Mpro} and mCherry) was transfected into HEK293
cells, followed by addition and incubation of the drugs. Levels of green fluorescence
normalized to red fluorescence were calculated. A volcano plot revealed ~120 drugs that
showed $\geq$ 50% reduction of Mpro activity with a p-value < 0.001 (**Fig. 2a**). To confirm
this result, we re-screened the identified ~120 drugs under similar conditions (Extended
Data Fig. 1). We further examined those top 15 drugs at a lower concentration (10 μM)
and found that 12 drugs showed $\geq$50% reduction of FlipGFP^{Mpro} fluorescence
(normalized by mCherry) at 10 μM concentration (**Fig. 2b**). We finally calculated an IC$_{50}$
for each of the 12 drugs. Six drugs were at 2 – 6 μM (**Fig. 2c**), and the rest were above 6
μM (Extended data Fig. 2).
145
Antiviral activity of identified drugs
We next investigated antiviral activity of selected drugs in Vero E6 cells. The monolayer
cells were pretreated with the drugs for 3 hours and cultivated in the presence of the
compounds at a concentration of 5 μM. After 16 hours of incubation, the culture media
samples were collected, and the amount of infectious particles were estimated by plaque
assay (**Fig. 3a, b** & Extended Data Fig. 3). Our data revealed that nine of the 12 drugs
showed significant antiviral activity at 5 μM. Strong inhibition was detected for
ethacridine with 5 – 6 logs reduction in viral titer, simeprevir ~4-log reduction, ABT-199
~2-log reduction, hydroxyprogesterone ~1-log reduction, cinacalcet ~1-log reduction.
Two of the 12 drugs (ivermectin and verteporfin) were cytotoxic at 5–13 μM in Vero E6
cells and, thus, were excluded from the further analysis (Extended Data Fig. 4). As a
comparison, we tested the antiviral effect of a reported Mpro inhibitor, ebselen, which
showed ~2-log reduction in viral titer. And the RdRp inhibitor remdesivir showed ~4-log
reduction.
Next, we determined dose-response curves for top five selected drugs. First, our
data revealed that the EC$_{50}$ of four drugs (simeprevir, cinacalcet, ABT-199 and
hydroxyprogesterone) was 1–3 μM (**Fig. 3c – g**), within a range similar to their IC$_{50}$ in
inhibiting Mpro (**Fig. 2c**). This was consistent with the expectation that SARS-CoV-2
replication is inhibited by inhibiting Mpro activity. Indeed, as we were finalizing this
study, a preprint report showed that simeprevir inhibits Mpro activity and SARS-CoV-2
[15]. On the other hand, ethacridine showed outstanding antiviral activity (EC$_{50}$ ~ 0.08 μM,
Fig. 3e), which is 40-fold lower (i.e. stronger) than its Mpro-inhibiting activity (IC$_{50}$ ~
3.54 μM, **Fig. 2c**). This suggests that the superior antiviral activity of ethacridine is not
mainly accounted for by its Mpro-inhibiting activity. Lastly, for comparison, we also
determined the EC$_{50}$ of remdesivir ~0.52 μM in a side-by-side manner (**Fig. 3h**), which
indicates that ethacridine is more potent than remdesivir.
172
173 **Ethacridine inhibits SARS-CoV-2 by inactivating viral particles**

Fig. 2. High-throughput screening and drug identification using FlipGFP^{Mpro} in living cells. (a) Volcano plot of 1622 FDA-approved drugs (20 μM) in inhibiting Mpro. HEK293T cells were transfected with FlipGFP^{Mpro}, mCherry and Mpro. FlipGFP fluorescence was normalized to co-expressed mCherry. **(b)** Normalized ratio of Mpro activity in drug (10 μM) vs DMSO-incubated cells for the third-round validation. The Mpro activity was determined as FlipGFP fluorescence normalized to mCherry. The ratio of Mpro activity was calculated by normalizing Mpro activity with that of cells treated with DMSO. Data are mean $\pm$ SD (n = 5). The 15 drugs were identified from a second-round imaging of the 120 identified drugs (20 μM, Extended data Fig. 1). (c) Dose-response curve of top six drugs in inhibiting Mpro. Inhibition ratio was calculated as (1-(ratio of Mpro activity)) $\times$ 100%. IC$_{50}$ was represented as mean $\pm$ SEM (n = 5). See Extended data Fig. 2 for the other six drugs.

Fig. 3. Antiviral activities of the identified drugs against SARS-CoV-2. (**a**) Schematic showing the experimental design, see methods for details. (**b**) Quantitative analysis of viral titer from plaque assays on Vero E6 cells treated with each drug at 5 μM. Data are mean $\pm$ SEM (n = 3). *** $p < 0.001$. (**c–h**) Dose-response and cell-toxicity curve of each drug against SARS-CoV-2 by plaque assays. The percentage of relative infection was determined by the ratio of infection rate of SARS-CoV-2 treated with each drug divided by that of DMSO control. EC_{50} and CC_{50} are represented as mean $\pm$ SEM (n = 3).

To determine how ethacridine inhibits SARS-CoV-2, we tested infectivity of the virus
particles after ethacridine treatment using plaque assay, and we also measured viral RNA
levels using qRT-PCR. We examined antiviral effect of ethacridine on different stages of
the viral lifecycle of SARS-CoV-2, including before virus-cell binding, after cell entry,
and after budding. In particular, we pre-incubated SARS-CoV-2 particles with
ethacridine (5 µM) or DMSO for 1 hr. The mixture was then added to Vero E6 cells for
viral adsorption at a multiplicity of infection (MOI) at 0.5. Next, we removed the solution
and added fresh medium containing ethacridine (5 µM) or DMSO. Sixteen hours later,
we collected supernatant and conducted plaque assay with overlaid agar without
ethacridine or DMSO, measured viral titer of the supernatant. We also conducted qRT-
PCR and measured viral RNA levels in the supernatant and in the cells. By this way, we
developed three conditions (**Fig. 4a**): 1) Control (DMSO + DMSO): the virus and cells
were not exposed to the drug but DMSO; 2) The virus and cells were exposed to the drug
at all stages, including 1 hour before infection, during replication, and after viral budding
(i.e. Eth. + Eth.); 3) The virus and cells were exposed to the drug only after viral entry,
during replication, and after budding (i.e. DMSO + Eth.). Lastly, we used a fourth
condition (**Fig. 4b**): we conducted plaque assay right after pre-treatment of SARS-CoV-2
with ethacridine for 1 hr (i.e. Eth. [1 hr]), which determines antiviral activity of the drug
on the viral particles (equivalent to addition of the drug after viral budding).
When ethacridine was present all of the time (Eth. + Eth.), viral titers were
reduced 3 – 4 logs, compared with the control (DMSO + DMSO) (**Fig. 4a**, lower left
panel). When ethacridine was added to the Vero cells after viral entry (DMSO + Eth.),
similar level of reduction in infectivity (3 – 4 logs reduction in viral titer) was observed
(**Fig. 4a**). Interestingly, when SARS-CoV2 was pre-incubated with ethacridine for 1 hr
(Eth. [1 hr]) and followed by plaque assay without drug, we also observed 3 – 4 logs
reduction in infectivity (**Fig. 4b**). This result suggests that the drug inactivates SARS-
CoV2 particles infectivity. Because of similar-level reduction in infectivity in all of the
three conditions, our data strongly suggests that ethacridine inhibits SARS-CoV-2 by
mainly inactivating viral particles.
We next examined viral RNA accumulation in infected cells. qRT-PCR
measurement revealed no change of viral RNA (vRNA) levels when the drug was added
after viral binding and cell entry (DMSO + Eth.) in both the supernatant and the cells
(**Fig. 4a**, lower middle and right panels), compared with the control (DMSO + DMSO).
This indicates that the drug has no effect on vRNA replication. In contrast, if ethacridine
is present all of the time (i.e. Eth. + Eth.), we observed a reduction of 4 – 5 folds in
vRNA accumulation (**Fig. 4a**, lower middle and right panels). Because plaque assay-
based measurement of the same conditioned sample (Eth. + Eth.) showed 2400-fold
reduction in viral titer, the effect of ethacridine on viral replication (4 – 5 folds reduction)
is about 500-fold smaller than its effect on viral infectivity. This further supports that
ethacridine inhibits SARS-CoV-2 by inactivating the viral particles. The 4 – 5 folds
reduction of vRNA copies is likely due to partial inactivation of virus particles during
preincubation and thus a reduced number of infected cells (see below). Therefore, our
data suggests that ethacridine inhibits SARS-CoV-2 by mainly inactivating viral
particles, including those before infection, as well as new viruses after budding, with no
or little effect on vRNA replication (**Fig. 4c**).

Fig. 4. Ethacridine bocks SARS-CoV-2 by inactivating viral particles.
(**a**) Upper panel: schematic showing the experimental design for plaque assay and qRT-PCR. The virus was pre-incubated with ethacridine or DMSO for 1 hr. The mixture was added to Vero cells for adsorption at 4°C for 1 hr. Details can be found in the text. Lower panel: quantitative analysis of viral titer from plaque assay (left), and viral RNA (vRNA) copies by qRT-PCR in Vero cells (middle) and supernatant (right). (**b**) Quantitative analysis of viral titer by plaque assay. (**c**) Proposed mode of action of ethacridine by mainly inactivating viral particles of the coronavirus with no or little effect on viral RNA replication. (**d**) Schematic showing the experimental design for immunostaining. (**e**, **f**) Representative images of immunostaining against nucleocapsid protein (N) and spike (S) in Vero E6 cells after infection with the virus that was pre-treated with ethacridine (5 μM) or DMSO (control), or no infection (mock). (**g**) Quantitative analysis of the immunofluorescence. Data are mean $\pm$ SEM (n = 3 or biological replicas). **: p value < 0.01. *: p value < 0.05. Scale bar, 20 μm.

To further investigate the mechanism of ethacridine-based inactivation of the viral
particles, we conducted immunostaining and fluorescence imaging to determine whether
the ethacridine-treated SARS-CoV-2 can bind to and enter the cells. We treated SARS-
CoV-2 with ethacridine (5 μM) or DMSO for 1 hour. Then the virus was added to cells
for adsorption (37°C, 1 hour) at a MOI = 100. Cells were then quickly washed and fixed
with 4% PFA (**Fig. 4d**). Immunostaining with antibodies against the nucleocapsid protein
(N) of SARS-CoV-2 showed strong anti-N fluorescence on the plasma membrane of the
cells infected with virus+DMSO mixture, but little anti-N fluorescence in cells treated
with virus+ethacridine mixture (**Fig. 4e**). Immunostaining against Spike (S) of SARS-
CoV-2 also revealed fluorescence signals on control cells, but little fluorescence on cells
infected with the virus+ethacridine mixture (**Fig. 4f**). Quantification of fluorescence
revealed that ethacridine treatment led to a significant reduction of the anti-N and anti-S
fluorescence (**Fig. 4g**). These results indicate that ethacridine-treated SARS-CoV-2
cannot bind cells to initiate infection.
We also tested the dependency of the viral-inactivation effect of ethacridine on
dose, incubation time and incubation temperature with a plaque assay. For the conditions
tested, the viral-inactivation effect showed dose-dependency but was comparable to a 1-
or 2-hour incubation at room temperature or 37°C (Extended data Fig. 5).
237
Further validation in human cells
We further evaluated the anti-viral effect of ethacridine in human cells, including a
human lung epithelial A549 cell line stably expressing human ACE2 (A549[ACE2]) and
human primary nasal epithelial (HNE) cells to ensure that the antiviral effect is not
restricted to the Vero E6 cells. A plaque assay in A549[ACE2] cells revealed that
ethacridine-treated virus showed a dramatic decrease in infectivity (Extended data Fig.
6), indicating that the inhibitory effect of ethacridine is cell-type independent. To further
support this and validate the antiviral effects, primary HNE cells were infected with
ethacridine-treated virus and incubated with the media containing 5 μM ethacridine for
48 hours (**Fig. 5a**). Immunostaining using anti-N and anti-S antibodies in HNE cells
showed that, in the presence of ethacridine, the proportion of infected cells was
dramatically decreased (**Fig. 5b, c**), suggesting that ethacridine is also anti-viral in HNE
primary cells. Thus, ethacridine is effective in a cell culture model of primary nasal
epithelial cells at the air-liquid interface.
252
253 **DISCUSSION**
254
Several small molecule-based inhibitors have been identified that inhibit major targets
involved in the viral lifecycle of SARS-CoV-2, including the main protease Mpro and the
SARS-CoV-2 replicase RdRp. For example, α-ketoamide inhibitors of Mpro and the
FDA-approved drug ebselen have been reported to inhibit Mpro activity with EC50 at
0.53 and 4.67 μM, respectively [8-10]. The replicase RdRp inhibitor, remdesivir, has been
designed and very recently approved by FDA to treat severe COVID-19 patients [6,16].
However, very recent data from WHO showed no or little effect on hospital COVID-19
patients, including overall mortality, initiation of ventilation and duration of hospital stay
[17]. Thus, therapeutic efficacy of remdesivir is far from satisfactory, and new therapies are
urgently needed.

Fig. 5. Ethacridine blocks SARS-CoV-2 in primary human nasal epithelial cells. (**a**) Schematic showing the experimental design, see methods for details. (**b**) Representative images of immunostaining against spike (S) and nucleocapsid protein (N) in HNE cells 48 hours after viral infection in the presence ethacridine (5 μM) or DMSO (control), or no infection (mock). (**c**) Quantification of infection ratio based on anti-S or anti-N staining. Data are mean ± SEM (three biological replicates). **: p value < 0.01. Scale bar, 20 μm.

The current work started with developing a target-specific cell-based screening
assay. We developed the Mpro sensor, and our screen of 1622 FDA-approved drugs led
to identification of nine drugs that inhibit Mpro activity and show anti-SARS-CoV-2
activity. We characterized the antiviral property of five drugs further. Four of them,
including simeprevir, ABT-199, hydroxyprogesterone and cinacalcet effectively inhibited
Mpro and blocked SARS-CoV-2. This suggests that these four drugs inhibit SARS-CoV-
2 mainly by inhibiting Mpro activity. These drugs have similar or better EC50 than
ebselen, and thus they may be repurposed as potential therapies against COVID-19.
Furthermore, previous computational docking predicted that simeprevir and venetoclax
(ABT-199) bind and inhibit Mpro [18,19]. Hydroxyprogesterone inhibits SARS-CoV-2, but
its mode of action is unclear [20]. Our data suggests that hydroxyprogesterone may block
SARS-CoV-2 by inhibiting Mpro activity. Thus, our FlipGFP-based HTS approach
enable screening Mpro inhibitors in the cellular context, can be expanded to visualize
activity of key proteases of other viruses at BSL2 and to identify their inhibitors.
The most potent antiviral drug discovered in current work, ethacridine, showed
stronger antiviral potency than remdesivir and very little cell toxicity (**Fig. 3e**). It blocks
SARS-CoV-2 mainly by inactivating the viral particles. First, unlike the other identified
drugs that show similar range of IC_{50} in inhibiting Mpro and EC50 against SARS-CoV-2,
ethacridine has much higher anti-viral potency ($EC_{50} \sim 0.08$ μM) than its Mpro-inhibiting
activity ($IC_{50} \sim 3.5$ μM). This indicates that ethacridine's main mode of action is not via
inhibiting Mpro. Second, we provide direct evidence that after ethacridine treatment,
SARS-CoV-2 particles lose their infectivity and cannot infect or bind host cells. On the
other hand, ethacridine does not block viral RNA replication as shown by the qRT-PCR
data. Therefore, our data suggest that ethacridine blocks SARS-CoV-2 mainly by
inactivating the viral particles with no effect on viral RNA replication. The detailed
mechanisms of how ethacridine inactivates the viral particles will require further
investigation, such as potential ultrastructural changes of viral particles and binding of
ethacridine to viral RNA and/or protein.
Our findings reveal a new approach against SARS-CoV-2 via inactivating viral
particles, of which the efficacy is expected to be cell type-independent. Indeed,
ethacridine blocks SARS-CoV-2 in human cell line A549[ACE2] cells and also in primary
HNE cells. Furthermore, ethacridine is not toxic in various animal models including rat,
mice and rabbits. For example, mice treated with 20 mg/kg ethacridine by i.p. injection
showed no toxicity [21]. While currently ethacridine is mainly used as a topical wound
disinfectant, it has been applied to patients for treating puerperal sepsis via intravenous
injection [22]. Thus, it is plausible to validate its antiviral effect in animal models and
COVID-19 patients. Moreover, because it blocks the coronavirus by directly inactivates
the viral particles, ethacridine may be complemented with other pathway-targeted drugs
such as the replicase inhibitor remdesivir in treating COVID-19 patients for a better
clinical outcome.
305

Methods

Plasmid construction
Plasmid constructs were created by standard molecular biology techniques and confirmed by exhaustively sequencing the cloned fragments. To create pcDNA3-Mpro-flipGFP-T2A-mCherry, the GFP10-E5-GFP11-Mprosubstrate-K5 fragment was first generated by PCR with an overlapping primer with coding sequence of the MPro substrate, AVLQSGFR. This fragment was then digested with NheI and AflII and inserted into the pcDNA3-TEV-flipGFP-T2A-mCherry [11].

Characterization of GFP-based Mpro activity sensor FlipGFP^{Mpro}
To check the FlipGFP^{Mpro} signal in the presence of wildtype or mutant MPro of SARS-CoV-2, HEK293T cells were seeded onto eight-well plate and 24 hours later, 40 ng of pcDNA3-FlipGFP^{Mpro}-T2A-mCherry was transfected with 40 ng of pLVX-EF1alpha-nCoV2019-nsp5-2xStrep-IRES-Puro (expressing wildtype Mpro of SARS-CoV-2) or pLVX-EF1alpha-nCoV2019-nsp5-C145A-2xStrep-IRES-Puro (expressing C145A MPro mutant). To check the FlipGFP^{Mpro} signal after SARS-CoV-2 infection, HEK293T on 10-mm coverglasses were transfected with 200 ng of pcDNA3.1-hACE2 (Addgene, #145033) 24 hours later. Those 293T cells were further infected with SARS-CoV-2 at an MOI of 0.5. At 24 hours after infection, 293T cells were fixed and processed for immunostaining against dsRNA (Scicons, 10010500) and visualized with goat anti-mouse IgG H&L (Cy5 ®) (Abcam, ab6563). Images were taken using Nikon Eclipse Ti-E Spinning Disk under 20X and 60X. The cell pixel intensity in the GFP and mCherry channels were scored using Analyze Particle function in ImageJ. The ratio of GFP pixel intensity against mCherry pixel intensity was then compared in cells with wildtype and mutant MPro or in cells with SARS-CoV-2 or mock infection.

High-throughput screen (HTS) against the FDA-approved library
An FDA-approved drug library (MedChemExpress) containing 1622 compounds was used for HTS. For initial screening, 293T cells were seeded onto a 96-well plate, incubated overnight and transfected with 20 ng of pcDNA3-Mpro-flipGFP-T2A-mCherry and 20ng pLVX-EF1alpha-nCoV2019-nsp5-2xStrep-IRES-Puro encoding MPro using calcium-phosphate transfection. At 3 hours after transfection, compounds were added to individual wells at a 100X dilution for a final concentration of 20 μM. Verification of the 120 hits from the initial screening and further IC$_{50}$ testing of the 12 hits followed a similar protocol except that a decreasing concentration series was used. Images were acquired 20 hours after transfection in the GFP and mCherry channels with a Nikon Eclipse Ti-E Spinning Disk. The cells' pixel intensity in the GFP and mCherry channels were scored using Analyze Particle function in imageJ. MPro activity was calculated as the ratio of the GFP pixel intensity versus the mCherry pixel intensity.

Antiviral assay
The African green monkey kidney Vero E6 cell line was obtained from the American Type Culture Collection (ATCC, no. 1586) and maintained in Minimum Essential Medium (MEM; Gibco Invitrogen) supplemented with 10% fetal bovine serum (FBS; Gibco Invitrogen), 1% penicillin-streptomycin-glutamine (Gibco Invitrogen), at 37°C in a

humidified 5% CO_2 incubator. A clinical isolate of SARS-CoV-2 (USA-WA1/2020, BEI
Cat No: NR-52281) was propagated in Vero E6 cells. Viral titers were quantified with a
plaque assay. All the infections were performed at biosafety level-3 (BSL-3).

   To assess the antiviral activity, ~70% confluent monolayers of Vero E6 cells
(3×10^5 cells/well in 24-well plates) were pretreated with drugs at different concentration
for 3 hours (pretreatment) and then infected with SARS-CoV-2 (MOI=0.5) at 37ºC for 1
hour. The virus solution was removed, cells were further cultured with fresh medium
containing drugs at different concentrations. At 16 hours post-infection, viral titers of the
supernatants were detected with a plaque assay.

Plaque assay
Confluent monolayers of Vero E6 cells grown in six-well plates were incubated with the
serial dilutions of virus samples (250 µl/well) at 37ºC for 1 hour. Next, the cells were
overlayed with 1% agarose (Invitrogen) prepared with MEM supplemented with 2%
FBS. Three days later, cells were fixed with 4% formaldehyde for 2 hours, the overlay
was discarded, and samples were stained with crystal violet dye.

Viral RNA quantification
Viral RNA (vRNA) was extracted from cell pellets or supernatants with Tri-reagent
(Ambion), following the manufacturer's instructions. The supernatant samples were
pretreated with RNAse A for 2 hours at 37ºC to remove non-encapsided RNA. These
samples were also spiked with equivalent volumes of *Drosophila* C Virus (DCV) to
enable normalization, based on variation in RNA extractions. RNA from supernatants
was used directly to make cDNA using the iScript RT Supermix (BioRad). For cell
pellets, 1–2 mg of RNA was treated with DNAse I (NEB), x µL and 2 µL of this reaction
was used to make cDNA. qPCR was done using the Luna Universal qPCR Master Mix
(NEB) and run on a CFX connect qPCR Detection System (BioRad). To determine the
number of vRNA copies per ml, plasmids containing the nucleocapsid gene of SARS-
CoV-2 (cloned from the USA-WA1/2020 isolate) or the DCV full genome were used as
standards and diluted serially 10-fold to determine target copy numbers. Threshold cycle
(Ct) values were plotted against the number of target-copies, and the resultant standard
curve was used to determine the number of genome equivalents of vRNA in the samples.
For cell pellet samples, the vRNA copy number was normalized to the housekeeping
gene *huel,* and supernatant samples were normalized to spiked DCV RNA. All samples
were within the range of linearity of a standard curve and primers efficiencies were 100%
+/- 5%. The primers used for SARS-CoV-2 are 5′-TCCTGGTGATTCTTCTTCAGG-3′
and 5′-TCTGAGAGAGGGTCAAGTGC-3′, DCV 5-CAGCAAAGAAACAGCGTGAG-
3' and 5′-CACTTGCGCAACAATACGAG-3′, huel 5′-
TCAGACGACGAAGTCCCCATGAAG-3′ and 5′-
TCCTTACGCAATTTTTTCTCTCTGGC-3′

Cytotoxicity assay
Cytotoxicity of the identified drugs on Vero E6 cell was determined with WST-1 cell
proliferation assays (ROCHE, 5015944001). Twenty thousand cells were seeded into a
96-well plate and incubated for 20–24 h at 37 °C, and 1 µL of each compound at
decreasing concentrations was added. After 18 hrs incubation at 37 °C, WST-1 assays

398　were performed according to manufacturer's protocols. All experiments were performed
399　in triplicate.
400
Adsorption assay
The virus sample was supplemented with 5 μM ethacridine or with DMSO and incubated
for 1 hour at 37ºC. Samples were added to ~70% Vero E6 cells monolayers on coverslips
with an MOI of 100. The cells were immediately placed on ice for 1 hour, and then the
virus suspension was quickly removed, and cells were washed three times with ice-cold
PBS. Cells were immediately fixed with 4% PFA for 30 min at room temperature. PFA
was further washed with PBS and quenched with 1 M glycine in PBS. Immunostaining
was further carried out using an antibody against nucleocapsid (Genetex, SARS-CoV-2
(COVID-19) nucleocapsid antibody, GTX135357) or an antibody against spike (Genetex,
SARS-CoV-2 (COVID-19) spike antibody [1A9], GTX632604) and visualized with goat
anti-rabbit IgG H&L (Alexa Fluor® 488) (Abcam, ab150077) or goat anti-mouse IgG
H&L (Alexa Fluor® 555) (Abcam, ab150114). Images were acquired using Nikon
Eclipse Ti-E Spinning Disk under 60X and processed in ImageJ. Fluorescence signals
were calculated by pixel intensity using Analyze Particle function in Image J.
415
Statistics
All statistics were performed in GraphPad Prism. IC_{50} or EC_{50} of MPro inhibition,
cytotoxicity and antiviral activity was calculated using the non-linear fit function
(Variable slope). Non-paired t-test was used to compare differences between groups. One
way-ANOVA and Dunnett's multiple comparisons test was used to compare differences
among multiple groups.
422

423　**Acknowledgments:** We thank David Gordon and Nevan Krogan for sharing the cDNA
424　of Mpro of SARS-CoV-2.
425
426　**Funding:** This work was supported in part by NIH (R35 GM131766) to X.S., NIH (R01
427　AI36178, AI40085, P01 AI091575), the Bill and Melinda Gates Foundation and the
428　DARPA Intercept program (Contract No. HR0011-17-2-0027) to R.A.

429
Author contributions: X.L., X.S. initiated the project and designed the fluorescent
assays. X.L., J.Y. prepared the drug library. X.L. performed screening and characterized
dose response of Mpro inhibition. P.L., Y.X., R.A. designed the antiviral activity assays.
P.L., Y.X. conducted plaque assay. P.L., Y.X., X.L. performed quantitative analysis.
M.K. performed RT-qPCR. X. L. performed toxicity assays. P.L., Y.X., X.L. performed
and analyzed immuno-staining. P.L., Y.X., T.N., J.V.N., C.W. P. K. J. and R.A. designed
the HNE cell infection experiments. P.L. and Y.X. performed viral infection, C.W., T.N.
performed cell culturing, C.W. conducted immunostaining and imaging, X.L. performed
quantitative analysis of the HNE cells. X.S. wrote initial manuscript, X.L., P.L., Y.X.,
M.K., C.W. wrote the methods. All authors edited and contributed to the final draft.
440
441
442　**Competing interests:**
443　X.L., P.L., Y.X., R.A., X.S. have filed a patent on a new use of the identified compounds.

444
445 **Data and materials availability:** All data are available in the main text or in the
446 supplementary information.
447

448 **Supplementary Information:**

449 Extended data Figs. 1–6

References

1.   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of
probable bat origin. *Nature* 1–20 (2020). doi:10.1038/s41586-020-2012-7
2.   Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C. & Garry, R. F. The
proximal origin of SARS-CoV-2. *Nat Med* 1–3 (2020). doi:10.1038/s41591-020-
0820-9
3.   Gorbalenya, A. E. *et al.* The species Severe acute respiratory syndrome- related
coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nature*
Microbiology 1–9 (2020). doi:10.1038/s41564-020-0695-z
4.   Liu, C. *et al.* Research and Development on Therapeutic Agents and Vaccines for
COVID-19 and Related Human Coronavirus Diseases. *ACS Cent. Sci.*
acscentsci.0c00272–17 (2020). doi:10.1021/acscentsci.0c00272
5.   Gordon, D. E. *et al.* A SARS-CoV-2 protein interaction map reveals targets for
drug repurposing. *Nature* 1–13 (2020). doi:10.1038/s41586-020-2286-9
6.   Yin, W. *et al.* Structural basis for inhibition of the RNA-dependent RNA
polymerase from SARS-CoV-2 by remdesivir. *Science* eabc1560–10 (2020).
doi:10.1126/science.abc1560
7.   van Hemert, M. J. *et al.* SARS-Coronavirus Replication/Transcription Complexes
Are Membrane-Protected and Need a Host Factor for Activity In Vitro. *PLoS*
Pathog **4,** e1000054–10 (2008).
8.   Jin, Z. *et al.* Structure of Mpro from COVID-19 virus and discovery of its
inhibitors. *Nature* 1–28 (2020). doi:10.1038/s41586-020-2223-y
9.   Zhang, L. *et al.* Crystal structure of SARS-CoV-2 main protease provides a basis
for design of improved α-ketoamide inhibitors. *Science* **368,** 409–412 (2020).
10.   Dai, W. *et al.* Structure-based design of antiviral drug candidates targeting the
SARS-CoV-2 main protease. *Science* eabb4489–10 (2020).
doi:10.1126/science.abb4489
11.   Zhang, Q. *et al.* Designing a Green Fluorogenic Protease Reporter by Flipping a
Beta Strand of GFP for Imaging Apoptosis in Animals. *J Am Chem Soc* **141,**
4526–4530 (2019).
12.   Tsien, R. Y. The green fluorescent protein. *Annu Rev Biochem* **67,** 509–544
(1998).
13.   Szymczak, A. L. *et al.* Correction of multi-gene deficiency in vivo using a single
'self-cleaving' 2A peptide-based retroviral vector. *Nat Biotechnol* **22,** 589–594
(2004).
14.   Zhang, J., Chung, T. & Oldenburg, K. A Simple Statistical Parameter for Use in
Evaluation and Validation of High Throughput Screening Assays. *Journal of*
Biomolecular Screening **4,** 67–73 (1999).
15.   Lo, H. S. *et al.* Simeprevir suppresses SARS-CoV-2 replication and synergizes
with remdesivir. *bioRxiv* **395,** 1054–20 (2020).
16.   MD, Y. W. *et al.* Remdesivir in adults with severe COVID-19: a randomised,
double-blind, placebo-controlled, multicentre trial. *The Lancet* **395,** 1569–1578
(2020).
17.   WHO Solidarity trial consortium *et al.* Repurposed antiviral drugs for COVID-19
–interim WHO SOLIDARITY trial results. *medRxiv* **17,** e1003252–17 (2020).

18. Hosseini, F. S. & Amanlou, M. Anti-HCV and anti-malaria agent, potential
candidates to repurpose for coronavirus infection_ Virtual screening, molecular
docking, and molecular dynamics simulation study. *Life Sci.* **258,** 118205 (2020).
19. Chen, Y. W., Yiu, C.-P. B. & Wong, K.-Y. Prediction of the SARS-CoV-2 (2019-
nCoV) 3C-like protease (3CLpro) structure: virtual screening reveals velpatasvir,
ledipasvir, and other drug repurposing candidates. *F1000Res* **9,** 129–18 (2020).
20. Jeon, S. *et al.* Identification of Antiviral Drug Candidates against SARS-CoV-2
from FDA-Approved Drugs. *Antimicrobial Agents and Chemotherapy* **64,** 270–9
(2020).
21. Rotin, L. E. *et al.* Ibrutinib synergizes with poly(ADP-ribose) glycohydrolase
inhibitors to induce cell death in AML cells via a BTK-independent mechanism.
Oncotarget **7,** 2765–2779 (2016).
22. Oginz, P. & (null). Intravenous injections of rivanol in puerperal sepsis. *AJS* 647–
650 (1929). doi:10.1016/S0002-9610(29)90444-X
514

Extended data Fig. 1. Verification of the selected top 120 drugs using the Mpro activity reporter FlipGFP^{Mpro}. Ratio of Mpro activity was calculated based on FlipGFP^{Mpro} fluorescence of drug-incubated HEK293 cells, divided by that of DMSO-treated HEK293 cells. FlipGFP^{Mpro} fluorescence was normalized by mCherry in HEK293 cells, which co-expressed FlipGFP^{Mpro}, mCherry, and Mpro. Data are mean $\pm$ SD (n = 5).

Extended data Fig. 2. Dose-response curve of Mpro inhibition. The Mpro activity was determined as FlipGFP fluorescence normalized by mCherry. The ratios of Mpro activity were calculated by normalizing Mpro activity with that of cells treated with DMSO. Inhibition ratio was calculated as (1-(ratio of Mpro activity)) X100%. Data are mean $\pm$ SD (n = 5). IC_{50} was represented as mean $\pm$ SEM (n = 5).

Extended data Fig. 3. Antiviral activities of the identified drugs. Antiviral activities of five drugs (5 μM) were quantified by a plaque assay with SARS-CoV-2 in Vero E6 cells. Data are mean $\pm$ SD (n = 3). *: p value < 0.05; ***: p value < 0.001. ns: not significant. PFU: plaque-forming unit.

Extended data Fig. 4. Cytotoxicity of the identified drugs. The cytotoxicities of the indicated compounds were determined in Vero E6 cells with the WST-1 assay. CC_{50} is represented as mean $\pm$ SEM (n = 3).

Extended data Fig. 5. Virucide effect of ethacridine on SARS-CoV-2. Effects of ethacridine on the infectivity of SARS-CoV-2 were examined using plaque assay at 37°C (**a**, **b**) or in the room temperature (RT) (**c**, **d**). SARS-CoV-2 was mixed with ethacridine for 1 or 2 hours before being added to infect Vero E6 cells. Data are mean ± SEM (n = 3).

Drug(5uM)

Adsorption

Plaque Assay

3 hrs

1 hr

16 hrs

DMSO   5 uM

Ethacridine

+

1 hr

one bar 37 for 1hour

Plaque assay (b, c)

Immunostaining (d-j)

Pre-incub. SARS-CoV-2 w/ eth. (1hr) → Plaque assay

A549-ACE2

Viral titer (PFU/ml)   **

10^7
10^6
10^5
10^4
10^3
10^2
10^1
10^0
0

DMSO   Ethacridine

Plaque assay

RT-PCR

Viral titre (pfu/ml)   ***

DMSO   5 uM

Plaque assay (b, c)

Immunostaining (d-j)

Ethacridine   1 hr

Extended data Fig. 6. Quantitative analysis of viral titer by plaque assay in the human cells A549 that stably express ACE2. SARS-CoV-2 was pre-incubated with ethacridine (5 µM) for 1hr, followed by plaque assay on the human A549 cells stably expressing ethacridine

Ethacridine   Ethacridine

Viral titre (pfu/ml)   ***

37 for 1hour

A549 cell

Plaque assay

RT-PCR

16 hrs

DMSO   ethacridine

capsid

Viral titres
Total viral RNA copies

P<0.05

Normalized Intensity   ***

1.5
1.0
0.5
0.0

SPIKE

DMSO   ethacridine

Normalized Intensity   ***

1.5
1.0
0.5
0.0

DMSO   ethacridine

P value   0.0006

N=3
P value   <0.0001

RT for 2hour

number   **

10^7
10^6
10^5
10^4

25