A genome-wide CRISPR/Cas9 knock-out screen identifies the DEAD

box RNA helicase DDX42 as a broad antiviral inhibitor

Boris Bonaventure[1], Antoine Rebendenne[1]*, Francisco Garcia de Gracia[1]*, Joe McKellar[1]*,

Ségolène Gracias[2], Emmanuel Labaronne[3], Marine Tauziet[1], Ana Luiza Chaves Valadão[1], Eric

Bernard[1], Laurence Briant[1], Nathalie Gros[4], Wassila Djilli[1], Mary Arnaud-Arnould[1], Valérie

Courgnaud[5], Hugues Parrinello[6], Stéphanie Rialle[6], Reiner Schulz[7], Emiliano Ricci[3], Nolwenn

Jouvenet[2], Olivier Moncorgé[1] and Caroline Goujon[1][#]

1 IRIM, CNRS, Université de Montpellier, France

2 Institut Pasteur, Virus Sensing and Signaling Unit, Department of Virology, CNRS UMR 3569,

Paris, France

3 LBMC, Université de Lyon, ENS de Lyon, CNRS, INSERM, Lyon, France

4 CEMIPAI, CNRS, Université de Montpellier, France

5 IGMM, CNRS, Université de Montpellier, France

6 Montpellier GenomiX (MGX), Biocampus, CNRS, INSERM, Université de Montpellier, France

7 Department of Medical & Molecular Genetics, King's College London, United Kingdom

* Contributed equally to the work

[#] Corresponding and lead author: Caroline Goujon, IRIM UMR9004, 1919 route de Mende, 34293

Montpellier cedex 5. Phone: +33 4 34 35 94 33, e-mail: caroline.goujon@irim.cnrs.fr, ORCID iD:

0000-0001-8571-1108

Abstract

Genome-wide CRISPR/Cas9 knock-out genetic screens are powerful approaches to unravel new regulators of viral infections. Here, we took advantage of the ability of interferon (IFN) to restrict HIV-1 infection, in order to create an environment hostile to replication and reveal new inhibitors through a CRISPR screen. This approach led to the identification of the RNA helicase DDX42 as an intrinsic inhibitor of HIV-1. Depletion of endogenous DDX42 increased HIV-1 DNA accumulation and infection in several human cell lines and primary cells, irrespectively of IFN treatment. DDX42 overexpression inhibited HIV-1, whereas a dominant-negative mutant of DDX42 increased infection. Importantly, DDX42 restricted retrotransposition of long interspersed elements-1 (LINE-1), infection with other retroviruses and positive-strand RNA viruses, including Chikungunya virus (CHIKV) and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). However, DDX42 did not inhibit infection with negative-strand RNA viruses such as influenza A virus (IAV), arguing against a general, unspecific effect on target cells. In line with this, RNA-seq analysis did not reveal changes upon DDX42 depletion that could explain the observed phenotypes. Proximity ligation assays showed that DDX42 was found in the vicinity of viral elements during infection, and RNA immunoprecipitation confirmed DDX42 interaction with RNAs from LINE-1, CHIKV and SARS-CoV-2, but not IAV. This strongly suggested a direct mode of action of DDX42 on viral ribonucleoprotein complexes. Taken together, our results identify DDX42 as a new, broadly active intrinsic antiviral inhibitor.

Introduction

The intrinsic and innate immunity are at the frontline against viral invasion and provide a rapid and global defence. The innate immunity relies on viral sensing by Pathogen Recognition Receptors (PRRs) inducing the production of type 1 and 3 interferons (IFNs). Secreted IFNs bind to specific receptors and activate the JAK-STAT signalling cascade, which leads to the expression of hundreds of IFN-stimulated genes (ISGs). The cellular reprogramming induced by ISG expression allows the establishment of an antiviral state that efficiently limits viral replication. Some ISGs are indeed direct antiviral effectors harbouring powerful antiviral activities[1]. In addition to the IFN response, antiviral proteins that are constitutively expressed are able to immediately counteract incoming virus replication and are referred to as intrinsic inhibitors; they are part of the so-called intrinsic immunity.

Intensive efforts have been made over the past decades to identify genes able to limit viral replication. Several ISGs were identified a long time ago as major players of innate immunity against viruses, such as the myxovirus resistance protein 1 (MX1) Dynamin Like GTPase, 2′,5′-oligoadenylate synthetases (OASs) and ribonuclease L (RNASeL), or protein kinase R (PKR)[2–4]. More recently, gain-of-function and loss-of-function screens have identified new IFN-induced antiviral effectors[5–7]. A growing list of cellular proteins with various functions has hence been identified as capable of limiting different steps of virus life cycles[8–10]. Viruses have often evolved to counteract the action of these so-called restriction factors. However, type 1 IFNs (e.g. IFN-alpha and -beta) induce, through the expression of ISGs, an antiviral state particularly efficient at inhibiting HIV-1 when cells are pre-exposed to IFN[9]. The dynamin-like GTPase MX2, and, recently, the restriction factor TRIM5α, have both been shown to participate in this IFN-induced inhibition[6,11–13]. While numerous antiviral ISGs have been identified, less is probably known about the extent of the intrinsic, antiviral inhibitor repertoire. The recent identification of TRIM7 as an enterovirus inhibitor illustrates the fact that important intrinsic, antiviral inhibitors most certainly remain to be revealed[14].

With the hypothesis that additional HIV-1 inhibitors remained to be identified, we took advantage of the hostile environment induced by IFN to develop a whole-genome, CRISPR/Cas9 screen

strategy in order to reveal intrinsic and innate inhibitors. This strategy led us to identify DDX42 as

a new intrinsic inhibitor of HIV-1. We reveal that endogenous DDX42 is antiviral in various cell

types, including primary targets of HIV-1, and impairs the accumulation of viral DNA. Moreover,

our data show a broad activity against lentiviruses and the retrovirus Murine Leukemia Virus

(MLV). Reminiscent of other HIV-1 inhibitors such as APOBEC3G, DDX42 also blocks LINE-1

spread by interacting with their RNAs. Interestingly, while three different negative strand RNA

viruses were found insensitive to DDX42, several positive strand RNA viruses, including the

flavivirus Zika (ZIKV), the alphavirus Chikungunya (CHIKV) and the severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2), were inhibited by this non-processive RNA helicase.

Finally, RNA immunoprecipitation assays showed that DDX42 specifically binds to viral RNAs

from sensitive viruses, suggesting a direct mode of action. Overall, our study sheds light on a new

intrinsic, antiviral function of a so far poorly studied DEAD-box RNA helicase, and provide new

insights on a broad-spectrum antiviral inhibitor.

Results

The Genome-Scale CRISPR Knock-Out (GeCKO) sgRNA library[15–17] was used to generate

knock-out (KO) populations in the glioblastoma T98G cell line. This cell line is both highly

permissive to HIV-1 infection and potently able to suppress infection following type 1 IFN

treatment (Fig. S1A). The screen strategy is depicted in Figure 1A. Cas9-expressing T98G cells

were independently transduced with lentiviral vectors (LVs) coding the two-halves of the GeCKO

library, at a low multiplicity of infection (MOI). Next-generation sequencing showed more than

94% sgRNA coverage for each sub-library (not shown). Cells were pre-treated with type 1 IFN

(IFN-alpha) and incubated with VSV-G-pseudotyped, HIV-1 based LVs coding for an antibiotic

resistance cassette. The cells which were successfully infected despite the IFN treatment were

selected by survival in the presence of antibiotics. In order to enrich the populations with mutants

of interest and to limit the presence of false-positives, two additional rounds of IFN treatment,

infection and selection (with different antibiotics) were performed. As expected, the cells enriched

after each round became less refractory to HIV-1 infection following IFN treatment (Fig. S1B).

Figure 1. A whole-genome CRISPR/Cas9 screen to identify new HIV-1 inhibitors.

A. Screen strategy. GeCKO cell populations (obtained by transduction of T98G/Cas9 cells with GeCKO v2 LV library) were IFN-treated, challenged with HIV-1 LVs coding for an antibiotic resistance gene and selected. Three rounds of IFN treatment, infection and selection were performed. Genomic DNAs of initial GeCKO and 3-time selected populations were extracted, the sgRNA-coding sequences amplified and sequenced.

B. Candidate gene identification. MAGeCK computational statistical tool[18] was used to establish a Robust Rank Aggregation (RRA) score for each gene based on sgRNA enrichment and number of sgRNAs per gene. Genes belonging to type 1 IFN response pathway (in blue) and DDX42 (in red) are shown (respective ranks into brackets) for 2 independent screens (the results of which were merged in the analysis).

C. Candidate validation. T98G/Cas9/CD4/CXCR4/Firefly KO populations were generated for the 25 top hits of each screen. The control (CTRL) condition represents the mean of 4 negative CTRL populations, obtained with 4 non-targeting sgRNAs; *IFNAR1* and *MX2* KO populations were used as positive controls. KO cell populations were treated with IFN and infected with HIV-1 Renilla and luciferase signals were measured 24 h later (Renilla signals were normalized to Firefly). IFN inhibition (i.e. ratio of untreated / IFN-treated conditions) was calculated and set at 100% inhibition for CTRL. A representative experiment is shown (mean and standard deviation from technical duplicates).

The differential sgRNA abundance between the initial GeCKO populations and selected populations was analysed by next-generation sequencing (NGS) and the MAGeCK algorithm was used to rank the gene candidates (Fig. 1B). An enrichment was observed for 200 genes (RRA score > 0,01), with the best hits being *IFNAR1*, *JAK1* and *STAT2* (Fig. 1B). The crucial mediators of type 1 IFN signalling cascade were among the top hits in both screens (with the notable

exception of *STAT1*), validating our approach and confirming the identification of relevant genes.

Interestingly, most of the other candidates displayed unknown functions, or functions that were *a*

priori unrelated to innate immunity. Of note, very little overlap was observed between the two

independent screens, performed with two sub-libraries. However, such a poor overlap between

biological replicates has been observed before and does not preclude obtaining valid data[19]. The

top 25 candidate genes from each screen were selected for further validation. T98G/Cas9 cells

expressing HIV-1 CD4 and CXCR4 receptors, as well as Firefly luciferase as an internal control

(T98G/Cas9/CD4/CXCR4/Firefly cells), were transduced with sgRNA-expressing LVs to generate

individual KO populations, using the identified sgRNA sequences. Four irrelevant, non-targeting

sgRNAs, as well as sgRNAs targeting *IFNAR1* and *MX2*, were used to generate negative and

positive control populations, respectively. The KO cell populations were pre-treated with IFN and

infected with an HIV-1 reporter virus expressing Renilla luciferase and bearing HIV-1 envelope[20]

(hereafter called HIV-1 Renilla). Infection efficiency was analysed 30 h later (Fig. 1C). As

expected[11,21,22], *IFNAR1* and *MX2* KO fully and partially rescued HIV-1 infection from the

protective effect of IFN, respectively. The KO of two candidate genes, namely *WARS2* and

DDX42, allowed a partial rescue of HIV-1 infection from the IFN-induced inhibition, suggesting a

potential role of these candidate genes in HIV-1 inhibition.

DDX42 is a member of the DEAD box family of RNA helicases, with RNA chaperone activities[23]

and, as such, retained our attention. Indeed, DEAD box helicases are well-known to regulate HIV-

1 life cycle[24]. However, to our knowledge, the impact of DDX42 on HIV-1 replication had never

been studied. In order to validate the effect of *DDX42* KO on HIV-1 infection in another model cell

line, two additional sgRNAs were designed (sgRNA-2 and -3) and used in parallel to the one

identified in the GeCKO screen (sgDDX42-1) (Fig. 2A). U87-MG/CD4/CXCR4 cells were used

here, as we previously extensively characterized the IFN phenotype in these cells[11]. Control and

DDX42 KO cell populations were generated. Control and DDX42 KO cell populations were pre-

treated or not with IFN prior to infection with HIV-1 Renilla. Of note, CRISPR/Cas9 KO of *DDX42*

induced only a partial decrease of DDX42 protein levels (Fig. 2A) and cell populations derived

quite rapidly (not shown), suggesting an essential role for DDX42 in cell survival or proliferation.

We observed however that DDX42 partial depletion with all 3 sgRNAs improved HIV-1 infection,

confirming that endogenous DDX42 had a negative impact on HIV-1 replication. Interestingly, the

increase in infection efficiency induced by *DDX42* KO was observed independently of the IFN

treatment. DDX42 is not an ISG, as shown in several cell lines (e.g. U87-MG, T98G, HEK293T)

and in primary T cells and monocyte-derived macrophages (Fig. S2A and GSE46599[11]). The fact

that the IFN-induced state is at least partially saturable (Fig. S1A) explains why an intrinsic

inhibitor of HIV-1, which is not regulated by IFN, could be identified by our approach. Indeed,

removing one barrier to infection presumably rendered the cells more permissive and, in this

context, IFN had less of an impact on viral replication.

Figure 2. DDX42 is an intrinsic inhibitor of HIV-1.

A. Top: *DDX42* KO and CTRL KO U87-MG/CD4/CXCR4/Cas9/Firefly cells were generated using 3 sgRNAs and

4 non-targeting sgRNAs, respectively (for CTRL, the average of data obtained with 4 cell populations is shown).

Cells were treated or not with IFN 24 h prior to infection with HIV-1 Renilla. Relative luminescence results for

IFN-treated and -untreated conditions are shown. Two-way ANOVA on log-transformed data with Sidak's test.

Bottom: Immunoblot analysis of DDX42 levels is shown for 1 CTRL and DDX42-depleted populations; Actin

served as a loading control.

B. DDX42 silencing efficiency measured by RT-qPCR (top) and immunoblot (bottom) in parallel samples from C.

C. siRNA-transfected U87-MG/CD4/CXCR4 cells were treated or not with IFN for 24 h prior to infection with HIV-

1 Renilla. Relative luminescence results for IFN-treated and -untreated conditions are shown. Multiple linear

regression analysis.

D. DDX42-depleted cells were infected with HIV-1, and infection efficiency was measured by CA p24^{Gag}

intracellular staining and flow cytometry analysis. When indicated, cells were treated with azidothymidine (AZT)

and lamivudine (3TC). Two-way ANOVA on log-transformed data with Dunnett's test.

E. siRNA-transfected MDMs were infected with a CCR5-tropic version of HIV-1 Renilla. Relative luminescence

results from independent experiments performed with cells from 3 donors are shown. Two-way ANOVA on log-

transformed data with Dunnett's test.

F. Primary CD4+ T cells were electroporated with Cas9-sgRNA RNPs using 2 non-targeting sgRNAs (sgCTRL1

and 2) and 5 sgRNAs targeting DDX42. Top: Cells were then infected with HIV-1 Renilla and relative infection

efficiencies obtained with cells from three donors are shown. Two-way ANOVA on log-transformed data with

Dunnett's test. Bottom: DDX42 protein levels were determined by immunoblot, Actin served as a loading control.

A representative immunoblot is shown.

G. Firefly- or DDX42-expresssing U87-MG/CD4/CXCR4 cells were infected with HIV-1 Renilla. Relative infection

efficiencies are shown. Multiple linear regression analysis.

A-G Data represent the mean ± S.E.M of three independent experiments.

In order to confirm DDX42's effect on HIV-1 infection with an independent approach, we used

three different siRNAs to knockdown DDX42 expression. We observed that depleting DDX42

(with ~90% efficiency both at the mRNA and protein levels, Fig. 2B) increased HIV-1 Renilla

infection efficiency by 3 to 8-fold in U87-MG/CD4/CXCR4 cells, irrespectively of the presence of

IFN (Fig. 2C). Of note, wild-type HIV-1 infection was also significantly impacted by DDX42

silencing, as measured by Capsid (CA p24^{Gag}) intracellular staining 30 h post-infection in U87-

MG/CD4/CXCR4 cells (Fig. 2D). We then investigated whether DDX42 had an impact in HIV-1

primary target cells. In monocyte-derived macrophages (MDMs), we observed that HIV-1 infection

was increased by about 2-fold following DDX42 silencing (Fig. 2E), whereas DDX42 mRNA

abundance was decreased by only 40% (Fig. S2B). Electroporation of pre-assembled Cas9-

sgRNA ribonucleoprotein complexes (RNPs) was used to efficiently deplete DDX42 in primary

CD4+ T cells (Fig. 2F). DDX42 depletion increased HIV-1 infection by 2- to 3-fold, showing a role

of DDX42 as an intrinsic inhibitor of HIV-1 in primary CD4+ T cells. Next, we analysed the

consequences of DDX42 overexpression on HIV-1 infection. An irrelevant control (Firefly) or

DDX42 were ectopically expressed in U87-MG/CD4/CXCR4 and the cells were challenged with

HIV-1 (Fig. 2G). DDX42 overexpression induced a substantial inhibition of HIV-1 infection (~5-

fold decrease in infection efficiency in comparison to the control). Interestingly, the expression of

K303E DDX42 mutant, which is unable to hydrolyse ATP and may supposedly act as a dominant

negative[25,26], increased HIV-1 infection by 3-fold, reminiscent of the impact of DDX42 depletion

(Fig. S2C). Altogether, these data showed that endogenous DDX42 is able to intrinsically inhibit

HIV-1 infection.

In order to determine which step of HIV-1 life cycle was affected by DDX42, we quantified HIV-1

DNA accumulation over time in DDX42-silenced and control cells (Fig. 3A; silencing efficiency is

shown in Fig. 3B). DDX42 depletion increased accumulation of early and late reverse transcript

products (by 2.5- to 8-fold), as well as proviral DNA and 2-long terminal repeat (2-LTR) circles at

48h post-infection (by 2.5- to 4.5-fold). Importantly, DDX42 silencing did not impact HIV-1 entry

(Fig. S3A). These data suggested that endogenous DDX42 could inhibit reverse transcription

and/or impact genome stability, leading to a decrease in viral DNA accumulation. We

hypothesized that if that was the case, DDX42 should be found in close proximity to HIV-1 reverse

transcription complexes during infection. In agreement with this, proximity ligation assay (PLA)

performed on HIV-1 infected MDMs showed that DDX42 was indeed in close vicinity of Capsid

(Fig. 3C).

We next examined the ability of DDX42 to inhibit infection by various primate lentiviruses,

including lab-adapted strains of HIV-1 (NL4-3, IIIB), transmitted/founder strains[27] (CH077.t,

CH106.c, REJO.c), HIV-2 and simian immunodeficiency virus from rhesus macaque (SIV_{MAC}).

DDX42 was depleted or not in TZM-bl reporter cells prior to infection with VSV-G-pseudotyped

lentiviruses, and infection efficiency was monitored 24h later (Fig. 3D; silencing efficiency is

shown in Fig. 3E). DDX42 depletion increased infection levels similarly with all the tested strains

of HIV-1 (i.e. 3- to 5-fold). HIV-2rod10 and SIV_{MAC} infection efficiencies were also slightly improved

in the absence of DDX42 (~2-fold). The analysis was then extended to two non-primate

lentiviruses, equine infectious anemia virus (EIAV) and feline immunodeficiency virus (FIV), using

GFP-coding LVs in U87-MG cells (Fig. 3F). DDX42 depletion appeared to increase HIV-1, HIV-2

and FIV LV infection to the same extent (~2-fold), whereas EIAV infection was less impacted. Of

note, DDX42 antiviral activity appeared less potent on HIV-1 LVs compared to full-length HIV-1,

which might suggest that genome length or cis-acting elements could play a role in DDX42

inhibition. We also observed that DDX42 depletion led to a significant increase in infection with

GFP-coding, MLV-derived vectors (Fig. 3G). These results strongly support a general antiviral

activity of DDX42 against retroviruses.

DDX42 can be found in the cytoplasm but is predominantly located in the nucleus in various cell

types, including monocyte-derived macrophages[28,29] (Fig. S3B). Considering that DDX42 showed

a broad activity against retroviruses and seemed to act at the level of reverse transcription, we

sought to investigate whether DDX42 could inhibit retrotransposons. Long interspersed nuclear

elements (LINE)-1 are non-LTR retrotransposons, which have been found to be active in the germ

line and some somatic cells[30]. Interestingly, DDX42 was identified among the suppressors of

LINE-1 retrotransposition through a genome-wide screen in K562 cells, although not further

characterized[31]. To confirm that DDX42 could inhibit LINE-1 retrotransposition, HEK293T cells

were co-transfected with GFP-expressing LINE-1 plasmids (RPS or LRE3) or an inactive LINE-1

(JM111) together with a DDX42- or a control (Firefly)-expressing plasmid[32]. LINE-1

retrotransposition was quantified by flow cytometry 7 days later (Fig. 3H). As the GFP cassette is

cloned in antisense and disrupted by an intron, GFP is only expressed after LINE-1 transcription,

splicing, Orf2p-mediated reverse transcription, and integration[32]. Successful retrotransposition

events were observed in >1.25% of control cells, but in only <0.25% of DDX42-expressing cells

(i.e. a percentage similar to what observed with the non-active LINE-1), showing that DDX42

ectopic expression significantly suppressed LINE-1 retrotransposition. Next, we investigated

whether DDX42 could physically interact with LINE-1 RNAs. Cells were co-transfected with GFP-

expressing LINE-1 RPS plasmid and either flag-tagged-Firefly or -DDX42. The cells were lysed 4

days later and the flagged proteins immunoprecipitated. The immunoprecipitation eluates were

then divided in two; the immunoprecipitated proteins were analysed by immunoblot (Fig. 3I) and their associated RNAs were extracted and analysed by RT-qPCR using LINE-1 specific primers (Fig. 3J). A significant enrichment of LINE-1 RNAs was observed with DDX42 immunoprecipitation as compared to the Firefly negative control, showing that DDX42 could interact with LINE-1 RNAs.

Figure 3. Characterization of DDX42 inhibitory activity against retroviruses and retroelements.

A. siRNA-transfected U87-MG/CD4/CXCR4 cells were infected with HIV-1 and relative amounts of Minus Strand Strong Stop (MSSS), 1st and 2nd Strand Transfer DNAs, and nuclear forms of HIV-1 DNA (proviral DNA, and 2-LTR circles) were quantified by qPCR. DNAs from cells infected for 48 h in the presence of AZT and 3TC were used as a control. Mixed-effects analysis on log-transformed data with Dunnett's test.

B. Silencing efficiency in parallel samples from A.

C. PLAs were performed in MDMs infected with HIV-1 or not (N.I. CTRL), using anti-Capsid and anti-DDX42 antibodies (nuclei stained with Hoechst). Images were acquired using a LSM880 Airyscan microscope. Left: representative images, scale-bar: 10 μm. Right: Average punctae quantified per cell in 3 independent assays done on MDMs from different donors with mean ± SD (n>65 cells per condition). Mann-Whitney test.

D. siRNA-transfected TZM-bl cells were infected and β-galactosidase signals measured 24 h later. The ratio of the signal in DDX42-depleted versus CTRL cells is shown.

E. Silencing efficiency in parallel samples from D.

F. siRNA-transfected U87-MG/CD4/CXCR4 cells were infected with HIV-1- HIV-2- FIV- EIAV-based, GFP-coding LVs and infection efficiency was scored 24 h later by measuring the percentage of GFP expressing cells by flow cytometry.

G. siRNA-transfected U87-MG/CD4/CXCR4 cells were infected with GFP-coding B-MLV and infection efficiency measured 24h later by flow cytometry. Simple linear regression analysis.

H. HEK293T were co-transfected with GFP-coding LINE-1 plasmids (RPS-GFP or LRE3-GFP) or with an inactive LINE-1 plasmid (JM111) together with either a Firefly- or DDX42-coding plasmid. GFP expression was measured by flow cytometry 7 days later. Two-way ANOVA on log-transformed data with Sidak's test.

I. HEK293T were co-transfected with pRPS-GFP and a Flag-Firefly- (negative control) or Flag-DDX42-coding plasmid, followed by Flag immunoprecipitation and immunoblot analysis. A representative immunoblot is shown.

J. Left, RNA extraction and LINE-1 RT-qPCR on parallel samples from H. Two-way ANOVA on log-transformed data with Sidak's test. Right, Percentage of immunoprecipitated RNA from I. T-test on log-transformed data.

A-H. Data represent the mean ± S.E.M of 3 independent experiments. **J**. Data represent the mean ± S.E.M of 5 independent experiments.

Finally, we sought to determine whether DDX42's inhibitory activity was specific towards retroviruses and retroelements, or could be extended to other viruses, as observed for many other anti-HIV-1 proteins, such as ZAP or BST-2/Tetherin[1,8]. To this aim, we tested the impact of DDX42 depletion on eight RNA viruses from five different families: the orthomyxovirus influenza A virus (IAV), the rhabdovirus vesicular stomatitis virus (VSV), the paramyxovirus measles virus (MeV), the flaviviruses ZIKV, Dengue virus serotype 2 (DENV-2) and yellow fever virus (YFV), the alphavirus CHIKV, and the coronavirus SARS-CoV-2, which is responsible for the current coronavirus disease (COVID)-19 pandemic (Fig. 4). Strikingly, DDX42 depletion had no significant effect on IAV, VSV and MeV replication in U87-MG and Huh-7 cells, respectively (Fig. 4A-C; silencing efficiency is shown in Fig. S4A), thereby strongly suggesting that manipulating DDX42 expression did not have a broad and unspecific impact on target cells. Interestingly, depletion of endogenous DDX42 had a modest but significant, positive impact on ZIKV in U87-

MG cells (Fig. 4D), whereas the impact in Huh-7 cells was less important (Fig. S4B and A).

Similarly, DDX42 silencing had little effect on 2 other flaviviruses (DENV-2 and YFV) in Huh-7

cells, as measured by the number of cells positive for the viral protein E (Fig. S4C-D). By contrast,

DDX42 depletion had a profound effect on both CHIKV (Fig. 4E) and RC-2020-00558 SARS-

CoV-2 replication (Fig. 4F-G) (up to 1 log- and 3 log-increase in infection efficiency with CHIKV

and SARS-CoV-2, respectively; silencing efficiencies in the different cell lines used are shown in

Fig. S4A). Plaque assays confirmed a strong impact of DDX42 depletion on infectious SARS-

CoV-2 production (Fig. 4G). In agreement with this, DDX42 was recently identified as a potential

inhibitor of SARS-CoV-2 replication in a CRISPR screen in simian cells[33]. Next, we used PLA to

determine whether DDX42 was in the vicinity of SARS-CoV-2 components in infected cells. To

this aim, PLA was performed with either anti-double strand (ds)RNA or anti-Nucleoprotein (N)

antibody, together with anti-DDX42 antibody, followed by an immunofluorescence staining to

identify the infected cells (Fig. 4H-I). In the latter, there was a significantly higher number of

dsRNA-DDX42 and N-DDX42 PLA punctae than in control cells. This suggested a potential

interaction between DDX42 and SARS-CoV-2 viral components. To test whether DDX42 RNA

helicase could interact with viral RNAs, RNA immunoprecipitation experiments were conducted

following viral infection of U87-MG and A549-ACE2 cells expressing either Flag-DDX42 or

negative control Flag-Firefly. A significant enrichment of the viral RNAs recovered with Flag-

DDX42 was observed in comparison to the negative control, and this was observed for the

sensitive viruses SARS-CoV-2 and CHIKV (Fig. 4J-K), but not for IAV, which was insensitive to

DDX42 antiviral activity (Fig. 4L and 4A).

Figure 4. DDX42 exerts a broad antiviral activity on positive strand viruses and interacts with viral RNAs from targeted viruses.

A. Relative IAV infection efficiency in siRNA-transfected U87-MG cells (Nanoluciferase activity 16 h post infection).

B. Relative VSV infection efficiency in siRNA-transfected U87-MG cells (Firefly activity 24 h post infection).

C. Relative MeV infection efficiency in siRNA-transfected Huh-7 cells (GFP+ cells scored 24h post infection). Multiple linear regression analysis.

D. Relative ZIKV infection efficiency in siRNA-transfected U87-MG cells (Nanoluciferase activity 24 h post infection). Multiple linear regression analysis.

E. Relative CHIKV infection efficiency in siRNA-transfected U87-MG cells (Nanoluciferase activity 24 h post infection). Multiple linear regression analysis.

F. Relative SARS-CoV-2 infection in siRNA-transfected A549-ACE2 cells (RdRp RT-qPCR 48 h post-infection). Mixed-effects analysis on log-transformed data with Dunnett's test.

G. Viral production in cell supernatants from F (48 h post-infection, MOI 0,05) measured by plaque assays.

H. A549-ACE2 cells were infected or not with SARS-CoV-2 for 24 h prior to PLA using mouse anti-dsRNA (J2) and rabbit anti-DDX42 antibodies, followed by additional immunofluorescence (IF) staining with anti-mouse Alexa Fluor 546 antibody (PLA in green, IF in magenta). Representative Z-stack projection images are shown; scale bar: 15 μm. Average punctae were quantified in 3 independent PLA assays with mean ± SD (n>75 cells per condition). Mann-Whitney test.

I. Identical to H but using an anti-N antibody instead of anti-dsRNA antibody.

J. Left, Quantification of SARS-CoV-2 RNA by RT-qPCR in RNA from total cell lysates (input) and in Flag-Firefly (negative control) and Flag-DDX42 immunoprecipitation (IP). Multiple linear regression analysis on log-transformed data.

Right, Percentage of immunoprecipitated RNA. Paired t-test on log-transformed data.

K. Identical to J following CHIKV infection.

L. Identical to J following IAV infection.

Data represent the mean ± S.E.M. of three (**A-E, J-L**), four (**F**) or two (**G**) independent experiments.

As mentioned above, DDX42's lack of effect of negative strand RNA viruses argued against a global, indirect effect on the target cells. However, to confirm this, we performed RNA-seq analysis on siRNA-treated U87-MG and A549-ACE2 cells. The results showed that DDX42 depletion didn't have a substantial impact on global cellular RNA expression (Supplementary File 1 and Fig. S5). Of note, only 63 genes were commonly found differentially expressed upon DDX42 depletion with the 3 different siRNAs in U87-MG cells, and only 23 genes were identified in common in U87-MG and A549-ACE2 cells (differentially expressed genes, DEGs, with a fold increased ≥ 2 and an adjusted p-value < 0.05). Importantly, no known restriction factors were identified among the DEGs (Supplementary File 1). Taken together, these data strongly suggested that the DEAD-box RNA helicase DDX42 directly impacted viral replication, by interacting with viral RNAs.

Discussion

Here, we identified for the first time the RNA helicase DDX42 as an intrinsic inhibitor of HIV-1, capable of limiting the efficiency of viral DNA accumulation. Moreover, our study revealed broad activity of endogenous DDX42 against retroviruses and retroelements, which was observed in various cell types, including primary CD4+ T cells. Strikingly, we observed that DDX42 was able to inhibit viruses from other families, which possess different replication strategies, including SARS-CoV-2 and CHIKV. However, DDX42 did not have an impact on all the viruses we tested, as three different negative-strand RNA viruses were found insensitive to DDX42 antiviral activity. This is reminiscent of other broad-spectrum antiviral inhibitors such as MX1, which show some specificity despite being able to inhibit viruses from various families[34]. Further work is now warranted to explore in depth the breadth of DDX42 antiviral activity and determine whether it is truly specific of positive strand RNA viruses.

Interestingly, our PLA assays showed a close proximity between DDX42 and HIV-1 Capsid, which is a viral protein recently shown to remain associated with reverse transcription complexes until proviral DNA integration in the nucleus[35–37]. We also observed a close proximity between DDX42 and SARS-CoV-2 N or dsRNA. Furthermore, LINE-1 RNAs, as well as SARS-CoV-2 and CHIKV RNAs, were specifically pulled-down when DDX42 was immunoprecipitated. Taken together, these observations strongly suggest a direct mode of action of DDX42, which could interact with target viral RNAs. DDX42 is known to be a non-processive helicase, which also possesses RNA annealing activities and the ability to displace RNA-binding proteins from single-stranded RNAs[23]. Moreover, DDX42 binds G-quadruplexes[38], which are secondary structures found in cellular and viral nucleic acids and involved in various processes, such as transcription, translation and replication[39,40]. All these known activities of DDX42 would be consistent with a potential role in RNP remodeling[23,41]. Nonetheless, further investigation will be needed to determine whether DDX42 acts directly by altering viral RNPs, and, if that's the case, what are the determinants for viral RNP recognition.

In conclusion, this work highlights the importance of understanding the mechanism of action of

DDX42 RNA helicase and its contribution to the control of RNA virus replication, an understanding

which may contribute to the development of future antiviral interventional strategies.

Methods

Plasmids. The pLentiCas9-Blast, pLentiGuide-Puro vectors and the GeCKO sub-library A and B

plasmids were a gift from Prof. F. Zhang (Addgene #52962, #52963, and #1000000048,

respectively[17]). LVs coding for sgRNAs targeting the candidate genes and control genes were

obtained by cloning annealed oligonucleotides in BsmBI-digested pLentiGuide-Puro, as

described (Addgene). Control sgRNAs and sgRNAs targeting the candidate genes, *MX2* and

IFNAR1, were designed with the Optimized CRISPR Design tool (not available anymore), or with

Chopchop (chopchop.cbu.uib.no). The sgRNA coding sequences used were as follow: MX2 5'-

CCGCCATTCGGCACAGTGCC-3', IFNAR1 5'-GACCCTAGTGCTCGTCGCCG-3', sgCTRL-1 5'-

AGCACGTAATGTCCGTGGAT-3', sgCTRL-2 5'-CAATCGGCGACGTTTTAAAT-3', sgCTRL-3

5'-TTAATTTGGGTGGGCCCTGC-3', sgCTRL-4 5'-TTGGATATTAATTAGACATG-3', sgDDX42-

1 5'-TCCTGAACCACACCAGCAGT-3', sgDDX42-2 5'-GGTGGTCCTGGCACTAAGCG-3',

sgDDX42-3 5'-AGGCACTGTGGGACTGCTGT-3'. All the other sgRNA sequences are available

upon request. In order to produce the HIV-1 based LVs used to perform the different steps of the

screen (pRRL.sin.cPPT.CMV/NeomycinR.WPRE, pRRL.sin.cPPT.CMV/HygromycinR.WPRE

and pRRL.sin.cPPT.CMV/ZeocinR.WPRE), neomycin, hygromycin and zeocin resistance genes

(i.e. the genes coding for Neomycin phosphotransferase II, Hygromycin B phosphotransferase,

and *Sh ble*) were amplified by PCR from pcDNA3.1 (ThermoFisher Scientific), pAHM[11], and

pcDNA3.1/Zeo (ThermoFisher Scientific), respectively, and cloned by replacement of GFP in

pRRL.sin.cPPT.CMV/GFP.WPRE[42] using BamHI and SalI restriction sites. The

pRRL.sin.cPPT.SFFV/E2-crimson-IRES-PuromycinR.WPRE has been described[43]. Human

DDX42 cDNA was amplified by RT-PCR using the SuperScript III™ (Invitrogen) from mRNAs of

MDMs using primers DDX42-forward 5'-

AATTAATTTAGGATCCATGAACTGGAATAAAGGTGGTCCTG and DDX42-reverse 5'-

AATTAATTTACTCGAGCTAACTGTCCCATCGACTTTTCTTGCG, and cloned by replacement of

E2-crimson in BamHI-XhoI-digested pRRL.sin.cPPT.SFFV/E2crimson-IRES-

PuromycinR.WPRE, in order to obtain pRRL.sin.cPPT.SFFV/DDX42-IRES-PuromycinR.WPRE.

The pRRL.sin.cPPT.SFFV/CD4-IRES-CXCR4.WPRE was obtained by replacement of E2-

crimson-IRES-PuroR in pRRL.sin.cPPT.SFFV/E2-crimson-IRES-PuromycinR.WPRE with a

BamHI/SalI fragment digested CD4-IRES-CXCR4 PCR fragment obtained from pMLV-CD4-

IRES-CXCR4 (a gift from Prof. N. Sherer, Wisconsin University, USA).

pRRL.sin.cPPT.SFFV/Firefly-IRES-PuromycinR.WPRE was obtained by amplification of Firefly

by PCR from pGL4 (Promega) and cloned into BamHI-XhoI-digested pRRL.sin.cPPT.SFFV/E2-

crimson-IRES-PuromycinR.WPRE. In some experiments, LVs without a selection marker were

used: the IRES-PuromycinR cassette was removed by XhoI-SalI digestion and subsequent

ligation, to obtain pRRL.sin.cPPT.SFFV/Firefly.WPRE and pRRL.sin.cPPT.SFFV/DDX42.WPRE.

DDX42 K303E mutant was obtained by site-directed mutagenesis (by overlapping PCR using the

aforementioned DDX42-forward and -reverse primers, respectively combined initially with reverse

primer 5'-GGCTGCAGTTTCCCCACTACCTGTTTTGGCAATACC and forward primer 5'-

GGTAGTGGGGAAACTGCAGCCTTCATTTGGCC). pRRL.sin.cPPT.SFFV/ACE2.WPRE has

been described[44] (Addgene 145842). Flag-DDX42 and Flag-Firefly were amplified by PCR from

the aforementioned LV plasmids and cloned into a NotI-XhoI-digested modified version of

pCAGGS[45] to obtain pCAGGS/flag-DDX42.WPRE and pCAGGS/flag-Firefly.WPRE. The NL4-

3/Nef–internal ribosome entry signal (IRES)-Renilla (NL4-3/Nef-IRES-Renilla) and the CCR5-

version of this proviral clone were gifts from Prof. Sumit Chanda[20]. Wild-type and Ba-L Env

bearing HIV-1 NL4-3, IIIB and HIV-2 proviral clones have been described[46–48], as well as the

transmitted founder HIV-1 molecular clones CH077.t, CH106.c, REJO.c (gifts from Prof. B.

Hahn[27]) and HIV-2$_{ROD10}$ and SIV$_{MAC239}$[49,50]. GFP-coding HIV-1 based LV system (i.e. p8.91 HIV-1

Gag-Pol, pMD.G, and GFP-coding minigenome), and HIV-2, FIV, and EIAV-derived, GFP coding

LVs, as well as MLV-derived, GFP coding retroviral vectors have all been described[51,52,53,54]. The

LINE-1 plasmid 99 RPS-GFP PUR (pRPS-GFP), 99 RPS-GFP JM111 PUR (pJM111) and

pLRE3-GFP were developed by Prof. Kazazian's lab[32,55,56].

Cell lines. Human cell lines HEK293T, A549, U87-MG, TZM-bl were obtained from the ATCC

and the AIDS reagent program, respectively. T98G cells were a gift from Prof. G. Kochs (Freiburg

University, Germany), MDCK cells a gift from Prof. W. Barclay (Imperial College London, UK), Vero E6 cells (Merck) were a gift from Christine Chable (CEMIPAI, CNRS). Human hepatocellular carcinoma Huh-7 cells[57] were kindly given by Annette Martin (Institut Pasteur, Paris). These cell lines were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum and 1 % penicillin-streptomycin (Thermofisher). T98G/Cas9 and U87-MG/Cas9 were obtained by transduction of T98G and U87-MG, respectively, with HIV-1-based LVs expressing the spCas9-P2A-Blasticidin cassette (pLentiCas9-Blast[17]). U87-MG/CD4/CXCR4 have been described [11] and were further modified to express Cas9 and Firefly using pLentiCas9-Blast and pRRL.sin.cPPT.SFFV/Firefly.WPRE, respectively. T98G/Cas9/CD4/CXCR4/Firefly were obtained by successive transductions of T98G/Cas9 with pRRL.sin.cPPT.SFFV/CD4-IRES-CXCR4.WPRE at a high MOI, and pRRL.sin.cPPT.SFFV/Firefly.WPRE, at a low MOI, respectively. Cell surface staining with anti-CD4 and CXCR4 antibodies (Miltenyi Biotec) confirmed than more than 95% cells were positive for both markers. A549 cells stably expressing ACE2 were generated by transduction with RRL.sin.cPPT.SFFV.ACE2.WPRE containing-vector. For antibiotic selection, cells were treated with 10 μg/mL Blasticidin (InvivoGen), 1 mg/mL Zeocin (InvivoGen), 2 μg/mL Puromycin (Sigma-Aldrich), 250 μg/mL Hygromycin (Sigma-Aldrich), 1 mg/mL G418 (Sigma-Aldrich). When indicated, universal type 1 IFN (PBL Interferon Source) was added at 1000 U/mL for 16-24h prior to virus infection or RNA extraction, and AZT and 3TC (AIDS reagent program) at 10 μM for 2 h prior to infection.

Primary cells. Blood from healthy donors was obtained from the Etablissement Français du Sang, under agreement n°21PLER2019-0106. Peripheral blood mononuclear cells (PBMCs) were isolated by centrifugation through a Ficoll® Paque Plus cushion (Sigma-Aldrich). Primary human CD4+ T cells and monocytes were purified by positive selection using CD3 and CD14 MicroBeads, respectively (Miltenyi Biotec), as previously described[11]. Monocytes were incubated for 3 hours in serum-free Roswell Park Memorial Institute (RPMI) 1640 medium and further differentiated into macrophages by culture for 5-7 days in RPMI 1640 supplemented with 10%

fetal calf serum, 1% penicillin-streptomycin and 100 ng/ml granulocyte-macrophage colony-

stimulating factor (GM-CSF; Miltenyi). CD4+ T cells were cultured in RPMI supplemented with

10% fetal bovine serum and 1% penicillin-streptomycin, and stimulated for 48h with 10 µg/ml

phytohemagglutinin (PHA) (Fisher Scientific) and 50 U/mL interleukin-2 (IL-2, Miltenyi Biotec)

prior to electroporation.

Genome-scale CRISPR/Cas9 screens. The plasmids coding GeCKO sub-libraries A and B were

amplified and prepared according to the provided guidelines (Lentiviral Crispr Toolbox, Addgene).

60 million T98G/Cas9 cells were transduced with GeCKO LVs at a MOI of 0,1 to cover about 100-

times the half-library complexity. After 48h, the cells were selected with puromycin, amplified for

12-15 days. 45 million cells were harvested and frozen down at -80°C for subsequent genomic

DNA extraction, using the QIAamp DNA Blood Maxi Kit according to manufacturer's instructions

(Qiagen). In parallel, 60 million cells from the initial GeCKO populations were used for the screen.

The cells were treated with 1000 U/mL IFN for 24h, infected with LVs coding a hygromycin

resistance cassette. 48h later the cells selected with hygromycin and the surviving cells amplified.

Two other rounds of IFN treatment, LV infection and antibiotic selection were subsequently

performed with LVs coding a neomycin resistance cassette and a zeocin resistance cassette,

respectively. The three time-selected populations were amplified and 45 million cells were

harvested and stored at -80°C for subsequent genomic DNA extraction, as previously. After

genomic DNA extraction, the sgRNA coding sequences integrated in the genomic DNA from the

initial and 3-times selected populations were amplified by touch-down PCR and sequenced by

Illumina deep sequencing. To this aim, 120 µg of genomic DNA was amplified using DNA

Herculase II Fusion DNA polymerase (Agilent) in the presence of 2% DMSO, 1 mM of dNTPs;

and 400 nM of the following primers: Forward-primer1: 5'-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCTTGTGGAAAGGACGAAACACC-3' for

screen A or Forward-primer2: 5'-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGATCTTGTGGAAAGGACGAAACACC-3'

used for screen B, together with reverse primer: 5'-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAAAGGTCCATTAGCTGCAAAGATTCCTCT

C-3'). Briefly, after 5 minutes at 95°C, 14 cycles of pre-amplification were performed with a

hybridization temperature decreasing by 0.5°C per cycle (30 sec at 95°C, 30 sec at 60°C, 30 sec

at 72°C), followed by 30 cycles of amplification (30 sec at 95°C, 30 sec at 53°C, 30 sec at 72°C).

50 ng of each amplicon was dual indexed in a 5-cycle PCR reaction using the PCR module and

indexed primers from the Nextera kit (Illumina). Resulting libraries were purified on AMPure XP

magnetic beads (Beckman Coulter) using a 0,8X ratio and verified on Fragment Analyzer using

the HS NGS fragment kit (Agilent). Libraries were quantified using microfluorimetry (QuBit,

Invitrogen), mixed with a PhiX library (Illumina) and sequenced on one single read 50nt lane of

Hiseq2500 using the rapid mode.

Image analyses and base calling were performed using the Illumina HiSeq Control Software and

Real-Time Analysis component (v1.18.66.3). Demultiplexing was performed using Illumina's

conversion software (bcl2fastq 2.20). The quality of the raw data was assessed using FastQC

(v0.11.5) from the Babraham Institute and the Illumina software SAV (Sequencing Analysis

Viewer). Potential contaminants were investigated with the FastQ Screen[58] (v0.11.4) software

from the Babraham Institute.

Sequencing reads were trimmed using Cutadapt[59] (v1.13), with options -g [primer sequence] -u

[length of remaining 3' bases] -e 0.2 -m 18 -l 20, to remove primer sequences and retrieve the 20

bases long sequences corresponding to sgRNAs. These retrieved sequences were then aligned

to the GecKOv2 Human Library (A or B) reference sequences (keeping only non-duplicated

sgRNA sequences, the duplicated ones being annotated) using Bowtie[60] (v1.2), with options -v 2

-norc -S. Resulting bam files were sorted and indexed using Samtools[61] (v1.5). Quantification of

sgRNAs was done using Samtools idxstats. MAGeCK[18] (v0.5.7) was used to normalize (total

count method) and identify enriched sgRNAs and genes in 3-times selected cell populations

versus starting GeCKO transduced cells (mageck test command).

Lentiviral and retroviral production. To produce lentiviral vector particles, HEK293T cells were

transfected by polyethylenimine (PEI) co-transfection with miniviral, HIV-1 based genome coding

plasmids (e.g. LentiCas9-Blast, LentiGuide-Puro or pRRL-SFFV), p8.91 (HIV-1 GagPol) and

pMD.G (VSV-G) at a ratio of 1:1:0.5, respectively. The medium was replaced after 6h and viral

particles were harvested 42h later, filtered, and directly used to transduce target cells (or stored

at -80°C). After 4 to 6 hours, the transduction medium was replaced with complete DMEM, and

the cells were treated 48h later with the relevant antibiotics. The HIV-2-, FIV-, EIAV-GFP coding

LVs were produced using GFP-coding HIV-2-, FIV-, EIAV-based miniviral genomes, together with

HIV-2-, FIV-, EIAV- GagPol, expression constructs and pMD.G at a ratio of 1:1:0.5. MIGR1 MLV-

derived retroviral vectors were obtained with B-MLV Gag-Pol-expressing plasmid pCIG3B, the

GFP-expressing minigenome pMIGR1 and pMD.G. at a ratio of 1:1:0.5, respectively and

harvested as previously described.

HIV-1 Renilla and NL4-3 HIV-1 were produced by standard PEI transfection of HEK293T. When

indicated, pMD.G was cotransfected with the provirus at a 3:1 ratio. The culture medium was

changed 6h later, and virus-containing supernatants were harvested 42h later. Viral particles were

filtered, purified by ultracentrifugation through a sucrose cushion (25% weight/volume in Tris-

NaCl-EDTA buffer) for 75 min at 4°C and 28,000 rpm using a SW 32 TI rotor (Beckman Coulter),

resuspended in serum-free RPMI 1640 or DMEM medium and stored in small aliquots at -80°C.

Viral particles were titrated using an HIV-1 p24^{Gag} Alpha-Lisa kit and an Envision plate reader

(Perkin Elmer) and/or by determining their infection titers on target cells.

Lentiviral and retroviral infections. For infections with replication-competent HIV-1 Renilla or

wild-type and/or VSV-G pseudotyped-HIV-1, target cells were plated at 2.5 x 10^4 cells per well in

96-well plates or at 2 x 10^5 cells per well in 12-well plates and infected for 24-48 h before lysis

and Renilla (and Firefly) luciferase activity measure (Dual-Luciferase® Reporter Assay System

Promega) or fixation with 2% paraformaldehyde (PFA)-PBS, permeabilization (Perm/Wash buffer,

BDBiosciences) and intracellular staining with the anti-p24^{Gag} KC57-FITC antibody (Beckman

Coulter), as described previously[62]. For TZM-bl assays, the β-galactosidase activity was

measured using the Galacto-Star™ system (ThermoFisher Scientific). For infections with lentiviral

and retroviral vectors, target cells were plated at 2.5 x 10^4 cells per well in 96-well plates the day

prior to infection with vectors at the indicated MOIs, and the percentages of infected cells were

scored by flow cytometry 24h later. For primary CD4+ T cell infections, 10^5 cells were infected

with 100 ng p24^{Gag} of HIV-1 Renilla for 24 h prior to lysis and luciferase activity measure. For

MDM infections, 8 x 10^4 cells were infected with 100 ng p24^{Gag} of a CCR5-tropic version of HIV-

1 Renilla for 30 h prior to lysis and luciferase activity measure.

Retrotransposon assays. For GFP-based retrotransposon assays, HEK293T cells (2 x 10^5 cells)

were co-transfected with either 1 µg of pJM111 (a negative control with two point mutations in

ORF1 that abolish retrotransposition), pRPS-GFP or pLRE3-GFP with either 1 µg of pCAGGS-

Flag-Firefly or pCAGGS-Flag-DDX42. At 7 days post-transfection, the percentage of GFP-

expressing cells was scored by flow cytometry.

CRISPR/Cas9 knock-out. For CRISPR/Cas9 knock-out in cell lines, Lentiguide-Puro LVs coding

sgRNAs targeting the indicated genes or non-targeting sgRNAs were produced, and U87-MG

Cas9/CD4/CXCR4/Firefly were transduced for 6 h before replacing the supernatants with fresh,

complete medium. The transduced cells were selected with puromycin two days later and

amplified for 12-15 days. For CRISPR/Cas9 knock-out in activated primary CD4+ T cells, 2 million

cells per condition were washed with PBS1X, and electroporated using the 4d-Nucleofector®

(Lonza) and the Amaxa P3 primary cell kit with 183 pmol of crispr/tracr RNA duplex (Alt-R

CRISPR-Cas9 crRNA XT and tracrRNA XT, IDT®) and 61 pmol of Cas9 (Alt-R® S.p. Cas9

Nuclease V3, IDT®). After electroporation, the cells were incubated for 4 days at 37°C in X-

VIVO15 medium (Lonza) supplemented with 1% pen/strep and IL-2 at 500 U/ml prior to cell

counting and infection. The crRNA sequences of the sgDDX42-1, -2, and -3 were identical to the

ones cloned in pLentiguide-Puro, and the crRNA of the sgDDX42-4 and sgDDX42-5 were pre-

designed by IDT®, as follow: sg4-DDX42 5'-CGGAGATCTATTAACTGCTG-3', sg5-DDX42 5'-

GAGTTGGTGAGTTTTCAGC-3'.

siRNA transfection. DDX42 and control knockdowns were achieved by transfecting the indicated siRNAs at 44nM, 14.2nM, and 100nM final in U87-MG and Huh-7 cells, TZM-bl cells and MDMs, respectively, with lipofectamine RNAimax (Thermofisher Scientific) according to the manufacturer's instructions. The scramble siRNA controls used were universal siCTRL1 (SIC001) and siCTRL2 (SIC002) (Sigma-Aldrich) and the sequences of the siRNAs targeting DDX42 were siDDX42-1: 5'-CAGAAUGCCUGGUUUCGGA-3' (SASI_Hs01_00119846, Sigma-Aldrich®), siDDX42-2: 5'-CUUACCUUGUGUUUGAUGA-3' (SASI_Hs01_00119845, Sigma-Aldrich®), siDDX42-4: 5'-AUCUCGAAUACCCUUUACG-3' (ID :136410, Ambion®).

RNA immunoprecipitation. For LINE-1 RNA immunoprecipitation, HEK293T cells were co-transfected with equal amounts of pRPS-GFP and pCAGGS-Flag-DDX42 or -Flag-Firefly. For viral RNA immunoprecipitation, U87-MG cells were transduced with either Flag-Firefly or Flag-DDX42 coding lentiviral vectors (pRRL.sin.cPPT.SFFV/Flag-Firefly.WPRE and pRRL.sin.cPPT.SFFV/Flag-DDX42.WPRE, respectively) and infected with CHIKV at MOI 0.1, SARS-CoV-2 at MOI 0.13, or A/Victoria/3/75 IAV at MOI 0.1 for 24 h. 4 days post LINE-1-transfection or 24 h post-infection, cells were washed twice in PBS1X, incubated for 10 min with 0,1% formaldehyde in PBS1X, for 5 min in 250 mM Glycine and washed twice in cold PBS1X. Cells were lysed in RIPA buffer (50 mM Tris/HCl pH 7.5, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 1 mM EDTA, protease inhibitor cocktail and 40 U/mL RNasin). The lysates were clarified by centrifugation at 16,000*g*, for 10 min at 4 °C. Fractions of cell lysates were harvested at this stage to serve as controls for protein and RNA inputs (15%) and the rest was incubated with Flag-magnetic beads (ThermoFisher Scientific) for 2 h at 4 °C. The beads were washed 5 times in RIPA buffer and the immunoprecipitated proteins eluted using 150 µg/mL 3x Flag peptide (Sigma-Aldrich) in elution buffer (50 mM Tris/HCl pH 7.5, 75 mM NaCl, 1mM DTT, protease inhibitor cocktail and 40 U/mL RNasin) for 2 h. Fractions of eluates were harvested for immunoblot analysis (1/6th) and the rest subjected to RNA extraction (5/6th). RNA extractions were then performed using TRIzol (ThermoFisher Scientific).

RNA quantification by RT-qPCR. To check silencing efficiency or measure gene induction after IFN treatment, 0,5-2 x 10^6 cells were collected 2-3 days after siRNA transfection or 24h after IFN treatment or no treatment, and total RNAs were isolated using the RNeasy kit with on-column DNase treatment (Qiagen). cDNAs were generated using 250 ng RNA (High-Capacity cDNA Reverse Transcription Kit, Applied Biosystem, ThermoFisher Scientific, catalogue number 4368814) and analysed by quantitative (q)PCR using TaqMan gene expression assays (Applied Biosystem) specific for ACTB (Hs99999903_m1), GAPDH (Hs99999905_m1), and DDX42 (Hs00201296_m1). Triplicate reactions were run according to the manufacturer's instructions using a ViiA 7 Real-Time PCR system. For relative quantification, samples were normalized to both ACTB and GAPDH mRNA expression and ΔΔCt analysis was performed.

For the measure of LINE-1 RNAs, 100 ng RNA (from cell extracts) or 25 μl of RNA extracted from the IP eluates (i.e. ~60% of the total amount of immunoprecipitated RNA) were reverse transcribed and analysed by qPCR using primers and probe specific for *ORF2*: ORF2-forward 5'-CACCAGTTAGAATGGCAATCATTAAA-3', ORF2-reverse 5'-GGGATGGCTGGGTCAAATGG-3' with ORF2-probe 5'-[FAM]-AGGAAACAACAGGTGCTGGAGAGGATGC-[TAMRA]-3. Absolute quantification was performed using a pRPS-GFP standard curve.

For the measure of SARS-CoV-2 replication, 3 x 10^5 cells were harvested and total RNA was extracted using the RNeasy kit (Qiagen) employing on-column DNase treatment. 125 ng of cellular RNAs were used to generate cDNAs that were analysed by qPCR using RdRp primers and probe, as follow: RdRp_for 5'-GTGARATGGTCATGTGTGGCGG-3', RdRp_rev 5'-CAAATGTTAAAAACACTATTAGCATA-3', and RdRp_probe 5'-[FAM]-CAGGTGGAACCTCATCAGGAGATGC-[TAMRA]-3'[63]. pRdRp (which contains an RdRp fragment amplified from SARS-CoV-2-infected cell RNAs[44]) was diluted in 20 ng/ml salmon sperm DNA to generate a standard curve to calculate relative cDNA copy numbers and confirm the assay linearity (detection limit: 10 molecules of RdRp per reaction).

For the measure of the amounts of viral RNAs in the RNA immunoprecipitation experiments, 100 ng RNA from cell lysates (input) or 25 μl of RNA extracted from the IP eluates (i.e. ~60% of the total amount of immunoprecipitated) were reverse transcribed using the High-Capacity cDNA

Reverse Transcription Kit as above. For SARS-CoV-2, the cDNAs were analysed by RdRp RT-

qPCR. For CHIKV RNA, the following primers and probe were used for the qPCR reactions: E1-

C21-forward   5'-ACGCAGTTGAGCGAAGCAC-3',   E1-C21-reverse   5'-

CTGAAGACATTGGCCCCAC-3'[64],   and   E1-C21-probe   5'-[FAM]-

CTCATACCGCATCTGCATCAGCTAAGCTCC-[TAMRA]-3'. pE1 (which contains an E1 fragment

amplified from CHIKV-infected cell RNAs using primers E1-C21 forward and E1-C21 reverse and

cloned into pPCR-Blunt II-TOPO) was used to generate a standard curve and ensure the linearity

of the assay (detection limit: at least 10 molecules per reaction). For A/Victoria/3/75 IAV RNA, the

following primers and probe, specific for the PA segment, were used, as follow: PA-forward 5'-

TTGCTGCACAGGATGCATTA-3', PA-reverse 5'- AGATTGGAGAAGACGTGGCT-3' and PA-

probe 5'-[FAM]- TGGCTCTGCAATGGGACACCTCTGC-[TAMRA]-3'.   pPolI-RT-Victoria-PA [65]

was used to generate a standard curve and ensure the linearity of the assay (detection limit: at

least 10 molecules per reaction).

Quantification of HIV-1 DNAs. To measure HIV-1 cDNAs, 2×10^5 cells transfected with a control

siRNA or siRNAs targeting DDX42 were plated in 24-well plates, and treated or not with 10 μM

AZT and 3TC 1-2 h prior to infection. The cells were infected with NL4-3 HIV-1 (60 ng p24^{Gag}) for

2 h at 37°C, washed with PBS1X and incubated in complete DMEM before being harvested at

the indicated times. Cell pellets were frozen at -80°C after two washes in PBS1X. Total DNA

extraction was performed using the DNeasy kit (Qiagen) according to the manufacturer's

instructions, and a DpnI-treatment step was performed prior to qPCR. Strong stop reverse

transcription   products   were   detected   using   forward   primer   oHC64   5'-

TAACTAGGGAACCCACTGC-3' and reverse primer oHC65 5'-GCTAGAGATTTTCCACACTG-

3',   2nd   strand   transfer   product   using   oHC64   and   oSA63R   5'-

CTGCGTCGAGAGATCTCCTCTGGCT-3',   together   with   oHC66   probe   5'-[FAM]-

ACACAACAGACGGGCACACACTA-[TAMRA]-3'. 2-LTR circular forms were detected using

2LTR-forward   5'-GTAACTAGAGATCCCTCAG-3'   and   2LTR-reverse   5'-

TGGCCCTGGTGTGTAGTTC-3'   together   with   2LTR-probe   5′-[FAM]-

CTACCACACACAAGGCTACTTCCCTGAT-[TAMRA]-3'. Integrated viral DNA was analysed

using an Alu qPCR as described before[11]. Briefly, a preamplification of 16 cycles was performed

(15 sec at 94°C, 15 sec at 55°C, 100 sec at 68°C) with Platinum Taq DNA High Fidelity

polymerase (Invitrogen) using 100 nM of genomic Alu forward primer 5′-

GCCTCCCAAAGTGCTGGGATTACAG and 600 nM of U3-reverse primer 5'-

CTTCTACCTTATCTGGCTCAAC-3'. The pre-amplification step was performed on serial dilutions

of all the DNA samples, as well as of a positive control (total DNA from U87-MG infected with a

high input of NL4-3), to ensure the linearity of the assay. Background levels were assessed using

linear, one-way amplification by performing the pre-amplification PCR with the U3-reverse primer

alone. Then a qPCR was performed on pre-amplification products using U3-forward primer 5'-

TCTACCACACACAAGGCTAC-3' and U3-reverse primer with the U3 probe 5'-[FAM]-

CAGAACTACACACCAGGGCCAGGGGTCA-[TAMRA]-3'. qPCR reactions were performed in

triplicates, in Universal PCR master mix using 900nM each primer and 250nM probe with the

following program: 10 min at 95°C followed by 40 cycles (15 sec at 95°C and 1 min at 60°C).

pNL4-3 or pTOPO-2LTR (generated by pTOPO cloning of a 2-LTR circle junction amplified from

NL4-3 infected cells, using oHC64 and U3-reverse primers into pCR™2.1-TOPO™) were diluted

in 20 ng/ml of salmon sperm DNA to create dilution standards used to quantify relative cDNA

copy numbers and confirm the linearity of all assays.

Proximity Ligation assays (PLAs). The proximity ligation assays were performed using the

Duolink® in situ Detection Reagents (Sigma-Aldrich, DUO92014). For PLA with HIV-1, MDMs

were plated in 24-well plates with coverslips pre-treated with poly-L-lysin (Sigma-Aldrich) and

infected with 1 μg p24^{Gag} of HIV-1 NL4-3 (Ba-L Env) or mock-infected. For PLA with SARS-CoV-

2, A549-ACE2 cells were plated in 24-well plates with coverslips and infected at an MOI of 0,1.

24 h later, the cells were fixed with 2-4% paraformaldehyde in PBS1X for 10 min, washed in

PBS1X and permeabilized with 0.2% Triton X-100 for 10 min. After a couple of washes in PBS1X,

either NGB buffer (50 mM NH$_4$Cl, 2% goat serum and 2% bovine serum albumin in PBS) or

Duolink® blocking solution was added for 1h. Cells were incubated with mouse AG3.0 anti-HIV-

1 Capsid antibody (National Institutes of Health (NIH) AIDS Reagent Program #4121), or J2 anti-

dsRNA antibody (SCICONS), or anti-SARS-CoV-2 Nucleoprotein (N; BioVision), and rabbit anti-

DDX42 antibody (HPA023571, Sigma-Aldrich) diluted in NGB buffer or in Duolink® blocking

solution for 1h. After 2 washes in PBS1X, the cells were incubated with the DUOLINK® in situ

PLA® Probe Anti-rabbit minus (DUO92006) and DUOLINK® in situ PLA® Probe Anti-mouse plus

(DUO92001) for 1h at 37°C. After 2 washes in PBS1X, the ligation mix was added for 30 min at

37°C. After 2 washes in PBS1X, the cells were incubated with the amplification mix for 100 min

at 37°C followed by 3 washes in PBS1X. In the case of SARS-CoV-2 infection, an additional

staining was performed by incubating cells in an anti-mouse Alexa Fluor secondary antibody.

Finally, the cells were washed twice with PBS1X and stained with Hoechst at 1 μg/mL for 5 min,

washed again and the coverslips mounted on slides in Prolong mounting media (ThermoFisher

Scientific). Z-stack images were acquired using an LSM 880 confocal microscope (ZEISS) using

a 63x lens. PLA punctae quantification was performed using the FIJI software[66]. Briefly, maximum

z-projections were performed on each z-stack and the number of nuclei per field were quantified.

Then, by using a median filter and thresholding, PLA punctae were isolated and quantified

automatically using the Analyse Particles function. To obtain a mean number of dots per cell, the

number of PLA dots per field were averaged by the number of nuclei. In the case of SARS-CoV-

2 infection, the infected cells were identified using N or dsRNA immunofluorescence staining. For

representative images, single cells were imaged using a LSM880 confocal microscope coupled

with an Airyscan module. Processing of the raw Airyscan images was performed on the ZEN

Black software.

Immunoblot analysis. Cell pellets were lysed in sample buffer (200 mM Tris-HCl, pH 6.8, 5.2%

SDS, 20% glycerol, 0.1% bromophenol blue, 5% β-mercaptoethanol), resolved by SDS–PAGE

and analysed by immunoblotting using primary antibodies specific for human DDX42

(HPA023571, Sigma-Aldrich), Flag (M2, Sigma-Aldrich) and Actin (A1978, Sigma-Aldrich),

followed by secondary horseradish peroxidase-conjugated anti-mouse or anti-rabbit

immunoglobulin antibodies and chemiluminescence (Bio-Rad). Images were acquired on a

ChemiDoc™ gel imaging system (Bio-Rad).

IAV production and infection. We have described previously IAV NanoLuciferase reporter virus

generation[43]. Stocks were titrated by plaque assays on MDCK cells. IAV challenges were

performed in serum-free DMEM for 1 h and the medium was subsequently replaced with DMEM

containing 10%. IAV infection experiments were performed in triplicates in 96-well plates with

cultures maintained for 16 h post-challenge. NanoLuciferase activity was measured with the

Nano-Glo assay system (Promega), and luminescence was detected using a plate reader

(Infinite® 200 PRO, Tecan).

VSV production and infection. A VSV-G pseudotyped-VSV-Δenv reporter virus, coding both

GFP and Firefly Luciferase, was obtained from Gert Zimmer . The virus was amplified on pMD.G

transfected HEK293T and titrated thanks to the GFP reporter gene. For infection, 2.5×10^4 cells

per well in 96-well plates were infected at the indicated MOIs. 24h after infection, cells were lysed

and Firefly luciferase activity was measured (Firefly luciferase Assay System Promega).

Measles virus production and infection. Measles virus GFP strain (MeV-GFP), which was

kindly provided by F. Tangy (Institut Pasteur, Paris), was previously described[68]. Viral stocks were

produced on Vero NK cells. After 4 days of infection, supernatant was collected and then

centrifugated to eliminate dead cells or fragments. Stocks were tittered using median tissue

culture infectious dose assays ($TCID_{50}$) on Vero NK cells. Cells were infected with 10-fold serial

dilutions of viral stocks and incubated for 7 days. Cells were then washed with PSB and fixed with

3% formaldehyde crystal violet during 30 min and finally rinsed with water. For infections, Huh-7

cells were infected at the indicated multiplicity of infection (MOI) in DMEM without FBS for 2 h in

small volume of medium to enhance contacts with the inoculum and the cells. After 2 h, the viral

inoculum was replaced with fresh DMEM 10% FBS 1% P/S. 24 h post-infection the cells were

harvested and samples separated in half for Western blot and flow cytometry analysis.

ZIKV production and infection. The nanoluciferase expressing ZIKV construct has been described[69]. The corresponding linearized plasmid was transcribed in vitro using the SP6 mMESSAGE mMACHINE™ (Thermofischer Scientific) and HEK293T cells were transfected with the transcribed RNA. After 7 days, supernatants were harvested, filtered and stock titers were determined by plaque assays on Vero cells. For infections, 2.5×10^4 cells per well in 96-well plates were infected, at the indicated MOIs. 24h after infection, cells were lysed and Nanoluciferase activity was measured using the Kit Nano Glo luciferase Assay (Promega).

CHIKV production and infection. The Nanoluciferase luciferase coding CHIKV construct was a gift from Andres Merits. The linearized plasmid coding CHIKV genome was transcribed with the T7 mMESSAGE mMACHINE kit (Thermofischer Scientific) and 5×10^5 HEK293T were transfected with 1-4 μg of transcribed RNA, using Lipofectamine 2000 (Thermofischer Scientific). After 24h, supernatants were harvested, filtered and viruses were then amplified on baby hamster kidney (BHK21) cells. Stock titers were determined by plaque assays on Vero cells. For infections, 2.5×10^4 cells per well in 96-well plates were infected at the indicated MOIs. 24h after infection, cells were lysed and Nanoluciferase activity was measured.

SARS-CoV-2 production and infection. The SARS-CoV-2 BetaCoV/France/IDF0372/2020 isolate was supplied by Pr. Sylvie van der Werf and the National Reference Centre for Respiratory Viruses hosted by Institut Pasteur (Paris, France). The patient sample from which strain BetaCoV/France/IDF0372/2020 was isolated was provided by Dr. X. Lescure and Pr. Y. Yazdanpanah from the Bichat Hospital, Paris, France. The virus was amplified in Vero E6 cells (MOI 0,005) in serum-free media supplemented with 0,1 μg/ml L-1-p-Tosylamino-2-phenylethyl chloromethylketone (TPCK)-treated trypsin (Sigma-Aldrich). The supernatant was harvested at 72 h post infection when cytopathic effects were observed (with around 50% cell death), cell debris were removed by centrifugation, and aliquots stored at -80C. Viral supernatants were titrated by plaque assays in Vero E6 cells. Typical titers were 5.10^6 plaque forming units (PFU)/ml. Infections of A549-ACE2 cells were performed at the indicated multiplicity of infection (MOI; as

calculated from titers obtained in Vero E6 cells) in serum-free DMEM and 5% serum-containing

DMEM, respectively. The viral input was left for the duration of the experiment and cells lysed at

48 h post-infection for RT-qPCR analysis.

Statistical analyses. Statistical analyses were performed with GraphPad Prism. Analysis types

are mentioned in Fig. legends and all comparisons are relative to the indicated controls. For data

with experimental factors greater than two, multiple linear regression was performed. For data

with two categorical factors, ANOVA was used, and repeated measures ANOVA when a pairing

factor was present. Simple linear regression was used when the relationship between a

continuous factor and a continuous response variable was investigated. P values are denoted as

follow: ns not significant, $p<0.05$ *, $p<0.01$ **, $p<0.001$ ***, $p<0.0001$****.

Data availability

The datasets generated during and/or analysed during the current study are available from the

corresponding authors on reasonable request.

Requests for materials

Requests for material should be addressed to Caroline Goujon at the corresponding address

above.

Acknowledgements

We wish to thank Tom Doyle and Chad Swanson for their useful comments on the manuscript,

and Matthieu Lewis, Nadine Laguette, Nathalie Arhel, Juliette Fernandez, Jean-Luc Battini, Georg

Kochs, Andres Merits, Frédéric Tangy, Sylvie van de Werf and Nathan Sherer for the generous

provision of reagents, protocols and/or for helpful discussions. We are grateful to Nicolas Manel

and Helena Izquierdo-Fernandez for sharing their protocol for efficient knockdown in primary

CD4+ T cell using CRISPR/Cas9 RNP electroporation. We are grateful to Myriam Boyer,

Stéphanie Viala, Baptiste Monterroso and Orestis Faklaris from the imaging and flow cytometry

facility MRI, for advice with flow cytometry and confocal microscopy, respectively. This work was

supported by the Institut National de la Santé et de la Recherche Médicale (INSERM) (to CG),

the ATIP-Avenir programme (to CG), institutional funds from the Centre National de la Recherche

Scientifique (CNRS) and Montpellier University, France REcherche Nord&Sud Sida/HIV et

Hépatites (ANRS) (ECTZ21792 to CG; ECTZ35478 to OM), Sidaction (to CG), a Starting Grant

from the European Research Council (759226, ANTIViR) (to CG), PhD studentships from the

Ministry of Higher Education and Research (to BB, JM and AR), and a 4th year PhD studentship

from the Fondation pour la Recherche Médicale (to BB). SR and HP acknowledge financial

support from the France Génomique National infrastructure, funded as part of "Investissement

d'Avenir" program managed by Agence Nationale de la Recherche (contract ANR-10-INBS-09).

We acknowledge the imaging facility MRI, member of the national infrastructure France-

BioImaging supported by the French National Research Agency (ANR-10-INBS-04) and the BSL-

3 facility CEMIPAI (UAR 3725 CNRS Montpellier University).

Author contributions

B.B. and CG designed the study, analysed the data and wrote the manuscript. B.B. and C.G.

performed the whole-genome screens and candidate validation. B.B. carried out most of the

experiments, with technical assistance from A.R., F.G.d.G, J.M., M.T., W.D., A.M., A.L.C.V.,

M.A.A., and O.M.; V.C. provided some of the lentiviral vector stocks; E.B and L.B. performed

CHIKV infections, N.G. performed ZIKV-Nluc infections, S.G. and N.J. performed MeV, ZIKV

PF13, DENV-2 and YFV infections; M.L and E.R analysed the RNA-seq data; R.S. performed the

statistical analyses; H.P. and S.R. performed the Illumina sequencing and MaGECK analyses,

respectively. All authors have read and approved the manuscript.

Conflicts of interest statement

The authors have no conflicts of interest to declare in relation to this manuscript.

References

1.   Schoggins, J. W. Interferon-Stimulated Genes: What Do They All Do? *Annu. Rev. Virol.* **6**, 567–584 (2019).

2.   Lindenmann, J. Resistance of mouse to mice adapted influenza A virus. *Virology* **16**, 203–204 (1962).

3.   Zhou, A. Expression cloning of 2-5A-dependent RNAase: A uniquely regulated mediator of interferon action.

Cell **72**, 753–765 (1993).

4.   Zilberstein, A., Kimchi, A., Schmidt, A. & Revel, M. Isolation of two interferon-induced translational inhibitors:

a protein kinase and an oligo-isoadenylate synthetase. *Proc. Natl. Acad. Sci.* **75**, 4734–4738 (1978).

5.   Kane, M. *et al.* Identification of Interferon-Stimulated Genes with Antiretroviral Activity. *Cell Host Microbe* **20**,

392–405 (2016).

6.   OhAinle, M. *et al.* A virus-packageable CRISPR screen identifies host factors mediating interferon inhibition

of HIV. *eLife* **7**, e39823 (2018).

7.   Schoggins, J. W. *et al.* A diverse range of gene products are effectors of the type I interferon antiviral

response. *Nature* **472**, 481–485 (2011).

8.   Chemudupati, M. *et al.* From APOBEC to ZAP: Diverse mechanisms used by cellular restriction factors to

inhibit virus infections. *Biochim. Biophys. Acta Mol. Cell Res.* **1866**, 382–394 (2019).

9.   Doyle, T., Goujon, C. & Malim, M. H. HIV-1 and interferons: who's interfering with whom? *Nat. Rev. Microbiol.*

13, 403–413 (2015).

10.   Ghimire, D., Rai, M. & Gaur, R. Novel host restriction factors implicated in HIV-1 replication. *J. Gen. Virol.*

99, 435–446 (2018).

11.   Goujon, C. *et al.* Human MX2 is an interferon-induced post-entry inhibitor of HIV-1 infection. *Nature* **502**,

559–562 (2013).

12.   Kane, M. *et al.* MX2 is an interferon-induced inhibitor of HIV-1 infection. *Nature* **502**, 563–566 (2013).

13.   Jimenez-Guardeño, J. M., Apolonia, L., Betancor, G. & Malim, M. H. Immunoproteasome activation enables

human TRIM5α restriction of HIV-1. *Nat. Microbiol.* (2019) doi:10.1038/s41564-019-0402-0.

14.   Fan, W. *et al.* TRIM7 inhibits enterovirus replication and promotes emergence of a viral variant with

increased pathogenicity. *Cell* **184**, 3410-3425.e17 (2021).

15.   Shalem, O. *et al.* Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* **343**, 84–87

(2014).

16.   Shalem, O., Sanjana, N. E. & Zhang, F. High-throughput functional genomics using CRISPR-Cas9. *Nat.*

Rev. Genet. **16**, 299–311 (2015).

17.   Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide libraries for CRISPR screening.

Nat. Methods **11**, 783–784 (2014).

18. Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9

knockout screens. *Genome Biol.* **15**, 554 (2014).

19. Doench, J. G. Am I ready for CRISPR? A user's guide to genetic screens. *Nat. Rev. Genet.* **19**, 67–80

(2018).

20. Goujon, C. *et al.* Evidence for IFNα-induced, SAMHD1-independent inhibitors of early HIV-1 infection.

Retrovirology **10**, 23 (2013).

21. Bulli, L. *et al.* Complex interplay between HIV-1 Capsid and MX2-independent IFNα-induced antiviral factors.

J. Virol. (2016) doi:10.1128/JVI.00458-16.

22. Xu, B., Pan, Q. & Liang, C. Role of MxB in Alpha Interferon-Mediated Inhibition of HIV-1 Infection. *J. Virol.*

92, (2018).

23. Uhlmann-Schiffler, H., Jalal, C. & Stahl, H. Ddx42p--a human DEAD box protein with RNA chaperone

activities. *Nucleic Acids Res.* **34**, 10–22 (2006).

24. Taschuk, F. & Cherry, S. DEAD-Box Helicases: Sensors, Regulators, and Effectors for Antiviral Defense.

Viruses **12**, (2020).

25. Granneman, S., Bernstein, K. A., Bleichert, F. & Baserga, S. J. Comprehensive Mutational Analysis of Yeast

DEXD/H Box RNA Helicases Required for Small Ribosomal Subunit Synthesis. *Mol. Cell. Biol.* **26**, 1183–

1194 (2006).

26. Rocak, S. Characterization of the ATPase and unwinding activities of the yeast DEAD-box protein Has1p

and the analysis of the roles of the conserved motifs. *Nucleic Acids Res.* **33**, 999–1009 (2005).

27. Ochsenbauer, C. *et al.* Generation of transmitted/founder HIV-1 infectious molecular clones and

characterization of their replication capacity in CD4 T lymphocytes and monocyte-derived macrophages. *J.*

Virol. **86**, 2715–2728 (2012).

28. Uhlmann-Schiffler, H., Kiermayer, S. & Stahl, H. The DEAD box protein Ddx42p modulates the function of

ASPP2, a stimulator of apoptosis. *Oncogene* **28**, 2065–2073 (2009).

29. Zyner, K. G. *et al.* Genetic interactions of G-quadruplexes in humans. *eLife* **8**, (2019).

30. Faulkner, G. J. & Billon, V. L1 retrotransposition in the soma: a field jumping ahead. *Mob. DNA* **9**, 22 (2018).

31. Liu, N. *et al.* Selective silencing of euchromatic L1s revealed by genome-wide screens for L1 regulators.

Nature **553**, 228–232 (2018).

32. Moran, J. V. *et al.* High frequency retrotransposition in cultured mammalian cells. *Cell* **87**, 917–927 (1996).

33. Wei, J. *et al.* Genome-wide CRISPR screens reveal host factors critical for SARS-CoV-2 infection. *Cell*

(2020) doi:10.1016/j.cell.2020.10.028.

34. Haller, O., Staeheli, P., Schwemmle, M. & Kochs, G. Mx GTPases: dynamin-like antiviral machines of innate

immunity. *Trends Microbiol.* **23**, 154–163 (2015).

35. Burdick, R. C. *et al.* HIV-1 uncoats in the nucleus near sites of integration. *Proc. Natl. Acad. Sci.* **117**, 5486–

5493 (2020).

36. Dharan, A., Bachmann, N., Talley, S., Zwikelmaier, V. & Campbell, E. M. Nuclear pore blockade reveals that

HIV-1 completes reverse transcription and uncoating in the nucleus. *Nat. Microbiol.* (2020)

doi:10.1038/s41564-020-0735-8.

37. Peng, K. *et al.* Quantitative microscopy of functional HIV post-entry complexes reveals association of

replication with the viral capsid. *eLife* **3**, e04114 (2014).

38. Zyner, K. G. *et al.* Genetic interactions of G-quadruplexes in humans. *eLife* **8**, e46793 (2019).

39. Fay, M. M., Lyons, S. M. & Ivanov, P. RNA G-Quadruplexes in Biology: Principles and Molecular

Mechanisms. *J. Mol. Biol.* **429**, 2127–2147 (2017).

40. Ruggiero, E. & Richter, S. N. G-quadruplexes and G-quadruplex ligands: targets and tools in antiviral

therapy. *Nucleic Acids Res.* **46**, 3270–3283 (2018).

41. Will, C. L. *et al.* Characterization of novel SF3b and 17S U2 snRNP proteins, including a human Prp5p

homologue and an SF3b DEAD-box protein. *EMBO J.* **21**, 4978–4988 (2002).

42. Goujon, C. *et al.* Characterization of Simian Immunodeficiency Virus SIVSM/Human Immunodeficiency Virus

Type 2 Vpx Function in Human Myeloid Cells. *J. Virol.* **82**, 12335–12345 (2008).

43. Doyle, T. *et al.* The interferon-inducible isoform of NCOA7 inhibits endosome-mediated viral entry. *Nat.*

Microbiol. **3**, 1369–1376 (2018).

44. Rebendenne, A. *et al.* SARS-CoV-2 triggers an MDA-5-dependent interferon response which is unable to

control replication in lung epithelial cells. *J. Virol.* (2021) doi:10.1128/JVI.02415-20.

45. Moncorge, O. *et al.* Investigation of Influenza Virus Polymerase Activity in Pig Cells. *J. Virol.* **87**, 384–394

(2013).

46. Adachi, A. *et al.* Production of acquired immunodeficiency syndrome-associated retrovirus in human and

nonhuman cells transfected with an infectious molecular clone. *J. Virol.* **59**, 284–291 (1986).

47. Simon, J. H., Southerling, T. E., Peterson, J. C., Meyer, B. E. & Malim, M. H. Complementation of vif-

defective human immunodeficiency virus type 1 by primate, but not nonprimate, lentivirus vif genes. *J. Virol.*

69, 4166–4172 (1995).

48. Schaller, T. *et al.* HIV-1 capsid-cyclophilin interactions determine nuclear import pathway, integration

targeting and replication efficiency. *PLoS Pathog.* **7**, e1002439 (2011).

49. Ryan-Graham, M. A. & Peden, K. W. C. Both Virus and Host Components Are Important for the Manifestation

of a Nef- Phenotype in HIV-1 and HIV-2. *Virology* **213**, 158–168 (1995).

50. Gaddis, N. C. *et al.* Further Investigation of Simian Immunodeficiency Virus Vif Function in Human Cells. *J.*

Virol. **78**, 12041–12046 (2004).

51. Naldini, L. *et al.* In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector.

Science **272**, 263–267 (1996).

52. Bainbridge, J. W. *et al.* In vivo gene transfer to the mouse eye using an HIV-based lentiviral vector; efficient

long-term transduction of corneal endothelium and retinal pigment epithelium. *Gene Ther.* **8**, 1665–1668

(2001).

53. O'Rourke, J. P., Newbound, G. C., Kohn, D. B., Olsen, J. C. & Bunnell, B. A. Comparison of gene transfer

efficiencies and gene expression levels achieved with equine infectious anemia virus- and human

immunodeficiency virus type 1-derived lentivirus vectors. *J. Virol.* **76**, 1510–1515 (2002).

54. Saenz, D. T., Teo, W., Olsen, J. C. & Poeschla, E. M. Restriction of feline immunodeficiency virus by Ref1,

Lv1, and primate TRIM5alpha proteins. *J. Virol.* **79**, 15175–15188 (2005).

55. Ostertag, E. M., Prak, E. T., DeBerardinis, R. J., Moran, J. V. & Kazazian, H. H. Determination of L1

retrotransposition kinetics in cultured cells. *Nucleic Acids Res.* **28**, 1418–1423 (2000).

56. Goodier, J. L., Cheung, L. E. & Kazazian, H. H. MOV10 RNA Helicase Is a Potent Inhibitor of

Retrotransposition in Cells. *PLoS Genet.* **8**, e1002941 (2012).

57. Nakabayashi, H., Taketa, K., Miyano, K., Yamane, T. & Sato, J. Growth of human hepatoma cells lines with

differentiated functions in chemically defined medium. *Cancer Res.* **42**, 3858–3863 (1982).

58. Wingett, S. W. & Andrews, S. FastQ Screen: A tool for multi-genome mapping and quality control.

F1000Research **7**, 1338 (2018).

59. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*

17, 10–12 (2011).

60. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short DNA

sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

61. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**, 2078–2079

(2009).

62. Goujon, C. & Malim, M. H. Characterization of the alpha interferon-induced postentry block to HIV-1 infection

in primary human macrophages and T cells. *J. Virol.* **84**, 9254–9266 (2010).

63. Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill.*

Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull. **25**, (2020).

64. Meertens, L. *et al.* FHL1 is a major host factor for chikungunya virus infection. *Nature* **574**, 259–263 (2019).

65. Zürcher, T., de la Luna, S., Sanz-Ezquerro, J. J., Nieto, A. & Ortín, J. Mutational analysis of the influenza

virus A/Victoria/3/75 PA protein: studies of interaction with PB1 protein and identification of a dominant

negative mutant. *J. Gen. Virol.* **77 (Pt 8)**, 1745–1749 (1996).

66. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682

(2012).

67. Rentsch, M. B. & Zimmer, G. A Vesicular Stomatitis Virus Replicon-Based Bioassay for the Rapid and

Sensitive Determination of Multi-Species Type I Interferon. *PLoS ONE* **6**, (2011).

68. Combredet, C. *et al.* A molecularly cloned Schwarz strain of measles virus vaccine induces strong immune

responses in macaques and transgenic mice. *J. Virol.* **77**, 11546–11554 (2003).

69. Mutso, M. *et al.* Reverse genetic system, genetically stable reporter viruses and packaged subgenomic

replicon based on a Brazilian Zika virus isolate. *J. Gen. Virol.* **98**, 2712–2724 (2017).

SUPPLEMENTARY INFORMATION

Supplemental methods

Only the methods specific for the supplemental Figures are described here.

Plasmids. pBlaM-Vpr and pAdVAntage have been described[70].

Lentiviral production. β-lactamase-Vpr (BlaM-Vpr)-carrying viruses, bearing the wild-type Env,

were produced by co-transfection of HEK293T cells with the NL4-3/Nef-IRES-Renilla provirus

expression vector, pBlaM-Vpr and pAdVAntage at a ratio of 4:1:0.5, as previously described[70].

Viral particles were titrated using an HIV-1 p24^{Gag} Alpha-Lisa kit and an Envision plate reader

(Perkin Elmer).

BlaM-Vpr assay for HIV-1 entry. This assay was performed as described previously[62]. Briefly, 2

x 10^5 U87-MG/CD4/CXCR4 cells were plated in 24-well plates and incubated with BlaM-Vpr

carrying NL4-3 particles (31, 62, 125 ng p24^{Gag}) or mock-infected for 3 h at 37°C. The cells were

then washed once in CO_2-independent medium and loaded with CCF2-AM substrate-containing

solution (ThermoFisher Scientific) for 1 h at room temperature before 2 washes and incubation at

room temperature for 16 h in development medium (CO_2-independent medium containing 2,5 mM

probenecid). Finally, the cells were trypsinized, washed and fixed in 1% paraformaldehyde (PFA)-

PBS1X before analysis with a FACSCanto™ II (Becton Dickinson).

1004

Viruses and infection. The Yellow Fever Virus (YFV) Asibi strain was provided by the Biological

resource Center of Institut Pasteur. Stocks were produced on Vero NK cells. After 3 days of

infection, viruses were concentrated by polyethylene glycol 6000 (PEG) precipitation and purified

by centrifugation in a discontinuous gradient of sucrose. The Dengue 2 strain Malaysia SB8553

(DENV-2) was obtained from the Centro de Ingeniería Genética y Biotecnología (CIGB), Cuba.

Stocks were generated on Vero NK cells. After 4 days of infection, viruses were concentrated by

PEG 6000 precipitation. The Zika strain PF13 (kindly provided by V. M. Cao-Lormeau and D.

Musso, Institut Louis Malardé, Tahiti Island, French Polynesia) was isolated from a viremic patient

in French Polynesia in 2013. Stocks were produced on C6-36 cells. After 2 days of infection,

viruses were concentrated by PEG 6000 precipitation and purified by centrifugation in a

discontinuous gradient of sucrose. YFV Asibi and ZIKV titers were assessed by plaque assays

using Vero NK cells, as described previously[71]. DENV-2 was tittered by in cell western assays on

Vero cells. Cells were fixed with PFA 4% during 30 min at room temperature (RT), then washed

in PBS and permeabilized with 0,5% triton in PBS (Sigma-Aldrich) during 10 min at RT. Cells

were then incubated 0,1% Tween in PBS (Sigma-Aldrich) containing 5% BSA (Sigma-Aldrich)

during 1 h at RT prior to incubation with mouse anti-Env 4G2 antibodies overnight at 4°C. After

1h of incubation with the secondary antibodies, cells were revealed with an Odyssey CLx infrared

imaging system (Li-Cor Bioscience).

Cells were infected at the indicated multiplicity of infection (MOI) in DMEM without FBS for 2 h in

small volume of medium to enhance contacts with the inoculum and the cells. After 2 h, the viral

inoculum was replaced with fresh DMEM 10% FBS 1% P/S. 24 hours post-infection the cells were

harvested and samples separated in half for Western blot and flow cytometry analysis. For the

latter, cells were fixed and permeabilized using BD Cytofix/Cytoperm (Fisher scientific) for 30 min

on ice (all the following steps were performed on ice and centrifuged at 4°C) and then washed

tree times with wash buffer. Cells infected with YFV, ZIKV and DENV-2 were incubated with the

pan-flavivirus anti-Env 4G2 antibody for 1 h at 4°C and then with Alexa 488 anti-mouse IgG

secondary antibodies (Thermo Fisher) for 45 min at 4°C in the dark. Data were acquired with an

Attune NxT Acoustic Focusing Cytometer (Life technologies) and analyzed using FlowJo

software.

1034

Quantification of mRNA expression. For all Figures except Fig. S4, see the main material and

methods; For Fig. S4: a TaqMan gene expression assays (Applied Biosystem) specific for ISG15

was used: Hs00192713_m1.

For the RNA extraction and RT-qPCR analysis specifically in Huh-7 cells, total RNAs were

extracted from cell lysates using the NucleoSpin RNA II kit (Macherey-Nagel), following the

manufacturer's protocol. Equal amounts of purified total RNA were used to synthetized first strand

cDNA using random hexamers (Thermo Fisher) and the ReverAid H Minus Moloney murine

leukemia virus (M-MuLV) reverse transcriptase (Thermo Fisher). Quantitative real-time PCR was

performed on a real-time PCR system (Quant Studio 6 Flex from Applied Biosytems) with SYBR

green PCR master mix (Life Technologies). Data were analyzed with the $\Delta\Delta$ CT method. All the

samples were performed in technical triplicate and normalized to GAPDH (glyceraldehyde-3-

phosphate dehydrogenase) as endogenous reference control. The primer sequences used for

Fig. S4 were as follow: GAPDH Forward: 5'-GGTCGGAGTCAACGGATTTG-3', reverse: 5'-

ACTCCACGACGTACTCAGCG-3'; DDX42 Forward: 5'-GGCCTATACCCTACTCACTCCC-3',

reverse: 5'-CCACCAATGTTCAGCTTTTTTCC-3'.

1050

Preparation of RNA-seq libraries. siRNA transfected U87-MG/CD4/CXCR4 and A549-ACE2

RNA extracts from three independent experiments were used for RNA-seq library preparation.

After determining sample RNA integrity numbers using a 2100 Bioanalyzer (Agilent), ribosomal

RNAs were depleted using the QIAseq FastSelect -rRNA HMR Kit (Qiagen) and libraries were

prepared according to manufacturer's instructions using the QIAseq Stranded Total RNA Lib Kit

(Qiagen). Libraries were quantified using a TapeStation D1000 ScreenTape. Equimolar amounts

of each library were then mixed and sequenced on 2 lanes (2x150 bp) on the Illumina HiSeq

3000/4000 platform (GENEWIZ).

1059

Analysis of high-throughput sequencing reads. Sequenced reads were filtered by quality and

sequence adaptors removed using fastp v0.20.1 (https://github.com/OpenGene/fastp)[72] with

following parameters "fastp --qualified_quality_phred 20 --disable_length_filtering --

detect_adapter_for_pe". Reads were pseudo-mapped against human cDNA sequenced

downloaded from Gencode database (https://www.gencodegenes.org/) and transcripts

abundance estimated with Kallisto v0.46.2 (https://pachterlab.github.io/kallisto/about)[73] with

parameters "--bias --bootstrap-samples 100".

1067

1068   **Differential analysis with DESeq2.** Differential expressed genes upon siRNA transfection were

1069   obtained using DESeq2[74] (version 1.32.0) in R (version 4.1.0). Briefly, transcript estimations were

1070   transformed in gene counts with tximport package[75] and the differential expression obtained with

1071   the model design "~ condition + replicate".

1072

1073   **Supplementary Information References**

1074

70. Cavrois, M., de Noronha, C. & Greene, W. C. A sensitive and specific enzyme-based assay detecting

HIV-1 virion fusion in primary T lymphocytes. *Nat. Biotechnol.* **20**, 1151–1154 (2002).

71. Beauclair, G. *et al.* Retinoic Acid Inducible Gene I and Protein Kinase R, but Not Stress Granules,

Mediate the Proinflammatory Response to Yellow Fever Virus. *J. Virol.* **94**, (2020).

72. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*

34, i884–i890 (2018).

73. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification.

Nat. Biotechnol. **34**, 525–527 (2016).

74. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol.* **15**, 550 (2014).

75. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-level estimates

improve gene-level inferences. *F1000Research* **4**, 1521 (2015).

1087

1088 **Supplementary Figures**

1089

Figure S1. Related to Figure 1.

A. IFN pre-treatment potently inhibits HIV-1 infection in T98G cells and is, at least partially, saturable.

T98G/Cas9/CD4/CXCR4/ Firefly cells were pre-treated with IFN for 24 h prior to infection with increasing doses

of NL4-3 Renilla (indicated in ng p24^{Gag}). Renilla activity was normalized to Firefly activity and the relative

infection efficiencies are shown. Data represent the average of 3 independent experiments and one standard

deviation from the mean.

B. GeCKO screen validation. GeCKO control cells and enriched cells from 3 successive rounds of selection

(Round 1, 2, and 3, as indicated) were treated with IFN or not and infected with GFP-expressing lentiviral vectors.

The percentage of infected cells was evaluated by flow cytometry 2 days post-infection. Data represent the

average of 2 independent experiments and one standard deviation from the mean.

1100

1101

1102

1103
Figure S2. Related to Figure 2.

A. DDX42 is not an ISG. The indicated primary cells or immortalized cell cultures were treated with IFN for 24 h
or left untreated. RNA was subsequently extracted and DDX42 and ISG15 (a prototype ISG) mRNA levels were
quantified by RT-qPCR; Actin B and GAPDH were used as endogenous controls. The bar chart shows the relative
levels of expression of DDX42 and ISG15 in the presence or absence of IFN treatment. Data represent the mean
± S.E.M of 3 independent experiments.

B. DDX42 silencing efficiency in monocyte-derived macrophages. siRNA-transfected MDMs were harvested
48h post-transfection for RNA extraction and quantification of DDX42 mRNA levels by RT-qPCR. Actin and
GAPDH were used as endogenous controls. Data represent the mean ± S.E.M of 3 independent experiments
performed with cells from 3 different blood donors (parallel samples from Fig. 2E).

C. Ectopic expression of DDX42 K303E mutant increases HIV-1 infection. U87-MG/CD4/CXCR4 cells were
transduced with lentiviral vectors expressing either Firefly (negative control), WT DDX42 (DDX42) or a motif I
mutant DDX42 K303, which has an impaired ATPase activity (DDX42 K303E). Transduced cells were infected
with NL4-3 Renilla and the infection efficiency was assessed 24h later by measuring Renilla activity. Data
represent the mean ± S.E.M of 4 independent experiments.

1119
1120

Figure S3. Related to Figure 3.

A. DDX42 depletion does not affect HIV-1 entry. DDX42-depleted U87-MG/CD4/CXCR4 cells were infected with the indicated amounts of viruses carrying the β-lactamase (BlaM)-Vpr fusion protein for 3 h. The cells were subsequently loaded with CCF2-AM substrate dye for 2h, washed extensively and incubated for another 16 h for the reaction to develop. Cells positive for the CCF2-AM product were scored by flow cytometry. Data represent the mean ± S.E.M of 3 independent experiments.

B. DDX42 localization in MDMs. MDMs were fixed, endogenous DDX42 and the nuclei were visualized using DDX42-specific antibodies and Hoechst staining, respectively, and confocal microscopy. Representative images are shown. Scale bar, 10 μm.

1131

Figure S4. Related to Figure 4.

A. DDX42 silencing efficiency in U87-MG, A549-ACE2 and Huh-7 cells.

B. Relative ZIKV PF13 infection efficiency in siRNA-transfected Huh-7 cells analysed by flow cytometry.

C. Relative DENV-2 infection efficiency in siRNA-transfected Huh-7 cells analysed by flow cytometry.

D. Relative YFV infection efficiency in siRNA-transfected Huh-7 cells analysed by flow cytometry.

A-D. Mean ± SEM of 3 independent experiments (4 for silencing efficiency in A549-ACE2 cells). Two-way

ANOVA with Sidak's test on log10 transformed data.

1139

1140

1141 **Figure S5.**

1142 Venn diagrams showing the Differentially Expressed Genes (DEGs) overlap between siRNA conditions in U87-

1143 MG cells, A549 cells and DEGs overlapping between U87-MG and A549 cells using cutoff criteria of log2 fold

1144 change (log2FC) >1 and p value <0.05.

1145 See **Supplemental File 1** for the identity of the DEGs.