CK2 alpha prime and alpha-synuclein pathogenic functional interaction mediates inflammation and transcriptional dysregulation in Huntington's disease

Dahyun Yu[1,*], Nicole Zarate[1,*], Francesco Cuccu[1,3], Johnny S. Yue[1,4], Taylor G. Brown[1], Wei Tsai[1], Rachel Mansky[1], Kevin Jiang[1], Hyuck Kim[1#], Angel White[1], Carmen Nanclares[1], Tessa Nichols-Meade[1], Sarah N. Larson[2], Katie Gundry[2], Ying Zhang[5], Cristina Tomas-Zapico[6#], Jose J. Lucas[6,7], Michael Benneyworth[1], Gülin Öz[2], Marija Cvetanovic[1], Alfonso Araque[1] and Rocio Gomez-Pastor[1±].

[1]Department of Neuroscience, School of Medicine, University of Minnesota, Minneapolis, MN, United States. [2]Center for Magnetic Resonance Research. Department of Radiology, School of Medicine, University of Minnesota, Minneapolis, MN, United States. [3]Department of Life and Environment Sciences, University of Cagliari, Cagliari, Italy. [4] Mounds View High School, Arden Hills, Minnesota. [5]Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, MN, United States. [6] Centro de Biología Molecular 'Severo Ochoa' (CBMSO) CSIC/UAM, Madrid, Spain. [7]Networking Research Center on Neurodegenerative Diseases (CIBERNED), Instituto de Salud Carlos III, Madrid, Spain

#Current address: HK, MEPSGEN, Seoul 05836, South Korea. CTZ, Department of Functional Biology, Physiology, University of Oviedo, Asturias 33006, Spain; Health Research Institute of the Principality of Asturias (ISPA), Asturias 33011, Spain.

*These authors contributed equally to the manuscript

±Correspondence should be addressed to Rocio Gomez-Pastor, University of Minnesota, 321 Church St. SE, Jackson Hall Room 6-145, Minneapolis, MN 55455, rgomezpa@umn.edu

Keywords: Huntington's disease, polyglutamine, CK2 alpha prime, alpha-synuclein, neuroinflammation, protein aggregation.

Abstract

Huntington's Disease (HD) is a neurodegenerative disorder caused by a CAG trinucleotide repeat expansion in the Htt gene for which no therapies are available. This mutation causes HTT protein misfolding and aggregation, preferentially affecting medium spiny neurons (MSNs) of the basal ganglia. Transcriptional perturbations in synaptic genes and neuroinflammation are key processes that precede MSN dysfunction and motor symptom onset. Understanding the interplay between these processes is crucial to develop effective therapeutic strategies to treat HD. We investigated whether protein kinase CK2α', a kinase upregulated in MSNs in HD and previously associated with Parkinson's disease (PD), participates in the regulation of neuroinflammation and synaptic function during HD progression. We used the heterozygous knock-in zQ175 HD mouse model that lacks one allele of CK2α' and performed cytokine proteome profiling, RNA-seq, electrophysiological recordings, and behavioral analyses. We showed that CK2α' haploinsufficiency in zQ175 mice ameliorated neuroinflammation, HTT aggregation, transcriptional alterations, excitatory synaptic transmission deficits, and motor dysfunction. Our RNA-seq analyses also revealed a connection between α-syn, a protein associated with PD, and the transcriptional perturbations mediated by CK2α' in HD. We found that CK2α' increased α-syn serine 129 phosphorylation (pS129-α-syn), a post-translational modification linked to α-synucleinopathy, and its accumulation in the nucleus of MSNs in zQ175 mice and in patients with HD. Our data demonstrated that CK2α' negatively contributes to HD by promoting striatal synucleinopathy, transcriptional alteration of synaptic genes and neuroinflammation. We propose CK2α' as a potential therapeutic target to treat HD.

Significance statement

Upregulation of Protein kinase CK2α' mediates neuroinflammation and transcriptional dysregulation of synaptic genes in the zQ175 mouse model of HD and its depletion improved several HD-like phenotypes in this mouse model. These effects are mediated, at least in part, by the CK2α'-dependent phosphorylation of α-syn serine 129 in the striatum of HD mice, a post-translational modification associated with synucleinopathy and PD. Our study highlights a potential convergent mechanism of neurodegeneration between HD and PD and suggests targeting CK2α' as a therapeutic strategy to ameliorate neuroinflammation and synaptic dysfunction in HD and perhaps other neurodegenerative diseases.

Introduction

Huntington's disease (HD) is a neurodegenerative disorder that manifests with progressive motor, cognitive, and psychiatric deficits for which there is no cure. HD is caused by a poly-glutamine (polyQ) expansion in exon 1 of the Huntingtin (*HTT*) gene. This mutation results in progressive misfolding and aggregation of mutant HTT protein (mtHTT) and preferentially affects GABAergic medium spiny neurons (MSNs) in the striatum (1-3). Transcriptional perturbations in synaptic genes and neuroinflammation are key processes that precede MSN death and motor symptom onset (4). However, our understanding of the interplay between these processes, mtHTT aggregation, and their contribution to MSN susceptibility in HD is still incomplete.

Protein kinase CK2 is at the crossroads between neuroinflammation, protein aggregation, and synaptic activity, and has recently emerged as a potential therapeutic target of neurodegeneration (5, 6). CK2 is a highly conserved serine/threonine kinase composed of two regulatory beta (CK2β) subunits and two catalytic subunits, alpha (CK2α) and alpha prime (CK2α') (7, 8). The two catalytic subunits share high structural homology, but they differ in their tissue distribution and their ability to phosphorylate different substrates (9, 10). Our previous work showed that CK2α', but not CK2α, is induced in HD MSNs and is responsible for the phosphorylation and degradation of the heat shock transcription factor HSF1, the master regulator of chaperones and protein quality control systems (11, 12). CK2α'-dependent degradation of HSF1 contributes to chaperone dysregulation and HTT aggregation in cells and mouse models of HD (11). However, other studies conducted *in vitro* have suggested a protective role of CK2 in HD (13, 14), imposing the necessity to clarify the specific involvement of CK2α' in HD pathogenesis and its potential as a therapeutic target in HD.

CK2 is involved in the phosphorylation and aggregation of HTT (11, 14), as well as the phosphorylation and aggregation of other pathological proteins like microtubule associated protein tau (MAPT) and alpha-synuclein (α-syn), proteins involved in Alzheimer's (AD) and Parkinson's disease (PD) (15-17). Tau and α-syn contribute to the activation of neuroinflammatory processes, transcriptional dysregulation, and synaptic deficits in AD and PD [18, 19].These pathophysiological alterations are also observed in HD as well as the association of tau and α-syn in HD pathology (20, 21). Specific tau modifications and cleavage products associated with cognitive decline in AD were found in the brains of HD patients (21, 22), while increased levels of α-syn were observed in the plasma of patients with HD (23, 24). α-syn colocalized with mtHTT aggregates in the brains of R6/1 HD mice, and its deletion resulted in amelioration of motor deficits (20, 25). However, the mechanisms by which these proteins are altered in HD and the extent to which they contribute to HD pathophysiology is still unknown.

In this study, we characterized the role of CK2α' in HD *in vivo* by using the heterozygous zQ175 HD mouse model that lacks one allele of CK2α'. We showed that CK2α' haploinsufficiency decreased the levels of pro-inflammatory cytokines and improved astrocyte health, restored synaptic gene expression and excitatory synapse function, and improved motor behavior in zQ175 mice. These neuropathological and phenotypic changes correlated with alterations in α-syn serine 129 phosphorylation

(pS129-α-syn) in the striatum, a post-translational modification involved in α-
synucleinopathy, establishing a novel connection between CK2α' function and
synucleinopathy in HD. Collectively, our data demonstrated that CK2α' plays a negative
role in HD and indicates the therapeutic potential of modulating CK2α' to achieve
enhanced neuronal function and neuroprotection.

Results

Increased CK2α' levels in the striatum of zQ175 mice parallel progressive HTT
aggregation and NeuN depletion

Increased CK2 activity has been associated with detrimental effects in protein
homeostasis and neuroinflammation in different neurodegenerative diseases, but its
role in HD is still controversial (11, 13, 14). To determine whether CK2α' plays a
negative role during HD pathogenesis, we first evaluated the relationship between HTT
aggregation, neuronal loss, and CK2α' levels in the striatum over time for the
heterozygous zQ175 mouse model at 3 (pre-symptomatic), 6 (early symptomatic), 12
(symptomatic), and 22 months (late-stage disease) of age (26, 27). We observed an
age-dependent increase of HTT aggregates (EM48$^+$ puncta) and a decreased number
of NeuN$^+$ neurons (pan-neuronal marker) in the striatum of zQ175 mice (**Fig. 1A-D,**
S1A). We confirmed that the depletion of NeuN$^+$ cells correlated with decreased Ctip2$^+$
neurons (MSN marker) (28)(**Fig. S1B, C**). However, we did not observe a significant
difference in the total number of neurons, measured by cresyl violet (**Fig. S1D-F**), or in
striatum volume (**Fig. S1G, H**), suggesting that changes in NeuN and Ctip2 reactivity
may reflect transcriptional dysregulation and/or neuronal dysfunction rather than
neuronal loss.

The levels of CK2α' increased over time in zQ175 mice in the striatum but not in the
cortex (**Fig. 1E-G**), coinciding with HTT aggregation and preceding NeuN depletion.
Regression analysis demonstrated that CK2α' levels had a significant positive
relationship with HTT aggregation (Pearson r(22)=0.87, p value<0.001) (**Fig. 1H**) and a
significant negative relationship with the number of NeuN$^+$ cells (Pearson r(22)=-0.78, p
value<0.001) (**Fig. 1I**).

CK2α' haploinsufficiency reduced production of inflammatory cytokines in zQ175

CK2 regulates signaling pathways involved in inflammation (29). Inhibition of CK2 has
previously shown to deplete pro-inflammatory cytokines such as IL-6 (6). IL-6 and other
cytokines are increased in HD and are often used as a marker of disease progression
(30). To determine whether the expression of CK2α' is directly related to neuronal-
mediated inflammation in HD, we first analyzed the levels of IL-6 in the striatal-derived
mouse cells ST*Hdh*$^{Q7/Q7}$ (control) and ST*Hdh*$^{Q111/Q111}$ (HD cells) (**Fig. 2A**). RT-qPCR
analysis showed an upregulation of CK2α' in HD cells which paralleled an upregulation

of IL-6. CK2α' knock-down in HD cells reduced mRNA levels of IL-6 when compared to
non-targeting siRNA (scramble) (**Fig. 2B**).

To further explore the role of CK2α' in modulating inflammation in HD, we used the
zQ175 mouse model lacking one allele of CK2α' (zQ175:CK2α'[(+/-)]) (**Fig. 2C, D**) which
has previously shown reduced HTT aggregation [11]. We analyzed the levels of 40
different cytokines and chemokines in the striatum of 12-14 month old mice (**Fig. 2E,**
S2A-C) and found seven cytokines (IFN-γ, IL-1α, IL-6, IL-7, IL-16, IL-17, IL-23, and
TNF-α) were decreased in zQ175:CK2α'[(+/-)] compared to zQ175 (**Fig. 2F**). This
suggests CK2α' contributes to the characteristic inflammatory profile of HD.

Inflammatory cytokines can be produced by both neurons and glial cells. Microglia are
pivotal regulators of CNS inflammatory processes and their activation has been
reported in numerous HD mouse models and in patients with HD (31-33). We evaluated
the levels of the microglial marker Iba1 (Ionized calcium binding adaptor molecule 1) by
immunoblotting (IB) and immunofluorescence (IF) in striatum extracts at 12-14 months
(**Fig. 2G, H, S2D-F**). Total Iba1 levels were significantly increased in zQ175 and
zQ175:CK2α'[(+/-)] mice compared to WT, but no significant differences were observed
between zQ175 and zQ175:CK2α'[(+/-)] mice. In addition, no changes were observed in
the number or area size of Iba1[+] cells across all genotypes (**Fig. S2D-F**) and no
expression of IL-6 was seen in Iba1[+] cells (**Fig. S3**), indicating that CK2α'-mediated
changes of pro-inflammatory cytokine levels may not be attributed to changes in
microglia. We then assessed whether astrocytes could contribute to inducing cytokines
in HD by co-staining IL-6 with different astrocyte markers (S100b, S100 Calcium
Binding Protein B; GFAP, Glial fibrillary acidic protein; and GS, glutamine synthetase)
(**Fig. S3**). Besides Ctip2, only GS[+] astrocytes showed IL-6 immunoreactivity in zQ175
mice.

CK2α' contributes to astrocyte pathology in HD

Increases in the number of astrocytes and astrocytic dysfunction are reported in several
neurodegenerative diseases(34), including HD (35). As such, we tested whether the
changes in cytokine levels due to CK2α' haploinsufficiency were related to changes in
GS[+] astrocytes during HD pathogenesis. We observed a progressive increase in the
number of GS[+] astrocytes in the striatum of zQ175 mice compared to WT, and a
decrease in zQ175:CK2α'[(+/-)] mice (**Fig. 3A, B**). To confirm the changes observed in
astrocyte number in brain slices, we performed proton magnetic resonance
spectroscopy ([1]H-MRS) *in vivo* at 22 months and measured the concentration of the
astroglia marker *myo*-inositol (Ins) (36, 37) and 14 additional brain metabolites (**Fig. 3C,**
D, S4A, B). The concentration of different neuronal metabolites including N-
acetylaspartate (NAA) and glutamate, which are dysregulated in HD (26, 38, 39), did not
differ between zQ175 and zQ175:CK2α'[(+/-)] mice (**Fig. S4A, B**). However, the levels of
myo-inositol (Ins) showed a significant reduction in zQ175:CK2α'[(+/-)] compared to zQ175
mice (**Fig. 3D**).

Cytokines like IL-1α, TNF-α, and complement component 1q (C1q) can induce the
accumulation of the complement component C3d protein, a protein involved in innate
immunity, altering astrocyte function and promoting astrocyte-dependent cytotoxicity
during aging and neurodegeneration (40, 41). Increased depositions of C3d have been
observed in the striatum of patients with HD and are considered a marker of astrocyte
pathology (40, 42, 43). Since depletion of CK2α' decreased IL-1α and TNF-α levels, we
evaluated the levels of C3d by IF and observed a significant age-dependent increase in
the intensity of C3d in zQ175 mice compared to WT mice which was lower in
zQ175:CK2α'$^{(+/-)}$ mice (**Fig. 3E, F**). C3d is expressed by different striatal components
other than astrocytes ^{42} so we determined the fraction of GS$^+$ astrocytes that showed
C3d depositions (GS$^+$C3d$^+$) (**Fig. 3E, E', G**). The percentage of GS$^+$C3d$^+$ astrocytes
showed an age-dependent increase for all genotypes. The levels of C3d progressively
increased in zQ175 mice (**Fig. 3G**), coinciding with the timing of CK2α' accumulation in
MSNs (**Fig. 1C**). In contrast, zQ175:CK2α'$^{(+/-)}$ showed a significant reduction in
GS$^+$C3d$^+$ astrocytes compared to zQ175 mice at 12 and 22 months and no significant
differences compared to WT mice (**Fig. 3G**), suggesting that neuronal accumulation of
CK2α' contributes to astrocyte pathology in HD.

Depletion of CK2α' improves neuronal function and motor behavior

Astrocyte pathology contributes to progressive loss of striatal excitatory synaptic
transmission and behavioral deficits associated with HD $^{(44, 45)}$. This is supported by
studies where conditional mutant HTT knock-out in astrocytes improved astrocyte
function and restored both synaptic protein expression and MSN excitability in the
BACHD mouse model (46). We assessed the impact of CK2α' depletion in AMPA-
mediated excitatory transmission by conducting whole-cell patch clamp recordings from
acute dorsolateral striatum coronal slices at 12 months (**Fig. 4A**). MSNs from all
genotypes showed similar profiles in the analysis of basal synaptic transmission,
including input/output curves, paired-pulse facilitation, and synaptic fatigue (**Fig. 4B-D**).
We observed a trend towards increased normalized excitatory postsynaptic currents
(EPSCs) in zQ175:CK2α'$^{(+/-)}$ mice compared to the other two genotypes, but the data
did not reach statistical significance (**Fig. 4D**). Spontaneous neurotransmitter release
and synaptic activity via miniature EPSC (mEPSC) recordings showed that mEPSC
amplitude, reflecting postsynaptic AMPA receptor function, was comparable among the
3 genotypes (**Fig. 4E**). However, mEPSC frequency, which reflects the probability of
neurotransmitter release from presynaptic vesicles and also correlates with the number
of synapses, was reduced in zQ175 mice (**Fig. 4F, G**), as previously reported (47), and
rescued in zQ175:CK2α'$^{(+/-)}$. CK2α' haploinsufficiency did not alter the number of MSNs
(Ctip2$^+$ cells) or the mRNA levels of the MSN markers (Drd1 and Drd2), but increased
the levels of synaptic proteins like the scaffold protein Dlg4 (PSD-95) and Ppp1rb1
(dopamine- and cAMP-regulated neuronal phosphoprotein DARPP-32), a key regulator
of the electrophysiological responses in striatal neurons (48, 49)(**Fig. S5A-C**). These
data supported that astrocyte health can influence neuronal synaptic protein levels and
excitability.

Glutamatergic synaptic transmission is often related to motor and cognitive function in
HD mouse models (45, 49). We conducted a series of motor tests including accelerating
rotarod and beam walk in WT, zQ175, and zQ175:CK2α'$^{(+/-)}$ mice at 3, 6, and 12 months
(**Fig. 5**). We also conducted cylinder and open field assessments on a different cohort
at 12 months comparing zQ175 and zQ175:CK2α'$^{(+/-)}$ (**Fig. S6**). We did not observe
significant differences between WT and zQ175 or between zQ175 and zQ175:CK2α'$^{(+/-)}$
at any tested age in the accelerating rotarod test (**Fig. 5A**), open field, or cylinder test
(**Fig. S6**). However, when we evaluated fine motor coordination and whole body
balance in the beam test, we observed a significant increase in foot slips of zQ175 mice
compared to WT at 3 months, but only with the most challenging beam (small round),
indicating early subtle motor deficits (**Fig. 5B**). At 12 months, zQ175 mice showed
increased foot slips in both the small round and small square beams compared to WT,
highlighting a worsening motor deficit. zQ175:CK2α'$^{(+/-)}$ mice showed a significant
reduction in foot slips compared to zQ175 mice at all tested ages and no significant
differences compared to WT.

We also performed tests to evaluate associative learning (fear conditioning), spatial
learning and memory (Barnes maze, BM), cognitive flexibility (BM reversal), and spatial
working memory (Y radial arm maze) by comparing zQ175 and zQ175:CK2α'$^{(+/-)}$ mice at
12 months of age, but no significant differences were observed between the two groups
(**Fig. S7**). This observation suggests that the positive effects of CK2α' depletion on
motor behavior may not additionally translate to improved cognitive functions.

CK2α' depletion rescued gene expression of synaptic genes connected to alpha-
synuclein

To elucidate the mechanism by which CK2α' contributes to HD, we investigated which
molecular pathways associated with HD could be specifically affected by CK2α'
haploinsufficiency. RNA-seq in the striatum of WT, zQ175, CK2α'$^{(+/-)}$, and
zQ175:CK2α'$^{(+/-)}$ mice at 12-14 months, followed by Weighted Gene Co-Expression
Network Analysis (WGCNA), revealed that the mouse transcriptome could be clustered
into 20 gene co-expression modules given the sample phenotypes (**Fig. S8, Table S1**).
Out of the 20 modules, 9 modules showed a significant difference in eigengene
expression between zQ175 and WT mice in a Kruskal-Wallis test (p value < 0.05) (**Fig.**
S9A, Table S2). Further evaluation of those 9 modules revealed 2 modules
(Greenyellow: 255 genes, and Red: 639 genes) were also significantly different between
zQ175 and zQ175:CK2α'$^{(+/-)}$ mice (p value < 0.05) (**Fig. 6A, Table S2**). Plots of module
eigengene expression confirmed that Greenyellow (**Fig. 6B**) and Red modules (**Fig. S8**)
were more consistent in the expression contrast between zQ175 mice and the rest of
the genotypes. Cook's distance (DESeq2) analyses revealed that these differences
were not due to the presence of outliers in our data set (**Fig. S9B**). Since the
Greenyellow module demonstrated a higher significance over the Red module (**Fig.**
6A), we decided to focus our analyses on the Greenyellow module. Ingenuity pathway
analysis (IPA) indicated that the five most significant canonical pathways in the
Greenyellow module were synaptogenesis signaling pathways (p-value 1.68E-06),

Ephrin A signaling (p-value 7.84E-05), glutamate receptor signaling (p-value 1.98E-04),
axonal guidance signaling (p-value 7.13E-04), and G-protein coupled receptor (GPCR)
signaling (p-value 1.14E-03) (**Fig. 6C, Fig. S9C**), all of which have been previously
implicated in HD pathogenesis [50]. Additional Gene Ontology (GO) annotation of cellular
components indicated that Greenyellow genes are significantly enriched in synaptic
components (**Fig. 6D**).

Connectivity analyses (showing only 15% of the most highly connected members of the
Greenyellow module) (**Fig. 6E**) revealed that the two most connected genes within the
hub were Slit1 (Slit Guidance Ligand 1), associated with "poor" behavior and a worse
prognosis in the R6/1 mouse mode (51), and Ncald (Neurocalcin delta), which regulates
multiple endocytosis-dependent neuronal functions and is situated on a locus that has
been associated with earlier clinical onset of HD [52, 53]. To better understand which
specific genes were more impacted in their expression by CK2α' depletion, we
examined the Differential Gene Expression (DGE) between control mice (WT and
CK2α'$^{(+/-)}$) and HD mice (zQ175 and zQ175:CK2α'$^{(+/-)}$) (**Table S3**). The DGE between
WT and zQ175 mice confirmed a large transcriptional dysregulation (n=885 genes,
Q<0.1) (**Fig. S9D, E Table S3**), as previously reported [50]. Interestingly, the DGE
between zQ175:CK2α'$^{(+/-)}$ and WT mice only reported 123 genes, in contrast to the 885
genes between zQ175 and WT (**Fig. S9F, G**). Since our sample distribution was not
equally sex balanced, we then used R package variance Partition and confirmed that
these expression changes were driven only by 'genotype' and not by 'sex' (**Fig. S9H**).

CK2α' depletion decreased inflammation and improved astrocyte health. Therefore, we
searched for microglial and astrocytic inflammatory RNA signatures in our data set (40).
As Castro-Diaz et al., showed (43), we did not observe significant changes in the
expression of these gene signatures even when comparing WT and zQ175 mice (**Fig.
S10A, B, Table S4, S5**). The disconnection between changes in protein levels of
inflammatory cytokines and the absence of transcriptional alterations in those genes
could be explained by post-translational events potentially regulated by CK2α'.
However, we recapitulate some transcriptional changes for genes of the HD-associated
astrocyte molecular signature in the zQ175 mice (43, 54), indicative of astrocytic
dysfunction (**Fig. S10C**). Notably, these changes were not observed when comparing
zQ175:CK2α'$^{(+/-)}$ and WT mice.

The DGEs analyzed between zQ175 and zQ175:CK2α'$^{(+/-)}$ mice revealed 39 specific
and significant genes (FDR<0.1) (**Fig. 6F, Table S6**), including Csnk2a2 (CK2α' gene)
as a positive control. Three genes (Ncald, Nrp2, and Slc30a3) were also among the
15% most highly connected members of the Greenyellow module (**Fig. 6E**). At least
40% of the DGEs (n=16) were related to synaptic functions (**Fig. 6F, Table S6**). We
performed an IPA on the 39 genes to query for biological meaning and found that the
most significant canonical pathway was for glutamate receptor signaling (p-value 2.09E-
03) (**Fig. 6A**). These pieces of data confirm the contribution of CK2α' to the
dysregulation of genes related to excitatory synaptic transmission in HD.

When looking at the most significant upstream regulators identified by IPA of both the
Greenyellow module and the 39 gene set identified by DGE, we found SNCA (alpha-

synuclein) (p-value 9.10E-11 and 1.03E-07, respectively). α-syn is a neuronal protein
that regulates multiple processes, including synaptic vesicle trafficking and
neurotransmitter release and transcription, and is associated with onset and
progression of Parkinson's disease (55, 56). α-syn was also shown to exert a toxic
effect in R6/1 mice, but its mechanism of action has remained unknown(20, 25). IPA
connected α-syn with some of the most differentially dysregulated genes in the set of 39
genes associated to synaptic functions including Ttr (Transthyretin), Grm2 (Glutamate
Metabotropic Receptor 2), Slc17a7 (Solute Carrier Family 17 Member 7; alias VGlut1),
C1ql3 (Complement Component 1, Q Subcomponent-Like), Cckbr (cholecystokinin B
receptor), Nrp2 (Neuropilin 2), and the transcription factors Tbr1 (T-Box Brain
Transcription Factor 1) and Nr4a2 (Nuclear Receptor Subfamily 4 Group A Member 2;
alias Nurr1) (**Fig. 6G, F**).

To determine the extent to which α-syn participates in the regulation of genes identified
in our IPA analyses (**Fig. 6G**), we used R6/1:SNCA^{KO} mice (**Fig. S11**), previously
shown to improve in motor behavior compared to R6/1 mice [20]. RT-qPCR showed
significant expression changes for SNCA, Nrp2, Cckbr, and Ttr between WT and R6/1
mice. Lack of SNCA significantly decreased the expression of Slc30a3 and Grm2, and
showed a trend toward decreased expression for Ttr and Cckbr in R6/1:SNCA^{KO}
compared to R6/1 mice. In addition, nearly all tested genes decreased their expression
when comparing R6/1:SNCA^{KO} to WT:SNCA^{KO}. Altogether, this data suggested that
transcriptional dysregulation of some synaptic genes in HD could be mediated by α-syn
dysfunction. Although some gene expression changes differed between R6/1 and
zQ175 mice, the effects mediated by SNCA^{KO} demonstrated a role of α-syn in the
regulation of the genes identified in our IPA analyses in HD.

Striatal synucleinopathy is found in zQ175 mice and is reduced by CK2α'
depletion

We next explored whether CK2α' was involved in the regulation of α-syn in HD. We
observed that the total amount of α-syn was similar between WT and zQ175 (**Fig. 7A-C**)
mice. zQ175:CK2α'$^{(+/-)}$ mice showed a trend towards increased α-syn, but did not reach
statistical significance (**Fig. 7B, C**). Nuclear and cytoplasm fractionation confirmed the
presence of α-syn in nuclear fractions from striatum samples (55, 57), and showed a
modest but significant increase in nuclear α-syn in zQ175 mice (**Fig. 7D, E**). IF analyses
for α-syn and HTT (EM48) also confirmed the colocalization between these two
proteins, as previously shown in R6/1 mice (20)(**Fig. 7F, G**). To determine if there was a
difference in the number and distribution of co-localized α-syn/HTT, we first analyzed
the number of EM48$^+$ puncta in both the nucleus and cytoplasm between zQ175 and
zQ175:CK2α'$^{(+/-)}$ mice. Cytoplasmic HTT aggregates were reduced in zQ175:CK2α'$^{(+/-)}$
compared to zQ175 mice, consistent with previous studies (11), although no significant
differences were observed in the number of nuclear HTT aggregates (**Fig. 7H, I**).
Despite the decrease in cytoplasmic HTT aggregates in zQ175:CK2α'$^{(+/-)}$ mice, no
significant differences were observed in the number of nuclear and/or cytoplasmic α-
syn/HTT colocalized puncta between zQ175 and zQ175:CK2α'$^{(+/-)}$ mice (**Fig. 7J**).

We then evaluated whether pS129-α-syn, a marker of synucleinopathy (58, 59), was
altered in HD and whether CK2α' could influence its levels. We observed that the levels
of pS129-α-syn increased in the striatum of zQ175 mice at 12 months compared to WT
(tested with 3 different pS129-α-syn antibodies: 81A, EP1536Y and D1R1R), and in the
striatum of patients with HD (**Fig. 8A-E, Fig. S12A**), indicating signs of synucleinopathy.
The levels of pS129-α-syn were significantly reduced in zQ175:CK2α'$^{(+/-)}$ mice
compared to zQ175, while no significant differences were observed with WT mice (**Fig.**
8D, E, Fig. S12A-C). pS129-α-syn was detected in both the cytoplasm and the nucleus
of zQ175 striatal cells, while no nuclear presence was detected in zQ175:CK2α'$^{(+/-)}$ mice
(**Fig. 8F, G**). In addition, we observed that pS129-α-syn colocalized with both
cytoplasmic and nuclear HTT puncta in zQ175 mice, while only cytoplasmic
colocalization was observed in zQ175:CK2α'$^{(+/-)}$ mice (**Fig. 8F, G, Suppl. Video 1**).

Discussion

Increased protein kinase CK2 activity has been associated with detrimental effects in
protein homeostasis and neuroinflammation in different neurodegenerative diseases,
including AD and PD (5). However, the role of CK2 in HD remained unclear. This study
demonstrated the adverse effects of the catalytic subunit CK2α' on neuroinflammation,
protein aggregation, neuronal function, and motor behavior in the zQ175 HD mouse
model and consolidated the important contribution of CK2α' to HD pathogenesis. We
also found this contribution is mediated, at least in part, by the ability of CK2α' to
phosphorylate α-syn and induce striatal synucleinopathy and synaptic gene
dysregulation (**Fig. 8H**).

We previously showed that CK2α' levels were induced in the striatum of patients with
HD, in MSN-like human iPSCs derived from patients with HD, and in zQ175 mouse
MSNs (11). CK2α' upregulation was associated with HD pathogenesis since CK2α'
genetic knockdown in different HD cell models resulted in decreased HTT aggregation
and increased cell viability(11). However, other studies using CK2 inhibitors suggested
CK2 induction as a protective mechanism in HD (13, 14). These conflicting reports
could be explained by the poor selectivity and high toxicity of previously used CK2
inhibitors (60). Our study demonstrated that reducing levels of CK2α' by removing one
allele of the Csnk2a2 (CK2α') gene in the zQ175 heterozygous HD mouse model
(zQ175:CK2α'$^{(+/-)}$) had significant and long-term benefits in several HD-related
phenotypes. CK2α' depletion significantly altered the production of several pro-
inflammatory cytokines in the striatum of symptomatic zQ175 mice. The benefits of
reducing neuroinflammation in HD have been shown in R6/2 mice intracranially injected
with a TNF-α inhibitor, which resulted in a significant increase in neuronal density and
improved motor function [61]. We obtained similar results when reducing the expression
of CK2α', suggesting CK2α' as a potential upstream regulator of neuroinflammation in
HD.

Neuroinflammation is a coordinated response to brain toxicity, involving the activation of
microglia and astrocytes. Increased reactive microglia were reported in numerous HD

mouse models [31, 33, 62] and in pre-symptomatic patients with HD (31, 32), and are often
associated with increased pro-inflammatory cytokines (4). However, our microglia
analyses failed to show robust microgliosis in zQ175 mice, consistent with previous
studies (43), and did not indicate microglia as a relevant cell type responsible for the
inflammatory profile observed in zQ175 or for the changes mediated by CK2α'
depletion. On the contrary, we observed increased astrogliosis in zQ175 mice,
manifested by an increased number of astrocytes and astrocytic pathology, which
decreased in zQ175:CK2α'$^{(+/-)}$. Although it is unknown how CK2α' influences astrocyte
proliferation, it has been shown that specific cytokines like TNF-α and IL-6 have a
proliferative effect on astrocytes (63) and therefore, CK2α'-mediated changes in these
cytokines could be responsible for these phenotypes in HD mice.

Accumulation of C3 and the C3 cleaved product, C3d, in HD astrocytes is canonically
considered a marker of astrocytic pathology associated with neuroinflammation (40, 42,
43). Liddelow et al., reported that astrocytes react to the presence of specific cytokines,
such as IL-1α, TNF-α, and C1q, by changing their transcriptional program and adopting
a neurotoxic phenotype (A1), which is also characterized by the accumulation of C3d
(40). Transcriptomic studies in purified HD astrocytes [43] and our own RNA-seq data
have failed to identify transcriptional changes characteristic of A1 astrocytes, but
instead revealed a transcriptional signature related to astrocytic dysfunction that is
associated with the activation of C3d. In AD, expression of C3d in astrocytes is induced
by neuroinflammatory processes triggered by amyloid-beta, and its activation
contributes to protein aggregation and cell death(64). CK2 phosphorylates C3 and
modulates its cleavage and activation(65). Therefore, CK2α' levels, as well as CK2α'-
mediated neuroinflammation, could influence C3d accumulation in astrocytes and
astrocytic pathology.

Increased pro-inflammatory cytokines can alter synaptic strength as well as
glutamatergic transmission and are also associated with structural and functional
disruption of synaptic compartments (66). Similarly, astrocyte dysfunction in HD
contributes to reduced striatal glutamatergic transmission and spine density, ultimately
decreasing MSN excitability(35, 46). We showed that CK2α'-dependent decrease of
inflammatory cytokines and astrogliosis correlated with increased expression of synaptic
proteins (PSD-95 and Darpp-32) and improved AMPA-mediated synaptic transmission.
Similarly, conditional deletion of mtHTT in astrocytes of BACHD mice rescued the
expression of synaptic proteins and synaptic activity in the striatum and improved
astrocyte health(46). However, whether decreased astrocytic pathology contributes to
improved neuronal activity in zQ175:CK2α'$^{(+/-)}$ mice is yet to be determined.

In contrast to CK2α, which is an essential protein with hundreds of targets, CK2α' has
very few identified substrates (9, 67). We previously showed the stress protective
transcription factor HSF1 as a substrate of CK2α' (11). HSF1 is the master regulator of
protein homeostasis in mammalian cells and has also been linked to the regulation of
anti-inflammatory and synaptic processes (12). CK2α' phosphorylates HSF1, signaling
HSF1 for proteasomal degradation and influencing chaperone expression and HTT
aggregation in cells and in HD mice (11). Our RNA-seq analysis validated the increased
expression of chaperones like Hsp70 and Hsp25 in the zQ175:CK2α'$^{(+/-)}$ mice,

consistent with previous findings (11). However, WGCNA and DGE did not reveal global
changes in specific transcriptional pathways associated with protein quality control
networks between zQ175 and zQ175:CK2α'$^{(+/-)}$, but instead showed two different
modules (Greenyellow and Red) primarily corresponding to signaling pathways related
to synaptogenesis and glutamate receptor signaling. SNCA (α-syn) was indicated by
IPA as one of the top putative upstream regulators of the CK2α'-mediated
transcriptional changes observed by WGCNA and DGE analyses. Although α-syn is not
a transcription factor, several reports have shown α-syn has the ability to modulate the
expression of transcription factors like Nurr1, which is involved in the regulation of
inflammation and is differentially expressed between zQ175 and zQ175:CK2α'$^{(+/-)}$ mice
(55, 56). α-syn aggregation is thought to be a key process in PD and other
synucleinopathies and is promoted by the presence of high levels of α-syn
phosphorylation and ubiquitylation (59). In particular, pS129-α-syn phosphorylation
regulates neurotransmitter uptake, notably dopamine as well as glutamate release,
thereby playing an important role at synaptic terminals and modulating synaptic
plasticity (59, 68).

The specific contribution of α-syn dysregulation in HD has yet to be determined. It is
known that α-syn participates in HD pathogenesis because α-syn KO mice have
demonstrated decreased mtHTT aggregation and attenuated body weight loss and
motor symptoms in R6/1 mice (20, 25). Interestingly, aggregation of α-syn and
consequent synucleinopathy in PD has been linked to the activation of CK2 and CK2-
dependent phosphorylation, although it is unknown which CK2 subunit participates in
this process (69). α-syn has also been shown to influence cytokine production (70).
Therefore, it is reasonable to hypothesize that CK2α' could contribute to HD by
increasing HTT aggregation via dysregulation of protein quality control mechanisms
and/or increasing α-syn phosphorylation. Here, we showed increased pS129-α-syn in
the striatum of symptomatic HD mice and patients with HD as well as an increased
localization of this phosphorylated form in the nucleus. Furthermore, we demonstrated
that CK2α' is involved in this process. Decreased phosphorylation and the accumulation
of pS129-α-syn in the nucleus could be responsible for the rescue of α-syn-associated
transcriptional alterations and the improvement of synaptic function, cytokine
expression, and astrocytic pathology seen in zQ175:CK2α'$^{(+/-)}$ mice. Overall, these
results provide strong evidence for CK2α' playing an important role in many aspects of
HD pathogenesis, supporting its potential as a therapeutic target in HD. Further
experiments will be necessary to decipher the mechanism by which CK2α'-mediated α-
synucleinopathy contributes to HD and to tease apart the differential contribution of HTT
aggregation and α-syn pathology to the symptomatology, onset, and progression of HD.

Materials and Methods

Cell lines

Mammalian cell lines used in this study were the mouse-derived striatal cells STHdh^{Q7}
and STHdh^{Q111} (Coriell Cell Repositories). Cell lines were authenticated by

immunoblotting for the detection of mutant HTT protein. Cells were grown at 33ºC in
Dulbecco's modified Eagle's medium (DMEM, Genesee) supplemented with 10% fetal
bovine serum (FBS), 100 U ml^{-1} penicillin/streptomycin and 100 ug ml^{-1} G418 (Gibco),
as previously described (11).

Mouse strains
All reported data abides to the ARRIVE guidelines for reporting animal research. For
this study we used a full-length knock-in mouse model of HD known as zQ175 on the
C57BL/6J background (B6J.zQ175 KI mice (Stock No. 027410)), which harbors a
chimeric human/mouse exon 1 carrying an expansion of ~188 CAG repeats and the
human poly-proline region (27). WT (C57BL/6) animals were used as controls. CK2α′
heterozygous mice (CK2α′$^{(+/-)}$) on the 129/SvEv-C57BL/6J background (Taconic
Biosciences TF3062) were originally obtained from Dr. Seldin (Boston University) (71).
CK2α′$^{(+/-)}$ were crossbred with WT and zQ175 for more than 10 generations prior to
selecting the founder animals of the study. All mice used have the C57BL/6J
background. All mice were housed under standard SPF conditions in a conventional
12:12 *light*/*dark* cycle. All experiments were conducted during the light cycle. Mice were
bred as previously described [11] to generate the following genotypes: WT (CK2α′$^{(+/+)}$
HTT$^{(0/0)}$), CK2α′$^{(+/-)}$ (CK2α′$^{(+/-)}$ HTT$^{(0/0)}$), zQ175 (CK2α′$^{(+/+)}$ HTT$^{(Tg/0)}$), zQ175;CK2α′$^{(+/-)}$
(CK2α′$^{(+/-)}$ HTT$^{(Tg/0)}$). Animals were analyzed at 3, 6, 12, and 22 months of age. We also
used samples from 5 month WT (mixed background CBA x C57BL/6), R6/1, SNCA^{KO},
and R6/1SNCA^{KO} obtained from Dr. Lucas. Sample size was set to a minimum of three
animals per genotype for every analysis. When possible, at least two females and two
males were used for each genotype and age. Littermates of the same sex were
randomly assigned to experimental groups. All animal care and sacrifice procedures
were approved by the University of Minnesota Institutional Animal Care and Use
Committee (IACUC) in compliance with the National Institutes of Health guidelines for
the care and use of laboratory animals under the approved animal protocol 2007-
38316A.
529

siRNA transfection

For CK2α' knock-down, ST*Hdh* cells were transfected with FlexiTube siRNA (5 nmol)
from Qiagen (Mm_Csnk2a2; SI00961051; SI00961058; SI00961065; SI00961072)
using DharmaFECT1 per manufacturer's guidelines. As a negative control, ON-
TARGETplus control Non-targeting pool siRNA (Dharmacon) was used. Cells were
collected 24 h after transfection for RNA extraction and RT-qPCR. All siRNAs have
been validated by RT-qPCR and immunoblotting for knockdown efficiency of the target
gene.

RNA preparation and RT-qPCR
RNA was extracted from STHdh cells and mouse striatal tissues by using the RNeasy
extraction kit (Qiagen) according to the manufacturer's instructions. cDNA for all

samples was prepared from 1 μg RNA using the Superscript First Strand Synthesis
System (Invitrogen) according to the manufacturer's instructions. SYBR green based
PCR was performed with SYBR mix (Roche). The primer sequences for target genes
were as follows: CK2α' forward: 5'- CGACTGATTGATTGGGGTCT-3' reverse: 5'-
AGAATGGCTCCTTTCGGAAT-3', IL-6 forward: 5'- AGTTGCCTTCTTGGGACT-3'
reverse: 5'-TCCACGATTTCCCAGAGAAC-3', PSD-95 (Dlg4) forward: 5'-
CCGACAAGTTTGGATCCTGT-3', reverse: 5'-ACGGATGAAGATGGCGATAG, Drd1
forward: 5'-AAGATGCCGAGGATGACAAC-3', reverse:5'-
CCCTCTCCAAAGCTGAGATG-3', Drd2 forward: 5'-TATGCCCTGGGTCGTCTATC-3',
reverse: 5'-AGGACAGGACCCAGACAATG-3', Darpp32(PPP1R1B) forward: 5'-
CCACCCAAAGTCGAAGAGAC-3', reverse: 5'-GCTAATGGTCTGCAGGTGCT-3',
GAPDH (used as an internal control gene) forward: 5'-AACTTTGGCATTGTGGAAGG-
3' reverse: 5'-ACACATTGGGGGTAGGAACA-3'. The qPCR amplification was
performed using the LightCycler 480 System (Roche). Each sample was tested in
triplicate and normalized to GAPDH levels. For analysis, the $2^{\Delta\Delta Ct}$ method was used to
calculate the relative fold gene expression of samples.
558

Immunoblot analysis

Sample preparation and immunoblotting condition were performed as previously
described (11). Striatum protein extracts from one hemisphere of mice were prepared in
cell lysis buffer (25 mM Tris pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% Triton-X100 and
0.1% SDS). Extra SDS was added to the suspension to a final concentration of 2% (w
per v) and lysates heated at 95°C for 5 min to solubilize tissue. Total tissue homogenate
was then centrifuged at 12,000 r.p.m. for 10 min. Protein samples were separated on 4–
20% SDS Criterion TGX Stain-Free gels (BioRad) at 110 V. Proteins were transferred to
a nitrocellulose membrane (BioRad 0.2 μm) and blocked with 5% non-fat dry milk in
TBS containing 0.25% Tween-20 (TBST) for 1 h at room temperature. Primary
antibodies were anti-CK2α' (Rabbit, Novus NB100-379 and Proteintech 10606-1-AP,
both 1:1000), anti-Iba1 (Rabbit, FUJIFILM Wako 019-19741, 1:1000), α-syn (Mouse,
Biolegend 834303, 1:1000, clone 4D6), pS129-α-syn (Mouse, Millipore MABN826,
1:500, clone 81A and Rabbit, EP1536Y Abcam ab51253, 1:1000), GAPDH (Mouse,
Santacruz sc-365062, 1:10000). Quantitative analyses were performed using ImageJ
software and normalized to GAPDH controls.

Immunohistochemistry

Mice were anesthetized with Avertin (250 mg/kg Tribromoethanol) and perfused
intracardially with 25mM Tris-base, 135mM Nacl, 3mM KCl, pH 7.6 supplemented with
7.5 mM heparin. Brains were dissected, fixed with 4% PFA in TBS at 4°C for 4-5 days,
cryoprotected with 30% sucrose in TBS for 4-5 days and embedded in a 2:1 mixture of
30% sucrose in TBS:OCT (Tissue-Tek). Immunostaining was performed as previously
described (11). Fluorescent images acquired on an epi-fluorescent microscope (Echo
Revolve) or confocal microscope (Olympus FV1000). Primary antibodies used and

dilutions are as follows: α-syn (Mouse, Biolegend 834303, 1:1000), pS129-α-syn
(Mouse, Millipore MABN826, 1:500 and Rabbit, D1R1R Cell signaling technology
23076S, 1:200), CK2α' (Rabbit, Proteintech 10606-1-AP, 1:500), Ctip2 (Rat, Abcam
ab18465, 1:500), complement component C3d (Goat, R&D Systems AF2655,
1:200), Glial fibrillary acidic protein GFAP (Rabbit, Invitrogen PA1-10019, 1:500), S100
Calcium Binding Protein B S100b (Rabbit, Abcam ab41548, 1:500), glutamine
synthetase GS (Mouse, BD Biosciences 610517, 1:1000 and Rabbit, Abcam 49873,
1:500), HTT (Mouse, Millipore, clone mEM48 Mab5374, 1:100, and Rabbit, Abcam
ab109115, 1:500), Iba1 (Rabbit, FUJIFILM Wako 019-19741, 1:200), NeuN (Mouse,
Millipore MAB377, 1:1000), IL-6 (Mouse, Santa Cruz Bio sc-32296, 1:100). For cell
number (Ctip, GS, NeuN, Iba1, DAPI) the Cell counter plugin from ImageJ software was
used and normalized to the image area ($300 \mu m^2$). EM48$^+$ aggregates were counted
using annotations in the Echo Revolve software and validated using the Puncta
Analyzer plugin in ImageJ. Number of pS129-α-syn/EM48 puncta was calculated using
the Puncta Analyzer plugin in ImageJ.

Cytokine and chemokine array

The Proteome Profiler Mouse Cytokine Array Panel (ARY006, R&D Systems) was used
to detect the levels of cytokine/chemokine of the mouse brain as per manufacturer's
instructions. Frozen striatum from 12 - 14 months was homogenized in PBS containing
Halt protease inhibitor cocktail and phosphatase inhibitors (Fisher Scientific) and 1%
triton X-100 (Sigma). Samples were stored at -80°C for 15 min, thawed and centrifuged
at 10,000 × g for 5 min to remove cell debris. A total of n=6 mice/genotype with a
female(F)/male(M) ratio (3F/3M WT, 5F/1M zQ175, and 4F/2M zQ175:CK2α'$^{(+/-)}$) were
analyzed. Samples were grouped in three different pools for each genotype: pools 1
and 2 contained three different mice per genotype, pool 3 contained a randomized
selection of 3 mice/genotype out the n=6 mice cohort. Similar numbers of male and
female mice were evaluated. Data was analyzed using ImageJ software and presented
as an average signal of three independent pool assays, calculating pairs of duplicate
spots corresponding to 40 different cytokines or chemokines.

Nuclear/Cytoplasm fractionation
616 Frozen striatum samples (~20 mg) were fractionated using the Minute^{TM} Cytosolic and
617 Nuclear Extraction Kit for Frozen/Fresh tissues (Invent Biotechnologies INC, Cat NT-
618 032) as per Manufacturer's instructions.

Cresyl Violet staining

Coronal brain slices were mounted on Fisherbrand Superfrost Plus Microscope Slide
and dried at 37 ºC overnight. Dried sections were incubated with 0.2% Cresyl violet
acetate (Sigma #C5042) pH 3.7 for 8 minutes. Samples were then dehydrated in 80%,

95%, and 100% Ethanol solution for two minutes each followed by Xylene incubation.
Slides were dried and mounted with permount (#SP15-500) and imaged on a Leica
DM2500. Neurons were identified as round light blue cells and manually counted.

Electrophysiological analyses

Acute dorsolateral striatum coronal slices (350 µm thick) were obtained from
approximately 12 months old mice using a vibratome, and processed as previously
described (72). Researchers were blind to the mouse genotype. The brain was quickly
removed after decapitation and placed in ice-cold artificial cerebrospinal fluid (ACSF).
Slices were incubated for at least 1h before use in a chamber at room temperature (21–
24 ºC) in ACSF containing (in mM): NaCl 124, KCl 2.69, KH_2PO_4 1.25, $MgSO_4$ 2,
$NaHCO_3$ 26, $CaCl_2$ 2, ascorbic acid 0.4, and glucose 10, and continuously bubbled with
carbogen (95% O_2 and 5% CO_2) (pH 7.4). Slices were then transferred to an immersion
recording chamber and superfused at 2 mL/min with gassed ACSF at 30-32 ºC and
visualized under an Olympus BX50WI microscope (Olympus Optical; Japan). To study
excitatory postsynaptic currents (EPSCs) picrotoxin (50 µM) and CGP54626 (1 µM)
were added to the solution to block the $GABA_A$ and $GABA_B$ receptors, respectively.
Whole-cell electrophysiological recordings were obtained from medium spiny neurons
(MSNs) using patch electrodes (3–10 MΩ) filled with an internal solution containing (in
mM): $KMeSO_4$ 135, KCl 10, HEPES-K 10, NaCl 5, ATP-Mg 2.5, GTP-Na 0.3 (pH 7.3).
Recordings were obtained with a PC-ONE amplifier (Dagan Instruments; Minneapolis,
MN, USA). Membrane potentials were held at −70 mV. Signals were filtered at 1 kHz,
acquired at a 10 kHz sampling rate, and fed to a Digidata 1440A digitizer (Molecular
Devices; San Jose, CA, USA). pCLAMP 10.4 (Axon Instruments, Molecular Devices;
San Jose, CA, USA) was used for stimulus generation, data display, data acquisition,
and data storage. To record evoked EPSCs, theta capillaries filled with ACSF were
used for bipolar stimulation and placed in the vicinity of the cell patched in the
dorsolateral striatum. Input–output curves of EPSCs were made by increasing stimulus
intensities from 0 to 100 µA. Paired-pulse facilitation was done by applying paired
pulses (2 ms duration) with 25, 50, 75, 100, 200, 300, and 500 ms inter-pulse intervals.
The paired-pulse ratio was calculated by dividing the amplitude of the second EPSC by
the first (PPR=EPSC-2/EPSC-1). Synaptic fatigue was assessed by applying 30
consecutive stimuli in 15 ms intervals. For miniature EPSCs (mEPSCs) tetrodotoxin
(TTX; 1 µM) was added to the solution.

Magnetic resonance imaging and spectroscopy

Animal Preparation for MR Scanning: Researchers were blinded to the genotype of the
mice during testing. Animals were induced with 3% isoflurane in a 1:1 mixture of
O_2:N_2O. Mice were secured in a custom-built mouse holder and physiological status was
monitored (SA Instruments) and recorded. Anesthesia was maintained with 1.5-2%
isoflurane to achieve a respiration rate of 70-100 breaths per minute. Body temperature

was maintained at 36-37ºC with a circulating warm water system and a heating fan
controlled by feedback received from a fiber-optic rectal thermometer. The scan session
was approximately 50 minutes for each animal. MR Protocol: All experiments were
performed on a 9.4T/31 cm scanner (Agilent), as described previously (73). See
Supplementary Methods for details.

Measurement of striatal volume

For volumetric analysis of striatum, coronal RARE images were used. The images had
an in-plane resolution of 0.125 mm x 0.125 mm and 1 mm slice thickness. All volumetric
quantifications were performed using ImageJ. The perimeter of the striatum was traced
using a polygon selection tool and the volume measurement was performed by running
Measure Stack plugin on ImageJ. All quantitative analyses were conducted blindly with
respect to mouse genotype.

Behavioral assays

Researchers at the Mouse Behavioral core at University of Minnesota were blind to the
genotypes of the mice during testing. Mice were transported from their colony room in
their home cages to the behavioral testing room, where they were allowed to acclimate
for at least one hour prior to the beginning of the experiment. See **Supplementary**
Methods for a complete description of all behavioral tests conducted in the study.

Beam Walk: The deficits of motor coordination were assessed by a beam walking test
which measures a mouse's ability to maintain balance while traversing a narrow beam
to reach a safe platform. For this, 19-mm (medium-round) or 10-mm (small-round)
diameter and 16-mm (medium-Square) or 10-mm (small-Square) width of 3 feet long
wood beams (made in house) were used and placed 19 inches above the test bench by
metal supports. Each mouse was placed on the beam at one end and allowed to walk to
the goal box. Foot slips were recorded manually when the hind paws slipped off the
beam. Testing included 3 training days with 4 consecutive trials. The performance was
measured by recording the time of traversing the beam and the number of paw faults or
slips from the beam. Maximum length of time per trial was 60 s; trials that exceeded 60
s (e.g. a mouse failed to cross the finish line within a minute) were recorded as 60 s.

Rotarod: It is a standard test of motor coordination and balance in rodents. Mice were
tested over 3 consecutive days. Each daily session included a single training trial of 15
min at 5 RPM on the rotarod apparatus (Ugo Basile). On day 3, the animals were tested
for 3 consecutive accelerating trials of 5 min with the rotarod speed changing from 5 to
50 RPM over 300 s, with an inter-trial interval of at least 15 min. The latency to fall from
the rod was recorded for each trial with mice remaining on the rod for more than 300 s
removed and scored at 300 s.

RNA-Seq Analyses

Gene expression analysis was carried out using the CHURP pipeline
(HTTps://doi.org/10.1145/3332186.3333156) using n=5 mice/genotype for WT, zQ175,
and zQ175:CK2α'$^{(+/-)}$ and n=3 mice for CK2α'$^{(+/-)}$, with a female (F)/male (M) ratio:
4F/1M WT, 1F/2M CK2α'$^{(+/-)}$, 2F/3M zQ175, 4F/1M zQ175:CK2α'$^{(+/-)}$. Read quality was
assessed using FastQC (HTTp://www.bioinformatics.bbsrc.ac.uk/projects/fastqc). Raw
reads were then trimmed to remove low quality 3' ends and adaptor contamination
using Trimmomatic with the following parameter:
ILLUMINACLIP:all_illumina_adapters.fa:4:15:7:2:true LEADING:3 TRAILING:3
SLIDINGWINDOW:4:15 MINLEN:18 (74). Processed reads were aligned to mouse
reference genome GRCm38 via HiSat2 (PMID: 31375807) using default parameters.
Post-alignment cleaning removed duplicated mapping, unmapped reads, and reads with
MAPQ<60. Gene-level expressions were quantified using Subread (PMID: 24227677),
and differential gene expression was determined with DESeq2 using default setting
(PMID: 25516281). Genes with a q < 0.1 were considered significant. Outliers'
identification was performed using Cook's distance (DESeq2), which is a measure of
how much a single sample is influencing the fitted coefficients for a gene. Driver factors
of gene expression variance (genotype and/or sex) were evaluated using R package
variance Partition. Pathway and clustering analysis were completed with Ingenuity
Pathway Analysis (Ingenuity Systems: RRID: SCR_008653) and gProfiler2 (PMID:
31066453). Data visualization was done using various R graphic packages, including
ggplo2, ggraph, and DESeq2 visualization functions. The RNA-seq data set generated
in this manuscript has been deposited at GEO (accession number GSE160586).
Reviewer token "**gpqrigisbxgprqf**".

WGCNA Analysis

Genes with less than 10 counts in more than 90% of samples were removed from
WGCNA analysis (75). The count-based gene expressions were first transformed using
a variance stabilizing method via DESeq2 vst function. The WGCNA R package (v1.69)
was used to construct an unsigned gene co-expression network with a soft threshold
power [beta] of 6. Nineteen tightly connected modules were detected, while genes that
were not connected to the 19 modules were collected in the 20th "grey" module.
WGCNA defines the expression of a given module by averaging module gene
expressions, which is also called module eigengene expression. Using a non-
parametric Kruskal-Wallis test (p value < 0.05), we can identify modules that differ
significantly among mouse samples with different genotypes. For example, the
"Greenyellow" module differs significantly between zQ175 mice and zQ175:CK2α'$^{(+/-)}$
mice. Data for the Greenyellow module were exported using a Cytoscape format for
visualization. Network figures are limited to the top 10% of genes with the strongest
network connections (the topological overlap threshold was raised until only 10% of
genes remained). The network modules are color coded by the differential expression
between zQ175 mice and zQ175:CK2α'$^{(+/-)}$ mice: blue, downregulated in zQ175
compared to ZQ175:CK2α'$^{(+/-)}$ mice; red, upregulated in zQ175 compared to

747 ZQ175:CK2α'$^{(+/-)}$. The size of the circles is scaled by the absolute value of the mean
748 log2 fold change between zQ175 and ZQ175:CK2α'$^{(+/-)}$ mice.

Quantification and Statistical analyses

Data are expressed as Mean ± SEM, Mean ± SD, or percentage, analyzed for statistical
significance, and displayed by Prism 8 software (GraphPad, San Diego, CA) or Excel
software (Microsoft). Pearson correlation tests were applied to test the normal
distribution of experimental data. Normal distributions were compared with t-test (two-
tailed) or ANOVA with appropriate post-hoc tests (Sidak's, Dunn's, or Tukey's) for
multiple comparisons. Non-normal distributions were compared with the non-parametric
Kruskal-Wallis test with an appropriate post-hoc test, as indicated. The accepted level of
significance was $p \leq 0.05$, therefore samples indicated as n.s (no significant) did not
reach $p \leq 0.05$. Statistical analyses for electrophysiological experiments were performed
with SigmaPlot 13.0 software. No statistical methods were used to predetermine sample
sizes, but sample sizes were chosen to be similar to those reported in previous
publications [11]. In the figures, we show the mean as an estimator of central tendency
including when we have used a non-parametric test, to maintain consistency with other
figures in the paper and because it is more intuitive to compare the mean values.

Acknowledgements

768 We are grateful to Drs. Sylvain Lesne and Michael Lee for sharing his expertise on
769 alpha-synuclein and sharing reagents, Maha Syed and Joyce Meints for technical
770 assistance, Jason Mitchell for assistance with confocal microscopy and Erin Greguske
771 for proofreading.

Authors' contributions

R.G.P obtained funding for the study and designed the experiments. D.Y and N.Z,
harvested mice, conducted behavioral analyses, performed IF and conducted alpha-
synuclein related experiments. F.C conducted cytokine proteome profiling and astrocyte
IF analyses. J.Y performed EM48 and Neu analyses. T.B conducted microglia analyses.
W.T conducted IL6-related experiments and qRT-PCR of SNCA^{KO} mice. K.J performed
striatal qRT-PCR. H.K conducted striatal volume analyses. C.N conducted
electrophysiological experiments. T.S.M conducted behavioral analyses. K.G and S.L
performed MRS analyses. A.W conducted a-syn IB and nuclear/cytoplasma
fractionation. R.M assisted in IF image analyses and a-syn/EM48 puncta analyses.

C.T.Z prepared SNCA^{KO} tissues. Y.Z conducted RNA-seq analyses. D.Y, N.Z, R.M, S.L,
K.G, C.N, W.T and R.G.P prepared and analyzed the data. G.O supervised the MR data
acquisition and analysis. M.B supervised mouse behavioral data acquisition and
analysis. A.A supervised electrophysiological recordings. M.C supervised microglia
analyses. J.J.L supervised SNCA^{KO} tissue preparation. R.G.P wrote the first draft of the
manuscript and all authors edited subsequent versions. All authors read and approved
the final version of the manuscript.

792 Funding

This work was supported by R.G.P's startup funds from University of Minnesota, the
Biomedical Research Awards for Interdisciplinary New Science BRAINS (to R.G.P) and
the National Institute of Health NINDS (R01 NS110694-01A1) (to R.G.P). The Center
for Magnetic Resonance Research is supported by the National Institute of Biomedical
Imaging and Bioengineering (NIBIB) grant P41 EB027061, the Institutional Center
Cores for Advanced Neuroimaging award P30 NS076408 and the W.M. Keck
Foundation. F.C. received the mobility scholarship program Global Links: an
Opportunity to Build a University Strategy 2018/2019 (GLOBUS Placement), Università
degli Studi di Cagliari, funded through the Regione Autonoma della Sardegna. National
Institute of Health NINDS (R01 NS197387) (to M.C.) and National Institute of Health
NINDS R01 MH119355 and R01 NS108686 (to A.A). Grants from Fundación Ramón
Areces, MICINN (SAF2009-08233) and MCIU/AEI/FEDER-UE (RTI2018-096322-B-100)
to JJL.

807 Data availability

808 RNA-seq data sets have been deposited at GEO. Mouse models and reagents
809 regenerated during the execution of this work are available upon request.

811 Declaration of interest

812 The authors declare no competing interests.
813

814 References

1.   T. H. s. D. C. R. Group, A novel gene containing a trinucleotide repeat that is
expanded and unstable on Huntington's disease chromosomes. *Cell* **72**, 971-983
(1993).
2.   M. DiFiglia *et al.*, Aggregation of huntingtin in neuronal intranuclear inclusions
and dystrophic neurites in brain. *Science* **277**, 1990-1993 (1997).

3.   R. J. Ferrante, N. W. Kowall, E. P. Richardson, Proliferative and degenerative
changes in striatal spiny neurons in Huntington's disease: a combined study using the
section-Golgi method and calbindin D28k immunocytochemistry. *J Neurosci* **11**, 3877-
3887 (1991).
4.   A. Crotti, C. K. Glass, The choreography of neuroinflammation in Huntington's
disease. *Trends Immunol* **36**, 364-373 (2015).
5.   J. Castello, A. Ragnauth, E. Friedman, H. Rebholz, CK2-An Emerging Target for
Neurological and Psychiatric Disorders. *Pharmaceuticals (Basel)* **10** (2017).
6.   A. F. Rosenberger *et al.*, Increased occurrence of protein kinase CK2 in
astrocytes in Alzheimer's disease pathology. *J.Neuroinflammation.* **13**, 4 (2016).
7.   L. A. Pinna, Protein kinase CK2: a challenge to canons. *J.Cell Sci.* **115**, 3873-
3878 (2002).
8.   D. W. Litchfield, Protein kinase CK2: structure, regulation and role in cellular
decisions of life and death. *Biochem.J.* **369**, 1-15 (2003).
9.   Y. Bian *et al.*, Global screening of CK2 kinase substrates by an integrated
phosphoproteomics workflow. *Sci.Rep.* **3**, 3460 (2013).
10.   I. Ceglia, M. Flajolet, H. Rebholz, Predominance of CK2α over CK2α' in the
mammalian brain. *Mol Cell Biochem* **356**, 169-175 (2011).
11.   R. Gomez-Pastor *et al.*, Abnormal degradation of the neuronal stress-protective
transcription factor HSF1 in Huntington's disease. *Nat Commun* **8**, 14405 (2017).
12.   R. Gomez-Pastor, E. T. Burchfiel, D. J. Thiele, Regulation of heat shock
transcription factors and their roles in physiology and disease. *Nat Rev Mol Cell Biol*
(2018).
13.   M. M. Fan, H. Zhang, M. R. Hayden, S. L. Pelech, L. A. Raymond, Protective up-
regulation of CK2 by mutant huntingtin in cells co-expressing NMDA receptors. *J*
Neurochem **104**, 790-805 (2008).
14.   R. S. Atwal *et al.*, Kinase inhibitors modulate huntingtin cell localization and
toxicity. *Nat.Chem.Biol.* **7**, 453-460 (2011).
15.   J. A. Greenwood, C. W. Scott, R. C. Spreen, C. B. Caputo, G. V. Johnson,
Casein kinase II preferentially phosphorylates human tau isoforms containing an amino-
terminal insert. Identification of threonine 39 as the primary phosphate acceptor. *J Biol*
Chem **269**, 4373-4380 (1994).
16.   E. Masliah *et al.*, Casein kinase II alteration precedes tau accumulation in tangle
formation. *Am.J.Pathol.* **140**, 263-268 (1992).
17.   E. A. Waxman, B. I. Giasson, Specificity and regulation of casein kinase-
mediated phosphorylation of alpha-synuclein. *J Neuropathol Exp Neurol* **67**, 402-416
(2008).
18.   C. Ising *et al.*, NLRP3 inflammasome activation drives tau pathology. *Nature* **575**,
669-673 (2019).
19.   E. Masliah *et al.*, Dopaminergic loss and inclusion body formation in alpha-
synuclein mice: implications for neurodegenerative disorders. *Science* **287**, 1265-1269
(2000).
20.   C. Tomás-Zapico *et al.*, α-Synuclein accumulates in huntingtin inclusions but
forms independent filaments and its deficiency attenuates early phenotype in a mouse
model of Huntington's disease. *Hum Mol Genet* **21**, 495-510 (2012).

21.   M. Fernández-Nogales *et al.*, Huntington's disease is a four-repeat tauopathy
with tau nuclear rods. *Nat Med* **20**, 881-885 (2014).
22.   P. Liu *et al.*, A soluble truncated tau species related to cognitive dysfunction is
elevated in the brain of cognitively impaired human individuals. *Sci Rep* **10**, 3869
(2020).
23.   A. Shalash *et al.*, Elevated Serum α-Synuclein Autoantibodies in Patients with
Parkinson's Disease Relative to Alzheimer's Disease and Controls. *Front Neurol* **8**, 720
(2017).
24.   M. Breza *et al.*, Elevated Serum α-Synuclein Levels in Huntington's Disease
Patients. *Neuroscience* **431**, 34-39 (2020).
25.   S. Corrochano *et al.*, α-Synuclein levels affect autophagosome numbers in vivo
and modulate Huntington disease pathology. *Autophagy* **8**, 431-432 (2012).
26.   T. Heikkinen *et al.*, Characterization of neurophysiological and behavioral
changes, MRI brain volumetry and 1H MRS in zQ175 knock-in mouse model of
Huntington's disease. *PLoS One* **7**, e50717 (2012).
27.   L. B. Menalled *et al.*, Comprehensive behavioral and molecular characterization
of a new knock-in mouse model of Huntington's disease: zQ175. *PLoS One* **7**, e49838
(2012).
28.   P. Arlotta, B. J. Molyneaux, D. Jabaudon, Y. Yoshida, J. D. Macklis, Ctip2
controls the differentiation of medium spiny neurons and the establishment of the
cellular architecture of the striatum. *J Neurosci* **28**, 622-632 (2008).
29.   N. N. Singh, D. P. Ramji, Protein kinase CK2, an important regulator of the
inflammatory response? *J Mol Med (Berl)* **86**, 887-897 (2008).
30.   F. B. Rodrigues *et al.*, Cerebrospinal Fluid Inflammatory Biomarkers Reflect
Clinical Severity in Huntington's Disease. *PLoS One* **11**, e0163479 (2016).
31.   N. Pavese *et al.*, Microglial activation correlates with severity in Huntington
disease: a clinical and PET study. *Neurology* **66**, 1638-1643 (2006).
32.   Y. F. Tai *et al.*, Imaging microglial activation in Huntington's disease. *Brain Res
Bull* **72**, 148-151 (2007).
33.   J. C. Savage *et al.*, Microglial physiological properties and interactions with
synapses are altered at presymptomatic stages in a mouse model of Huntington's
disease pathology. *J Neuroinflammation* **17**, 98 (2020).
34.   K. Li, J. Li, J. Zheng, S. Qin, Reactive Astrocytes in Neurodegenerative
Diseases. *Aging Dis* **10**, 664-675 (2019).
35.   B. S. Khakh *et al.*, Unravelling and Exploiting Astrocyte Dysfunction in
Huntington's Disease. *Trends Neurosci* **40**, 422-437 (2017).
36.   M. Castillo, L. Kwock, J. Scatliff, S. K. Mukherji, Proton MR spectroscopy in
neoplastic and non-neoplastic brain disorders. *Magn Reson Imaging Clin N Am* **6**, 1-20
(1998).
37.   A. Brand, C. Richter-Landsberg, D. Leibfritz, Multinuclear NMR studies on the
energy metabolism of glial and neuronal cells. *Dev Neurosci* **15**, 289-298 (1993).
38.   I. Tkac *et al.*, Homeostatic adaptations in brain energy metabolism in mouse
models of Huntington disease. *J Cereb Blood Flow Metab* **32**, 1977-1988 (2012).
39.   Q. Peng *et al.*, Characterization of Behavioral, Neuropathological, Brain
Metabolic and Key Molecular Changes in zQ175 Knock-In Mouse Model of Huntington's
Disease. *PLoS One* **11**, e0148839 (2016).

40.   S. A. Liddelow *et al.*, Neurotoxic reactive astrocytes are induced by activated
microglia. *Nature* **541**, 481-487 (2017).
41.   L. E. Clarke *et al.*, Normal aging induces A1-like astrocyte reactivity. *Proc Natl
Acad Sci U S A* **115**, E1896-E1905 (2018).
42.   S. K. Singhrao, J. W. Neal, B. P. Morgan, P. Gasque, Increased complement
biosynthesis by microglia and complement activation on neurons in Huntington's
disease. *Exp Neurol* **159**, 362-376 (1999).
43.   B. Diaz-Castro, M. R. Gangwani, X. Yu, G. Coppola, B. S. Khakh, Astrocyte
molecular signatures in Huntington's disease. *Sci Transl Med* **11** (2019).
44.   M. M. Zeron *et al.*, Increased sensitivity to N-methyl-D-aspartate receptor-
mediated excitotoxicity in a mouse model of Huntington's disease. *Neuron* **33**, 849-860
(2002).
45.   A. I. Smith-Dijak, M. D. Sepers, L. A. Raymond, Alterations in synaptic function
and plasticity in Huntington disease. *J Neurochem* **150**, 346-365 (2019).
46.   T. E. Wood *et al.*, Mutant huntingtin reduction in astrocytes slows disease
progression in the BACHD conditional Huntington's disease mouse model. *Hum Mol
Genet* **28**, 487-500 (2019).
47.   T. Indersmitten, C. H. Tran, C. T. Cepeda, M. S. Levine, Altered Excitatory and
Inhibitory Inputs to Striatal Medium-Sized Spiny Neurons and Cortical Pyramidal
Neurons in the Q175 Mouse Model of Huntington's Disease. *J.Neurophysiol.*, jn (2015).
48.   A. A. Fienberg *et al.*, DARPP-32: regulator of the efficacy of dopaminergic
neurotransmission. *Science* **281**, 838-842 (1998).
49.   E. Vezzoli *et al.*, Inhibiting pathologically active ADAM10 rescues synaptic and
cognitive decline in Huntington's disease. *J Clin Invest* **129**, 2390-2403 (2019).
50.   P. Langfelder *et al.*, Integrated genomics and proteomics define huntingtin CAG
length-dependent networks in mice. *Nat Neurosci* **19**, 623-633 (2016).
51.   A. Gallardo-Orihuela *et al.*, Transcriptional correlates of the pathological
phenotype in a Huntington's disease mouse model. *Sci Rep* **9**, 18696 (2019).
52.   G. M. o. H. s. D. G.-H. Consortium, Identification of Genetic Factors that Modify
Clinical Onset of Huntington's Disease. *Cell* **162**, 516-526 (2015).
53.   M. Riessland *et al.*, Neurocalcin Delta Suppression Protects against Spinal
Muscular Atrophy in Humans and across Species by Restoring Impaired Endocytosis.
Am J Hum Genet **100**, 297-315 (2017).
54.   O. Al-Dalahmah *et al.*, Single-nucleus RNA-seq identifies Huntington disease
astrocyte states. *Acta Neuropathol Commun* **8**, 19 (2020).
55.   D. Davidi *et al.*, α-Synuclein Translocates to the Nucleus to Activate Retinoic-
Acid-Dependent Gene Transcription. *iScience* **23**, 100910 (2020).
56.   M. Decressac *et al.*, α-Synuclein-induced down-regulation of Nurr1 disrupts
GDNF signaling in nigral dopamine neurons. *Sci Transl Med* **4**, 163ra156 (2012).
57.   M. W. Rousseaux *et al.*, TRIM28 regulates the nuclear accumulation and toxicity
of both alpha-synuclein and tau. *Elife* **5** (2016).
58.   A. Oueslati, Implication of Alpha-Synuclein Phosphorylation at S129 in
Synucleinopathies: What Have We Learned in the Last Decade? *J Parkinsons Dis* **6**,
39-51 (2016).
59.   H. Fujiwara *et al.*, alpha-Synuclein is phosphorylated in synucleinopathy lesions.
Nat.Cell Biol. **4**, 160-164 (2002).

60.   M. Mikula *et al.*, Halogenated imidazole derivatives block RNA polymerase II
elongation along mitogen inducible genes. *BMC Mol Biol* **11**, 4 (2010).
61.   H. Y. Hsiao *et al.*, Inhibition of soluble tumor necrosis factor is therapeutic in
Huntington's disease. *Hum Mol Genet* **23**, 4328-4344 (2014).
62.   S. Franciosi *et al.*, Age-dependent neurovascular abnormalities and altered
microglial morphology in the YAC128 mouse model of Huntington disease. *Neurobiol*
Dis **45**, 438-449 (2012).
63.   K. W. Selmaj, M. Farooq, W. T. Norton, C. S. Raine, C. F. Brosnan, Proliferation
of astrocytes in vitro in response to cytokines. A primary role for tumor necrosis factor. *J*
Immunol **144**, 129-135 (1990).
64.   H. Lian *et al.*, Astrocyte-Microglia Cross Talk through Complement Activation
Modulates Amyloid Pathology in Mouse Models of Alzheimer's Disease. *J Neurosci* **36**,
577-589 (2016).
65.   K. Nilsson Ekdahl, B. Nilsson, Phosphorylation of complement component C3
after synthesis in U937 cells by a putative protein kinase, casein kinase 2, which is
regulated by CD11b: evidence that membrane-bound proteases preferentially cleave
phosphorylated C3. *Biochem J* **328 (Pt 2)**, 625-633 (1997).
66.   A. Mottahedin *et al.*, Effect of Neuroinflammation on Synaptic Organization and
Function in the Developing Brain: Implications for Neurodevelopmental and
Neurodegenerative Disorders. *Front Cell Neurosci* **11**, 190 (2017).
67.   C. Franchin *et al.*, Re-evaluation of protein kinase CK2 pleiotropy: new insights
provided by a phosphoproteomics analysis of CK2 knockout cells. *Cell Mol Life Sci* **75**,
2011-2026 (2018).
68.   S. Hara *et al.*, Serine 129 phosphorylation of membrane-associated α-synuclein
modulates dopamine transporter function in a G protein-coupled receptor kinase-
dependent manner. *Mol Biol Cell* **24**, 1649-1660, S1641-1643 (2013).
69.   G. Lee *et al.*, Casein kinase II-mediated phosphorylation regulates alpha-
synuclein/synphilin-1 interaction and inclusion body formation. *J Biol Chem* **279**, 6834-
6839 (2004).
70.   C. Y. Chung, J. B. Koprich, H. Siddiqi, O. Isacson, Dynamic changes in
presynaptic and axonal transport proteins combined with striatal neuroinflammation
precede dopaminergic neuronal loss in a rat model of AAV alpha-synucleinopathy. *J*
Neurosci **29**, 3365-3373 (2009).
71.   X. Xu, P. A. Toselli, L. D. Russell, D. C. Seldin, Globozoospermia in mice lacking
the casein kinase II alpha' catalytic subunit. *Nat.Genet.* **23**, 118-121 (1999).
72.   A. Cavaccini, C. Durkee, P. Kofuji, R. Tonini, A. Araque, Astrocyte Signaling
Gates Long-Term Depression at Corticostriatal Synapses of the Direct Pathway. *J*
Neurosci **40**, 5757-5768 (2020).
73.   G. Öz *et al.*, Assessing recovery from neurodegeneration in spinocerebellar
ataxia 1: Comparison of in vivo magnetic resonance spectroscopy with motor testing,
gene expression and histology. *Neurobiol Dis* **74**, 158-166 (2015).
74.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
75.   P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation
network analysis. *BMC Bioinformatics* **9**, 559 (2008).

1003

Figure Legends

Figure 1. **CK2α' levels progressively increase in the striatum of zQ175 and correlate with HTT aggregation and NeuN depletion. *A*, *B*,** Immunostaining and quantification of HTT puncta detected with anti-HTT EM48 antibody, *C*, *D*, NeuN⁺ cells and *E*, *F*, CK2α' levels in zQ175 compared with WT mice at 3, 6, 12 and 22 months (*n*=6 mice/genotype). *G*, CK2α' mRNA levels analyzed by RT-qPCR in striatum and cortex of 6 month old mice. Data was normalized to GAPDH and WT striatum (n=3 mice/genotype). *H*, Linear regression analysis between CK2α' levels and HTT puncta, and *I*, between CK2α' levels and number of NeuN⁺ cells in zQ175 mice. The Pearson correlation coefficient (ρ) and R^2 are indicated. Data are mean ± SD with significance determined by one-way ANOVA Dunnett's post-hoc test in *B* and *F* and two-way ANOVA Tukey post-hoc test in *D* and *G*. p-values <0.05 are indicated. n.s = not significant. Scale bar, 50 μm.

Figure 2. **Expression of pro-inflammatory cytokines in the striatum of zQ175 is decreased by reduction in CK2α'**. *A*, RT-qPCR analysis for CK2α' and IL-6 in immortalized embryonic striatal cells ST*Hdh*$^{Q7/Q7}$ (control) and ST*Hdh*$^{Q111/Q111}$ (HD) (n = 5 independent experiments). Data were normalized to GAPDH and using control cells as a reference. *B*, siRNA knockdown of CK2α' for 24 h in ST*Hdh*$^{Q111/Q111}$ cells and RT-qPCR. Data were normalized with GAPDH and relativized to non-targeting control siRNA-treated cells (scramble), (*n* = 6 independent experiments). Data are mean ± SEM with significance determined by paired Student's t-test. *C*, *D*, Representative IF images and quantitative analysis of CK2α' in striatal MSNs immunostained for Ctip2, a specific MSN marker, in WT, zQ175 and zQ175:CK2α'$^{(+/-)}$ mice at 12 months of age (*n*=6 mice/genotype). Scale bar, 50 μm. *E*, *F*, Representative images of mouse cytokine array panels in striatum extracts of WT, zQ175, and zQ175:CK2α'$^{(+/-)}$ at 12-14 months of age (*n* = 6 mice/genotype). *G-I*, Iba1 IB in the striatum of 12 months old WT, zQ175 and zQ175:CK2α'. Quantification was measured by image analyses using Image J software. Scale bar, 100 μm. Error bars denote mean ± SD, values were analyzed by Student's t test in *F*, and one-way ANOVA and Tukey post-hoc test in *A, B, D*, and *H*. **p* represent p-values comparing zQ175 and WT, #*p* are p-values comparing zQ175 and zQ175:CK2α'$^{(+/-)}$.

Figure 3. **Decreased CK2α' ameliorated astrocyte pathology in zQ175 mice**. *A*, Representative images of striatum sections from 12-month-old mice immunostained for CK2α' and glutamine synthetase (GS, an astrocytic marker) in WT, zQ175 and zQ175:CK2α'$^{(+/-)}$. *B*, Quantification of GS⁺ cells from images in *A* (*n* = 5 mice/genotype). *C*, Representative coronal images of brain scans in 9.4T magnet showing the striatum voxel (green box) for MRS acquisition from each genotype at 22 months of age. *D*, *myo*-inositol (Ins) quantification in WT, zQ175 and zQ175:CK2α'$^{(+/-)}$ (*n* = 4-6 mice/genotype). *E*, Representative IF analysis of C3d co-stained with GS in the striatum of 12-month-old mice. *E'*, Higher magnification of the outlined section in *E*

shows a representative view of a single astrocyte immunolabeled with C3d and GS. ***F,***
Quantification of C3d signal from images in ***E***. ***G***, Quantification of C3d⁺ GS⁺ double-
positive cells (*n* = 5 mice/genotype). DAPI is used for nuclear staining. Scale bars, 50
µm. Error bars denote mean ± SD. *p*-values for differences between groups are
indicated in each graph and calculated using two-way ANOVA and Tukey post-hoc
tests. n.s. = not significant.

1053

Figure 4. **CK2α' haploinsufficiency increased the frequency of AMPA-mediated**
miniature excitatory postsynaptic currents (mEPSC) in the dorsolateral striatum
of zQ175 mice. ***A***, Image shows whole-cell patch-clamp recording diagram in acute
dorsolateral striatum slices, where Ctip2 labeled MSNs from 12-month-old mice. Scale
bar 500 µm, Ctr: Cortex; Str: Striatum. ***B***, Input–output curve (WT, n = 8; zQ175, n = 9;
zQ175:CK2α'$^{(+/-)}$ n = 13). Representative traces are shown in the top inset. ***C***, Short-
term potentiation measured via paired-pulse facilitation (WT, n = 8; zQ175, n = 9;
zQ175:CK2α'$^{(+/-)}$ n = 11). Representative traces of two consecutive stimuli delivered at
25 ms time intervals are shown in the top inset. ***D***, Short-term depression analyzed
through synaptic fatigue (WT, n = 7; zQ175, n = 9; zQ175:CK2α'$^{(+/-)}$ n = 12).
Representative traces are shown in the top inset. Values were analyzed using two-way
ANOVA with Tukey's post-hoc analysis. ***E-F***, Spontaneous recordings of mini excitatory
postsynaptic currents (mEPSCs). Amplitude (in pA; left panel) and frequency (in Hz;
right panel) were analyzed (WT, n = 10; zQ175, n = 9; zQ175:CK2α'$^{(+/-)}$ n = 12). ***G***,
Representative mEPSC traces. Values were analyzed using one-way ANOVA with
Dunn´s post-hoc analysis. P values <0.05 are indicated. Error bars represent mean ±
SEM from at least 3 mice/genotype.

1071

Figure 5. **Genetic deletion of CK2α' improved motor deficits in zQ175. *A***, Number
of foot slips recorded while walking on four different types of beams with different
degrees of difficulty from less to more challenging; medium-square, medium round,
small-square and small-round (Beam test), and ***B***, Latency to fall off the rod Rotarod
test) at 3, 6 and 12 months of age (n =16-18 mice/genotype in 3 months, n= 12-14 for 6
months and n = 5-6 for 12 months). Error bars denote mean ± SEM, values were
analyzed by two-way ANOVA with *Sidak's* post-hoc test. p-values <0.05 are indicated,
n.s = not significant.

1080

Figure 6. **Depletion of CK2α' restored the expression of synaptic genes**
associated with α-syn-dependent regulation in the striatum of symptomatic zQ175
mice. ***A***, Kruskal-Wallis test of module expressions between zQ175 (HD) mice and
zQ175:CK2α'$^{(+/-)}$ mice. The y-axis is the negative log transformed p-values. ***B***,
Expressions of module "Greenyellow" in each mouse sample. ***C***, IPA canonical pathway
analysis of module genes for module "Greenyellow". ***D***, Enrichment analysis of GO
terms in CC (cellular component) for module genes of module "Greenyellow". ***E***,
Network visualization of top 15% connected genes in "Greenyellow" Module. The size of
the circles was scaled by the absolute value of the mean log2-fold change between

zQ175 and zQ175:CK2α'$^{(+/-)}$ mice. **F,** Marker genes and their mean log2 fold change
between zQ175 and zQ175:CK2α'$^{(+/-)}$ mice compared to WT. **G,** IPA network analysis of
DGEs in zQ175:CK2α' mice showing α-syn as the most significant upstream regulator.

1093

Figure 7. α-syn differentially accumulates in the nucleus of symptomatic zQ175
mice and colocalized with mtHTT. **A,** α-syn (4D6 antibody) IB in the striatum of WT,
zQ175 and SNCA^{KO} and **B** in WT, zQ175 and zQ175:CK2α'$^{(+/-)}$ mice at 12 months old.
GAPDH used as loading control. **C,** α-syn protein levels analyzed by Image J from IB
analyses (n= 5-6 mice/genotype). **D,** Nuclear/cytoplasmic fractionation of striatum
samples from 12-month-old WT, zQ175 and SNCA^{KO} mice. **E,** Quantification of nuclear
α-syn from images in D (n=4 mice/genotype). **F,** Representative IF images of dorsal
striatum sections from 12 month old WT, zQ175 and zQ175:CK2α'$^{(+/-)}$ (n=3
mice/genotype) for α-syn and HTT (EM48 antibody). White arrows indicate α-syn/HTT
colocalization. Scale bar, 10 μm. **G,** Magnification of images from F showing nuclear
and cytoplasmic α-syn and HTT colocalization. Grey circles represent nuclei. White
arrows indicate α-syn/HTT colocalization. Scale bar, 2 μm. **H** Number of cytoplasmic
and **I** nuclear EM48$^+$ puncta calculated using Image J Puncta analysis plugin. A total of
three images per brain section and three brain sections per genotype were analyzed
(n=27 images, n=3 mice/ genotype). **J,** Number of colocalized α-syn and EM48$^+$ puncta
calculated using Image J Puncta analysis plugin (n=3 mice/genotype). Error bars denote
mean ± SEM, values were analyzed by Student's t-test.

1111

Figure 8. CK2α' regulates phosphorylation of S129-α-syn and nuclear
accumulation in symptomatic zQ175 mice. **A,** pS129-α-syn (EP1536Y antibody) IB in
the striatum of 12-month-old WT, zQ175 and SNCA^{KO} (n=4 mice/genotype)**. B** pS129-α-
syn (81A antibody) IB in the striatum of patients with HD (Vonsattel grade 3 and 4)
compared to age and sex matched controls. GAPDH is used as loading control. **C**
pS129-α-syn protein levels analyzed by Image J from images in **B**. Samples from grade
3 and grade 4 HD were all grouped for pS129-α-syn quantification. **D,** Representative
pS129-α-syn IF images (81A antibody) in the dorsal striatum of 12-month-old WT,
zQ175 and zQ175:CK2α'$^{(+/-)}$ (n=3 mice/genotype), Scale bar, 20 μm. **E,** pS129-α-syn
fluorescence signal was calculated using Image J from images in D (n=3
mice/genotype, n=27 images per mouse). **F,** Magnification of images in D showed
pS129-α-syn and EM48 colocalization in zQ175 and zQ175:CK2α'$^{(+/-)}$ (n=3 mice/group).
G, Quantification of pS129-α-syn and EM48 colocalized puncta using Image J puncta
plug in. All data are mean ± SEM. Statistical analyses were conducted by one-way
ANOVA. **H,** Working model for the role of CK2α' in the regulation of pS129-α-syn and
HD-like phenotype.

1128

1129

1130

1131

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8