Novel SARS-CoV-2 Whole-genome sequencing technique using Reverse Complement PCR enables easy, fast and accurate outbreak analysis in hospital and community settings

Femke Wolters[1*a], Jordy P.M. Coolen[1a], Alma Tostmann[1], Lenneke F.J. van Groningen[2], Chantal P. Bleeker-

Rovers[3], Edward C.T.H. Tan[4,5], Nannet van der Geest-Blankert[6], Jeannine L.A. Hautvast[7], Joost Hopman[1],

Heiman F.L. Wertheim[1], Janette C. Rahamat-Langendoen[1], Marko Storch[8], Willem J.G. Melchers[1].

[a]Shared first authorship

[1]Department of Medical Microbiology and Radboudumc Center for Infectious Diseases, Radboud university

medical center, Nijmegen, The Netherlands

[2]Department of Haematology, Radboud university medical center, Nijmegen, The Netherlands

[3]Department of Internal Medicine, division of Infectious Diseases, Radboud university medical center,

Nijmegen, The Netherlands

[4]Department of Emergency Medicine, Radboud university medical center, Nijmegen, The Netherlands

[5]Department of Surgery, Radboud University Medical Center, Nijmegen, The Netherlands

[6]Department of Occupational Health, Radboud university medical center, Nijmegen, The Netherlands

[7]Regional Public Health Service, Nijmegen, the Netherlands

[8]London Biofoundry, Imperial College Translation & Innovation Hub, White City Campus, 84 Wood Lane,

London, W12 0BZ, UK;

Corresponding author:

Femke Wolters, MD

Radboud university medical center, Geert Grooteplein Zuid 10, 6525 GA, Nijmegen, The Netherlands

Femke.wolters@radboudumc.nl

27
28

Abstract

Background: Current transmission rates of severe acute respiratory syndrome coronavirus 2 (SARS-

CoV-2) are still increasing and many countries are facing second waves of infections. Rapid SARS-

CoV-2 whole-genome sequencing (WGS) is often unavailable but could support public health

organizations and hospitals in monitoring and determining transmission links. Here we report the use

of reverse complement polymerase chain reaction (RC-PCR), a novel technology for WGS of SARS-

CoV-2 enabling library preparation in a single PCR saving time, resources and enables high

throughput screening. Additionally, we show SARS-CoV-2 diversity and possible transmission within

the Radboud university medical center (Radboudumc) during September 2020 using RC-PCR WGS.

Methods: A total of 173 samples tested positive for SARS-CoV-2 between March and September

2020 were selected for whole-genome sequencing. Ct values of the samples ranged from 16 to 42.

They were collected from 83 healthcare workers and three patients at the Radboudumc, in addition to

64 people living in the area around the hospital and tested by the local health services. For validation

purposes, nineteen of the included samples were previously sequenced using Oxford Nanopore

Technologies and compared to RC-PCR WGS results. The applicability of RC-PCR WGS in outbreak

analysis for public health service and hospitals was tested on six suspected clusters containing samples

of healthcare workers and patients with an epidemiological link.

Findings: RC-PCR resulted in sequencing data for 146 samples. It showed a genome coverage of up

to 98,2% for samples with a maximum Ct value of 32. Comparison to Oxford Nanopore technologies

gives a near-perfect agreement on 95% of the samples (18 out of 19). Three out of six clusters with a

suspected epidemiological link were fully confirmed, in the others, four healthcare workers were not

associated. In the public health service samples, a previously unknown chain of transmission was

confirmed.

Significance statement:

SAR-CoV-2 whole-genome sequencing using RC-PCR is a reliable technique and applicable for use

in outbreak analysis and surveillance. Its ease of use, high-trough screening capacity and wide

applicability makes it a valuable addition or replacement during this ongoing SARS-CoV-2 pandemic.

Funding: None

Research in context

Evidence before this study

At present whole genome sequencing techniques for SARS-CoV-2 have a large turnover time and are

not widely available. Only a few laboratories are currently able to perform large scale SARS-CoV-2

sequencing. This restricts the use of sequencing to aid hospital and community infection prevention.

Added value of this study

Here we present clinical and technical data on a novel Whole Genome Sequencing technology,

implementing reverse-complement PCR. It is able to obtain high genome coverage of SARS-CoV-2

and confirm and exclude epidemiological links in 173 healthcare workers and patients. The RC-PCR

technology simplifies the workflow thereby reducing hands on time. It combines targeted PCR and

sequence library construction in a single PCR, which normally takes several steps. Additionally, this

technology can be used in concordance with the widely available range of Illumina sequencers.

Implications of all the available evidence

RC-PCR whole genome sequencing technology enables rapid and targeted surveillance and response

to an ongoing outbreak that has great impact on public health and society. Increased use of sequencing

technologies in local laboratories can help prevent increase of SARS-CoV-2 spreading by better

understanding modes of transmission.

Introduction

76
In December 2019 China reported a group of patients with a severe respiratory illness caused by a thus

far unknown coronavirus. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was

identified as the causative agent.[1] Since its outbreak, the virus evolved into a pandemic with almost 37

million infections and over a million deaths worldwide by October 2020.[2] Many countries are

currently fighting second waves of infection whilst the healthcare systems are still under pressure from

the first wave. To reduce spread and mitigate workforce depletion, large scale testing of healthcare

workers (HCW) was implemented in the Netherlands early on.[3]

Current testing is based on RT-PCR detection of SARS-CoV-2 in nasopharynx or oropharyngeal

swabs. If tested SARS-CoV-2 positive, HCW are instructed to self-isolate at home, and source finding

and contact tracing is performed. These procedures enable us to identify patients and personnel at risk

of infection and to identify chains of transmission in the hospital. In the community setting, source

finding and contact tracing is performed by public health staff upon a notification of a SARS-CoV-2

positive individual. It facilitates the implementation of quarantine measures for high-risk contacts in

the community. Contact tracing is time consuming and with rising numbers of infections as currently

seen in the second wave, the public health capacity may reach the limits of feasibility of thorough

source and contact tracing investigations.[4] Routine sequencing the SARS-CoV-2 genome from

positive samples provides crucial insights into viral evolution and supports outbreak analysis.[5,6]

Current whole-genome sequencing (WGS) workflows often require cumbersome preparation, are

laborious to implement for high throughput screening or use less widely accessible sequencing

platforms, preventing widespread implementation. Here we present a novel strategy for fast, simple

and robust Next-Generation Sequencing (NGS) WGS library preparation. We show that the RC-PCR

method, which integrates tiled target amplification with Illumina library preparation has a simple

workflow with minimal hands-on time. We used this novel and practical method to I) validate and

compare it with another sequence technology to demonstrate its reliability and capacity and II) apply it

to a set of epidemiologically linked cases to illustrate its added value in detecting potential

transmission events in public health and hospital settings.

Material and Methods

In this study we conducted a validation to assess the performance and reproducibility of the novel RC-

PCR SARS-CoV-2 sequencing technology. Subsequently, we performed a clinical validation to assess

the potential added value in identifying chains of transmission in a hospital and public health setting.

Sample collection

Nasopharyngeal and oropharyngeal swabs collected in UTM or GLY medium of patients, healthcare

workers and samples for the local public health services that were tested for SARS-CoV-2 in our

laboratory. Samples collected between March 2020 and September 2020 were included in this study

and stored at -80°C. Detailed descriptions on included samples can be found in supplementary table 1.

A total of 173 SARS-CoV-2 positive and fifteen SARS-CoV-2 negative samples were tested.

Samples and selection of epidemiological clusters

Nineteen out of 188 samples were previously sequenced using Oxford Nanopore Technologies (ONT).

These nineteen samples were collected at the beginning of the pandemic, between March 9[th] and

March 20[th] and ONT sequencing data of these samples has been deposited at GISAID, a global

initiative curating sequenced SARS-CoV-2 genomes for public access (https://www.gisaid.org/).[6]

Hospital samples

Six epidemiolocal hospital clusters that were identified by the infection prevention and control (IPC)

team were included in this study. These clusters involved patients admitted at and healthcare workers

(HCW) employed by the Radboud university medical center. Of the identified clusters, three were

clusters of healthcare workers with an epidemiological link, and three involved a patient and several

healthcare workers with a suspected epidemiological link. To determine whether other HCW could be

linked to one of the clusters, samples of sporadic HCW (all other HCW who tested positive for SARS-

CoV-2 in September 2020) were included in the selection, as the second wave of infections in the

Netherlands started late August 2020. These consist of Radboud university medical center HCW and

the majority work in direct or indirect patient care. A minority of positive samples include employees

working at the medical faculty or research departments. Additionally, twenty samples were included

from patients and HCW who were tested between March and September 2020 and who were not

associated with any of these predefined clusters.

Community samples

We also included an additional 64 community samples that tested positive for SARS-CoV-2 in March

and April 2020 and that were tested by the local public health service. These were samples of persons

living in the defined public health region surrounding our hospital. See Table 1 for an overview of the

groups and clusters.

The Research Ethics Committee of the region Arnhem/Nijmegen reviewed the current study and

waived additional ethical approval. All personal data of patients, HCW and public health service

Table1: number of groups and clusters of samples that were sequenced for SARS-CoV-2.			
Groups	Samples (N)	Month SARS-CoV-2 PCR positive	IPC cluster information
Oxford Nanopore Technology (ONT)	19	March 2020	None
Cluster 1 – External outbreak link	6	September 2020	HCW linked to a known community outbreak and who had either visited the venue or had close contact to people (with positive test) who had visited the venue
Cluster 2 – Department C	5	September 2020	All HCW working at the same department in close proximity and who tested positive in the same week.
Cluster 3 – Patient ward E	2^	September 2020	A patient and an HCW; the HCW had contact with the patient without adequate personal protective equipment (PPE).
Cluster 4 – Patient ward H	3*	May 2020	Two HCW and one patient tested positive at the same department in a short time period. An epidemiological link was suspected since the employees came in contact with the patient.
Cluster 5 – Laboratory R	9	April 2020	All HCW working at the same department, tested positive in the same week.
Cluster 6 – Patient ward S	6	September 2020	One patient and 5 HCW, the HCW tested positive 5 days after being in contact with the positive patient, the event included an unexpected aerosol generating procedure and HCW were not protected with PPE.
Sporadic HCW September 2020	39	September 2020	none
Public Health services samples	64	March & April 2020	none
Other *(patients/employees tested up to September 2020)*	20^	March – September 2020	none
Negative	15	n.a.	n.a.
Total	188		

samples was anonymized. Cluster information was provided anonymously by the IPC team.

Real-Time Polymerase Chain Reaction

SARS-CoV-2 RT-PCR was performed on all samples during routine diagnostics. RNA was isolated

using Roche COBAS 4800 (Roche Diagnostics Corporation) with a CT/NG extraction kit according to

the manufacturers protocol. RT-PCR with primers targeting the envelope (E-gene) was used as

described by Corman *et al.* and performed on a LightCycler 480 (Roche Diagnostics Corporation)

using Roche Multiplex RNA Virus Mastermix.[7]

^1 sample was isolated and sequenced twice. *2 samples were isolated and sequenced twice

Reverse Complement Polymerase Chain Reaction

For all 188 selected samples, RNA isolation was repeated on the MagnaPure 96 (Roche Diagnostics

Corporation) using Small Volume isolate protocol with 200µl of sample and eluting isolated RNA in

50µl. cDNA-synthesis was performed using Multiscribe RT (Applied Biosystems) with 10µl of RNA

input (supplementary table 2). Four samples were replicates, RNA was isolated twice and tested in two

separate sequencing runs. They were randomly selected for the first run, but were also part of an IPC

identified cluster and therefore included in the second run.

Whole genome sequencing (WGS) was performed in 3 independent runs (96 samples each) using the

novel EasySeq™ RC-PCR SARS-CoV-2 WGS kit (NimaGen BV, Nijmegen, The Netherlands).

Figure 1 and 2 show a detailed description of the technology in which two types of oligo's are used to

start the targeted amplification. The RC-probe and the universal barcoding primer hybridize and start

the formation of specific SARS-CoV-2 primers with Unique Dual Index (UDI) and adapter sequences

already included. In contrast to other techniques where multiple steps are needed to add sequence

adapters and UDI's. This means a regular PCR-system can be used to produce SARS-CoV-2 specific

amplicons ready for sequencing. The kit uses 155 newly designed probes with a tiling strategy

previously implemented in the ARTIC protocol.[8] The probes are divided in two pools, A and B. Pool

A contains 78 probes and Pool B contains 77 probes. This strategy requires two separate RC-PCR

reactions but ensures there is minimal chance of forming chimeric sequences or other PCR artifacts

(See Figure 2). After the PCR, samples of each plate are pooled into an Eppendorf tube, resulting in

two tubes, for pool A and B, respectively. These are individually cleaned using AmpliClean™

Magnetic Bead PCR Clean-up Kit (NimaGen, Nijmegen, The Netherlands). Afterwards, quantification

using the Qubit double strand DNA (dsDNA) High Sensitivity assay kit on a Qubit 4.0 instrument

(Life Technologies) is performed and pool A and B are combined. The amplicon fragment size in the

final library will be around 435 bp. Next Generation Sequencing (NGS) was performed on an Illumina

MiniSeq® using a Mid Output Kit (2x150-cycles) (Illumina, San Diego, CA, USA) by loading 0.8 pM

on the flowcell. The first two runs (Run1 and Run1_new) were conducted to test the performance of

the RC-PCR on a large variety of Ct-values (Ct 16 – 41) using the standard protocol provided by

NimaGen. For sequencing Run1_new the RC-PCR product from Run1 was re-used and sequenced

with the exception that the final sequencing library was created by using a balanced library pooling

strategy based on estimated cDNA input (2 ul for Ct<20, 5 ul 20≤Ct<27 or 10ul Ct≥27). The final

sequence run (Run2) contains samples with a Ct range from 16 – 32, using the same Ct dependent

balanced library strategy.

Figure 1. Schematic representation of the RC-PCR technology to WGS SARS-CoV-2. The protocol consists of one single PCR-like reaction consisting of 2 steps. The schematic is adapted from Kieser *et al.* (Kieser *et al.*, 2020) **A.** Two types of oligo's are present, 1) the universal barcoding primer which includes a Unique Dual Index (UDI), sequence adapter, and universal tail. 2) the RC probe which contains an extension blocker, universal sequence, and the reverse complement of the SARS-CoV-2 genomic target sequence. **B.** The universal tail sequences anneal and form a SARS-CoV-2 specific PCR primer. **C - E.** A regular PCR in which the SARS-CoV-2 specific amplicons are created. **F.** The final amplicons are ready to sequence on an Illumina sequencer.

Figure 2. Schematic overview of the EasySeq SARS-CoV-2 WGS process. **A.** For 96 samples the kit provides a Pool A plate and a Pool B plate. These plates contain the RC-probe and the universal oligo's with an UDI per well. Identical UDI's between Pool A and B are used. **B.** For covering the full SARS-CoV-2 genome a tiling method similar to the ARTIC protocol is used.(DNA Pipelines R&D, 2020) The kit provides two distinct plates to separate into amplicon pools that do not overlap. This greatly enhances accurate sequencing output. **C.** The same wells of pool A and B share the same index and allow the combination of corresponding sequencing results to cover the full SARS-CoV-2 genome.

Data analysis

VirSEAK (JSI, Ettenheim, Germany) was used to map the Illumina paired-end reads to SARS-CoV-2

reference NC_045512.2. Consensus sequences were extracted for each sample using the virSEAK

export option, settings used can be found in supplementary table 3. All consensus sequences and

reference NC_045512.2 were aligned using MUSCLE (version 3.8.1551) using default settings.[9]

Sequence statistics were calculated using faCount (version 377). Mean read depth (RD) was calculated

using JSI/SEQUENCE PILOT (JSI, Ettenheim, Germany) to evaluate the amplicon depth of each of

the 155 amplicons. For the validation samples (ONT group Table 1) the sequence starts and ends were

trimmed to match RC-PCR region with Oxford Nanopore region. A maximum-likelihood phylogenetic

tree was inferred using IQ-TREE (version 2.0.3) under the GTR$+$F$+$I$+$G4 model with the

ultrafast bootstrap option set to 1,000. Phylogenetic tree visualization and annotation was performed

using iTOL (version 5.6.3) or FigTree (version 1.4.4) (http://tree.bio.ed.ac.uk/software/figtree/).[10] SNP

distances between samples was calculated using snp-dists (version 0.7.0)

(https://github.com/tseemann/snp-dists). From the genome alignments we calculated a minimum

spanning tree (MST) by applying the MSTreeV2 algorithm using GrapeTree (version 1.5.0).[11]

Visualization of the MST was performed using GrapeTree.

The clinical validation consisted of a comparison of the epidemiological information of the

community and hospital samples and the WGS findings to see whether sequencing confirmed or

dismissed the suspected links between the samples.

Results

Technical results RC-PCR

In this study we performed three Illumina MiniSeq Mid Output (2x150 bp) runs containing 96 samples

each that were prepared using the EasySeq^{TM} RC-PCR SARS-CoV-2 WGS kit. It has a turnaround

time of about 8.5 hours, consisting of 1-hour hands-on time for preparing 96 samples, 6.5 hours for

performing the RC-PCR, and 1-hour of hands-on time for pooling, sample clean-up. Run 2 had the

highest number of positive SARS-CoV-2 VirSEAK consensus retrievals (100%). Of Run1 65% was

retrieved, Run1_new 67%. Run2, containing samples with higher viral loads (Ct values 16-32),

reached an average coverage of 96.69%. Genome coverage for Run1_new was 88%. (Figure 3B)

Supplementary table 4 provides a detailed overview of the technical results of the three sequence runs.

Amplicon depth plots

The amplicon depth distribution highlights which parts of the SARS-CoV-2 genome are represented

and the number of reads for each of the amplicons. In essence this shows how well the individual parts

of the SARS-CoV-2 genome are represented in the results. To illustrate the amplicon distribution on

the SARS-CoV-2 genome, for each of the 155 amplicons a sequencing depth was calculated and

plotted per run (See Figure 3A). Most amplicons are centered around a Mean read depth (RD) of 100-

1000. While some amplicons show less depth, in most cases they still result in a consensus sequence.

Additionally, for Run2 the interquartile range of the Mean RD is smaller compared to the two other

runs. When comparing the amplicon depth obtained per probe, boxplots are made for each run divided

in three Ct groups (Ct<20, 20≤Ct<27, and Ct≥27) (see Figure3C). We see a decline in depth for

samples with Ct above 27. For Run1_new and Run2 samples with a Ct between 20 and 27 perform

slightly better than the Ct<20 group this is probably an effect of the balanced library input strategy

applied for these runs. To evaluate if the impact of amplicon sequencing depth affects SARS-CoV-2

genome completeness, boxplots with the Ct groups are displayed to the effect on genome coverage

(see Figure 3D). Here we notice a decline in genome coverage with increasing Ct values for Run1 and

Run1_new. Run2 maintains high genome coverages, however does not contain samples with Ct values

above 32.

Figure 3. Graphical representation of performance of the RC-PCR Illumina sequence runs. **A.** Boxplots of the interquartile range of the Mean read depth (RD) of the Amplicons on a $\log_{10}$ scale for all 155 probes sorted on the SARS-CoV-2 genome. **B.** Boxplot of the Ct value as determined by RT-PCR to illustrate the differences of viral load of the sample per run. **C.** Boxplots of the Mean RD of the amplicons ($\log_{10}$ scale used) grouped per run and by Ct value. **D.** Boxplots of the SARS-CoV-2 genome coverage as achieved by RC-PCR grouped per run and Ct value.

Regions of low sequencing coverage

In a detailed analysis of the coverage of the SARS-CoV-2 genome obtained by RC-PCR 5 missing

genomic regions were observed (Table 2). The largest missing region has a length of 186 bp and is

part of the Open Reading Frame 1a (ORF1a). A further two regions are the start (1-54 bp) and the end

(46-165bp) of the genome. We observed that region 14585-14725 is missing in the VirSEAK

consensus output but not in the JSI/SEQUENCE PILOT and at the time of writing the manuscript the

VirSEAK algorithm was updated to improve the consensus output. Overall, without this update, the

maximum SARS-CoV-2 genome coverage that can be achieved using RC-PCR is between 97,8% and

98,2%. In version 1 of the EasySeq™ RC-PCR SARS-CoV-2 WGS kit three probe pairs do not

produce amplicons, 6258_6426, 9504_9752, and 21241_21420, respectively. No data on these

genomic regions will be obtained (Table 2).

Table 2: missing regions in VirSEAK consensus output				
VirSEAK consensus output		**JSI/SEQUENCE PILOT**		
Genomic location	**Length (bp)**	**Probes**	**Genomic location**	**Length (bp)**
1 - 54	54	No Probe		
6309 - 6407	99	6258_6426	6204 - 6372	169
9554 – 9739	186	9504_9753	9450 - 9699	250
14585 – 14725	141			
21322 - 21331	10	21241_21420	21187 - 21366	180
29739/29756/29858 – 29903	165/148/46	29630_29857	29576 - 29803	228
Total base-pairs	655/638/536			

Validation of RC-PCR reproducibility

All samples from Run1 that obtained a consensus (n=57) were compared to the same 57 samples from

Run1_new to determine whether results are reproducible when repeating sequencing with the RC-PCR

product. Results in supplemental figure 1 show that 50 of the 57 clusters fully align between Run1 and

Run1_new. There are 7 samples in which the phylogenetic distance is larger. For those samples in

which the phylogenetic distance is larger than expected, alignments were analyzed. The samples from

Run1_new show a lower genome coverage, explaining larger phylogenetic distances in these cases.

This is in line with the results observed in table 3 with average genome coverage of 88% in Run1_new

versus 93% in Run1. Which is either caused by RC-PCR product storage or the influence of the

balanced library pooling strategy based on Ct values of the samples.

Four sample pairs were tested in both Run1 and Run2 to serve as biological replicates. The entire

process from RNA isolation to sequence analysis was performed twice on these four samples.

Phylogenetic analysis depicted in Figure 4 (Illumina biological replicates) shows perfect agreement

between these repeats and confirms the specificity and reproducibility of RC-PCR.

Figure 4. Validation of RC-PCR by comparing the results for the same 19 samples sequenced previously by ONT.(Munnink *et al.*, 2020) Additionally, reproducibility was tested by applying RC-PCR on 4 samples as biological replicates (beige).

Validation of RC-PCR with Oxford Nanopore Technologies® (ONT)

Nineteen out of the 188 samples were tested using both ONT and Illumina® sequencing. The ONT

sequences were available in the GISAID database and compared to the results of RC-PCR sequencing.

All nineteen samples provided sequencing results on both platforms (Figure 4, ONT in red and RC-

PCR in blue). Fourteen out of nineteen samples provided perfect pairs, four samples show a small

divergence in the phylogenetic tree. Single nucleotide polymorphism (SNP) distance was calculated to

identify the number of nucleotides discrepant between samples. This in combination with manual

inspection showed that they have identical sequences but RC-PCR samples miss certain genomic

regions compared to ONT which results in the phylogenetic differences. One pair does not match, the

ONT sample shows a large distance (EPI ISL 422891). Manual inspection of the alignment revealed a

wrongly placed ambiguous region in the ONT sample.

Clinical validation

Of the 188 tested samples, 173 were SARS-CoV-2 positive of which sequencing results were obtained

for 146 (57 in Run1 and 89 in Run2). All samples, excluding nineteen ONT and four duplicate

samples used for validation, are depicted in the phylogenetic tree of Figure 5. Only HCW and patients

are included in the minimum spanning tree of Figure 6. Figure 5 shows the genetic diversity of the

samples at different time points during the pandemic. Those collected during the first months of March

and April (community samples from public health service) are clearly separated from the other

samples, especially compared to the samples from September 2020 (Cluster 1,2,3,6, and the HCW).

In Figure 6 it is clear the epidemiological link between the samples three of the six clusters was

completely confirmed by the sequencing results. Clusters two, five, and six contained HCW that were

not related. In cluster one, linked to a venue outside the hospital, five samples group together with no

SNP distances, one sample has a distance of a single SNP suggesting the possibility of linked cases.

However, multiple "sporadic HCW" tested in September and two HCW previously linked to cluster

two and five also group within cluster one.

In cluster two only two samples group together, two others are genetically unrelated samples and one

samples has a SNP distance of 2 which could still be within the transmission chain. Cluster three, a

patient and HCW show a distance of only 1 SNP. Sample collection was performed on one occasion,

twelve days apart, which could account for the SNP difference. In cluster four two HCW and a patient

group together, confirming the suspected link. Cluster five, an outbreak at a laboratory, eight HCW

samples have identical SARS-CoV-2 genomes, only one sample is phylogenetically linked. Cluster six

originated from a SARS-CoV-2 positive patient seen at a department, where at that time multiple

HCW had close contact to the patient. At the time of presentation, no symptoms were present that

were indicative of SARS-CoV-2 and screening using a questionnaire was negative. Five of the HCW

tested SARS-CoV-2 positive in the following weeks. In four HCW a genetically similar SARS-CoV-2

virus was detected. Surprisingly multiple other HCW group in this same cluster with minimal

differences (0-3 SNPs), which could mean the outbreak was larger than anticipated or the patient was

not the source of the infection.

Even though no new clusters were identified among the "sporadic HCW", they do group with

previously identified clusters. Additional information about these HCW revealed that many of them

had a direct or indirect link to the community source that was known by the public health services,

Cluster one.

Sequencing of the 64 community samples showed seven people clustered together in the phylogenetic

tree of Figure 5. There was no prior information available on these tested persons, but additional

information provided by the Local Public Health Service indicated that two of the seven worked at the

same location, two were their partners, the others lived in the same neighbourhood at the initial four

people, although they had no known epidemiological link to these people other than the area of

residence. Of other public health service samples no contact tracing information was available and

other samples clustering could not be confirmed with an epidemiological link.

Figure 5. Phylogenetic tree of 123 RC-PCR WGS SARS-CoV-2 genomes rooted at the NC 045512.2 reference genome. These samples are obtained from SARS-CoV-2 positive tested patients, HCW, and samples provided by public health services. Six clusters of samples were identified by the hospital Infection prevention control team. Sample groups are indicated by the colored blocks. Additional Ct values and genome coverage are plotted in barplots to illustrate the diversity in viral load between the samples and the high genome coverage that can be achieved by RC-PCR, respectively.

Figure 6. Minimum spanning tree of all samples being part of a cluster as defined by the infection prevention team and all HCW September samples. The tree was calculated using GrapeTree with the MSTreeV2 algorithm. This figure clearly illustrates the relationship between the samples and the clusters.

Discussion

In this study we present the application of a novel method called Reverse Complement-PCR to

sequence the SARS-CoV-2 genome which combines target amplification and indexing in a single

procedure, directly creating a sequencing ready Illumina library. We applied this method to 173

hospital and community samples that tested positive for SARS-CoV-2 with RT-PCR. Most

epidemiological clusters from the hospital and the community were confirmed by phylogenetic

clustering. Based on our data, RC-PCR is a reproducible technology, it correlates well with Oxford

Nanopore sequencing, is able to sequence samples with Ct values up to 32 determined by RT-PCR and

within these samples retrieves a high SARS-CoV-2 genome coverage. Optimization of the protocols is

expected to increase coverage in samples with lower viral loads even further.

Previous studies showed the benefit of using WGS of SARS-CoV-2 for outbreak investigation

purposes and to study transmission routes.[6,12-16] Several methods have been optimized for this purpose.

The ARTIC Illumina method, a tiling multiplex PCR approach, was the first that enabled WGS of

SARS-CoV-2 using Illumina sequencers.[17] The technique has subsequently been optimized and

analysis, albeit in small sample numbers, concluded that it delivers sufficient quality to perform

phylogenetic analysis.[18-20] It had been used as targeted and random RT-PCR screening with

subsequent sequencing of the population in order to study the spread through the community.[12] More

recently Sikkema *et al.* were the first to describe the use of SARS-CoV-2 sequencing in healthcare

associated infections and identify multiple introductions into Dutch hospitals through community-

acquired infections.[5]

SARS-CoV-2 has an estimated mutation rate of $1.12 \times 10{-3}$ substitutions per site per year, which

results in 2.8 mutations every month.[21] The minimum spanning tree of Figure 6 shows several samples

with a genetic distance of only a single SNP. With the mutation rate in mind, it is unclear how to relate

these clusters since extensive contact tracing information is lacking and interpretation on SNP

regarding outbreak management is unknown. Since community samples of September were

unavailable, we are unable to determine whether the genetic diversity in the community was low

resulting in genetically similar SARS-CoV-2 strains in a hospital setting. However, since sequencing

of samples in March and April 2020 clearly resulted in a larger diversity of SARS-CoV-2, and this

was early on in the pandemic, it seems more likely that a common source of infection, in- or outside

the hospital is the cause. Further research is needed to determine the accepted SNP distance for the use

in outbreak analysis.[22] Although we know minimum spanning trees are often used in outbreak

analysis.[5] It is a simplification of the phylogeny which could result in erroneous conclusions in

outbreak analysis. Care should be taken in interpreting these results.

It should be noted that some of the amplicons result in lower coverage than others (See Figure 3).

Currently, developments are under way in which a better distribution of the amplicon depth will be

achieved resulting in genome coverage that could increase to almost 100%. The difference in genome

coverage between Run1 and Run1_new is most likely caused by storage of the library and subsequent

pooling on the basis of Ct value of the individual samples, nonetheless, repeated testing at higher Ct

values will be needed to confirm this.

With current increase in infections in many countries including the Netherlands and additional

measures being put in place to reduce SARS-CoV-2 spreading, real-time sequencing of public health

service samples could be used to target infection prevention measures nationwide and locally.[23] Its

application can range from incidental cluster analysis in the case of uncertain epidemiological links to

real-time surveillance in the community or health care institutes. Additionally, correlation between

specific SARS-CoV-2 strains or mutations and clinical outcome could be identified, supporting

clinical decision making to improve outcomes for patients.[24,25]

In conclusion, here we implemented for the first time, RC-PCR in the field of medical microbiology

and infectious diseases thereby showing it to be a robust method which requires only minimal hands-

on time compared to current sequencing methods and can be used for high throughput sequencing of

SARS-CoV-2. Moreover, RC-PCR and sequence analysis can support epidemiological data with

genomic data to identify, monitor, and screen clusters of samples to help identify chains of

transmission of SARS-CoV-2, enabling a rapid, targeted and adaptive response to an ongoing outbreak

that has great impact on public health and society.

Author contributions

F.W. and J.P.M.C. conducted the research, performed analysis, wrote manuscript and created the

figures. L.F.J.vG., C.P.B-R., E.C.T.H.T., N.vdG-B., J.L.A.H. proofreading and provided clinical

information and samples of patients and HCWs. A.T.and J.H. conducted the contact tracing and

proofreading of the manuscript. H.F.L.W., J.C.R-L., M.S., and W.J.G.M supervised the study and

drafted the manuscript.

Conflict of interest disclosures:

The authors have no conflict of interest to disclose.

Funding/support:

The EasySeq™ RC-PCR SARS-CoV-2 WGS kit was supplied by NimaGen B.V and sequencing of

the Illumina libraries was performed by NimaGen B.V.. Validation was performed by the Department

of Medical Microbioly at the Radboud university medical center for the purpose of using the

technology in routine diagnostics. Therefore, no other funding was applied for.

Role of funder/sponsor:

NimaGen B.V. had no role in the design and conduct of the study; collection, management, data

analysis; preparation or approval of the manuscript.

References

1.   Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China,

2019. *New England Journal of Medicine* 2020.

2.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real

time. *The Lancet infectious diseases* 2020; **20**(5): 533-4.

3.   Reusken CB, Buiting A, Bleeker-Rovers C, et al. Rapid assessment of regional SARS-CoV-2

community transmission through a convenience sample of healthcare workers, the Netherlands, March

2020. *Eurosurveillance* 2020; **25**(12): 2000334.

4.   McLachlan S, Lucas P, Dube K, et al. The fundamental limitations of COVID-19 contact

tracing methods and how to resolve them with a Bayesian network approach. 2020.

5.   Sikkema RS, Pas SD, Nieuwenhuijse DF, et al. COVID-19 in health-care workers in three

hospitals in the south of the Netherlands: a cross-sectional study. *Lancet Infect Dis* 2020; **20**(11):

1273-80.

6.   Munnink BBO, Nieuwenhuijse DF, Stein M, et al. Rapid SARS-CoV-2 whole-genome

sequencing and analysis for informed public health decision-making in the Netherlands. *Nature*

medicine 2020; **26**(9): 1405-10.

7.   Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by

real-time RT-PCR. *Eurosurveillance* 2020; **25**(3): 2000045.

8.   DNA Pipelines R&D BF, Diana Rajan, Emma Betteridge, Lesley Shirley, Michael Quail,

Naomi Park, Nicholas Redshaw, Iraad F Bronner, Louise Aigrain, Scott Goodwin, Scott Thurston,

Stefanie Lensing, Charlotte Beaver, Ian Johnston. COVID-19 ARTIC v3 Illumina library construction

and sequencing protocol V.1. 2020. https://www.protocols.io/view/covid-19-artic-v3-illumina-library-

construction-an-beuzjex6?version_warning=no (accessed 10th september 2020.

9.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res 2004; **32**(5): 1792-7.

10.   Minh BQ, Schmidt HA, Chernomor O, et al. IQ-TREE 2: New Models and Efficient Methods

for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* 2020; **37**(5): 1530-4.

11.   Zhou Z, Alikhan N-F, Sergeant MJ, et al. GrapeTree: visualization of core genomic

relationships among 100,000 bacterial pathogens. *Genome research* 2018; **28**(9): 1395-404.

12.   Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS-CoV-2 in the Icelandic

Population. *N Engl J Med* 2020; **382**(24): 2302-15.

13.   Meredith LW, Hamilton WL, Warne B, et al. Rapid implementation of SARS-CoV-2

sequencing to investigate cases of health-care associated COVID-19: a prospective genomic

surveillance study. *Lancet Infect Dis* 2020; **20**(11): 1263-72.

14.   Quick J, Grubaugh ND, Pullan ST, et al. Multiplex PCR method for MinION and Illumina

sequencing of Zika and other virus genomes directly from clinical samples. *Nat Protoc* 2017; **12**(6):

1261-76.

15.   Richard M, Kok A, de Meulder D, et al. SARS-CoV-2 is transmitted via contact and via the

air between ferrets. *bioRxiv* 2020.

16.   Stefanelli P, Faggioni G, Lo Presti A, et al. Whole genome and phylogenetic analysis of two

SARS-CoV-2 strains isolated in Italy in January and February 2020: additional clues on multiple

introductions and further circulation in Europe. *Euro Surveill* 2020; **25**(13).

17.   Quick J. nCoV-2019 sequencing protocol 2020. *Publisher Full Text* 2020.

18.   Batty EM, Kochakarn T, Wangwiwatsin A, et al. Comparing library preparation methods for

SARS-CoV-2 multiplex amplicon sequencing on the Illumina MiSeq platform. *BioRxiv* 2020.

19.   Pillay S. Illumina Nextera DNA Flex library construction and sequencing for SARS-CoV-2:

Adapting COVID-19 ARTIC protocol. 2020. https://www.protocols.io/view/illumina-nextera-dna-

flex-library-construction-and-bhjgj4jw (accessed September 30th 2020.

20.   Pillay S, Giandhari J, Tegally H, et al. Whole Genome Sequencing of SARS-CoV-2: Adapting

Illumina Protocols for Quick and Accurate Outbreak Investigation during a Pandemic. *Genes (Basel)*

2020; **11**(8).

21.   Koyama T, Platt D, Parida L. Variant analysis of SARS-CoV-2 genomes. *Bull World Health*

Organ 2020; **98**(7): 495-504.

22.   Yin C. Genotyping coronavirus SARS-CoV-2: methods and implications. *Genomics* 2020.

23.   RIVM. Current information about COVID-19 (novel coronavirus). October 13th 2020 2020.

https://www.rivm.nl/en/novel-coronavirus-covid-19/current-information. (accessed October 15th 202.

24.   Gong YN, Tsao KC, Hsiao MJ, et al. SARS-CoV-2 genomic surveillance in Taiwan revealed

novel ORF8-deletion mutant and clade possibly associated with infections in Middle East. *Emerg*

Microbes Infect 2020; **9**(1): 1457-66.

25.   Wang C, Liu Z, Chen Z, et al. The establishment of reference sequence for SARS-CoV-2 and

variation analysis. *J Med Virol* 2020; **92**(6): 667-74.

461