Is degree of sociality associated with reproductive senescence? A comparative analysis across birds and mammals

Csongor I. Vágási[1,*], Orsolya Vincze[2], Jean-François Lemaître[3], Péter L. Pap[1], Victor Ronget[4] and Jean-Michel Gaillard[3,*]

[1]Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babe□-Bolyai University, Cluj-Napoca, Romania

[2]Department of Tisza Research, MTA Centre for Ecological Research-DRI, Debrecen, Hungary

[3]Laboratoire de Biométrie et Biologie Évolutive, CNRS, Université Lyon, Villeurbanne, France

[4]Unité Eco-anthropologie (EA), Muséum National d'Histoire Naturelle, CNRS, Université Paris Diderot, Paris, France

*Author for correspondence:

Csongor I. Vágási

e-mail: csvagasi@gmail.com

Jean-Michel Gaillard

e-mail: jean-michel.gaillard@univ-lyon1.fr

Running title: Sociality and reproductive senescence

ORCID: 0000-0002-8736-2391 (CIV), 0000-0001-5789-2124 (OV), 0000-0001-9898-2353 (JFL), 0000-0002-3659-7684 (PLP), 0000-0001-9991-0482 (VR), 0000-0003-0174-8451 (JMG)

Our understanding on how widespread reproductive senescence is in the wild and how the onset and

rate of reproductive senescence vary among species in relation to life histories and lifestyles is

currently limited. More specifically, whether the species-specific degree of sociality is linked to the

occurrence, onset and rate of reproductive senescence remains unknown. Here, we investigate these

questions using phylogenetic comparative analyses across 36 bird and 101 mammal species

encompassing a wide array of life histories, lifestyles and social traits. We found that female

reproductive senescence (1) is widespread and occurs with similar frequency (about two thirds) in

birds and mammals; (2) occurs later in life and is slower in birds than in similar-sized mammals; (3)

occurs later in life and is slower with an increasingly slower pace of life in both vertebrate classes;

and (4) is only weakly associated, if any, with the degree of sociality in both classes after

accounting for the effect of body size and pace of life. However, when allowing for species

differences in pace of life, a higher degree of sociality was associated with later and weaker

reproductive senescence in females, which suggests that degree of sociality is indirectly related to

reproductive senescence via the pace of life.

Keywords:

brain size, coloniality, cooperative breeding, life history, reproductive ageing, vertebrates

Subject Areas:

ecology, evolution

1. Introduction

Reproductive senescence (or reproductive ageing) – the decline in reproductive performance with increasing age – is widespread in nature [1,2], except for species with indeterminate growth that gain mass and thereby increase fecundity with age [3]. Recent studies have revealed that both the timing and the strength of reproductive senescence is highly variable across species [4,5], although our knowledge is still very limited about how ecological factors and species-specific life history shape variation in either the onset or the rate of reproductive senescence [1,6]. Among these factors, the possible role played by the species-specific degree of sociality has never been investigated.

Sociality is evolutionarily associated with a complex set of life-history traits. Most notably, social species might have longer lifespan and decreased actuarial senescence (see [7–9] for reviews). Indeed, social life in cooperative breeders and colonial species can buffer environmentally-driven mortality risks and might ultimately slow down actuarial senescence (e.g. [10] for a case study on cooperatively breeding Seychelles warblers, *Acrocephalus sechellensis*), even if the relationship between sociality and actuarial senescence is likely to be complex and might differ both within and among species [7]. However, the association between social life and the occurrence, onset and rate of reproductive senescence has never been investigated so far, although similar relationship with the intensity of senescence is expected for survival and reproduction. We aimed here to fill this knowledge gap using the most comprehensive comparative analyses performed to date across bird and mammal species.

Within populations, the evolutionary hypotheses explaining why social individuals should display a weaker senescence than solitary ones [7] are rooted in the principle of energy allocation [11]. This principle states that increased allocation of finite resources to a given biological function (e.g. reproduction) compromises allocation to a competing function (e.g. somatic maintenance that promotes survival) [12]. Increased allocation of resources to reproduction early in life, which is favoured by natural selection in growing populations [13], is expected to have detrimental

consequences in terms of actuarial and/or reproductive senescence [6]. This trade-off is predicted by

both antagonistic pleiotropy and disposable soma theories of ageing [14,15], and is well supported

by current empirical evidence [16,17]. For instance, male red deer (*Cervus elaphus*) allocating

substantial resources to sexual competition during early life show a steeper rate of reproductive

senescence in late life ([18]; see also [19] for examples in birds). How social lifestyle may buffer

against such costs? For instance, helpers in cooperative breeders reduce the workload of

reproducers according to the load-lightening hypothesis [20]. Thus, the principle of allocation, a key

concept of life-history evolution [12,21], explains how senescence can either increase due to a

delayed cost of high performance during early life [16,22] or decrease thanks to a reduced

reproductive effort allowed by a high degree of sociality (e.g. the presence of helpers [20]).

Assuming that these processes can explain the variance in senescence observed at the inter-specific

level, two hypotheses can be proposed to expect a negative covariation between the degree of

sociality and reproductive senescence:

(H1) Given the inevitable costs of reproduction [23,24], a high reproductive effort observed

at a given species should lead to an earlier and/or faster reproductive senescence [16]. For a

given reproductive effort, a higher degree of sociality in a given species might facilitate the

reproductive duties of individuals and therefore reduce directly the costs of reproduction [7]

and ultimately shape the senescence patterns of that species [7,17]. Thus, the mitigation of

reproductive cost by a social mode of life should lead to postponed onset and/or decelerated

rate of reproductive senescence of a given species.

(H2) The degree of sociality can drive the evolution of reproductive senescence in a given

species indirectly through decreasing adult mortality risk, thereby slowing down the pace of

life. Life-history theory postulates that a decreased rate of environmentally-driven mortality

should favour slower growth rate, longer time to maturation, older age at first reproduction

and reduced allocation to reproduction by young adults [12], as well as later onsets and

slower rates of both actuarial and reproductive senescence [6,25]. Indeed, sociality has been

shown to mitigate multiple forms of environmentally driven mortality risks (e.g. starvation,

predation). Thus, the presence of social partners in a given species might indirectly lead to

delayed or decelerated senescence via slowing down the pace of life of that species [26].

Under both hypotheses, reproductive senescence should be less pronounced in species with a higher

degree of sociality by involving either a direct response of the reproductive effort at each

reproductive attempt (H1) or indirectly through a slower pace of life selecting for a lower

reproductive effort early in life (H2). If the degree of sociality is directly associated with

reproductive senescence (H1), we predict a substantial effect of the degree of sociality even after

the effects of allometry and pace of life on reproductive senescence are accounted for. If the degree

of sociality is indirectly associated with reproductive senescence via the pace of life (H2), we

predict no detectable effect of the degree of sociality once the effects of allometry and pace of life

are accounted for.

Here, we modelled age-specific changes in reproductive traits at the species level and tested

whether the degree of sociality accounts for the variation in the occurrence, onset and rate of

reproductive senescence observed across birds and mammals ($n = 36$ and 101 species, respectively).

We accounted for the confounding effect of phylogenetic inertia, allometric constraints and species'

ranking on the slow–fast continuum of life histories (i.e. pace of life) in our phylogenetic

comparative analyses, as all these processes are known to shape variation in senescence [5].

2. Methods

(a) Female reproductive senescence data

As age-specific reproductive output is easier to measure in females than in males (e.g. due to extra-

pair offspring often produced by males; [27]) and has been reported in a much higher number of

vertebrate species, we focus on the reproductive ageing of females in both birds and mammals.

Reproductive senescence parameters of 101 wild or semi-captive mammal species were taken from

[28]. This data set includes the presence/absence of reproductive senescence and, for species with

evidence of senescence, the age at onset and the rate of reproductive senescence. All those

parameters were estimated from age-specific birth rates (i.e. number of female offspring alive at

birth that are produced by a female of age x, tabulated as m_x in a life table) extracted from published

life tables or graphs using WebPlotDigitizer (https://automeris.io/WebPlotDigitizer/). The

acquisition of age-specific reproductive data for mammals is fully detailed in [28]. In cooperatively

breeding mammals, age-specific reproductive data were collected for dominant females (e.g. [29]),

as subordinate females have – by definition – no access to reproduction.

In birds, we conducted a systematic literature search of age-specific changes in reproductive traits

in wild populations to extract data similar to those obtained for mammals (see Electronic

Supplementary Material, ESM for search methods). Unlike in mammals, age-specific birth rates

(i.e. the m_x parameter) were seldom reported in bird studies because the probability of breeding –

necessary for birth rate calculations – is often unknown. Therefore, to increase the number of

species, we also included age-specific number of hatchlings or number of fledglings per female

when birth rates could not be extracted or computed. Some studies reported standardized values (i.e.

normalized values or residuals from models) instead of raw values of age-specific reproduction. We

included those studies in our analyses and controlled for the effect of standardization (yes/no).

When reproductive data were reported for multiple populations of the same species, we only

included the study with the largest sample size, as done in mammals [28]. To estimate senescence

parameters, we accounted for differences in the age-specific sample sizes, as done in mammals

[28]. We used the original age-specific sample size when reported in the original studies, and we

calculated the number of females expected to be alive at age x from the observed age distribution of

females when sample sizes were not reported. We collected female reproductive data for 36 avian

species (see ESM 'Data set').

Age-dependent reproductive traits in birds followed similar distributions to mammalian ones.

Hence, we computed reproductive senescence parameters in birds using the same methods as in

mammals. Briefly (see [28] for further details), four different age-dependent models (i.e. constant

model, linear model, threshold model with one threshold and two linear segments, threshold model

with two thresholds and three linear segments) weighted by the age-specific sample size were fitted

on the reproductive data using the R package 'segmented' [30]. The final model was selected using

Akaike's Information Criterion (AIC) (see the Methods in the ESM for model selection procedure

and ESM 'Model selection' for the AIC values associated to each alternative senescence models;

see also ESM 'Segmented' for the segmented fits of the selected models plotted separately for each

bird species; similar table and plots for mammals can be found in [28]). Based on the selected

model, different procedures were used to infer reproductive senescence from the slope of the

different linear segments and their associated standard error. When reproductive senescence

occurred (i.e. slope of one of the segments < 0), the rate and the onset of reproductive senescence

were reported as the slope of the linear segment and the age corresponding to the beginning of the

segment, respectively. Using this procedure, we detected reproductive senescence for most of the

bird species for which it was observed in the original studies from which the data were extracted,

only minor discrepancies were found mostly due to the use of different statistical methods (see

ESM 'Occurrence' for a comparison of the results found on reproductive senescence using our

standardized procedure against the results found in the original studies; a similar comparison for

mammals can be found in [28]).

(b) Life-history traits

To assess the relationship between the degree of sociality and reproductive senescence, we first had

to account for inter-specific differences in body size and biological time [31], which structure most

life-history variation across vertebrates [32]. Body mass is a reliable measure of species-specific

size that shapes age-specific reproductive and survival rates via allometric effects. Thus, small bird

and mammal species display both earlier and steeper reproductive senescence than large ones [5].

Likewise, for a given size, slow-living species display both later and slower reproductive

senescence, an effect well illustrated by the comparison of similar-sized birds and mammals [5].

Generation time is the most appropriate metric to position species on the slow–fast continuum of

life histories [33]; however, data to accurately measure generation time were missing for many of

the species studied here [34]. Thus, instead of generation time, we used a compound of the age at

first reproduction and maximum longevity observed in the focal case study to measure species-

specific pace of life. In birds, we collected data on female body mass from [35], age at first

reproduction and longevity from the same papers including age-specific reproduction data (ESM

'Data set'), while in mammals data of the same traits were retrieved from [28].

(c) Sociality traits

The social environment varies considerably across species and this diversity can have vast

evolutionary consequences [36]. We use four simple sociality traits (i.e. cooperative breeding,

coloniality, parental cooperation and relative brain size; see also [37,38]) to assess the species-

specific degree of sociality (Table 1) and test whether these traits are associated or not with the

occurrence, rate and onset of reproductive senescence across birds and mammals. These four

proxies of sociality cover different ranges of degree of sociality. For instance, cooperative breeders

often live in social systems with more complex social interactions than colonial ones, and therefore

imply different costs and benefits to the individuals. The diversity of social traits we use in this

study makes possible to assess whether social lifestyle in general or specific social systems in

particular are associated with reproductive senescence, if any.

We used three sociality traits in birds (i.e. presence/absence of coloniality, parental cooperation and

relative brain size) and two sociality traits in mammals (i.e. presence/absence of cooperative

breeding and relative brain size). Both the degree of sociality and the use of social information are

higher in species breeding in large and dense colonies of non-kin individuals as compared with

solitarily breeding ones [39], and hence the degree of coloniality has been considered as a proxy of

the sociality degree in studies of longevity across bird species [37]. Because coloniality cannot be

defined with confidence in mammals, but in a few species only (e.g. in black-tailed prairie dogs,

Cynomys ludovicianus; [40]), we had to omit this trait in this vertebrate class. We used parental

cooperation as a metric of the degree of sociality, which reflects whether female-only, male-only or

shared female–male parental care is typical for a given bird species [41]. Family, where individuals

form short-term pair bonds during breeding and raise their offspring cooperatively (i.e. have

biparental care) is the simplest social system, and species with biparental care display a higher

degree of sociality than species in which only females care for their young [38]. This metric is

relevant in birds because it influences the reproductive costs of females and thus is likely to

modulate female reproductive senescence parameters. Biparental care is the most common form of

social behaviour between unrelated individuals in birds, with over 90% of all living birds being

biparental [42]. The degree of sociality is considered high (i.e. implying frequent and complex

social interactions among individuals) in species living in small cooperative breeding groups with

helpers as compared with non-cooperatively breeding ones. Cooperative breeders have the most

intense social system among mammals [43]. The presence/absence of cooperative breeding was not

considered in birds due to the low number of species with regular cooperative breeding in our data

set. Lastly, the relative brain size (i.e. brain size for a given body size) is higher in species with high

degree of social bonding (e.g. primates and whales/dolphins) or reproductive pair bonding (e.g.

monogamous carnivores and ungulates, bats and birds) [38], making possible its use to measure the

degree of sociality [38,44]. Quantifying the degree of sociality in comparative studies encompassing

species with a large range of life-history strategies is far from trivial, which leads most comparative

studies to use only proxies of sociality instead of accurate metrics.

Table 1. Sociality traits and their meaning in terms of degree of sociality.

sociality trait	degree of sociality	
	low	high
colonial breeding (birds)	no	yes
parental cooperation (birds)	female care	female & male care
cooperative breeding (mammals)	no	yes
relative brain size (birds and mammals)	small	large

In birds, we collected data on brain mass from [45], presence/absence of coloniality from [46] and

parental cooperation during breeding from [41] (see ESM 'Data set'). In the latter source, parental

cooperation was separately quantified for the pre- and post-hatching periods, which are highly

correlated (Pearson correlation, $r = 0.76$, df $= 29$, $t = 6.24$, $p < 0.0001$). We calculated the average

of these two periods (henceforth parental cooperation) reflecting the sex bias in parental care during

breeding. Values range from –1 (exclusive female care) to 1 (exclusive male care), with 0 reflecting

an equal share of parental duties between sexes. Coloniality had a perfect overlap with marine

environment in our bird data set, as all colonial species are seabirds and all solitary ones are

terrestrial, which reflects a strong phylogenetic bias and a limitation of our coloniality data. In

mammals, data on brain mass were obtained from [47], presence/absence of cooperative breeding

from [48] and we completed species with lacking information with additional sources (see ESM

'Data set').

(d) Statistical analyses

All analyses were performed in R version 4.0.1 [49]. To make meaningful inferences about the

effect of body size, pace of life and degree of sociality on reproductive senescence, all models were

controlled for phylogenetic inertia. In birds we used a rooted, ultrametric consensus tree built using

the SumTrees Python library [50] based on 1,000 trees. These trees were obtained from birdtree.org

[51] using the Hackett backbone tree [52]. For mammals, we used a published phylogenetic super-

tree (see also [53]).

Female body mass, age at first reproduction, longevity and brain mass were highly correlated across

both bird and mammal species (Table S1). Therefore, to avoid multicollinearity problems, we

conducted a phylogenetically-controlled principal component analysis (PPCA) as implemented in R

package 'phytools' [54] on the first three traits (all log-transformed) separately for birds and

mammals. We retained the first two phylogenetic principal components (PPCs), where the first PPC

is a size component (hereafter PPC size), which explained 69% and 79% of variation in birds and

mammals, respectively, and the second PPC is a pace of life component (hereafter PPC pace),

which explained additional 23% and 12% of variation in birds and mammals, respectively (Table

S2). Larger values indicate larger body mass (PPC size) and slower pace of life (PPC pace),

respectively (Table S2). PPC size and PPC pace were used in the subsequent analyses to control for

allometry and pace of life, respectively. Given that we were specifically interested in the effect of

relative brain size (a proxy measure of the degree of sociality) on reproductive senescence, we did

not include brain size in the PPCA. Nonetheless, to avoid collinearity of brain size with PPC size,

we estimated relative brain size as residuals of a standard major axis regression (as implemented in

R package 'lmodel2') between log-transformed brain size and PPC size and used this measure in

the multifactorial models.

To explore variation in reproductive senescence patterns, we used phylogenetic logistic regressions

for evidence of reproductive senescence and phylogenetic linear regressions separately for onset

and rate of reproductive senescence as implemented in R package 'phylolm' [55]. Age at onset and

the absolute value of the rate of reproductive senescence were log-transformed prior to the analysis.

In birds, for each senescence metric, the reproductive trait used to assess reproductive senescence

(i.e. birth rate m_x, number of hatchlings or number of fledglings) and the presence/absence of

coloniality were tested as fixed factors, while PPC size, PPC pace, residual brain size and parental

cooperation were included as covariates. We did not need to account for either the hunting status

(i.e. no bird species in the data set is hunted) or the data quality (i.e. all bird studies were based on

longitudinal data and only included known-aged individuals). Similarly to analysis in mammals (see

[28]), we tested whether the probability to detect reproductive senescence in birds was influenced

by the sample size (i.e. total number of reproductive records in the population; log-transformed in

the analysis; ESM 'Data set'). For the rate of reproductive senescence, the effect of data

standardization (yes/no) was also tested. In mammals, for each senescence metric, data quality

(transversal/longitudinal), hunting status (hunted/not hunted) and presence/absence of cooperative

breeding were included as fixed factors, while PPC size, PPC pace and residual brain size were

included as covariates. In both birds and mammals, the effect of age at onset of reproductive

senescence (log-transformed) was also tested in models of rate of reproductive senescence because

a negative correlation is expected to occur [28]. Non-linear effects of PPC size and PPC pace were

also modelled in both bird and mammal models using second-degree orthogonal polynomials, but

were only retained in the model when their inclusion decreased AIC values by > 2 compared with

the initial model without the polynomials. In no case where a quadratic model was selected over the

linear model did a cubic model outperform the quadratic model, meaning that a second-order

polynomial satisfactorily accounted for observed non-linear relationships. Sample size varied across

models because some variables (e.g. brain size) had missing values in certain species and rate as

well as onset of senescence were only analysed for species in which evidence of reproductive

senescence was detected. To test H2 according to which the effect of the degree of sociality acts

indirectly through slowing down the pace of life, we repeated all the above-mentioned models after

removing PPC pace from the models.

Due to the limited number of bird species, we adopted an AIC-based stepwise forward model

selection procedure to avoid over-parametrization of models. As a first step, an intercept model was

constructed for each dependent variable. In the second step, each explanatory variable (except

metrics of sociality) was added one by one to this model and the model with the smallest AIC value

(if ΔAIC < 2) was further elaborated until adding extra variables did not decrease AIC value by > 2.

If any of the single-predictor models had $\Delta AIC < 2$, the intercept model was used as the base

model. In the third step, to test the association between the degree of sociality and reproductive

senescence, the sociality traits were added one by one to the base model and the change in AIC was

checked. Given that relative brain size and parental cooperation had missing values for some

species, when testing their effect on reproductive senescence metrics, their corresponding base

models were refitted for the subset of species with the full set of available data. These models are

presented in birds (Table S3). Given the large sample size in mammals, we present the full models

with all explanatory variables entered simultaneously (Table S4). Consequently, the final sample

size is 88 mammalian species (out of 101 species) because brain size data were missing for 13

species; however, repeating the analyses by excluding brain size and keeping only cooperative

breeding as sociality trait, which is available for the entire species pool, the results of cooperative

breeding remain unchanged (results not shown).

3. Results

(a) Occurrence of senescence in birds and mammals

Reproductive senescence was detected in 61% (22 out of the 36 species) of bird species and 68%

(69 out of 101 species) of mammal species. The occurrence of reproductive senescence was similar

in birds and mammals (Chi-squared test $\chi^2 = 0.34$, df $= 1$, $p = 0.562$). The probability of detecting

reproductive senescence tended to increase with sample size in birds ($\beta \pm$ s.e. $= 0.43 \pm 0.31$, $p =$

0.16), but this effect was not significant (as opposed with mammals, see [28]).

(b) Allometry, pace of life and the degree of sociality in birds

Results of occurrence, rate and onset of reproductive senescence in birds are presented in Table S3.

The occurrence of reproductive senescence in birds was unrelated to body size and pace of life, and

was independent of the reproductive trait used to assess reproductive senescence. None of the

sociality traits was associated with the probability to detect reproductive senescence (Table S3).

The rate of reproductive senescence decreased non-linearly with increasing body size (linear term: β

$\pm$ s.e. $= -3.23 \pm 0.81$; quadratic term: $\beta \pm$ s.e. $= -1.41 \pm 0.69$), decreased linearly with an

increasingly slower pace of life ($\beta \pm$ s.e. $= -0.89 \pm 0.23$), and varied among reproductive traits used

to assess reproductive senescence. The rate of reproductive senescence was the slowest when using

birth rates, intermediate when using the number of hatchlings and fastest when using the number of

fledglings. Data standardisation did not explain substantial variation in the rate of reproductive

senescence either. The rate of reproductive senescence tended to decrease with increasingly later

onset of senescence, although this effect was not statistically significant. None of the sociality traits

was associated with the rate of reproductive senescence, which does not support H1. Once the

marked effect of pace of life was removed from the model, the rate of reproductive senescence was

slower in colonial birds than in solitary breeders ($\beta \pm$ s.e. $= -1.06 \pm 0.35$), in support of H2 (Fig. 1*a*,

Table S3).

The onset of reproductive senescence increased linearly with both body size ($\beta \pm$ s.e. $= 0.32 \pm 0.04$)

and slower pace of life ($\beta \pm$ s.e. $= 0.39 \pm 0.14$). None of the sociality traits was related to the age at

onset of reproductive senescence, which does not support H1. Once the strong effect of pace of life

was removed from the model, the onset of reproductive senescence was later in colonial birds than

in solitary species ($\beta \pm$ s.e. $= 0.41 \pm 0.21$), in support of H2 (Fig. 1*b*, Table S3).

These results do not support H1, but do support H2, which involves an indirect relationship between

degree of sociality and both the rate and onset of reproductive senescence via a slowing down of the

overall pace of life in species with higher degree of sociality.

Figure 1. (*a*) Predicted residual rate and (*b*) residual onset of reproductive senescence (± s.e.m.) in

function of coloniality in birds once PPC pace (i.e. pace of life) is removed from the models. The

residual rate was extracted from a linear regression including PPC size (polynomial) and

reproductive trait as explanatory variables. The residual onset was obtained from a linear regression

including PPC pace as an explanatory variable. The relationship between relative brain size and (*c*)

the residual rate and (*d*) the residual onset of reproductive senescence in mammals once PPC pace is

removed from the models. The residual rate and onset of senescence were extracted from linear

regressions including PPC size, but not including the otherwise statistically significant effect of

PPC pace. Slopes were obtained from a linear regression between residual brain size (explanatory)

and the residual rate and onset (dependent), respectively.

(c) Allometry, pace of life and the degree of sociality in mammals

Results of occurrence, rate and onset of reproductive senescence in mammals are presented in Table

S4.

Reproductive senescence was more likely to be detected when data originated from longitudinal

rather than transversal studies ($\beta \pm$ s.e. $= -1.29 \pm 0.52$). Larger-sized mammals were more likely to

experience reproductive senescence than smaller ones ($\beta \pm$ s.e. $= 0.23 \pm 0.10$). Neither relative brain

size, nor cooperative breeding was related to the probability to detect reproductive senescence in

mammals (Table S4).

The rate of reproductive senescence decreased linearly with increasing body size ($\beta \pm$ s.e. $= -0.43 \pm$

0.06) and non-linearly with increasingly slower pace of life (linear term: $\beta \pm$ s.e. $= -4.45 \pm 1.35$;

quadratic term: $\beta \pm$ s.e. $= -2.41 \pm 1.08$). Contrary to the H1 expectation, cooperative breeding

mammals had higher rates of reproductive senescence compared to non-cooperative species ($\beta \pm$

s.e. $= 1.3 \pm 0.43$), while relative brain size was unrelated to the rate of reproductive senescence.

When the marked effect of pace of life was removed from the model, species with larger relative

brain size had slower rate of reproductive senescence ($\beta \pm$ s.e. $= -0.64 \pm 0.32$; Fig. 1c), in support to

H2. When the pace of life was not controlled for, however, the relationship between cooperative

breeding and the rate of reproductive senescence disappeared, which does not support H1 (Table

S4).

The age at onset of reproductive senescence increased linearly with both body size and increasingly

slower pace of life ($\beta \pm$ s.e. $= 0.23 \pm 0.05$). Neither relative brain size, nor cooperative breeding was

related to the age at onset of reproductive senescence in mammals, which does not support H1.

Once the marked effect of pace of life was removed from the models, species with large relative

brain size showed a later onset of senescence than species with small relative brain size ($\beta \pm$ s.e. =

0.38 ± 0.19, Fig. 1d, Table S4), in support to H2.

As in birds, these results do not support H1, but support H2 that involves an indirect relationship

between degree of sociality and the rate and onset of reproductive senescence via a slowing down of

the overall pace of life in species with higher degree of sociality.

(d) Comparing reproductive senescence between birds and mammals

In both classes, the rate of reproductive senescence tended to decrease with increasingly later onset

of reproductive senescence, with the same apparent strength (Fig. 2a). However, the relationship

was statistically significant only in mammals likely because of a lack of power (smaller sample

size) in birds. When looking at the allometric relationships, the rate of reproductive senescence

decreased and the onset of reproductive senescence occurred later with increasing size in both birds

and mammals (Fig. 2b,c). Interestingly, for a given body mass, mammals displayed both steeper

and earlier reproductive senescence than birds did (Fig. 2b,c), which is in line with the common

view that birds senesce less than similar-sized mammals.

Figure 2. Association between (*a*) age at onset and rate of reproductive senescence, (*b*) female

body mass and rate of reproductive senescence, and (*c*) female body mass and age at onset of

reproductive senescence across bird species (* non-colonial, × colonial) and mammal species (▲).

Female body mass was used here to measure body size because it captures the differences in size

range between birds and mammals unlike PPC size, and it is very strongly correlated with PPC size

(Table S2). Slopes were obtained from single-predictor phylogenetic regressions between the

plotted variables (dashed line for birds, continuous line for mammals). Polynomial effects are only

plotted where statistically significant (see Tables S3 and S4).

4. Discussion

A previously published review revealed an increasing number of case studies reporting reproductive

senescence in the wild [2]. Here, we quantified the occurrence of this phenomenon on the largest

species-level data set so far compiled on birds and mammals. We found that the proportion of

species that display detectable reproductive senescence is similar in avian (0.61; present study) and

mammalian (0.68; [28]) species. Interestingly, these proportions are similar to those reported in a

previous comparative study of 19 species of birds and mammals (0.65; [5]). However, as the current

prevalence of reproductive senescence is likely to be under-estimated (see [28] for further

discussion), the biological meaning of these values is disputable. Nevertheless, these studies

together emphasize that reproductive senescence is the rule rather than the exception, at least in

endotherm vertebrates. The positive effect of sample size on the probability of detecting senescence

in mammals (see [28]) constitutes a limitation of our analyses, although this limitation is not

detectable for birds, likely due to the smaller data set in this class.

Our findings highlight that birds display a later onset and a slower rate of reproductive senescence

as compared with similar-sized mammals. Note that the strength of senescence in birds increases

with offspring development (i.e. from birth rate to number of fledglings), while for mammals

senescence was only computed for birth rates. Therefore, we expect that the differences in onset and

rate of reproductive senescence between the two classes would be even stronger if weaning success

were also considered in mammals. This notion is supported by a recent review showing that

maternal effect senescence (i.e. an increasing offspring mortality with mother age) is very common

in mammals [56]. The more intense reproductive senescence in mammals than in same-sized birds

we report matches the class differences reported in longevity (i.e. birds live c.a. 1.5 times longer

than similar-sized mammals [57]). Birds also display a much slower pace of life, and, for a given

pace of life, birds and mammals of a given size have similar senescence patterns [5]. This suggests

that the *modus operandi* of senescence has a deep evolutionary root and is mostly shaped by

allometric constraints and pace of life. To test whether the differences between the two classes are

explained by flight capacity in birds, and hence their lower environmentally driven mortality, a

comparison of reproductive senescence between birds and flying mammals would be promising (see

[58] for longevity).

In line with previous observations for other biological times (e.g. longevity, gestation length; see

[59]), we found strong effects of allometry and pace of life on both the rate and the onset of

reproductive senescence in both birds and mammals. The heavier and slower-paced a species is, the

more postponed and slower its senescence is. Both senescence metrics correspond to biological

times with a dimension of time for the onset of senescence and a dimension of frequency (i.e.

inverse of time) (sensu [31]) for the rate of senescence, which explains the negative relationship we

found between the rate and the onset of senescence across birds and mammals. Our analyses thus

provide a first evidence that these senescence metrics can be interpreted as life-history traits

describing the speed of the life cycle of a given species, alike development time [60], age at first

reproduction [61] or longevity [62], which have been much more intensively studied. Our results,

which are based on the largest number of bird and mammal species compiled to date, bring

convincing support that the process of reproductive senescence is embedded in the life-history

strategy of a given species [5,16,17] and has a role in the evolution of life histories.

The degree of sociality appears to have a very limited direct influence on reproductive senescence

when the effects of allometry and pace of life are accounted for, which supports the view expressed

above that reproductive senescence in a given species is mostly driven by the species size and

position on the slow-fast continuum of life histories. With the exception of cooperative breeding in

mammals, none of the sociality traits we analysed (i.e. relative brain size in birds and mammals,

colonial breeding and parental cooperation in birds) were associated with either the occurrence or

the rate and onset of reproductive senescence. These results support the conclusions reached about

the putative role of sociality in actuarial senescence and longevity [7].

One striking result of this work is that the degree of sociality was associated with a decreased

strength of senescence in terms of both rate and onset when species differences in pace of life were

not controlled for. As these associations vanished when we controlled for the pace of life, we

conclude that the social mode of life *per se* does not influence reproductive senescence. Instead, the

social lifestyle shapes the entire life-history strategy, which supports H2 and refutes H1. Thus,

cooperative breeders often display delayed dispersal and reproductive suppression of subordinates

[43], so that the age at first reproduction is also delayed and the number of breeding attempts is thus

decreased, which can lead to increased longevity [16]. Moreover, evidence suggests that a slower

pace of life is evolutionary linked to colonial breeding in birds [37] and to larger brain size in

mammals [63], and species displaying a high degree of sociality also display slower development,

delayed age at primiparity, better survival prospects and longer lifespan (reviewed in [7]). Whether

a large relative brain size is directly related to a slower pace of life (cognitive buffer hypothesis;

[64]) for a given degree of sociality or a large relative brain size is more likely to evolve in social

species (social brain hypothesis; [38]), which leads to slow down the pace of life is currently

unknown and requires further investigation.

It might be premature to conclude firmly that the degree of sociality has no direct effect on the

magnitude of reproductive senescence. Currently, we lack accurate metrics for measuring the

degree of sociality across a wide range of species and the metrics we used in this study have

limitations. For instance, cooperative breeding would require a more detailed typology based on

four classes (i.e. solitary, social, communal and cooperative; as per [48]) to describe accurately the

different levels of social complexity. Moreover, colonial breeding was confounded with habitat type

in our analyses of the bird data set, because all colonial species are seabirds and occupy thus marine

habitats. This redundancy cannot be separated completely by controlling for phylogenetic inertia,

and therefore future studies will be required to assess whether the association between coloniality

and reproductive senescence differs (with and without accounting for the pace of life) between

terrestrial and marine colonial species. Unfortunately, we failed to identify any data fulfilling our

selection criteria on reproductive senescence in terrestrial colonial birds. We also relied on the

social brain hypotheses, which proposes that relative brain size is larger in species with a higher

degree of social bonds [38] to justify our use of the relative brain size as a measure of the degree of

sociality. This hypothesis has received so far mixed support when assessing its plausibility in

animal taxa with a wide diversity of social systems [38,65]. However, the social brain hypothesis

holds for species displaying complex social interactions, such as cetaceans or primates [38,66]. Our

results based on relative brain size should also be treated with caution because brain size is only a

rough index of sociality and is related to other life-history traits that might influence senescence

(relationship with longevity [67,68]). Taken together, our conclusion that the degree of sociality has

no direct influence on reproductive senescence in birds and mammals will need to be investigated

more thoroughly when better measures of the degree of sociality will be available for a substantial

set of species. The recent development of social network analysis [69], which allows detailed

accounts of individual interactions within populations, should play a key role for doing that.

Interestingly, the only detectable direct effect of the degree of sociality on reproductive senescence

was opposite to our prediction. We found that cooperatively breeding mammals senesce faster, not

slower, than non-cooperative ones for a given size and pace of life. At first sight, this finding

contradicts within-population studies that showed almost consistently that helpers buffer the

demographic senescence of breeders [7,43]. However, cooperative breeding might have opposite

effect on reproductive senescence depending on the level of biological organization we consider.

For instance, getting the breeder tenure requires winning aggressive social interactions that increase

the level of physiological stress at the long term [70,71], which might exacerbate reproductive

senescence [8]. Additionally, the buffered effect of cooperative breeding on reproductive

senescence among individuals within a population of a given species can translate into an increased

reproductive senescence of cooperative breeding species compared to non-cooperative breeding

ones. Within a population of cooperative breeders, reproductively active individuals (i.e.

dominants) usually receive alloparental assistance from helpers (i.e. subordinates), which decreases

the cost of a given reproductive effort and leads thereby either to a postponed onset or to a

decelerated rate of reproductive or actuarial senescence (e.g. load-lightening hypothesis; [20]). For

instance, in Alpine marmots (*Marmota marmota*), individuals that have benefited from more

helping during their prime-age reproductive stage display a reduced actuarial senescence compared

to those that received less help [72], leading to increased individual heterogeneity in the amount of

senescence. At the population level, as considered in across-species analyses, the reproductive

suppression and associated physiological stress of individuals that help repeatedly before reaching a

dominant status and/or the paucity of substantial help when being breeders might lead to more

pronounced reproductive senescence. Overall, at the population level, increased costs of helping

when subordinate or lack of help when dominant for a large number of individuals might

counterbalance the benefits of having many helpers during breeding events that only a reduced

number of individuals enjoy. This strong individual heterogeneity in the strength of reproductive

senescence within populations of cooperative breeders might lead the average magnitude of

reproductive senescence to be higher in these species than in non-cooperatively breeding ones.

5. Conclusions

Our results indicate that degree of sociality is not directly associated with female reproductive

senescence. Instead, the positive covariation between the degree of sociality and a slower pace of

life has deeper evolutionary roots, which encompass both a later onset and a slower rate of

reproductive senescence.

Ethics. This study does not have ethical aspects.

Data accessibility. The data sets supporting this article have been uploaded as part of the Electronic Supplementary

Material (see ESM 'Data set').

Authors' contributions. C.I.V., J.-F.L. and J.-M.G. designed the study; C.I.V., J.-F.L., O.V., P.L.P., V.R. and J.-M.G.

provided data; O.V. and V.R. carried out the statistical analyses; C.I.V., O.V., J.-F.L. and J.-M.G. drafted the

manuscript with considerable help from P.L.P. and V.R.; all authors approved the final version of the manuscript.

Competing interests. We declare we have no competing interests.

Funding. C.I.V. was supported by the Hungarian National Research, Development and Innovation Office (#PD

121166), P.L.P. by a grant from the Romanian Ministry of Research and Innovation (#PN-III-P4-ID-PCE-2016-0404).

Acknowledgements. We are grateful to Hanna Kokko and three anonymous reviewers for insightful comments that

greatly improved our manuscript.

524 References

1. Lemaître J-F, Gaillard J-M. 2017 Reproductive senescence: new perspectives in the wild. *Biol. Rev.* **92**, 2182–

2199. (doi:10.1111/brv.12328)

2. Nussey DH, Froy H, Lemaître J, Gaillard J-M, Austad SN. 2013 Senescence in natural populations of animals:

widespread evidence and its implications for bio-gerontology. *Ageing Res. Rev.* **12**, 214–225.

(doi:10.1016/j.arr.2012.07.004)

3. Vaupel JW, Baudisch A, Dölling M, Roach DA, Gampe J. 2004 The case for negative senescence. *Theor.*

Popul. Biol. **65**, 339–351. (doi:10.1016/j.tpb.2003.12.003)

4.   Jones OR *et al.* 2014 Diversity of ageing across the tree of life. *Nature* **505**, 169–173.

(doi:10.1038/nature12789)

5.   Jones OR *et al.* 2008 Senescence rates are determined by ranking on the fast–slow life-history continuum. *Ecol.*

Lett. **11**, 664–673. (doi:10.1111/j.1461-0248.2008.01187.x)

6.   Gaillard J-M, Lemaître J-F. 2017 The Williams' legacy: a critical reappraisal of his nine predictions about the

evolution of senescence. *Evolution* **71**, 2768–2785. (doi:10.1111/evo.13379)

7.   Lucas ER, Keller L. 2020 The co-evolution of longevity and social life. *Funct. Ecol.* **34**, 76–87.

(doi:10.1111/1365-2435.13445)

8.   Bourke AFG. 2007 Kin selection and the evolutionary theory of aging. *Annu. Rev. Ecol. Evol. Syst.* **38**, 103–

128. (doi:10.1146/annurev.ecolsys.38.091206.095528)

9.   Snyder-Mackler N *et al.* 2020 Social determinants of health and survival in humans and other animals. *Science*

368, eaax9553. (doi:10.1126/science.aax9553)

10.   Hammers M, Kingma SA, Spurgin LG, Bebbington K, Dugdale HL, Burke T, Komdeur J, Richardson DS. 2019

Breeders that receive help age more slowly in a cooperatively breeding bird. *Nat. Commun.* **10**, 1301.

(doi:10.1038/s41467-019-09229-3)

11.   Cody ML. 1966 A general theory of clutch size. *Evolution* **20**, 174–184. (doi:10.2307/2406571)

12.   Stearns SC. 1992 *The Evolution of Life Histories*. London, UK: Oxford University Press.

13.   Cole LC. 1954 The population consequences of life history phenomena. *Q. Rev. Biol.* **29**, 103–137.

(doi:10.1086/400074)

14.   Williams GC. 1957 Pleiotropy, natural selection, and the evolution of senescence. *Evolution* **11**, 398–411.

(doi:10.2307/2406060)

15.   Kirkwood TBL, Rose MR. 1991 Evolution of senescence: late survival sacrificed for reproduction. *Philos.*

Trans. R. Soc. London B **332**, 15–24. (doi:10.1098/rstb.1991.0028)

16.   Lemaître J-F, Berger V, Bonenfant C, Douhard M, Gamelon M, Plard F, Gaillard J-M. 2015 Early–late life

trade-offs and the evolution of ageing in the wild. *Proc. R. Soc. London B* **282**, 20150209.

(doi:10.1098/rspb.2015.0209)

17.   Baudisch A, Vaupel JW. 2012 Getting to the root of aging. *Science* **338**, 618–619.

(doi:10.1126/science.1226467)

18.   Lemaître J-F, Gaillard J-M, Pemberton JM, Clutton-Brock TH, Nussey DH. 2014 Early life expenditure in

sexual competition is associated with increased reproductive senescence in male red deer. *Proc. R. Soc. London*

B **281**, 20140792. (doi:10.1098/rspb.2014.0792)

19.   Preston BT, Jalme M Saint, Hingrat Y, Lacroix F, Sorci G. 2011 Sexually extravagant males age more rapidly.

Ecol. Lett. **14**, 1017–1024. (doi:10.1111/j.1461-0248.2011.01668.x)

20.   Crick HQP. 1992 Load-lightening in cooperatively breeding birds and the cost of reproduction. *Ibis* **134**, 56–61.

(doi:10.1111/j.1474-919X.1992.tb07230.x)

21.   van Noordwijk AJ, de Jong G. 1986 Acquisition and allocation of resources: their influence on variation in life

history tactics. *Am. Nat.* **128**, 137–142. (doi:10.1086/284547)

22.   Nussey DH, Kruuk LEB, Donald A, Fowlie M, Clutton-Brock TH. 2006 The rate of senescence in maternal

performance increases with early-life fecundity in red deer. *Ecol. Lett.* **9**, 1342–1350. (doi:10.1111/j.1461-

0248.2006.00989.x)

23.   Speakman JR. 2008 The physiological costs of reproduction in small mammals. *Philos. Trans. R. Soc. London*

B **363**, 375–398. (doi:10.1098/rstb.2007.2145)

24.   Hamel S, Gaillard J-M, Yoccoz NG, Loison A, Bonenfant C, Descamps S. 2010 Fitness costs of reproduction

depend on life speed: empirical evidence from mammalian populations. *Ecol. Lett.* **13**, 915–935.

(doi:10.1111/j.1461-0248.2010.01478.x)

25.   Williams P, Day T, Fletcher Q, Rowe L. 2006 The shaping of senescence in the wild. *Trends Ecol. Evol.* **21**,

458–463. (doi:10.1016/j.tree.2006.05.008)

26.   Williams SA, Shattuck MR. 2015 Ecology, longevity and naked mole-rats: confounding effects of sociality?

Proc. R. Soc. London B **282**, 20141664. (doi:10.1098/rspb.2014.1664)

27.   Birkhead TR, Møller AP. 1995 Extra-pair copulation and extra-pair paternity in birds. *Anim. Behav.* **49**, 843–

848. (doi:10.1016/0003-3472(95)80217-7)

28.   Lemaître J, Ronget V, Gaillard J-M. 2020 Female reproductive senescence across mammals: a high diversity of

patterns modulated by life history and mating traits. *Mech. Ageing Dev.* **192**, 111377.

(doi:10.1016/j.mad.2020.111377)

29.   Thorley J, Duncan C, Sharp SP, Gaynor D, Manser MB, Clutton□Brock T. 2020 Sex□independent senescence

in a cooperatively breeding mammal. *J. Anim. Ecol.* **89**, 1080–1093. (doi:10.1111/1365-2656.13173)

30.   Muggeo VMR. 2017 Interval estimation for the breakpoint in segmented regression: a smoothed score-based

approach. *Aust. N. Z. J. Stat.* **59**, 311–322. (doi:10.1111/anzs.12200)

31.   Lindstedt SL, Calder WA. 1981 Body size, physiological time, and longevity of homeothermic animals. *Q. Rev.*

Biol. **56**, 1–16. (doi:10.1086/412080)

32.   Stearns SC. 1983 The influence of size and phylogeny on patterns of covariation among life-history traits in the

mammals. *Oikos* **41**, 173–187. (doi:10.2307/3544261)

33.   Gaillard J-M, Yoccoz NG, Lebreton J-D, Bonenfant C, Devillard S, Loison A, Pontier D, Allaine D. 2005

Generation time: a reliable metric to measure life-history variation among mammalian populations. *Am. Nat.*

166, 119–123. (doi:10.1086/430330)

34.   Staerk J, Conde DA, Ronget V, Lemaître J, Gaillard J, Colchero F. 2019 Performance of generation time

approximations for extinction risk assessments. *J. Appl. Ecol.* **56**, 1436–1446. (doi:10.1111/1365-2664.13368)

35.   Dunning Jr JB. 2008 *Body masses of birds of the world. CRC handbook of avian body masses.* Taylor and

Francis Group, Boca Raton, FL.

36.   Székely T, Moore AJ, Komdeur J, editors. 2010 *Social Behaviour: Genes, Ecology and Evolution.* Cambridge,

UK: Cambridge University Press. (doi:10.1017/CBO9780511781360)

37.   Møller AP. 2006 Sociality, age at first reproduction and senescence: comparative analyses of birds. *J. Evol.*

Biol. **19**, 682–689. (doi:10.1111/j.1420-9101.2005.01065.x)

38.   Dunbar RIM, Shultz S. 2007 Evolution in the social brain. *Science* **317**, 1344–1347.

(doi:10.1126/science.1145463)

39.   Evans JC, Votier SC, Dall SRX. 2016 Information use in colonial living. *Biol. Rev.* **91**, 658–672.

608   (doi:10.1111/brv.12188)

609   40.   Hoogland JL. 1995 *The Black-Tailed Prairie Dog: Social Life of a Burrowing Mammal*. Chicago, IL:

610   University of Chicago Press.

611   41.   Liker A, Székely T. 2005 Mortality costs of sexual selection and parental care in natural populations of birds.

612   *Evolution* **59**, 890–897. (doi:10.1111/j.0014-3820.2005.tb01762.x)

613   42.   Lack D. 1968 *Ecological Adaptations for Breeding in Birds*. London, UK: Methuen and Co.

614   43.   Clutton-Brock T. 2009 Structure and function in mammalian societies. *Philos. Trans. R. Soc. London B* **364**,

615   3229–3242. (doi:10.1098/rstb.2009.0120)

616   44.   Shultz S, Dunbar RI. 2007 The evolution of the social brain: anthropoid primates contrast with other

617   vertebrates. *Proc. R. Soc. London B* **274**, 2429–2436. (doi:10.1098/rspb.2007.0693)

618   45.   Sayol F, Lapiedra O, Ducatez S, Sol D. 2019 Larger brains spur species diversification in birds. *Evolution* **73**,

619   2085–2093. (doi:10.1111/evo.13811)

620   46.   del Hoyo J, Elliott A, Sargatal J. 2016 HBW Alive: Handbook of the Birds of the World. See

621   https://www.hbw.com/.

622   47.   Burger JR, George MA, Leadbetter C, Shaikh F. 2019 The allometry of brain size in mammals. *J. Mammal.*

623   **100**, 276–283. (doi:10.1093/jmammal/gyz043)

624   48.   Lukas D, Clutton-Brock T. 2012 Cooperative breeding and monogamy in mammalian societies. *Proc. R. Soc.*

625   *London B* **279**, 2151–2156. (doi:10.1098/rspb.2011.2468)

626   49.   R Core Team. 2020 *R: a language and environment for statistical computing*. Vienna, Austria. http://www.R-

627   project.org/: R Foundation for Statistical Computing.

628   50.   Sukumaran J, Holder MT. 2010 DendroPy: a Python library for phylogenetic computing. *Bioinformatics* **26**,

629   1569–1571. (doi:10.1093/bioinformatics/btq228)

630   51.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. 2012 The global diversity of birds in space and time.

631   *Nature* **491**, 444–448. (doi:10.1038/nature11631)

632   52.   Hackett SJ *et al.* 2008 A phylogenomic study of birds reveals their evolutionary history. *Science* **320**, 1763–

633   1768. (doi:10.1126/science.1157704)

634   53.   Bininda-Emonds ORP *et al.* 2007 The delayed rise of present-day mammals. *Nature* **446**, 507–512.

635   (doi:10.1038/nature05634)

636   54.   Revell LJ. 2012 phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol.*

637   *Evol.* **3**, 217–223.

638   55.   Tung Ho LS, Ané C. 2014 A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Syst.*

639   *Biol.* **63**, 397–408. (doi:10.1093/sysbio/syu005)

640   56.   Ivimey-Cook E, Moorad J. 2020 The diversity of maternal-age effects upon pre-adult survival across animal

641   species. *Proc. R. Soc. London B* **287**, 20200972. (doi:10.1098/rspb.2020.0972)

642   57.   Holmes D, Martin K. 2009 A bird's-eye view of aging: what's in it for ornithologists? *Auk* **126**, 1–23.

643   (doi:10.1525/auk.2009.1109)

644   58.   Healy K, Guillerme T. 2014 Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc.*

645   *R. Soc. London B* **281**, 20140298.

646   59.   Gaillard J-M, Lemaître J-F, Berger V, Bonenfant C, Devillard S, Douhard M, Gamelon M, Plard F, Lebreton J.

647   2016 Axes of variation in life histories. In *Encyclopedia of Evolutionary Biology* (ed R Kliman), pp. 312–323.

648   New York, NY: Elsevier. (doi:10.1016/B978-0-12-800049-6.00085-8)

649   60.   Reiss JO. 1989 The meaning of developmental time: a metric for comparative embryology. *Am. Nat.* **134**, 170–

650   189. (doi:10.1086/284974)

651   61.   Wootton JT. 1987 The effects of body mass, phylogeny, habitat, and trophic level on mammalian age at first

652   reproduction. *Evolution* **41**, 732–749. (doi:10.2307/2408884)

653   62.   Sacher G. 1959 Relationship of lifespan to brain weight and body weight in mammals. In *The Lifespan of*

654   *Animals* (eds G Wolstenholme, M O'Conner), pp. 115–141. Churchill, London: Ciba Foundation.

655   (doi:10.1002/9780470715253.ch9)

656   63.   Barton RA, Capellini I. 2011 Maternal investment, life histories, and the costs of brain growth in mammals.

657   *Proc. Natl. Acad. Sci. USA* **108**, 6169–6174. (doi:10.1073/pnas.1019140108)

658   64.   Sol D. 2009 The cognitive-buffer hypothesis for the evolution of large brains. In *Cognitive Ecology II* (eds R

659   Dukas, JM Ratcliffe), pp. 111–134. Chicago, IL: University of Chicago Press.

660   65.   Isler K, van Schaik CP. 2012 Allomaternal care, life history and brain size evolution in mammals. *J. Hum. Evol.*

661   **63**, 52–63. (doi:10.1016/j.jhevol.2012.03.009)

662   66.   Fox KCR, Muthukrishna M, Shultz S. 2017 The social and cultural roots of whale and dolphin brains. *Nat.*

663   *Ecol. Evol.* **1**, 1699–1705. (doi:10.1038/s41559-017-0336-y)

664   67.   Jiménez☐Ortega D, Kolm N, Immler S, Maklakov AA, Gonzalez☐Voyer A. 2020 Long life evolves in

665   large☐brained bird lineages. *Evolution*, in press. (doi:10.1111/evo.14087)

666   68.   González-Lagos C, Sol D, Reader SM. 2010 Large-brained mammals live longer. *J. Evol. Biol.* **23**, 1064–1074.

667   (doi:10.1111/j.1420-9101.2010.01976.x)

668   69.   Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting animal social network analysis. *J.*

669   *Anim. Ecol.* **84**, 1144–1163. (doi:10.1111/1365-2656.12418)

670   70.   Cohas A, Rey B, Federico V, Regis C, Lardy S, Bichet C. 2018 Stress levels of dominants reflect underlying

671   conflicts with subordinates in a cooperatively breeding species. *Behav. Ecol. Sociobiol.* **72**, 72.

672   (doi:10.1007/s00265-018-2484-8)

673   71.   Goymann W, Wingfield JC. 2004 Allostatic load, social status and stress hormones: the costs of social status

674   matter. *Anim. Behav.* **67**, 591–602. (doi:10.1016/j.anbehav.2003.08.007)

675   72.   Berger V, Lemaître J-F, Allainé D, Gaillard J-M, Cohas A. 2018 Early and adult social environments shape sex-

676   specific actuarial senescence patterns in a cooperative breeder. *Am. Nat.* **192**, 525–536. (doi:10.1086/699513)