Detecting significant components of microbiomes by random forest with forward variable selection and phylogenetics

Tung Dang[1] Hirohisa Kishino[1]

Abstract

A central focus of microbiome studies is the characterization of differences in the microbiome composition across groups of samples. A major challenge is the high dimensionality of microbiome datasets, which significantly reduces the power of current approaches for identifying true differences and increases the chance of false discoveries. We have developed a new framework to address these issues by combining (i) identifying a few significant features by a massively parallel forward variable selection procedure, (ii) mapping the selected species on a phylogenetic tree, and (iii) predicting functional profiles by functional gene enrichment analysis from metagenomic 16S rRNA data. We demonstrated the performance of the proposed approach by analyzing two published datasets from large-scale case-control studies: (i) 16S rRNA gene amplicon data for *Clostridioides difficile* infection (CDI) and (ii) shotgun metagenomics data for human colorectal cancer (CRC). The proposed approach improved the accuracy from 81% to 99.01% for CDI and from 75.14% to 90.17% for CRC. We identified a core set of 96 species that were significantly enriched in CDI and a core set of 75 species that were enriched in CRC. Moreover, although the quality of the data differed for the functional profiles predicted from the 16S rRNA dataset and functional metagenome profiling, our approach performed well for both databases and detected main functions that can be used to diagnose and study further the growth stage of diseases.

Introduction

A microbiome is the full collection of genes of all microbes in a community; for example, all bacteria in a sample from the gut of a healthy individual or from an individual with a disease. Identifying different microbiome compositions between two or more groups is one of the most important purposes of microbiome studies [1] [2]. High-throughput sequencing technologies have allowed the microbiome composition and function in different environments to be quantified correctly [3] [4].

Machine learning technology has been applied in microbiome studies, especially for predicting specific diseases and supporting medical diagnosis [5] [6]. Random forest (RF) algorithms are considered to be a promising approach [7] [8].

[1]Department of Agricultural and Environmental Biology, The University of Tokyo, Tokyo, Japan

The RF algorithm is a modification of bagging that aggregates a large collection of decision trees [9]. A main step in building an ensemble of decision trees is to perform random sampling of the available features to generate different subspaces of features at each node of each unpruned decision tree. This strategy can produce better estimation performances than a single decision tree because each tree estimator has low bias but high variance, whereas a bias–variance trade-off is achieved by the bagging process of RFs. RF methods have been applied successfully to genetic and microbiome data [10] [11] [12]. It is anticipated that RF methods and implemented importance measures will help in the identification of microbiome species that can be used to distinguish diseased and non-diseased samples. Identifying a core set of the most significant microbial species is of high interest, not only for diagnosis of certain diseases but also to gain valuable insights into the biological functionality and mechanisms of these species.

However, the performance of a decision tree and the diversity of decision trees in the ensemble significantly influence the performance of RF algorithms. The generalization error for RFs involves measures of how accurate the individual classifiers are and their interdependence. Therefore, the high dimensionality problems of microbiome datasets pose a number of

challenges. For example, microbiome datasets tend to contain a large number of microbiome species whose functions may not be related to the disease of interest. Common random sampling methods may select a sizeable number of subspaces that do not include the informative microbiome species and functions. As a consequence, the decision trees generated from these subspaces will have reduced average strength, thereby increasing the error bounds for the RF algorithm.

Although a number of different algorithms and tools have been developed for microbiome analysis [5], effective approaches require a lot of possible combinations of variables, which exponentially increases the computational burden as the number of involved features increases. Even though a small number of machine learning methods, including the RF algorithm, can be easily parallelized, building prediction models for thousands of microbiome species and functions can be very time consuming.

In this study, we developed a novel analysis pipeline that tackles the challenges described above. The core of our pipeline is an RF classifier coupled with forward variable selection (RF-FVS), which selects a minimal-size core set of microbial species or functional signatures that maximize the predictive performance of the RF classifier. To reduce the computational cost for the high-dimensional data, we designed a parallelized algorithm and integrated a prescreen algorithm. The RF-FVS pipeline is the basis for the integrated analysis of microbiome data, detecting significant species in the phylogenetic tree [13] and predicting functional capabilities of microbial communities based on 16S rRNA datasets [14]. To estimate the dependence of the pipeline on the quality of the microbiome functional profile data, we analyzed the predicted functional profiles obtained from a 16S rRNA dataset that were approximately 85% accurate and the functional profiles obtained from a shotgun metagenomic dataset. We found that our RF-FVS approach performed well, even for the predicted functional profile data.

Materials and methods

Forward variable selection for random forests

The RF approach is an ensemble method that combines a large number of individual binary decision trees. Two main randomization procedures have been implemented to reduce variance of individual decision trees, deliver diversity amongst decision trees, and thus improve prediction accuracy. First, randomly selected training samples for each of the individual trees are applied to construct sufficiently diverse trees. Second, at each node within a tree, a set of randomly chosen candidate predictor variables is identified for the split.

However, random feature subspace sampling may not be a good strategy to deal with high dimensional data because a large proportion of the features may not be informative of the class of an object in the high dimensional data. If a random sampling strategy is implemented to select the subset of eligible features at each node, almost all the subsets are likely to contain a large number of non-informative features. For example, the 16S rRNA gene amplicon data for *Clostridioides difficile* infection (CDI) that we used to evaluate the performance of our proposed approach contained a total of 3347 microbiome species, but only 96 of the species were informative. Therefore, if a subset of species, which is usually the square root of the total number of species, is selected by resampling randomly at any node within the decision tree, the mean number of informative species selected at each node will be two. Therefore, individual decision trees built using such nodes will have low accuracy and the performance of the RF algorithm will suffer. In our approach, we used forward variable selection to identify a small number of informative variables to improve the performance of individual decision trees in an ensemble.

A key idea behind our algorithm was to divide the total number of variables into two groups, a remaining variables group and a selected variables group. We started with an empty group for the selected variables. At each step, a variable from the remaining variables group was added to the selected variables group such that the specified criterion was improved (i.e., area under the receiver operating characteristic [ROC] curve [AUC], a weighted average of the precision and recall [F1 score] or predictive accuracy). Model selection for microbial signature identification also can be performed using our RF-FVS algorithm. At each forward iteration, given the selected variables, the randomized parameter optimization algorithm for RFs implements a randomized search over parameters, where each setting is sampled from a distribution over possible parameter values. Thus, the best RF model is specified by these selected variables.

Moreover, a high-speed computational strategy based on multi-processing architecture was developed to parallelize the forward variable selection algorithm at the single machine level and thus significantly reduce runtimes. Another key idea behind our algorithm was to create many subsets of variables, so that each subset had one of the variables from the remaining variables group added to the selected variables group. Because of the high dimensionality problems of microbiome data, the number of these subsets is usually significantly larger than the number of processors in a single computer system. Our solution was to create queues, so that subsets are assigned randomly, and each processor runs the computational processes from its own privately prepared queue. Figure 1 shows how all computational burdens for searching important feature relevance are appropriately decomposed, so that they can be computed in a parallel environment. The

Fig. 1 Massively parallel forward variable selection algorithm for the random forest (RF) classifier. The number of queues depends on the number of CPU cores available in the computer system.

RF model with the highest accuracy value for each subset of features is selected by a specific processor. Processors in symmetric multiprocessing communicate with each other through shared memory architecture to decide only the best feature among multiple candidates that should be added to the selected features. The algorithm stops when there are no additional variables that improve the current optimization parameters or when the maximum number of components to be included in the group of selected variables is achieved. The main parameters of the RF classifier that were optimized in our algorithm, were number of trees in the forest, maximum depth of the tree, minimum number of samples required to split an internal node, minimum number of samples required to be at a leaf node, and number of features to consider when looking for the best split.

Functional gene enrichment analysis

Functional profiles were predicted from the 16S rRNA gene data for CDI using the Tax4Fun prediction tool that was developed to analyze the enrichment of functional genes of microbiomes [14] [15]. The output from QIIME with a SILVA database extension (SILVA 119) was used to pre-process raw data for Tax4Fun. Tax4Fun transforms the SILVA-based operational taxonomic units (OTUs) into a taxonomic profile of KEGG organisms that is normalized by the 16S rRNA copy number (obtained from the NCBI genome annotations). The result is a table containing relative KEGG ortholog (KO) abundance levels. The KO profiles obtained from predictions or actually measured from shotgun metagenomic or metatranscriptomic data can be used for functional diversity profiling based on the KEGG (pathways, modules, or EC categories) annotation systems. Because one KO can be assigned to multiple groups, our approach detects the most important functional profiles associated with disease or healthy samples.

Phylogenetic transformation of microbiota data for random forest

To avoid spurious statistical analyses because of the relative nature of microbial abundance data in microbiota studies

Fig. 2 Workflow for the new approach developed and implemented in this study.

[16], we used the Phylogenetic Isometric Log-Ratio (PhILR) transformation [13]. The main idea behind the PhILR transformation is to consider the bacterial phylogenetic tree as a natural and informative sequential binary partition to construct an isometric log-ratio that converts compositional data into a real Euclidean space. This phylogenetically driven isometric log-ratio transformation can help to capture the hierarchical pattern of microbial community structure. Therefore, the RF-FVS algorithm can identify the main internal nodes of a phylogenetic tree that represent phylogenetically related bacterial groups (clades), thereby offering the opportunities for biological insights.

Pre-screening algorithm for random forest coupled with forward variable selection

We used the Boruta algorithm [17] as a relevant embedded feature selection algorithm that uses the RF classifier to detect all strongly and weakly relevant OTUs (or phylogenetic internal nodes, or functional profiles) to reduce the considerable data dimensionality. This improved the classification accuracies and significantly decreased the computation time. The main idea of this algorithm was to duplicate each OTU, thus creating "shadow" OTUs by randomly permuting the observations of duplicated OTUs at the first step. Then, the importance of all the OTUs are computed (calculated as Z-scores) and the maximum Z-score among the shadow bands are identified when the RF classifier is run. The number of times the importance of an OTU is higher than the maximum Z-score among the shadow OTUs is counted. An OTU is deemed "important" when the frequency is significantly higher than the expected value, otherwise the

OTU is deemed "unimportant" and removed. Basically, the Boruta algorithm works with Scikit-learn dependency and acts as an interface medium on the RF classifier, which uses the importance measure generated by the original algorithm. The Boruta algorithm depends on the implementation of the Scikit-learn RF classifier, which runs on a single code to identify all the important features in a dataset with respect to an outcome variable. Figure 2 shows the global framework of our new approach.

Two empirical datasets

To examine the performance of our RF-FVS approach, we analyzed two empirical datasets from large-scale case control studies as follows.

Clostridioides difficile infection

We collected a published case-control 16S rRNA gene amplicon sequencing gut microbiome dataset with 3347 OTUs for *C. difficile* infection (CDI) [18]) from 338 individuals that included disease metadata and sequencing data. To understand how clinical- and microbiome-based factors were associated with CDI, the gut microbiomes of individuals with CDI, individuals with diarrhea who tested negative for *C. difficile* infection (diarrheal control), and healthy individuals (non-diarrheal control) were characterized. The 183 diarrheal stool samples that were analyzed were from individuals with CDI (94 samples) and diarrheal controls (89 samples), and 155 non-diarrheal control stool samples were from the non-diarrheal controls. Taxonomic assignments were generated using a naive

Fig. 3 Confusion matrices comparing the performance of the random forest classifier (with and without forward variable selection) for clinical data and 16S rRNA gene amplicon data of *Clostridioides difficile* infection (CDI) dataset. **a** Random forest classifier for the clinical data; **b** Random forest classifier for the 16S rRNA gene amplicon dataset; **c** Random forest classifier coupled with forward variable selection (RF-FVS) approach for the 16S rRNA gene amplicon dataset.

Bayesian classifier trained against a 16S rRNA gene training set provided by the Ribosomal Database Project (RDP) [19]. We also collected a set of published 16S rRNA gene sequences [20], among which, 120 samples were from individuals with human colorectal cancer (CRC) and 172 samples were from tumor-free controls. Taxonomic assignments were generated using the Bayesian classifier as described above.

Human colorectal cancer

We also collected a published fecal shotgun metagenomic dataset [21], which included 290 samples from tumor-free controls and 285 samples from individuals with CRC. Taxonomic profiles were generated using the mOTU profiler tool [22] to guarantee consistency of the biological information. Unlike the 16S rRNA gene amplicon data, one of the most important advantages of fecal shotgun metagenomic data is that it provides functional abundance profiles, such as those for eggNOG gene family or KEGG module abundance profiles, that allowed a direct analysis of the functional potential of the gut microbiome to be conducted.

Results

Forward variable selection increased accuracy for the 16S rRNA gene amplicon data

We compared the predictive power of our approach to classify three groups (CDI case, diarrheal control, non-diarrheal control) between the microbiome data and the clinical data. The clinical data included age, sex, race, antibiotic use, antacid use, a vegetarian diet, surgery within the past 6 months, a history of CDI, residence with another person who had CDI, and residence with another person who works in health care. The clinical data were 51% accurate (AUC = 0.7). In contrast, the microbiome data had significantly higher accuracy (81%) and AUC (0.95). In particular, the accuracy was high for the microbiome data of the non-diarrheal control. Moreover, when treating the composition of the microbiota data by PhILR transformation, we found that the RF classifier achieved

high accuracy (85%) and AUC (0.95). However, it was still difficult to distinguish CDI cases and diarrheal controls in both the microbiome and phylogenetic transformation data. Overall, the accuracy of the diagnosis was significantly improved for the CDI and healthy groups for both the microbiome and clinical data. By focusing on 119 species, the accuracy of the RF classifier increased to 96% (AUC = 0.97) (Figure 3), and CDI cases were identified with an accuracy of 94% (AUC = 0.94) and diarrheal controls were identified with an accuracy of 93% (AUC = 0.97). For the phylogenetic transformation data, the accuracy was 95% (AUC = 0.97) when the forward variable selection approach detected the 36 phylogenetic internal nodes.

Significant species on the 16S rRNA tree
phylogenetic tree

Group C

Group A

Group B

Fig. 4 Phylogenetic tree of the 16S RNA microbiota (OTU) data from the CDI dataset. The red positives indicate the 119 important microbial species that remained after forward variable selection. Red, OTUs associated with CDI cases; blue, OTUs associated with the non-diarrheal controls.

The selected OTUs from nine families were clustered in the 16S RNA tree: *Peptostreptococcaceae, Verrucomicrobiaceae*, *Veillonellaceae, Ruminococcaceae*, *Lachnospiraceae*, *Bacteroides*, *Porphyromonadaceae, Lactobacillaceae* and *Enterobacteriaceae* (figure 4). They formed three main groups of species that were associated with the differentiation of CDI cases from the non-diarrheal controls. The species in Group B (such as denovo 54, 1326, 601, 2346, and 3486) that showed strong positive correlations with CDI cases belonged to a diverse number of families (Supplementary Fig. 1 and Table 1), including *Peptostreptococcaceae* [23], *Lactobacillaceae* (including

Fig. 5 Confusion matrices that compare the performance of random forest (with and without forward variable selection) for the 16S rRNA gene amplicon data and shotgun metagenomics data for the human colorectal cancer (CRC) datasets. **a** Random forest classifier for the 16S rRNA gene amplicon data; **b** Random forest classifier for the shotgun metagenomics data; **c** Random forest classifier coupled with forward variable selection (RF-FVS) for the shotgun metagenomics data.

the *Lactobacillus* genus), *Enterococcaceae* (including the *Enterococcus* genus) [24] [25], *Verrucomicrobiaceae*, and *Veillonellaceae* [26]. For example, species (such as denovo 54) in the *Clostridium* cluster XI genus in family *Peptostreptococcaceae* that showed high positive correlations with CDI cases (Supplementary Fig. 1) are main candidates for treatment development [27] and the study of new challenges that have arisen because of the discontinuation of therapy for CDI disease [28]. Pérez-Cobas *et al.* [29] reported that the most striking changes in the microbiome of CDI cases occurred in the *Lactobacillaceae* family, whose frequency increased from <1% at the beginning of antibiotic treatment to 83.3% and 70% on days 35 and 38 of an antibiotic course, then reduced to 15.5% after the antibiotic therapy. Most of the species in group C (such as denovo 127, 1399, and 788) that showed positive correlations with CDI cases belonged to the *Enterobacteriaceae* family (Supplementary Table 1 and Table 1.1). Studies [30] [31] [32] have shown that relative overgrowth of members of *Enterobacteriaceae* family was one of the main causes of significantly disturbed microbiota in CDI. Thus, *C. difficile* colonization may be facilitated by increased endotoxin production with increased intestinal permeability.

Most of species in group A (such as denovo 557, 302, 1888, 1987, 1983 and 610), some species in group C (such as denovo 26, 9, 1295, 447 and 347), and one species in group B (denovo 156) that were enriched in the non-diarrheal controls belonged to the *Ruminococcaceae*, *Lachnospiraceae*, *Bacteroides*, and *Porphyromonadaceae* families (Supplementary Fig. 1 and Table 1). Short-chain fatty acid (SCFA) production is known to play a principal role in the regulation of intestinal inflammatory processes [33] and intestinal barrier maintenance [34], and *C. difficile* infection was found to cause significant reductions in the *Ruminococcaceae* and/or *Lachnospiraceae* families that produce butyrate and SCFA [35] [36]. *Lachnospiraceae* and *Ruminococcaceae* sequences were dominant at 45.8% and 17.4% respectively in healthy fecal microbiota, whereas sequences from other families constituted <3% of all the pyrosequencing reads. However, the proportions of these species decreased significantly in CDI cases (*Lachnospiraceae* (11.2%), *Ruminococcaceae* (3.0%)) [36]. Moreover, the four species in group A (denovo 2019,

1773, 3205 and 2528) (Supplementary Fig. 1.1) showed weak associations with the CDI case.

When we applied our RF classifier to the clinical data, we found that antibiotic treatment contributed significantly to an increase in the accuracy of the prediction models [29] [37]

Forward variable selection for shotgun data

Our RF classifier successfully distinguished the CRC cases and tumor-free controls in both the 16S rRNA gene amplicon data and the shotgun metagenomics data (Figure 5). The average accuracy for the 16S rRNA gene amplicon data, which included 18 448 OTUs, was 68.18% (AUC = 0.73). For the shotgun metagenomics data, the accuracies of the RF classifier for CRC cases and tumor-free controls were 70% (AUC = 0.86) and 80% (AUC = 0.86), respectively. The forward variable selection detected 75 microbial species (out of 849 species) that were differentially abundant in the CRC microbiome, which increased the accuracy of the RF classifier to 88% (AUC = 0.89) for the CRC cases and to 92% (AUC = 0.89) for the tumor-free controls. These findings are consistent with previous reports of significant enrichment of novel species in the fecal microbiomes of patients with CRC. For example, we detected the three most important species, *Parvimonas micra*, *Flavonifractor plautii*, and *Gemella morbillorum*, that helped to improve the accuracy of our RF classifier (Supplementary Fig. 2). Yu *et al.* [38] and Feng *et al.* [39] identified *Parvimonas micra* as a key species that was consistently enriched in the microbiomes of patients with CRC, and we found this species had the most importance among our selected species. Gupta *et al.* [40] found that *Flavonifractor plautii* was associated significantly and enriched in CRC samples of Indian patients. *Flavonifractor plautii* was linked with the degradation of beneficial anticarcinogenic flavonoids, and this role was strongly correlated with enzymes and modules involved in flavonoid degradation in CRC samples of the Indian patents. In our study, *Flavonifractor plautii* was significantly associated with CRC samples of cohorts from France, Germany, China, United States, and Austria (Supplementary Fig. 3). Our analysis showed that the dysbiosis of fecal microbiota was characterized by

Fig.6 Correlation of the 23 main functional profiles with CDI cases, antibiotics, proton pump inhibitors, and non-diarrheal controls.

enrichment of potential pathogens and the reduction in butyrate-producing members such as *Fusobacterium*, in CRC samples [41] [42]. We also identified other main species (Supplementary Fig. 2), *Gemella morbillorum* [43], *Prevotella intermedia* [44], *Prevotella nigrescens* [45], which have been shown previously to be strongly associated with an increased risk of CRC. In the gut microbiome of the tumor-free controls, we detected *Bifidobacterium adolescentis* and *Lactobacillus ruminis*, which previously have been associated with healthy samples and negatively correlated with CRC cases (Supplementary Fig. 2). These species may have a protective role against the development of non-alcoholic fatty liver disease and obesity [46].

Influential bacterial functions predicted by 16S rRNA data.

The forward variable selection algorithm detected 119 OTUs out of 3347 OTUs. To predict functional profiles from these OTU candidates, we examined 5818 of the 21 620 functional profiles of the KEGG organisms in the Tax4Fun framework. The RF method gave an average accuracy of 81% (AUC = 0.93) for the 5818 predicted functional profiles. Specifically, the accuracies for the CDI cases, diarrheal controls, and non-diarrheal controls were 68%, 79%, and 93% (AUCs = 0.88, 0.92, and 0.97) respectively. To reduce the computational burden, we used the prescreen algorithm at the first step, which reduced the predicted functional profiles from 5818 to 2534. Then, we applied the RF-RVS algorithm, which identified 23 functional profiles (out of 2534) that were different for the CDI cases compared with the controls, which significantly increased the average accuracy to 90% (AUC = 0.95). Specifically, the accuracies for CDI cases, diarrheal controls, and non-diarrheal controls were 77%, 93%, and 98% (AUCs = 0.90, 0.95, and 0.97) respectively. Some of the 23 functional profiles were strongly associated with CDI cases, but a larger number of them were associated with healthy gut (non-diarrheal controls) (Figure 6). Our results confirmed that bacteria support human health with functions such as histidinol dehydrogenase, gluconate 5-dehydrogenase, lactaldehyde reductase, and alpha-1,3-rhamnosyltransferase [47] [48] [49]. However, these functions were absent in the microbiomes of patients with CDI, mainly because they were treated with antibiotics and proton pump inhibitors, which killed these bacteria (Figure 6). Therefore, *C. difficile*, which is resistant to these treatments, became dominant and

Data type	Analysis framework			Accuracy	Time computation
	Boruta alogirithm	Random forest	Forward algorithm		
OTUs data	✗	✓	✓	96.04%	1 week
	✓	✓	✓	99.01%	1 day
PhILR transformation	✗	✓	✓	95.05%	26 hours
	✓	✓	✓	95.05%	8.75 hours
Functional profile data	✗	✓	✓	90.1%	4 days
	✓	✓	✓	90.1%	13 hours

Table 1 Random forest classifier with forward variable selection (RV-FVS) with and without the prescreen algorithm for the CDI dataset. ✔ indicates that the prescreen algorithm was used; ✖ indicates that the prescreen algorithm was not used. All algorithms were run in a parallel environment. The properties of the parallel version were evaluated on a high-performance computer (Intel® Xeon® Gold 6230 Processor 2.10 GHz × 2, 40 cores, 2 threads per core, 93.1-Gb RAM), under Ubuntu 20.04.1 LTS.

increased the risk of CDI. Conversely, some of the functional profiles were strongly enriched in the CDI cases. They included alcohol dehydrogenase, L-iditol 2-dehydrogenase, and glucose 6-dehydrogenase, which are related to the growth of *C. difficile* [50] [51], and xanthine dehydrogenase, which is related to the high-level resistance of *C. difficile* to antibiotics treatments [52].

Influential bacterial functions predicted using the shotgun data

We used evolutionary genealogy of genes: Non-supervised Orthologous Groups (eggNOG) orthologous gene family abundances and KEGG module abundance profiles to detect functional profiles in the shotgun metagenomics data for CRC. Because the number of functions in the KEGG and eggNOG databases are very large (7955 and 31185, respectively), we used the prescreen algorithm before applying forward variable selection.

Our RF-RVS algorithm identified 29 out of 7955 functions in the KEGG database that significantly improved the performance of the RF classifier. Specifically, the accuracy increased from 60% (AUC = 0.79) to 84% (AUC = 0.87) for CRC cases and from 80% (AUC = 0.79) to 97% (AUC = 0.87) for tumor-free controls (Supplementary Fig. 4). A number of functions, such as HOMODA hydrolase (K10623), carbamoyl-phosphate synthase 1 (K01948), heptose-I-phosphate ethanolaminephosphotransferase (K19353), tartronate-semialdehyde synthase (K01608), ABC transporters (K11707), and biofilm formation - *Pseudomonas aeruginosa*/bacterial secretion system (K11903) had strong positive correlations with CRC cases (Supplementary Fig. 5). The contributions of some of these functions to the stage progress of CRC have been reported.

For example, carbamoyl-phosphate synthase 1, a metabolic enzyme that utilizes ammonia to produce carbamoyl phosphate, is encoded by one of four novel driver genes that were identified as hubs for stage-III progression of colorectal cancer [53].

Our RF-RVS algorithm also identified 53 out of 31185 functions in the eggNOG database that significantly improved the performance of the RF classifier. Specifically, the accuracy increased from 62% (AUC = 0.78) to 86% (AUC = 0.90) for CRC cases and from 80% (AUC = 0.78) to 97% (AUC = 0.90) for tumor-free controls (Supplementary Fig. 6). Although a large number of functions were significantly positive correlated with CRC cases, such as ENOG410Y6BY, ENOG410XYS8, ENOG411EMB, and ENOG410ZGTS (Supplementary Fig. 7), experimental information about their functions is lacking. These genes are likely to be good candidates for further studies.

Forward variable selection and the prescreening algorithm reduced the CPU time

Although the RF classifier achieved high accuracy in analyzing the microbiome data, the high dimensionality of the data meant the computational burden was high. The RF classifier took about one week to identify 119 species out of 3347 species in the 16S rRNA gene amplicon data for CDI. By using forward variable selection to identify a small number of informative variables, the 3347 species were reduced to 1008 species and the RF-FVS algorithm detected 96 species with an increased accuracy of 99% (AUC = 0.99) (Table 1). The time computation also was reduced from one week to one day. For the functional profile predictions, the RF-FVS algorithm took about 4 days to identify 65 out of 5818 functional profiles. The prescreening algorithm reduced the total number of functional profiles from 5818 to 2534, and the RF-FVS then needed only 13 hours to detect

23 out of the 2534 functional profiles. The accuracy of the RF classifier also increased to 90.1%.

Discussion

A number of publicly available databases contain information about microbial species and their functions that are associated with disease or health. The large number of microbial species and functional profiles in these databases significantly negatively influence the power of machine learning classifiers, including RF algorithms. Further, the computational cost of running these algorithms to detect a few informative features in the high-dimensional space of microbiome data is still very high. In this study, we developed a novel procedure that significantly enhances the RF classifier and substantially improves its performance in terms of high-speed computation and high accuracy. We tested its performance using two microbiome datasets that contained a large number of species and functional signatures (>30000 variables) but a very small proportion of significant variables.

Our RF-FVS approach was useful in several respects. Firstly, the RF-FVS algorithm identified the core set of microbial species and their functional profiles, which considerably increased the predictive accuracy of the RF classifier. The highest increase in the predictive accuracy was to 99% for the CDI cases classification. Moreover, because 16S rRNA sequencing data do not directly provide insights into the functional capabilities of the microbiome community, we integrated the Tax4Fun tool into our pipeline to predict the functional profiles of microbial communities based on the 16S rRNA datasets. Therefore, the RF-FVS approach could detect the minimal-size core set and optimal predictive subset of functional profiles of the selected microbial species and the linkages among them, which will provide insights into the ecological functioning of habitats. Some unknown species and genes within the dedicated taxa and functional profiles were detected in the group of selected features that made meaningful contributions to the predictive performance of the RF classifier. These species and genes are likely to be good candidates for future experimental studies.

Secondly, to overcome the computational burden of high-dimensional data that limits the implementation of existing machine learning approaches, we developed a parallel computational strategy algorithm for handling large-scale problems in the forward variable selection algorithm. This parallel strategy helps to equally divide the computational burden of the search processes among processors. Thus, the forward variable selection computation process is completely optimized and parallelized based on data partitioning. In datasets of tens of thousands of species and their functional profiles, selecting only a few hundred of the most significant samples can be a major problem. Our current strategy is focused on parallelly searching for the features of interest. In the future, the numbers of samples and features in microbiome datasets are likely to explode at a rapid pace. We anticipate that hybrid-partitioning strategies that partition the data both horizontally (over samples) and vertically (over features) will become essential to speed up the computational processes [54] [55].

Acknowledgment

We thank Margaret Biswas, PhD, from Edanz Group (https://en-author-services.edanzgroup.com/ac) for editing a draft of this manuscript. This study was supported by Grant-in-Aid for Scientific Research (B) 19H04070 from the Japan Society for the Promotion of Science.

References

[1] Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y. Y., Keilbaugh, S. A., ... & Sinha, R, "Linking long-term dietary patterns with gut microbial enterotypes," *Science,* vol. 334, no. 6052, pp. 105-108, 2011.

[2] Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., ... & Peng, Y, "A metagenome-wide association study of gut microbiota in type 2 diabetes," *Nature,* vol. 490, no. 7418, pp. 55-60, 2012.

[3] Arumugam, M., Raes, J., Pelletier, E., Le Paslier, D., Yamada, T., Mende, D. R., ... & Bertalan, M, "Enterotypes of the human gut microbiome," *Nature,* vol. 473, no. 7346, pp. 174-180, 2011.

[4] Turnbaugh, P. J., Ley, R. E., Hamady, M., Fraser-Liggett, C. M., Knight, R., & Gordon, J. I, "The human microbiome project," *Nature,* vol. 449, no. 7164, pp. 804-810, 2007.

[5] Knight, R., Vrbanac, A., Taylor, B. C., Aksenov, A., Callewaert, C., Debelius, J., ... & Melnik, A. V, "Best practices for analysing microbiomes," *Nature Reviews Microbiology,* vol. 16, no. 7, pp. 410-422, 2018.

[6] Zhou, Y. H., & Gallins, P, "A review and tutorial of machine learning methods for microbiome host trait prediction," *Frontiers in Genetics,* vol. 10, p. 579, 2019.

[7] Breiman, L, "Random forests," *Machine learning,* vol. 45, no. 1, pp. 5-32, 2001.

[8] Ho, T. K, "The random subspace method for constructing decision forests," *IEEE transactions on pattern analysis and machine intelligence,* vol. 20, no. 8, pp. 832-844, 1998.

[9] Dietterich, T. G, "An experimental comparison of three methods for constructing ensembles of decision trees: Bagging, boosting, and randomization," *Machine learning,* vol. 40, no. 2, pp. 139-157, 2000.

[10] Bureau, A., Dupuis, J., Falls, K., Lunetta, K. L., Hayward, B., Keith, T. P., & Van Eerdewegh, P, "Identifying SNPs predictive of phenotype using random forests," *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society,* vol. 28, no. 2, pp. 171-182, 2005.

[11] Díaz-Uriarte, R., & De Andres, S. A, "Gene selection and classification of microarray data using random forest," *BMC bioinformatics,* vol. 7, no. 1, p. 3, 2006.

[12] Knights, D., Costello, E. K., & Knight, R, "Supervised classification of human microbiota," *FEMS microbiology reviews,* vol. 35, no. 2, pp. 343-359, 2011.

[13] Silverman, J. D., Washburne, A. D., Mukherjee, S., & David, L. A, "A phylogenetic transform enhances analysis of compositional microbiota data," *Elife,* p. e21887, 2017.

[14] Aßhauer, K. P., Wemheuer, B., Daniel, R., & Meinicke, P, "Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data," *Bioinformatics,* pp. 2882-2884, 2015.

[15] Wemheuer, F., Taylor, J. A., Daniel, R., Johnston, E., Meinicke, P., Thomas, T., & Wemheuer, B, "Tax4Fun2: prediction of habitat-specific functional profiles and functional redundancy based on 16S rRNA gene sequences," *Environmental Microbiome,* pp. 1-12, 2020.

[16] Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J, "Microbiome datasets are compositional: and this is not optional," *Frontiers in microbiology,* p. 2224, 2017.

[17] Kursa, M. B., & Rudnicki, W. R, "Feature selection with the Boruta package," *J Stat Softw,* pp. 1-13..

[18] Alyxandria M. Schubert, Mary A. M. Rogers, Cathrin Ring, Jill Mogle, Joseph P. Petrosino, Vincent B. Young, David M. Aronoff, Patrick D. Schlossa, "Microbiome data distinguish patients with Clostridium difficile infection and non-C. difficile-associated diarrhea from healthy controls," *MBio,* p. 5(3), 2014.

[19] Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R, "Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy," *Applied and environmental microbiology,* no. 73, pp. 5261-5267, 2007.

[20] Baxter, N. T., Ruffin, M. T., Rogers, M. A., & Schloss, P. D, "Microbiota-based model improves the sensitivity of fecal immunochemical test for detecting colonic lesions," *Genome medicine,* no. 8, pp. 1-10,

2016.

[21] Wirbel, J., Pyl, P. T., Kartal, E., Zych, K., Kashani, A., Milanese, A., ... & Sunagawa, S, "Meta-analysis of fecal metagenomes reveals global microbial signatures that are specific for colorectal cancer," *Nature medicine,* vol. 4, no. 25, pp. 679-689, 2019.

[22] Milanese, A., Mende, D. R., Paoli, L., Salazar, G., Ruscheweyh, H. J., Cuenca, M., ... & Schmidt, T. S, "Microbial abundance, activity and population genomic profiling with mOTUs2," *Nature communications,* no. 10, pp. 1-11, 2019.

[23] Milani, C., Ticinesi, A., Gerritsen, J., Nouvenne, A., Lugli, G. A., Mancabelli, L., ... & Ferrario, C, "Gut microbiota composition and Clostridium difficile infection in hospitalized elderly individuals: a metagenomic study," *Scientific reports,* vol. 6, no. 1, pp. 1-12, 2016.

[24] Ross, C. L., Spinler, J. K., & Savidge, T. C, "Structural and functional changes within the gut microbiota and susceptibility to Clostridium difficile infection," *Anaerobe,* vol. 41, pp. 37-43, 2016.

[25] Ling, Z., Liu, X., Jia, X., Cheng, Y., Luo, Y., Yuan, L., ... & Xu, X, "Impacts of infection with different toxigenic Clostridium difficile strains on faecal microbiota in children," *Scientific reports,* no. 4, p. 7485, 2014.

[26] De Wolfe, T. J., Eggers, S., Barker, A. K., Kates, A. E., Dill-McFarland, K. A., Suen, G., & Safdar, N, "Oral probiotic combination of Lactobacillus and Bifidobacterium alters the gastrointestinal microbiota during antibiotic treatment for Clostridium difficile infection," *PLoS One,* vol. 13, no. 9, p. e0204253, 2018.

[27] Kolling, G. L., Wu, M., Warren, C. A., Durmaz, E., Klaenhammer, T. R., & Guerrant, R. L, "Lactic acid production by Streptococcus thermophilus alters Clostridium difficile infection and in vitro Toxin A production," *Gut Microbes,* vol. 3, no. 6, pp. 523-529, 2012.

[28] Gazzola, A., Panelli, S., Corbella, M., Merla, C., Comandatore, F., De Silvestri, A., ... & Cambieri, P, "Microbiota in Clostridioides difficile-Associated Diarrhea: Comparison in Recurrent and Non-Recurrent Infections," *Biomedicines,* vol. 8, no. 9, p. 335, 2020.

[29] Pérez-Cobas, A. E., Artacho, A., Ott, S. J., Moya, A., Gosalbes, M. J., & Latorre, A., "Structural and functional changes in the gut microbiota associated to Clostridium difficile infection," *Frontiers in microbiology,* p. 335, 2014.

[30] Tanaka, S., Kobayashi, T., Songjinda, P., Tateyama, A., Tsubouchi, M., Kiyohara, C., ... & Nakayama, J, "Influence of antibiotic exposure in the early postnatal period on the development of intestinal microbiota," *FEMS Immunology & Medical Microbiology,* vol. 56, no. 1, pp. 80-87, 2009.

[31] Prasad, N., Labaze, G., Kopacz, J., Chwa, S., Platis, D., Pan, C. X., ... & Kreiswirth, B. N, "Asymptomatic rectal colonization with carbapenem-resistant Enterobacteriaceae and Clostridium difficile among residents of a long-term care facility in New York City," *American journal of infection control,* vol. 44, no. 5, pp. 525-532, 2016.

[32] Seddon, M. M., Bookstaver, P. B., Justo, J. A., Kohn, J., Rac, H., Haggard, E., ... & Al-Hasan, M. N, "Role of early de-escalation of antimicrobial therapy on risk of Clostridioides difficile infection following Enterobacteriaceae bloodstream infections," *Clinical Infectious Diseases,* vol. 69, no. 3, pp. 414-420, 2019.

[33] Maslowski, K. M., Vieira, A. T., Ng, A., Kranich, J., Sierro, F., Yu, D., ... & Xavier, R. J, "Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43," *Nature,* vol. 461, no. 7268, pp. 1282-1286, 2009.

[34] Koruda, M. J., Rolandelli, R. H., Bliss, D. Z., Hastings, J., Rombeau, J. L., & Settle, R. G, "Parenteral nutrition supplemented with short-chain fatty acids: effect on the small-bowel mucosa in normal rats," *The American journal of clinical nutrition,* vol. 51, no. 4, pp. 685-689, 1990.

[35] Lawley, T. D., Clare, S., Walker, A. W., Stares, M. D., Connor, T. R., Raisen, C., ... & Deakin, L. J, "Targeted restoration of the intestinal microbiota with a simple, defined bacteriotherapy resolves relapsing Clostridium difficile disease in mice," *PLoS Pathog,* vol. 8, no. 10, p. e1002995, 2012.

[36] Antharam, V. C., Li, E. C., Ishmael, A., Sharma, A., Mai, V., Rand, K. H., & Wang, G. P, "Intestinal dysbiosis and depletion of butyrogenic bacteria in Clostridium difficile infection and nosocomial diarrhea," *Journal of clinical microbiology,* vol. 51, no. 9, pp. 2884-

2892, 2013.

[37] Theriot, C. M., Koenigsknecht, M. J., Carlson Jr, P. E., Hatton, G. E., Nelson, A. M., Li, B., ... & Young, V. B, "Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase susceptibility to Clostridium difficile infection," *Nature communications,* no. 5, p. 3114, 2015.

[38] Yu, J., Feng, Q., Wong, S. H., Zhang, D., yi Liang, Q., Qin, Y., ... & Wang, X, "Metagenomic analysis of faecal microbiome as a tool towards targeted non-invasive biomarkers for colorectal cancer," *Gut,* pp. 70-78, 2017.

[39] Feng, Q., Liang, S., Jia, H., Stadlmayr, A., Tang, L., Lan, Z., ... & Su, L., "Gut microbiome development along the colorectal adenoma–carcinoma sequence," *Nature communications,* p. 6528, 2015.

[40] Gupta, A., Dhakan, D. B., Maji, A., Saxena, R., PK, V. P., Mahajan, S., ... & Amato, K. R, "Association of Flavonifractor plautii, a flavonoid-degrading bacterium, with the gut microbiome of colorectal cancer patients in India," *mSystems,* vol. 4, no. 6, pp. e00438-19, 2019.

[41] Castellarin, M., Warren, R. L., Freeman, J. D., Dreolini, L., Krzywinski, M., Strauss, J., ... & Holt, R. A, "Fusobacterium nucleatum infection is prevalent in human colorectal carcinoma," *Genome research,* vol. 22, no. 2, pp. 299-306, 2012.

[42] Kostic, A. D., Gevers, D., Pedamallu, C. S., Michaud, M., Duke, F., Earl, A. M., ... & Baselga, "Genomic analysis identifies association of Fusobacterium with colorectal carcinoma," *Genome research,* vol. 22, no. 2, pp. 292-298, 2012.

[43] Kwong, T. N., Wang, X., Nakatsu, G., Chow, T. C., Tipoe, T., Dai, R. Z., ... & Chan, F. K, "Association between bacteremia from specific microbes and subsequent diagnosis of colorectal cancer," *Gastroenterology,* pp. 383-390, 2018.

[44] Yang, Y., Cai, Q., Shu, X. O., Steinwandel, M. D., Blot, W. J., Zheng, W., & Long, J, "Prospective study of oral microbiome and colorectal cancer risk in low‐income and African American populations," *International journal of cancer,* vol. 144, no. 10, pp. 2381-2389.

[45] Szafrański, S. P., Deng, Z. L., Tomasch, J.,

Jarek, M., Bhuju, S., Meisinger, C., ... & Wagner-Döbler, I, "Functional biomarkers for chronic periodontitis and insights into the roles of Prevotella nigrescens and Fusobacterium nucleatum; a metatranscriptome analysis," *npj Biofilms and Microbiomes,* pp. 1-13, 2015.

[46] Nobili, V., Putignani, L., Mosca, A., Del Chierico, F., Vernocchi, P., Alisi, A., ... & Drago, L, "Bifidobacteria and lactobacilli in the gut microbiome of children with non-alcoholic fatty liver disease: which strains act as health players?," *Archives of medical science: AMS,* vol. 14, no. 1, p. 81, 2018.

[47] Agyirifo, D. S., Wamalwa, M., Otwe, E. P., Galyuon, I., Runo, S., Takrama, J., & Ngeranwa, J, "Metagenomics analysis of cocoa bean fermentation microbiome identifying species diversity and putative functional capabilities," *Heliyon,* p. e02170, 2019.

[48] Xie, M., Wu, J., An, F., Yue, X., Tao, D., Wu, R., & Lee, Y, "An integrated metagenomic/metaproteomic investigation of microbiota in dajiang-meju, a traditional fermented soybean product in Northeast China," *Food Research International,* pp. 414-424, 2019.

[49] O'Callaghan, A., & van Sinderen, D, "Bifidobacteria and their role as members of the human gut microbiota," *Frontiers in microbiology,* p. 925, 2016.

[50] Nawrocki, K. L., Wetzel, D., Jones, J. B., Woods, E. C., & McBride, S. M., "Ethanolamine is a valuable nutrient source that impacts Clostridium difficile pathogenesis," pp. 1419-1435, 2018.

[51] Collins, J., Robinson, C., Danhof, H., Knetsch, C. W., Van Leeuwen, H. C., Lawley, T. D., ... & Britton, R. A, "Dietary trehalose enhances virulence of epidemic Clostridium difficile," *Nature,* vol. 553, pp. 291-294, 2018.

[52] Deshpande, A., Wu, X., Huo, W., Palmer, K. L., & Hurdle, J. G, "Chromosomal Resistance to Metronidazole in Clostridioides difficile can be Mediated By Epistasis Between Iron Homeostasis and Oxidoreductases," *Antimicrobial Agents and Chemotherapy,* 2020.

[53] Palaniappan, A., Ramar, K., & Ramalingam, S, "Computational identification of novel stage-specific biomarkers in colorectal cancer progression," *PloS one,* vol. 11, no. 5, p. e0156665, 2016.

[54] Xing, E. P., Ho, Q., Xie, P., & Wei, D, "Strategies and principles of distributed machine learning on big data," *Engineering,* vol. 2, no. 2, pp. 179-195, 2016.

[55] Lee, S., Kim, J. K., Zheng, X., Ho, Q., Gibson, G. A., & Xing, E. P, "On model parallelization and scheduling strategies for distributed machine learning," *In Advances in neural information processing systems,* pp. 2834-2842, 2014.

[56] Duvallet, C., Gibbons, S. M., Gurry, T., Irizarry, R. A., & Alm, E. J, "Meta-analysis of gut microbiome studies identifies disease-specific and shared responses," *Nature communications,* pp. 8(1), 1-10, 2017.

[57] Kanehisa, M., Goto, S., Sato, Y., Kawashima, M., Furumichi, M., & Tanabe, M, "Data, information, knowledge and principle: back to metabolism in KEGG," *Nucleic acids research,* vol. 42, no. D1, pp. D199-D205, 2014.

[58] Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M. C., ... & Jensen, L. J, "eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences," *Nucleic acids research,* vol. 44, no. D1, pp. D286-D293, 2016.