Sea anemone genomes reveal ancestral metazoan chromosomal macrosynteny

Bob Zimmermann[1,2,#], Juan D. Montenegro[1,2,#], Sofia M.C. Robb[3], Whitney J. Fropf[3], Lukas Weilguny[1], Shuonan He[3], Shiyuan Chen[3], Jessica Lovegrove-Walsh[1], Eric M. Hill[3], Cheng-Yi Chen[3], Katerina Ragkousi[3,4], Daniela Praher[1], David Fredman[1], Darrin Schultz[1], Yehu Moran[1,5], Oleg Simakov[1,2], Grigory Genikhovich[1], Matthew C. Gibson[3]*, Ulrich Technau[1,2,6]*

[1]Department of Neurosciences and Developmental Biology, Faculty of Life Sciences, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[2]Research platform SinCeReSt, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria
[3]Stowers Institute for Medical Research, Kansas City, MO 64110 USA
[4]Department of Biology, Amherst College, Amherst, MA 01002 USA
[5]*Current address:* The Alexander Silberman Institute of Life Sciences, Faculty of Science, The Hebrew University of Jerusalem, Jerusalem 9190401, Israel
[6]Max Perutz laboratories, University of Vienna, Dr. Bohrgasse 5, 1030 Vienna

[#] these authors contributed equally
*To whom correspondence should be addressed: ulrich.technau@univie.ac.at, mg2@stowers.org

Abstract

Draft genome sequences of non-bilaterian species have provided important insights into the evolution of the metazoan gene repertoire. However, there is little information about the evolution of gene clusters, genome architectures and karyotypes during animal evolution. In this regard, slowly evolving anthozoan Cnidaria, the sister group of Bilateria, are particularly informative. Here we report chromosome-level genome assemblies of two related cnidarians, the sea anemones *Nematostella vectensis* and *Scolanthus callimorphus*. We find a robust set of 15 chromosomes with a clear one-to-one correspondence between the two species. Both sea anemone genomes show remarkable chromosomal conservation with other cnidarians, several bilaterians and the sponge *Ephydatia muelleri*, allowing us to reconstruct ancestral cnidarian and metazoan chromosomal blocks, consisting of at least 19 and 16 ancestral linkage groups, respectively. We show that, in contrast to Bilateria, the Hox and NK clusters of investigated cnidarians are largely disintegrated, despite the presence of staggered *hox/gbx* expression in *Nematostella*. This loss of microsynteny conservation may be facilitated by shorter distances between cis-regulatory sequences and their cognate transcriptional start sites. In line with that, we find no clear evidence for topologically associated domains, suggesting fundamental differences in long-range gene regulation compared to vertebrates. These data suggest that large sets of ancestral metazoan genes have been retained in ancestral linkage groups of some extant lineages, yet, higher order gene regulation with associated 3D architecture may have evolved only after the cnidarian-bilaterian split.

Introduction

Comparative genomics and epigenomics have provided fundamental insights into the evolution of gene regulation in multicellular organisms. Among basally branching animals, this led to the surprising discovery of conserved gene repetoires[1–3], of microsyntenic gene clusters and patterns of DNA methylation and histone modification codes[4–6]. Recent applications of long-read sequencing technologies and proximity ligation methods such as Hi-C facilitated the genome assembly on chromosomal level of many organisms. These chromosome-level genome assemblies have presented the opportunity to compare the content and localization of homologous genes between distantly related species and the reconstruction of ancestral linkage groups. In vertebrates and several other bilaterians, topological analysis has revealed three-dimensional chromosomal architecture[7], organized as chromosomal compartments and topologically associated domains (TADs)[8]. TADs have been shown to act as boundaries of gene regulation[9–12]. Analyses on chromosome evolution and three-dimensional genome structure have mainly focused on bilaterians, and it remains unclear whether non-bilaterian animals share the topological signatures of bilaterian chromosomes. In this regard, studying Cnidaria, the sister clade to Bilateria, is a crucial step to understanding the evolution of animal genomes.

Cnidaria constitute a large clade of basally branching Metazoa, dating back between 590 and 690
Mya[13–15]. Their robust phylogenetic position as the sister group to Bilateria makes them the key
group to study the evolution of bilaterian features, such as axis organization, mesoderm formation
and central nervous system development[16]. The starlet sea anemone, *Nematostella vectensis*
(Figure 1a - Hexacorallia; Actiniaria; Edwardsiidae), has been developed into an important model
organism[17–19] and in 2007 became the first non-bilaterian animal to have a draft scaffold-level
genome assembled[1]. The *Nematostella* genome revealed uncanny conservation of gene content
to vertebrates as well as the first observations of large-scale macrosyntenic conservation by way
of comparisons with human chromosomes to *Nematostella* scaffolds[1].
92
Since the release of the *Nematostella* genome, genomes of the representatives of all five
cnidarian classes have become available[2,20–27] providing valuable insight into various aspects of
the cnidarian gene complement and genome organization. However, these genomes originated
from distantly-related species, and few genomic studies of Cnidaria have sought to search for
genomic conservation signals of recently-diverged species, a cornerstone of comparative
genomics[28]. For example, no genome sequence of an edwardsiid sea anemone has been
reported, aside from that of *Nematostella*. Another sea anemone in the family Edwardsiidae is
the "worm sea anemone" *Scolanthus callimorphus* (Figure 1b), dwelling in European intertidal
zones[29,30], which according to our molecular clock calculations has separated from *Nematostella*
vectensis approximately 174 Mio years ago, to date its closest sequenced relative (EDF1, see
Materials and Methods for details).
104
Here, we report chromosome-level genome assemblies and new gene models for the model
organism *Nematostella vectensis* and of the closely-related species *Scolanthus callimorphus*. We
show that the macrosyntenic localization of orthologous genes across the chromosomes are
highly conserved across many cnidarians and bilaterians, allowing us to reconstruct the minimal
set of ancestral eumetazoan chromosomal linkage blocks. Notably, the macrosyntenic
conservation contrasts with the loss of most of the microsyntenic blocks, including the integrity of
the Hox cluster. Moreover, in *Nematostella*, which like all non-bilaterians lacks the TAD boundary
protein CTCF[4,31], HiC analyses did not reveal any obvious TAD-like structures, as known in
Bilateria. We propose that the proximity of most cis-regulatory regions to their target genes is an
ancestral metazoan feature, whereas large-scale three-dimensional structures, such as TADs,
only became necessary in bilaterians with larger genomes and more distantly located enhancers.
116
117 **Results**
118
119 **High Quality Chromosome-Level Assemblies of Two Edwardsiid Genomes**
120
Using short-read sequencing and a *k*-mer coverage model, we estimated the genome length of
Nematostella at 244 Mb (EDF2), which is substantially shorter than previously suggested at 450
Mb[1]. This discrepancy could be in part attributed to the previous use of four haplotypes in
sequencing, lower coverage and read length. The genome of the sea anemone *Exaiptasia pallida*
is similar in length to *Nematostella*[32], while the estimated 414 Mb of the *Scolanthus* genome is at
present the largest sequenced sea anemone genome (Figure 1c). Using PacBio long-read

sequencing and high-throughput conformation capture (Hi-C), we then assembled chromosome-level *Nematostella* and *Scolanthus* genomes, which surpass the quality of the original *Nematostella* genome in terms of contiguity, accuracy and mappability (see Supplementary Text, EDF2,3 for details).

Gene completeness as measured by alignment of single-copy pan-metazoan genes to the assemblies using the BUSCO method[33] was comparable to the previous assembly (EDF2i), however in order to completely assess this, we generated new gene models. Using a combination of IsoSeq and RNASeq data, we identified 24,525 gene models and 36,280 transcripts (see Supplementary Text for details). BUSCO analysis showed that the gene set was more complete than a previously-generated gene set[34]. In addition, the new gene models showed a better mappability to a publicly available single-cell RNA-Seq data set[35,36] (EDF4).

To facilitate the usage of the newly assembled genomes, we established a publicly accessible genome browser. Both new genome assemblies and associated data are available for browsing, downloading, and BLAST at SIMRbase (https://simrbase.stowers.org). The *Nematostella vectensis* genome assembly, referred to as Nvec200, has an abundance of aligned track data, including the newly generated gene models, a large collection of published RNAseq and ChIP-seq analyses, as well as 145 ultra-conserved non-coding elements (UCNEs) shared between *Nematostella* and *Scolanthus* (Supplementary Text; EDT8).

Comparison of *Nematostella* Chromosomes to Metazoan Genomes

We identified 15 chromosomes in the new genome assemblies of *Nematostella* and *Scolanthus*, and we numbered these chromosomes from 1 to 15 according to decreasing size (Figure 1d,e). This is in line with the previous estimates based on the number of *Nematostella* metaphase plates[1] and the analysis of *Nematostella* chromosome spreads[37]. Most chromosomes, according to their homologous pair, corresponded in length but are much larger in *Scolanthus* (Figure 4d). This was accounted for by a large fraction of unclassified and potentially lineage-specific repeat sequences (EDT1, Supplementary text). No obvious heteromorphic sex chromosomes could be identified based on read mapping depth, or from the level of heterochromatin usually associated with non-recombinant sex chromosomes. Ascertaining the sex-determining regions of the genome will require more data and detailed analyses.

Next we wished to determine the extent to which the *Nematostella* and *Scolanthus* chromosomes exhibit conservation of gene content and order (micro- and macrosynteny). Indeed, each of the 15 chromosomes of both species shared a majority of orthologous genes with a single corresponding chromosome in the other species (Figure 2a). We found that 8117 of 8692 mutual best BLAST hits between *Nematostella* and *Scolanthus* were retained on their respective chromosomes, implying a one-to-one homology between all 15 chromosomes. However, gene order was largely lost from the most recent common ancestor (MRCA), which we estimate to have diverged approximately 174 Mya (EDF1).

To assess the macrosyntenic conservation between anthozoans we compared the *Nematostella*
chromosomes to those of the sea anemone *Exaiptasia pallida,* the stony coral *Acropora millepora*
and the soft coral *Xenia sp*. (Figure 2a). While both *Exaiptasia* and *Acropora* are only assembled
on the scaffold level, we observed that the gene content of the scaffolds suggest a similar
karyotype to *Nematostella*. The chromosome-level genome assembly of the octocoral soft coral
Xenia sp. also appears to have 15 chromosome-scale scaffolds[20]. However, despite the apparent
identical number of chromosomes, the chromosome-scale scaffolds of *Xenia* showed surprisingly
little homology to those of the hexacoral *Nematostella*. We discovered many translocations and
fusions, with only 3 chromosomes demonstrating a one-to-one relationship (Figure 2a).
179
Notably, the hydrozoan *Hydra vulgaris,* which belongs to the clade of Medusozoa, also has 15
chromosomes[38]. While our analyses revealed many ancestral linkage groups in the comparison
with *Hydra*, we also detected numerous chromosomal rearrangements, including splits and
fusions (Figure 2a, EDF5). Specifically, we observe that *Nematostella* chromosomes 2 and 3 are
partially or completely split in other cnidarian genomes. By comparison, the scyphozoan jellyfish
Rhopilema has 21 chromosomes, supported with karyotype images[21,39]. We observed that all
Rhopilema and *Nematostella* chromosomes show a clear 1-to-1, 1-to-2 or, in a single case, a 1-
to-3 macrosyntenic correspondence, which can be traced to the ancestral linkage groups
(EDF5,7). These comparisons among cnidarians suggest that the history of cnidarian
chromosomes may be more complex than previously envisioned. This is in line with recent
interpretations of cnidarian chromosome comparisons[40].
191
We then compared the chromosome-scale scaffolds of *Nematostella* with that of a bilaterian, the
cephalochordate *Branchiostoma floridae*. Cephalochordates are early branching chordates
lacking the two rounds of whole-genome duplication and allotetraploidization suggested for
vertebrates[41,42], as represented by their single Hox cluster[43] (Figure 2a). Strikingly,
Branchiostoma and *Nematostella* chromosomes retain extensive macrosynteny since their
divergence from the bilaterian-cnidarian MRCA. The large-scale macrosynteny retention detected
in *Branchiostoma* becomes less obvious once we focus on more recently-branching bilaterian
clades. Vertebrates, such as the early branching teleost fish *Lepisosteus oculatus[44]* and humans,
underwent two or more rounds of genome duplications and showed additional translocation
events from the MRCA (Figure 2a). Among protostomes, we observed that the lophotrochozoan
Ram's horn worm *Streplosbio benedicti* and the sea scallop *Patinopecten yessoensis[45]*, as well
as the ecdysozoan millipede *Trigoniulus corallinus[46],* retained many macrosyntenic links (Figure
2a)*.
205
The high degree of macrosynteny conservation with *Branchiostoma* motivated us to compare
Nematostella chromosomes to those of the representative of an even older lineage, the Porifera
(sponges), which branched off prior to the cnidarian-bilaterian split (EDF1) and is considered by
many to be the sister group to all other animals[47,48]*. Strikingly, the comparison with the recently-
assembled chromosomes of the sponge *Ephydatia muelleri* revealed many shared ancestral
linkage groups (Figure 2a).
212
213   **Reconstruction of Ancestral Linkage Groups**

214
The observed patterns of conserved macrosynteny between chromosomes of multiple distantly-
related species and the varying degrees of gene shuffling and chromosomal rearrangements in
others (e.g. *Drosophila, Caenorhabditis,* humans) prompted us to reconstruct the set of predicted
ancestral linkage groups for the MRCA of cnidarians, bilaterians and metazoans (Figure 2b). First,
we inferred gene orthology across all genomes investigated. For each of Metazoa, Bilateria and
Cnidaria, we determined ancestral linkage groups by determining genes with high chromosomal
linkage throughout each clade. Our method finds ALGs which maximize their "modularity", i.e. groups
with high chromosomal linkage within ALGs and low linkage between ALGs, by evaluating multiple
scenarios (see Methods and Supplemental Methods for details) to infer "ancestral linkage groups"
(ALGs). We can then color the oxford plots with the genes in these blocks suggesting a pattern
of splits and fusions in extant chromosomes from ancestral chromosomes.
226
We identified 17 ALGs present in the ancestor of cnidarians, although we could find similar
modularity in scenarios of 16-18 ALGs (EDF14). These correspond well to the extant cnidarian
chromosomes (Figure 2c, EDF5), however, while the *Rhopilema* and edwardsiid chromosomes
appear highly representative of the ancestral cnidarian karyotype, the soft coral *Xenia*
chromosomes appear to have undergone more chromosomal translocation events. Our results
also indicate that although both Hydrozoa and Edwardsiidae have a clear 15 chromosome
karyotype, their chromosomes originate from distinct fusions of cnidarian ALGs. For the ancestor
of Bilateria, we identified 16 ALGs (Figure 2c). In a previous study 17 ALGs were proposed[14],
which resulted in similarly high modularity in our analysis (EDF14). By comparison with the
sponge *Ephydatia,* we reconstructed a minimum of 16 metazoan ALGs for the metazoan
ancestor, which were maintained in the MRCA of Cnidaria, Bilateria and the sponge (Figure 2b,c).
238
In order to visualize how the chromosomes of extant species have undergone major splits and
fusions, we projected the reconstructed ALGs of the respective cnidarian, bilaterian and metazoan
ancestor to the chromosomes of the extant species. We found that many ALGs correspond to the
same chromosomes across multiple species and lineages (Figure 2c; EDF5-7). To determine the
extent of this, we further explored the relationships between metazoan ALGs and those of
Cnidaria and Bilateria (Figure 2c). Remarkably, many ancestral chromosomes exhibited a 1-to-1
correspondence across all predicted ancestral lineages (Figure 2c) and carried through to the
extant lineages (Figure 2d).
247
248 **Chromosomal Organization of the NK and extended Hox gene clusters**
249
The chromosome-level assembly of the *Nematostella* genome allowed us to address the evolution
of specific gene clusters. Prominent examples of clusters of homeodomain transcription factor
coding genes ancestral for Bilateria include the SuperHox cluster, the ParaHox cluster, the
NK/NK-like cluster as well as *NK2* group genes located separately[43,49,50]. It has been hypothesized
that all of them originated from a single gene cluster, which then disintegrated during evolution[50].
Our analysis revealed that *Nematostella* possesses a separate ParaHox cluster of two genes,
(*Gsx* and *Xlox/Cdx)* on chromosome 10, and a SuperHox cluster on chromosome 2 containing
Hox, *Evx*, *Mnx*, and *Rough*, as well as more distant *Mox* and *Gbx*[51] (Figure 3, EDF8, EDT6). We

identified an *NK* cluster on chromosome 5 containing *NK1*, *NK5*, *Msx*, *NK4*, *NK3*, *NK7*, *NK6*, a
more distant *Lbx*, a possible highly derived *Tlx-like* gene and, intriguingly, *Hex,* which is also
linked to the NK cluster in the hemichordate *Saccoglossus kowalevskii*[52] and in the
cephalochordate *Branchiostoma floridae*. Similar to Bilateria, the *NK2* genes were clustered
separately and found on the chromosome 2 (Figure 3, EDF8-9, EDT6). In contrast, in the earlier-
branching sponges, neither ParaHox nor extended Hox cluster genes exist, and only the NK
cluster is present with a single *NK2/3/4* gene, two *NK5/6/7* genes, an *Msx* ortholog, as well as
possible *Hex* and *Tlx* orthologs[53], (Figure 3, EDF10). Taken together, this allows us to propose
that the last common ancestor of Cnidaria and Bilateria possessed an NK-cluster on a
chromosome different from the one carrying the SuperHox cluster, and a separate NK2 cluster,
possibly on the same chromosome as the SuperHox cluster (Figure 3). The hypothesized
SuperHox-NK Megacluster[43], if it ever existed, must have both formed and broken apart during
the time after the separation of the sponge lineage, but before the origin of the cnidarian-bilaterian
ancestor (Figure 3a, Supplementary Text). The lack of selection pressure in favor of microsynteny
conservation is clearly illustrated by the comparison of the divergent Hox clusters of *Scolanthus*
and *Nematostella*. Although located on homologous chromosomes, the gene order, orientation,
and the number of intervening genes differs drastically between these two species (Figure 3b). In
contrast to the atomized Hox clusters of *Nematostella* and *Scolanthus*, the Hox cluster of their
very distant anthozoan relative *Xenia*, is compact and contains "anterior" Hox genes *HoxA* and
HoxB immediately next to the non-anterior Hox gene *HoxE* (Figure 3b) Aside from the tandem
duplications, this compact state likely represents the ancestral organization. Similarly, *HoxE* and
HoxB are immediate neighbors in the genome of the jellyfish *Rhopilema*, although the *Rhopilema*
Hox cluster shows some evidence of disintegration (Figure 3b).
281
282   **Topologically associating domains are not detected in either sea anemone genome**
283
In the past decade, high-resolution chromosome conformation capture has increased interest in
topologically associating domains (TADs), recurring chromosomal-folding motifs evidenced by
signals in Hi-C contact maps[54]. Flanking regions of TADs are positively correlated with CCCTC-
binding factor (CTCF) binding sites. Interestingly, no CTCF ortholog has been detected in non-
bilaterian animals[31,54], but previous studies of non-bilaterian animals have proposed the existence
of TAD-like structures[47].
290
Similar to previous studies, we sought to identify putative TADs within the edwardsiid genomes.
Initial inspections of the Hi-C maps of both *Scolanthus* and *Nematostella*, however, revealed that
both species appeared to lack TAD-like structures (Figure 4a; EDF11) and instead the Hi-C
contact frequency decayed smoothly as a function of read pair distance along the chromosomes
(EDF11). We quantified these findings by measuring the strength of topological boundaries,
measured by insulation score, and found the topological boundaries to be weaker in the
Nematostella genome compared to the fly (EDF12, p = 8.38×10^{-18}, Mann-Whitney *U* test) and
mouse genomes (p = 2.93×10^{-55}, Mann-Whitney *U* test). We also verified that there was a lack of
bilaterian-like TADs and weak topological domain boundaries in an independent *Nematostella*
chromosome-scale genome assembly and Hi-C dataset produced in parallel to ours (EDF11[55]).
301

TAD structures are often well-conserved around gene clusters that control spatio-temporal gene
expression, especially of developmental regulators, such as the TADs that separate the anterior
and posterior hox genes in the mouse HoxD cluster[56]. We therefore explored whether TAD-like
structures existed around Hox clusters in *Nematostella*. *Nematostella* has 6 Hox genes, one is on
chromosome 5, while the other 5 Hox genes are located on chromosome 2, but in two clusters,
separated by hundreds of genes between (Figure 4). We analyzed the genomic regions
surrounding both partial hox clusters but again we could not detect any significant TAD-like
structure.
310
311 **Correlation of gene regulation with three-dimensional genome organization**
312
We note that the *Nematostella* genome is fairly gene-dense (10±4 genes per 100kb), leaving
relatively little intergenic sequence. It is conceivable that most relevant cis-regulatory elements
are located in close proximity to the regulated gene. This is supported by the observation that
many *Nematostella* transgenic reporter constructs faithfully mimic endogenous expression
patterns with only 1-3 kilobases of upstream promoter sequence. Cis-regulatory regions are
reflected by ATAC-seq peaks, which detects open chromatin. We therefore wished to assess
whether there is a correlation between the distance of ATAC-seq peaks and the closest gene (i.e.
transcriptional start or end site) and the genome size. We compared several species, where
chromosome assemblies and ATACseq data are available and - as expected - we found that there
is indeed a correlation between distance and of ATAC-seq peaks to the next gene and the
genome size of the organism. However, there is also a correlation between small genomes and
the absence of clear TAD structures, for instance in *C. elegans*, sponges and cnidarians. Notably,
at least one cnidarian genome, *Hydra*, is relatively large (1.27 Gb), yet does not show typical TAD
structures, although smaller-scale 3D structures could be observed[57]. Thus, within bilaterians,
species with small genomes may show a tendency to lack TAD structures, while in non-
bilaterians, which lack CTCF, they may not organize into stable TAD-like structures. At present,
Hydra is the only available large non-bilaterian genome with relatively short distances between
ATAC-seq peaks to the next gene.
331
Since TADs not only assure maintenance and directionality of enhancers to the regulated gene,
they frequently also encompass more than one gene. In vertebrates, many TADs are conserved
between species, suggesting that their maintenance is under selection pressure. This predicts
that in organisms with TADs the number of microsyntenic genes is higher than in organisms
lacking TADs. To test this, we compared the number of microsyntenies in different cnidarians,
chordates and protostomes, roughly representing similar times of divergence. We find that, while
the amount of microsyntenies consisting mostly of two consecutive orthologous genes did not
show any difference between clades, microsyntenies consisting of at least three or four
orthologous genes were reduced in cnidarians. No microsyntenies of four and more genes were
found in any pair of species spanning the anthozoan-medusozoan split (e.g., *Nematostella-*
Rhopilema), whereas around a dozen such regions are shared among distantly branching
spiralian and chordate-vertebrate ("deuterostome") genomes (e.g., 11 blocks between the scallop
Pecten maximus bilaterian and the annelid *Streblospio benedicti*, and 14 blocks for amphioxus
Branchiostoma floridae to chicken *Gallus gallus* split) (Figure 4f). A similar trend was observed

for microsyntenies where intervening genes were allowed to be part of the blocks. For a commonly
used threshold of up to 5 intervening genes[58], *Nematostella-Rhopilema* species pair showed 7
blocks, whereas *Pecten-Streblospio* had 82, and amphioxus to chicken had 45 microsyntenic
blocks retained (Figure 4f, EDT12). Together, this data indicates higher retention and
evolutionary exploration of longer syntenic regions in bilaterian genomes, potentially allowed for
by the maintained distal topological architecture.

Discussion

The assembly of two high quality, chromosome-level edwardsiid sea anemone genomes has illuminated several intriguing aspects about chromosomal evolution, the NK and extended Hox clusters, the conservation of non-coding elements and the status of topologically associated domains in the common ancestor to cnidarians and bilaterians. In addition, the highly improved *Nematostella* genome and manually curated gene annotations will prove to be an invaluable resource for future studies of both coding and non-coding regions, structural variants among populations and continued development of functional tools for this model organism.

Nearly all members of the extended Hox cluster were distributed among distant, isolated microsyntenic blocks on chromosome 2 of *Nematostella* (pseudo-chromosome 4 in *Scolanthus*), with the single exception of *HoxF/Anthox1*, located on chromosome 5 (Figure 3; Supplementary Text). This indicates a lack of proximity constraint on the Hox genes in Cnidaria, contrasting with the situation in Bilateria. In addition, while a staggered spatiotemporal pattern of Hox expression along the secondary, directive axis of the *Nematostella* larva and polyp can be observed[59], unlike Bilateria, there is no correlation between expression and cluster position[60]. Notably, *HoxF/Anthox1* is not only located on a different chromosome but it is also the only Hox gene expressed in the (aboral) ectoderm, while all other Hox genes are expressed in the inner endomesodermal cell layer[61,62].

The dispersed NK and extended Hox clusters may be due to the diminishment or, possibe absence of higher-order chromosome organization at the level of microsynteny. In line with this, it was recently observed that the HoxD cluster boundaries in the mouse genome are marked by two TAD boundaries[56], and the cluster's intra-TAD gene order is deemed to be under selective pressure[63]. The lack of a CTCF gene in the *Nematostella* genome led us to hypothesize that the cnidarians might lack TADs, as TAD presence has been attributed to the appearance of CTCF[63]. Moreover, CTCF is absent not only in cnidarians but also in earlier branching ctenophores and sponges, which provides a possibility that the existence of TADs might represent a bilaterian-specific feature. While we were unable to detect any noticeable structure resembling the current definitions of TADs, it remains an open question as to whether larger or smaller structures, beyond the resolution of our data, could yet be detected. One study has suggested evidence for the higher-order chromosomal organization in a non-bilaterian, the sponge *Ephydatia*[47]. However, the contact maps resemble patterns we observe in our assemblies at the boundary of scaffolds or contigs, which can be the result of differential mappability from repetitive content or assembly issues. We therefore deliberately do not report any results from a TAD finder, since, after multiple rigorous rounds of manual assembly update, we can assert that the data we have generated do not qualitatively represent TAD boundaries per se, and most results would be likely false positives. While the precise definition of a TAD is still evolving[64,65], both data sets lack many characters of TADs identified in CTCF-containing genomes: hierarchical compartments, mammalian-specific "corner peaks" indicating strong interactions, and in our case, loop peaks and inter-contig compartments. This suggests that the presence of CTCF is necessary for the formation of TADs. The apparent absence of TADs in our analysis could also be explained by a higher degree of cell type variability. Therefore, we cannot exclude the possibility that performing the experiment with

402 a more homogenous cell population, or sequencing at a higher resolution, would reveal a signal
403 on a smaller scale.
404
While microsynteny analyses reveal little conservation of the local gene order in the genomes of
Nematostella and *Scolanthus*, macrosyntenic analysis of the edwardsiid chromosomes compared
to available cnidarian genomes revealed a high level of conservation. We identified a stable set
of 19 ALGs across all clades of sequenced cnidarian genomes. When compared to extant
genomes, we can trace a small number of recombination events from the ALGs since the common
cnidarian ancestor split an estimated 580 Mya. This stands in stark contrast to the history of, for
example, the 326 Mya old ancestral genome of Amniota, which is estimated to have 49 distinct
units, while the karyotypes of the extant amniote taxa consist of multiple translocated segments
and variable chromosomes[66]. However, far more remarkable is the macrosynteny maintained
between the edwardsiids, the early branching chordate *Branchiostoma*, and the sponge
Ephydatia. Our analyses revealed clear one-to-one, one-to-few or few-to-one conservation of the
chromosome-level linkages between cnidarians, sponges and early chordates, which suggests a
striking retention of macrosyntenies throughout evolution of these animal lineages. This in turn
allowed us to deduce a set of 16 ALGs of the last common metazoan ancestor, which was
maintained in the cnidarian-bilaterian ancestor and gave rise to the 19 ancestral cnidarian
chromosomes and the 17 ancestral bilaterian chromosomes[58]. The observed conservation of
macrosyntenic linkages can result from a strong selective pressure to maintain intact
chromosomes during meiosis, so that only local translocations within the chromosomes,
scrambling local gene order, are favored. Changes in population sizes or asexual reproduction or
selfing may facilitate chromosomal unit breakages. While such global chromosomal
rearrangements have been observed for some clades (most notably dipterans but also
nematodes and cephalopods), it still remains unclear whether additional gene regulatory
constraints may exist at the whole-chromosomal level[67]. It is tempting to speculate that the
emergence of the TADs in Bilateria may have restricted local rearrangements and at the same
time released the constraints on maintaining the ancestral macrosyntenies conserved all the way
back to the origin of multicellular animals.
431
The high level of macrosynteny maintaining ancestral chromosomal blocks over hundreds of
millions of years contrasts with the low level of microsynteny conservation, even between more
closely related cnidarians. Despite this reshuffling of genes within chromosomes, many
developmental regulators involved in axial patterning show a conserved expression pattern,
suggesting that their cis-regulatory elements have been maintained. For instance, in *Nematostella*
and *Hydra*, which are separated over 500 Million years, all 12 Wnt genes, *brachyury,* and *foxA*
are expressed at the oral pole, while *foxQ2* and *six3* are expressed aborally[68–71]. To ensure
conservation of spatio-temporal gene expression patterns cis-regulatory elements must remain
located in close proximity to the respective gene even when translocated to a new genomic
position. This is supported by our distance analyses of ATACseq peaks. By contrast, bilaterians
with large genomes, such as vertebrates, enhancers can be located up to several megabases
away from the gene they regulate, often with several genes in between. To maintain the gene
regulation by long distant enhancers, we predicted that the microsynteny conservation should be
under higher selective pressure than in non-bilaterians. Indeed, microsynteny conservation is

higher in vertebrates than in cnidarians, in line with recent comparisons between skates, mouse
and garfish[72]. We therefore propose a hypothetical evolutionary scenario, where close cis-
regulation is ancestral to metazoans and maintained in non-bilaterians, which lack CTCF. It
follows that CTCF evolved in the bilaterian common ancestor and was recruited to bind to the
genome to act as an insulator and, together with cohesin, as a structural component defining
recombination boundaries. This kept complex gene regulation by distant enhancers as a physical
boundary for recombination events, but also acted as an insulator in TADs. Bilaterians that evolve
a more compact genome would have a tendency to lose TADs and the dependence on CTCF.
This seems to be the case in the *C. elegans* autosomes[73,74] and is also predicted for other small
genomes, such as in urochordates.
456
457
458

Figure 1. a) *Nematostella vectensis*. photo credit: Patrick RH Steinmetz b) *Scolanthus callimorphus*. photo credit: Robert Reischl c) Genome size estimates of cnidarian genomes taken from[2,21,22,24–26,32,39,75–79].

Figure 2. a) Oxford plots of the macrosyntenic relationships of *Nematostella vectensis* (*y*-axis) to genomes
of various metazoans (*x*-axis). Pairwise orthologous are ordered by their position on the chromosome
labeled on the axis. Dots (genes) are colored according to their membership to the metazoan ancestral
linkage groups (ALGs) shown in (c). Species names are colored according to clade shown in (b). b)
Genomes represented in this figure and EDF5-7. c) Relationships between the ALGs colored according to
the metazoan ALG. Lines are drawn between ALGs to represent fissions. d) Graph representing the
relationships of metazoan (M), bilaterian (B) and cnidarian (C) ALGs to each other and extant genomes of
Ephydatia (em)*, Nematostella* (nv)*, Rhopilema* (re) and *Branchiostoma* (bf). Edges are colored by the
metazoan ALG in (b) from which its path has a source. Line width represents the fraction of the extant
genome or derived ALG (bilaterian or cnidarian) content of the ancestral genome. Faded edges represent
near one-to-one relationships (> .8).

Figure 3. Evolution of a selection of Antennapedia class homeobox gene clusters. a) Composition of the extended Hox, ParaHox and NK clusters in the sponge *Amphimedon*, the sea anemone *Nematostella*, in a chordate *Branchiostoma*, and the deduced cluster composition of the cnidarian-bilaterian ancestor (CBA). Greyed-out genes with question marks have uncertain orthology. Genes shown as borderless boxes have an uncertain position relative to neighboring cluster members. Stacked boxes represent clusters of paralogs of the indicated ancestral gene. Genes in immediate proximity are indicated by abutting boxes. Linked genes of the same class separated by 1 to 50 intervening genes are connected with solid lines, over 50, with dashed lines. Grey intergenic connectors in the CBA indicate that the distances and the number of the intervening genes between the cluster members cannot be estimated. *NK2* of the CBA may be linked to the extended Hox cluster. Since *Branchiostoma Gbx* remained unplaced in the chromosome-level assembly, its position was taken from the scaffold-level assembly of *Branchiostoma lanceolatum*[80]. A two-gene ParaHox CBA scenario is shown although a three-gene ParaHox CBA scenario is possible based on evidence from Scyphozoa[21]. b) Organization of the *Nematostella* Hox cluster in comparison to the Hox clusters of *Scolanthus*, the octocoral *Xenia* and a scyphozoan jellyfish *Rhopilema* indicates loss of microsynteny. c) Staggered expression of *Gbx* and Hox genes along the directive axis of *Nematostella* (oral view) partially reflects the position of the genes on the chromosome. Arrows show the direction of transcription for each of the genes. The number of intervening genes is indicated in white circles. (d-e) Chromosomal relationships, genomic content and locations of NK and extended Hox cluster and other landmark homeobox genes of (d) *Nematostella* and *Scolanthus* and (e) *Nematostella* and *Branchiostoma*. 1) Chord diagram of macrosyntenic relationships of chromosomes based on the inferred ancestral linkage groups. Chords represent the proportion of the chromosome's gene content in common with the compared genome, not the positional information. 2) Scaled and centered gene density relative to the respective genome. Red is high and blue is low. 3) Scaled and centered density of interspersed repeat elements relative to the respective genome. Red is high and blue is low. 4) Locations of the landmark homeobox genes.

Figure 4. Hi-C contact maps. Contact maps (a) Nematostella Hox clusters, (b) Mouse HoxD cluster and (c) Fly Hox cluster show striking differences in the patterns of intrachromosomal interactions; d) distribution of the distance from ATACseq peaks to their closest peak for human (Hs), mouse (Mm), Sea urchin (*Lytechinus variegatus*, Lv), mussel (*Patinopecten yessoensis*, Py), polychete (*Owenia fisuformis*, Of), fly (Dm), *Caenorhabditis* (Ce), *Hydra vulgaris* (Hv) and *Nematostella vectensis* (Nv); e) correlation of mean gene-ATACseq peak distance and genome size for the same species; f) number of microsyntenic blocks of 2, 3 and 4 genes in Cnidarians, Deterostomes and Spiralians. The number of syntenic blocks falls rapidly in Cnidarians compared to Deuterostomes and Spiralians as the synteny length increases.

Acknowledgements

We thank Matthew Nicotra for providing us with the HMW DNA extraction protocol used for *Scolanthus.* We thank Robert Reischl for the photo of *Scolanthus* and Patrick R.H. Steinmetz and Hanna Kraus for the photo of *Nematostella vectensis* (Figure 1). Special thanks to Tatiana

Lebedeva for the cartoon drawings of animals used in this study. We are grateful to the Stowers
Institute Molecular Biology Core facility, particularly Amanda Lawlor, Michael Peterson and Anoja
Perera. This work was supported by grants of the Austrian Science Fund FWF (P24858; P21108)
to U.T., support from the Stowers Institute for Medical Research to M.G. and an NIH Ruth L.
Kirschstein NRSA (F32 GM131522) to E.M.H.. We are also grateful for the support of the CNRS
Marine Station in Roscoff and the Assemble grant 227799 to U.T. for collecting *Scolanthus*.

Data Availability

All raw data is available via the National Center for Biotechnology Information under the accession
PRJNA667495. The assembled genomes can be downloaded, browsed and searched on publicly
available   browsers   at   https://simrbase.stowers.org/starletseaanemone   and
https://simrbase.stowers.org/wormanemone. Code used to generate the analyses is available
from the authors upon request. Publicly available data that were used in this manuscript include
SRR1771322-SRR1771324 (mouse Hi-C), SRR10512944 (*D. melanogaster* Hi-C), and
ERR8571699 (Wellcome Sanger *N. vectensis* Hi-C).

Materials and Methods

Animal Care and Source

546 *Nematostella vectensis* animals were cultured as previously described[81] at the University of
547 Vienna and the Stowers Institute. Adult male and female individuals were verified by induction in
548 isolation. *Scolanthus callimorphus* animals were collected at the Île Callot, Carantec, France. After
549 transport, they were kept in seawater at 20°C and fed freshly hatched *Artemia salina* weekly or
550 biweekly.

Sequencing

Short Read DNA-Seq

Genomic DNA samples were extracted from both adult male and female individual *Nematostella*
adults using the DNeasy Blood and Tissue Kit (Qiagen). After purification, approximately 5 ug of
genomic DNA was recovered from each sample. Following DNA extraction, samples were
sheared and size selected for ~500 bp using a Blue Pippin Prep machine (Sage Science).
Following size selection, sequencing libraries were created using a KAPA HTP Library Prep kit
(Roche) and subjected to paired-end sequencing on an Illumina NextSeq 500. *Scolanthus* DNA
samples for library preparation were aliquoted from high molecular weight extractions, described
below.

High Molecular Weight DNA Extraction and Library Prep

Nematostella high molecular weight DNA was extracted at Dovetail Genomics. Samples were
quantified using Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, USA). The
PacBio SMRTbell library (~20kb) for PacBio Sequel was constructed using SMRTbell Template
Prep Kit 1.0 (PacBio, Menlo Park, CA, USA) using the manufacturer recommended protocol. The

pooled library was bound to polymerase using the Sequel Binding Kit 2.0 (PacBio) and loaded
onto PacBio Sequel using the MagBead Kit V2 (PacBio). Sequencing was performed on the
PacBio Sequel SMRT cell, using Instrument Control Software Version 5.0.0.6235, Primary
analysis software Version 5.0.0.6236 and SMRT Link Version 5.0.0.6792, yielding 24.67 Gb over
3,050,403 subreads.

High molecular weight DNA from a single *Scolanthus callimorphus* adult animal was extracted
using a modified Urea-based DNA extraction protocol[82,83]. A whole animal was flash frozen and
ground with mortar and pestle. While frozen, drops of buffer UEB1 (7M Urea, 312.5 mM NaCl, 50
mM Tris-HCl pH 8, 20 mM EDTA pH 8.1% w:v N-Lauroylsarcosine sodium salt) were added and
crushed with the tissue. Tissue was incubated in a final volume of 10 mL UEB1 at RT for 10
minutes. Three rounds of phenol-chloroform extraction were performed followed by DNA
precipitation by addition of 0.7 volume isopropanol. The pellet was transferred to a fresh tube and
washed twice in 70% EtOH and twice more in 100% EtOH, dried, and resuspended in TE buffer.

A library for PacBio sequencing was then prepared from the high molecular weight sample using
the SMRTbell® Express Template Prep Kit v1. The libraries were then sequenced on a PacBio
Sequel machine over 3 SMRT Cells, yielding a total of 22.85 Gb over 1,474,285 subreads. An
aliquot of the same sample was used to prepare a library using the NEBNext® Ultra™ II DNA
Library Prep Kit for Illumina. This was then subjected to 50 cycles of single-end sequencing in
one flow cell lane using an Illumina HiSeq 2500 system.

Chicago libraries
2 Chicago libraries were prepared as described previously[84]. For each library, ~500 ng of HMW
gDNA (mean fragment length = 100 kbp) was reconstituted into chromatin in vitro and fixed with
formaldehyde. Fixed chromatin was digested with DpnII, the 5' overhangs filled in with biotinylated
nucleotides, and then free blunt ends were ligated. After ligation, crosslinks were reversed and
the DNA purified from protein. Purified DNA was treated to remove biotin that was not internal to
ligated fragments. The DNA was then sheared to ~350 bp mean fragment size and sequencing
libraries were generated using NEBNext Ultra enzymes and Illumina-compatible adapters. Biotin-
containing fragments were isolated using streptavidin beads before PCR enrichment of each
library. The libraries were sequenced on an Illumina HiSeq 2500 (rapid run mode). The number
and length of read pairs produced for each library was: 116 million, 2x101 bp for library 1; 35
million, 2x101 bp for library 2. Together, these Chicago library reads provided 125 x sequence
coverage of the genome (1-100 kb pairs).

Chromatin was extracted from a single *Nematostella vectensis* adult male and *Scolanthus*
callimorphus adult (unknown sex) nuclei using the Phase Genomics Proximo Hi-C animal
protocol. After proximity ligation and purification, 16 ng and 9 ng of DNA was recovered,
respectively. For library preparation 1 μl of Library Reagent 1 was added 12 PCR cycles were
performed. The final library was subjected to 150 total cycles of paired-end sequencing using an
Illumina NextSeq 550 machine yielding a total of 13.5 gigabases.

Hi-C sequencing, *Scolanthus callimorphus* PacBio library preparation and sequencing,
Scolanthus Illumina DNA library preparation and sequencing and adult *Nematostella vectensis*
RNA library preparation and sequencing was performed at the VBCF NGS Unit
(https://www.viennabiocenter.org/facilities). *Nematostella vectensis* DNA size selection, library
preparation, and sequencing were performed by the Molecular Biology Core at the Stowers
Institute for Medical Research.
618
Developmental and adult *Nematostella* RNA sequencing was performed as follows. *Nematostella*
were spawned and eggs were de-jellied and fertilized as previously described[81]. Spawning and
embryo development took place at 18°C. Eggs and embryos from different stages were collected
(300 per sample) in duplicate as indicated: eggs (within 30min of spawn), blastula (7.5hpf),
gastrula (23.5hpf) and planula (72hpf). Eggs and embryos were collected in eppendorf tubes and
centrifuged to a pellet at 21,000 x g for 1 min. All seawater was quickly removed and pellets were
resuspended in 150ml lysis buffer (RLT buffer supplied by the Qiagen RNeasy kit (#74104),
supplemented with -mercaptoethanol). The samples were homogenized with an electric pestle
(1 min continuous drilling) and further supplemented with 200 ml of the above lysis buffer.
Homogenized samples were then transferred into QIAshredder columns (Qiagen #79654) and
centrifuged at 21,000 x g for 2 min. The flow throughs were supplemented with 1 ml 70 % ethanol
and transferred to RNeasy columns and were processed according to the Qiagen RNeasy
protocol. Quality and integrity of the RNA was evaluated using the Agilent RNA 600 pico kit
(Agilent Technologies) and RNA samples were stored at -80°C until further processing. cDNA
libraries were then constructed for polyA stranded sequencing. The resulting libraries were
sequenced on Illumina HiSeq using paired end runs (RapidSeq- 2x150bp).
635
636 **Genome Assembly**
637
638 Size estimates for *Nematostella vectensis* and *Scolanthus callimorphus* were derived using
639 Genomescope[85], taking the result of the highest *k* (56 and 18) which converged under the model.
640
641 Initial assemblies based on PacBio sequencing of *Nematostella* and *Scolanthus* were generated
642 using canu version 1.8[86] with the parameters `rawErrorRate=0.3`
643 `correctedErrorRate=0.045`.
644
Nematostella haplotigs were removed using Purge Haplotigs[87]. First, the source PacBio reads
were aligned onto the canu assembly using minimap2[88] using the parameters `-ax map-pb --`
`secondary=no`. Following this a coverage histogram was generated using the Purge Haplotigs
script `readhist`. Per the documented Purge Haplotigs protocol, lower, mid and high coverage
limits were found by manual inspection of the plotted histogram to be 12, 57 and 130 respectively.
All initial contigs marked as suspect or artifactual were removed from further analysis with the
Purge Haplotigs script `purge`.
652
653 Due to lower sequencing coverage of *Scolanthus*, diploid per-scaffold coverage could not be
654 deconvolved from haploid, and therefore Purge Haplotigs could not be used. Removal of
655 redundant contigs was performed with Redundans version 0.14a[89] using the parameters `--`

656   `noscaffolding --nogapclosing --overlap 0.66.` Only contigs marked in the reduced
657   version of the genome were used in further analysis.
658
The input de novo assembly, shotgun reads, and Chicago library reads were used as input data
for HiRise, a software pipeline designed specifically for using proximity ligation data to scaffold
genome assembly[84]. Shotgun and Chicago library sequences were aligned to the draft input
assembly using a modified SNAP read mapper (http://snap.cs.berkeley.edu). The separations of
Chicago read pairs mapped within draft scaffolds were analyzed by HiRise to produce a likelihood
model for genomic distance between read pairs, and the model was used to identify and
break putative misjoins, to score prospective joins, and make joins above a threshold. After
scaffolding, shotgun sequences were used to close gaps between contigs.
667
668   **Repetitive DNA and Hi-C scaffolding**
669
Repetitive DNA was found using two strategies. First, known repeats found in repbase[90] were
searched in the assemblies using RepeatMasker[91] using the parameters `-s -align -e ncbi`
in addition to `-species nematostella` for *Nematostella* and `-species edwardsiidae` for
Scolanthus. Second, novel repeat sequences were found using RepeatModeler version 2.0[92].
After generating the repeat library, genomes' repeat regions were detected with the corresponding
library using the same parameters in RepeatMasker.
676
Hi-C sequences were aligned to the reduced and repbase masked genomes of *Nematostella* and
Scolanthus using `bwa mem`[93,94] using the parameters `-5SP`. For *Nematostella,* an additional
candidate assembly was generated by mapping Hi-C sequences to the Chicago library scaffolded
sequences using repbase masking (dovetail_standardmask) in addition to the contig-based
scaffolding (contig_standardmask). Duplicate reads were marked with the `samblaster` utility[95],
and duplicate, secondary and supplementary mappings were removed with `samtools`. These
mappings were used to generate initial chromosomal assemblies using Lachesis[96], specifying the
restriction site GATC. Assemblies were manually reviewed using Juicebox Assembly Tools
version 1.11.08[97]. Candidate assemblies were compared using the nucmer aligner with default
parameters and visualized using mummerplot[98]. Assemblies were converted over to Juicebox
format using juicebox_scripts (https://github.com/phasegenomics/juicebox_scripts). In the case
of *Scolanthus*, duplicate regions were clipped, and the resulting contigs were subjected to another
round of alignment, assembly and review.
690
Nematostella scaffold correctness was assessed using REAPR[99]. *Nematostella* assembly
nemVec1 was downloaded from the JGI website[1]. Sequences from the adult male and adult
female (see Sequencing) were aligned to nemVec1 and the *Nematostella* genome after
scaffolding with Chicago libraries using SMALT as well as the REAPR tool perfectmap using an
expected insert size of 400, as determined from fragment analysis. Error-free bases and contiguity
after breaking the genome were extracted from the results.
697
698   Genome and gene model set assembly and completeness was assessed using BUSCO version
699   3.0.2[33], using the gene set `metazoa_odb9` as the standard.

Gene Models

Nematostella, Scolanthus and *M. senile* paired end sequences obtained from previous studies[34,100] and data available on BioProject PRJNA430035 were used to generate *de novo* assembled transcripts.

Trinity version 5.0.2[101] was run on each library using the flags `--min_contig_length 200 --min_kmer_cov 2`. For those which had a strand-specific library preparation, the flag `--SS_lib_type RF` was applied. To reduce redundancy, cd-hit version 4.6.8[102,103] was applied with the flags `-M 0 -c 1`. Transdecoder version 5.0.6[104] was used to detect open reading frames in the resulting reduced set of transcripts. Transcript abundance was quantified using salmon version 1.2.1[105] using the flags `--seqBias --useVBOpt --discardOrphansQuasi --softclip`.

For PacBio Iso-seq, 12 *Nematostella* RNA samples were collected over the course of multiple developmental stages, adult tissues and regeneration time points. For developmental stages, zygotes spawned by a single batch of wildtype colony were kept at 22°C, and collected at 0 hpf, 24 hpf, 48 hpf, 72 hpf and 7 dpf. Adult tissues were collected from sex-sorted, sexually mature wildtype individuals kept at 22°C. The male and female mesenteries were harvested separately by surgically opening the body column and carefully peeling off the attached body column tissues. Adult oral discs were collected by surgical removal of tentacles as well as the attached pharyngeal regions. Regeneration was induced by amputating the oral part of a sexually mature individual at the mid-pharyngeal level. Regenerating tissues close to the wound were collected at 4 hpa and 12 hpa, respectively. All the samples were deep frozen and lysed using TRIzolTM reagent (Invitrogen). Phenol-chloroform extraction was performed to remove undissolved mesoglea from adult tissues. DirectzolTM RNA Miniprep Plus Kit (Zymo) was then used to purify total RNA from the aqueous phase. For each sample, 2 μg of total RNA with RIN > 7 was submitted to UC Berkeley for Iso-seq library construction.

RNA Libraries were sequenced at UC Berkeley using PacBio Sequel-II system. Raw subreads bams were processed and demultiplexed using PacBio's isoseq v3.2 conda pipeline. The steps include consensus generation, primer demultiplexing, polyA refinement and data clustering using default parameters. This resulted in the generation of 406,317 high quality HIFI reads and used to build Nvec200 transcriptome.

HIFI reads were mapped to the *Nematostella* genome using minimap2[88] using parameters (`-ax splice -uf --secondary=no`) to obtain the primary best alignments. Reads were then grouped and collapsed down to potential transcripts using PacBio's cDNA_Cupcake toolkit and TAMA[106]. Based on PacBio's guideline, transcripts with degraded 5' reads and have less than 10 FL counts were removed. Chimeric transcripts were then analyzed to find potential fusion genes. For reads that didn't map to the genome, de novo transcriptome assembly was performed using graph-based tool Cogent with kmer size equals 30. . Cupcake and TAMA results were merged into non-redundant gene models using stringtie v2[107]. Deep RNA-seq reads from 4 developmental

stages: egg, gastrula, pos-gastrula and planula were aligned to the genome using STAR v
2.7.3a[108]. Read alignments outside of the isoseq gene models were extracted with bedtools
v2.29.2[109] and used for reference-guided transcriptome assembly using Stringtie2. Final gene
models were obtained by merging Isoseq models and RNAseq models and manually corrected
using previously cloned full length CDS from Nematostella vectensis in NCBI (EDT7). Finally,
transdecoder v5.5.0 was used to produce CDS annotation using a minimum protein length of 50
amino acids and prioritizing ORFs with significant similarity to any family in the PFAM database[110].
Alignment of the protein candidates to the PFAM database was done using Hmmer v3.1b2[111].
RNAseq libraries from the *N. vectensis* developmental time series were downloaded and aligned
to the new genome using STAR v2.7.3a[108] with standard parameters. Mapping and assignment
efficiency was measured using featureCounts from the subread package[112] with the "-p" flag for
paired-end libraries.
756
Evidence for *Scolanthus* gene models were taken from RNA-sequencing and repeats. *Scolanthus*
RNA-seq reads (see Sequencing) were mapped to the *Scolanthus* contigs using STAR version
2.7.3a[108]. These mappings were used as evidence for intron junctions to generate putative gene
models and estimating hidden Markov model parameters using BRAKER2[113,114]. Gene models
were then refined using Augustus version 3.3.3[115] using extrinsic evidence from STAR splice
junctions and the location of repeats from RepBase (see Genome Assembly) as counter-evidence
for transcription. These models were filtered with the following criteria: 1) genes completely
covered by RepeatModeler repeats (see Genome Assembly) were removed 2) predicted gene
models were required to be either supported by external RNA-seq evidence as reported by
Augustus or have a predicted ortholog as reported by Eggnog-mapper[116]. This resulted in a set
of 24,625 gene models. Transcription factor identity was inferred by aligning the predicted protein
sequences to Pfam A domains version 32.0[117] using hmmer version 3.3[111]. Transcription factor
families were based on domains curated in a previous work[118].
770
771   Extended Hox cluster, NK cluster and ParaHox genes were found with BLAT[119] matches of
772   published models[51,61,120–125] to the nv1 genome, taking the best hits. If an NVE gene model[34]
773   corresponded to the matched genomic region, its location in the nv2 genome was then determined
774   for macrosynteny analysis. In cases where no published gene was known, reciprocal BLAST hits
775   between the bilaterian and cnidarian counterpart were taken as evidence for orthology.
776
777   **Divergence Estimates**
778
Single copy orthologs were detected by collecting common complete and duplicated BUSCO
genes present in the *Scolanthus* and *Nematostella* genomes. Where duplicated BUSCOs were
present, the transcript with the highest score was taken. This resulted in a total of 541 orthologs.
BUSCOs found in genomes obtained from previous studies[2,21,22,26,27,32,41,44,47,126–129] were used to
generate multiple alignments. Genes were aligned with mafft version 7.427 using the E-INS-i
model and a maximum 1000 refinement iterations[130]. Alignments were trimmed using trimAl
version 1.4.rev15 using the "gappyout" criteria[131]. A maximum likelihood tree was inferred using
iqtree version 2.0.6, using the model finder partitioned on each gene, constrained to nuclear
protein models[132]. Divergence estimates were determined using r8s version 1.8.1 using the

Langley-Fitch likelihood method[133]. Age ranges were estimated by fixing the split between
Bilateria and Cnidaria at 595.7 and 688.3 Mya[134].
790
791 Single copy orthologs were detected by collecting common complete BUSCO genes present in
792 the *A. millepora*, *A. digitifera*, *E. pallida*, *M. senile*, *Scolanthus* and *Nematostella* genomes. This
793 resulted in a total of 229 orthologs.
794
Ultraconserved Elements
795
796
In order to determine noncoding elements conserved between *Scolanthus* and *Nematostella*,
genomes repeat-masked from both *de novo* and repbase repeats were blasted using NCBI
BLAST+ version 2.10.0[135], using the flags `-evalue 1E-10 -max_hsps 100000000 -`
`max_target_seqs 100000000 -task megablast -perc_identity 0 -`
`template_length 16 -penalty -2 -word_size 11 -template_type`
`coding_and_optimal`. Additionally, the `-dbsize` parameter was set to the estimated genome
size. Candidate hits were then filtered using criteria loosely based on previous work[136]: for each
high-scoring pair, a sliding window method was used to determine subsections of the alignment
with at least 95 % identity, and extending these windows as long as the identity remains at this
level. *Nematostella* elements mapping to more than one locus in the *Scolanthus* genome were
reduced to the longest locus pair in both genomes. Elements mostly mapping to coding sequence
were removed, and the remaining elements were classified as intron or non-coding, depending
on location. Recurring UCE sequences that were not identified by RepeatModeler or
RepeatMasker were detected with blastclust version 2.2.26 requiring the length of hit to cover at
least 90 % of either sequence for linkage.
812
Macrosynteny Analysis
813
814
815 *Branchiostoma floridae* gene models and sequences were retrieved from the published study[41].
816 All against all comparisons were performed with OMA standalone version 2.5[137].Genomes were
817 downloaded from previous studies[2,3,20,22,26,32,44–47,76,126–129,138–141].
818
Ancestral genome reconstruction was carried out using a graph based approach. In brief, genes
were summarized into multi-species orthologous groups, and these comprised the nodes, and
orthology groups occurring on the same chromosome or scaffold of two different species were
linked together. A consensus approach to community detection based on the Leiden algorithm[142]
was used to determine ancestral linkage groups from this graph. For details, see the
Supplementary Text.
825
TAD Sliding Window Analysis
826
827
Hi-C maps were generated for multiple species' genomes (g:) and reads (r:), including those of
the ctenophore *Hormiphora californensis* (g: GCA_020137815.1, r: SRR13784181,
SRR13784182)[143], the cnidarian *Hydra vulgaris* strain 105 (g: GCA_022113875.1, r:
SRR14099165)[40], the cnidarian *Haliclystus octoradiatus* (g: GCA_916610825.1, r:

ERR6745733)[144], the *Nematostella* assembly and reads presented in this manuscript (also, Hi-C
reads from the Darwin Tree of Life project, ERR8571699[145]), the cnidarian *Diadumene lineata* (g:
GCA_918843875.1, r: ERR6688655)[146], the fire jellyfish *Rhopilema esculentum* (g:
http://doi.org/10.5524/100720[39], r: SRR11649085[21]), the fly *Drosophila melanogaster* (g:
assembly Release 6[128], r: SRR10512944[147]), the scallop *Pecten maximus* (g:
http://dx.doi.org/10.6084/m9.figshare.10311068, r: SRR10119404)[148], the cephalochordate
Branchiostoma floridae (g: GCA_000003815.2 Bfl_VNyyK, r: SRR12007919, SRR12059951)[41],
and *Mus musculus* (g: GRCm39[149], r: SRR1771322-SRR1771324[150]).
840
The Hi-C maps were generated by mapping the raw paired-end Hi-C reads to the genome
assemblies using chromap v0.2.3[151]. The pairs files were normalized using hicExplorer v3.6[152],
and balanced using Cooler v0.8.10[153]. The insulation scores were then calculated using FanC v
0.9.23b[73] using 100kb bin sizes. The peaks and valleys in the insulation scores were also called
with FanC v0.9.24[154]. We defined the delta between each peak and valley as the transition from
a highly-interacting region to a poorly-interaction region as the strength of transitions of topological
boundaries. We compared the distributions of these deltas with an uncorrected Mann-Whitney U
two-sided test to test for significant differences in the median values.
849
850   **Phylogenetic analysis of NK-like and SuperHox Genes**
851
Sequences of the NK class proteins were MUSCLE-aligned with default settings in MEGA11[155],
and trimmed using the Automated1 setting in TrimAl (v. 1.3)[131]. Trimmed sequences were used
for calculating the NJ, and ML trees. The NJ tree was calculated in MEGA11[155], and the ML tree
was calculated in IQ-TREE2[132] using the automatic model selection algorithm. Sequences of the
SuperHox cluster genes were aligned as above but not trimmed. NJ and ML trees were calculated
as above.
Nematostella gastrula ATAC-seq
859
Embryos were raised at 21°C until gastrula stage (24 hpf). Ten embryos were collected and
washed with cold PBS, then cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2,
0.1 % NP-40, 0.1% Tween-20, 0.01% Digitonin) was added. Embryos were disintegrated by
pipetting and incubated in lysis buffer for 6.5 minutes on ice. Nuclei were then pelleted by
centrifugation (0.5 rcf, 10 min, 4°C). The pellet was resuspended in 50 µl tagmentation mix (5x
TD buffer, 0.1% Tween-20, 0.01% Digitonin, 2.5 µl Tn5 transposase). TD buffer and Tn5
transposase were kindly provided by the David Garfield lab. The transposition reaction was
incubated at 37°C for 30 min with agitation at 1000 rpm and then cooled down on ice for 5 minutes.
DNA purification was carried out with the QIAquick PCR kit (QIAGEN, #28104) according to the
manufacturer's instructions, with an elution in 20 µl elution buffer. PCR amplification was
performed for 12 cycles according to Buenrostro(2015). A final purification step was performed
with the QIAquick PCR kit (QIAGEN, #28104), elution in 15 µl EB. The quality of the library was
validated with an agarose gel, from which DNA was extracted with a peqlab gel extraction kit,
eluted in 20 µl EB, and stored at -20°C. Library preparation and sequencing were done at
Novogene and Vienna Biocenter Core Facility (VBCF).

Comparative ATACseq analysis

878 ATACseq peak calling was performed for the following species using publicly available datasets:

Species	SRA	Stage	Reference genome	Reference
Owenia fusiformis	SRR15898385	Blastula	GCA_90381334 5.2	[156]
Lytechinus variegatus	SRR18841698 SRR18841699 SRR18841700	Gastrula	GCF_01814301 5.1	[157]
Patinopecten yessoensis	SRR18361867 SRR18361868 SRR18361869 SRR18361870 SRR18361871 SRR18361873	Gastrula	GCF_00211388 5.1	[158]
Nematostella vectensis	SRR24055088	Gastrula	This study	This study

All peak callings were done following the Encode guideline for ATACseq. *Owenia* and
Nematostella peak calls were done following the unreplicated data guidelines
(https://www.encodeproject.org/pipelines/ENCPL344QWT/) while *Lytechinus* and *Petinopectin*
following the replicated data guidelines
(https://www.encodeproject.org/pipelines/ENCPL787FUN/). Briefly, raw reads were trimmed,
aligned to their reference genome and filtered for mapq >=20. Psudoreplicates were produced for
each library as well as for the merged alignment of multiple libraries and peak calling was done
on the initial alignment, the merged alignment and all the pseudoreplicates. For unreplicated data,
peaks were filtered based on p-signal <=0.05 and similar peak calls between the original
alignment and at least one pseudoreplicate. For replicated data, final peaks were selected based
on IDR scores between the merged library peaks and each replicate library.

ATACseq peaks for Human (hg19), mouse (mm10), Caenohabditis (Ce11) and Drosophila (dm6)
were downloaded from the ChIP-atlas database (https://chip-atlas.org/, [159,160]). Only embryonic
stage datasets were used in this study. Repetitive regions were identified using RepeatModeler
and RepeatMasker for all genomes and peaks overlapping repeat regions were removed for
downstream analysis. A custom R script was used to find the closest gene for each ATAC peak
and measure the distances (available in https://github.com/jdmontenegro/NematostellaGenome).

All peaks overlapping genes were not considered in downstream analysis. Data points 1.5 times the interquartile range above the third quartile or below the first quartile were removed as outliers before plotting and regression. A simple linear regression was used to correlate genome size and mean intergenic distance for all species.

Microsynteny Analysis

Orthofinder version 2.5.4 was run to obtain orthogroups for the selected species. The microsynteny pipeline from Simakov et al[58] was run for each of the three clades (deuterostomes, spiralians, cnidarians) separately. For this a subset of orthogroups that had at least one gene per species in a given clade (e.g., every cnidarian species had an ortholog for the cnidarian clade micro-synteny analysis) was generated, to correct for any missing genes. The micro-synteny analysis was thus based on 6016, 4832, and 6539 orthogroups for cnidarians, spiralians, and deuterostomes, respectively. For the most strict profiling of gene block length, we run the micro-synteny pipeline without allowance for any intervening genes. The resulting blocks were then filtered to remove micro-syntenies composed of paralogous genes. The resulting total numbers of micro-syntenic blocks given different minimal required length (at least 2, 3, or 4 genes) are listed in EDT12.

References

1. Putnam, N. H. *et al.* Sea Anemone Genome Reveals Ancestral Eumetazoan Gene Repertoire and Genomic Organization. *Science* **317**, 86–94 (2007).

2. Chapman, J. A. *et al.* The dynamic genome of Hydra. *Nature* **464**, 592–596 (2010).

3. Srivastava, M. *et al.* The Amphimedon queenslandica genome and the evolution of animal complexity. *Nature* **466**, 720–726 (2010).

4. Schwaiger, M. *et al.* Evolutionary conservation of the eumetazoan gene regulatory landscape. *Genome Res.* **24**, 639–650 (2014).

5. Gaiti, F. *et al.* Landscape of histone modifications in a sponge reveals the origin of animal cis-regulatory complexity. *Elife* **6**, e22194 (2017).

6. Nanty, L. *et al.* Comparative methylomics reveals gene-body H3K36me3 in Drosophila predicts DNA methylation and CpG landscapes in other invertebrates. *Genome Res.* **21**,

935   1841–1850 (2011).

7. Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* **326**, 289–293 (2009).

8. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376–380 (2012).

9. Acemel, R. D., Maeso, I. & Gómez Skarmeta, J. L. Topologically associated domains: a successful scaffold for the evolution of gene regulation in animals. *WIREs Developmental Biology* **6**, e265 (2017).

10. Anderson, E., Devenney, P. S., Hill, R. E. & Lettice, L. A. Mapping the Shh long-range regulatory domain. *Development* **141**, 3934–3943 (2014).

11. Krefting, J., Andrade-Navarro, M. A. & Ibn-Salem, J. Evolutionary stability of topologically associating domains is associated with conserved gene regulation. *BMC Biol.* **16**, 87 (2018).

12. Rodríguez-Carballo, E., Lopez-Delisle, L., Yakushiji-Kaminatsui, N., Ullate-Agote, A. & Duboule, D. Impact of genome architecture on the functional activation and repression of Hox regulatory landscapes. *BMC Biol.* **17**, 55 (2019).

13. Cartwright, P. *et al.* Exceptionally Preserved Jellyfishes from the Middle Cambrian. *PLoS One* **2**, e1121 (2007).

14. Technau, U., Genikhovich, G. & Kraus, J. E. M. Cnidaria. in *Evolutionary Developmental Biology of Invertebrates 1* (ed. Wanninger, A.) 115–163 (Springer Vienna, 2015).

15. Zapata, F. *et al.* Phylogenomic Analyses Support Traditional Relationships within Cnidaria. *PLoS One* **10**, e0139068 (2015).

16. Technau, U. & Steele, R. E. Evolutionary crossroads in developmental biology: Cnidaria. *Development* **138**, 1447–1458 (2011).

17. Röttinger, E. Nematostella vectensis, an Emerging Model for Deciphering the Molecular and Cellular Mechanisms Underlying Whole-Body Regeneration. *Cells* **10**, (2021).

18. Rentzsch, F. & Technau, U. Genomics and development of Nematostella vectensis and

other anthozoans. *Curr. Opin. Genet. Dev.* **39**, 63–70 (2016).

19. Layden, M. J., Rentzsch, F. & Röttinger, E. The rise of the starlet sea anemone

Nematostella vectensis as a model system to investigate development and regeneration:

Overview of starlet sea anemone *Nematostella vectensis. Wiley Interdiscip. Rev. Dev. Biol.*

5, 408–428 (2016).

20. Hu, M., Zheng, X., Fan, C.-M. & Zheng, Y. Lineage dynamics of the endosymbiotic cell type

in the soft coral Xenia. *Nature* **582**, 534–538 (2020).

21. Nong, W. *et al.* Jellyfish genomes reveal distinct homeobox gene clusters and conservation

of small RNA processing. *Nat. Commun.* **11**, 3051 (2020).

22. Leclère, L. *et al.* The genome of the jellyfish Clytia hemisphaerica and the evolution of the

cnidarian life-cycle. *Nature Ecology & Evolution* **3**, 801–810 (2019).

23. Khalturin, K. *et al.* Medusozoan genomes inform the evolution of the jellyfish body plan.

Nature Ecology & Evolution **3**, 811–822 (2019).

24. Ohdera, A. *et al.* Box, stalked, and upside-down? Draft genomes from diverse jellyfish

(Cnidaria, Acraspeda) lineages: Alatina alata (Cubozoa), Calvadosia cruxmelitensis

(Staurozoa), and Cassiopea xamachana (Scyphozoa). *Gigascience* **8**, (2019).

25. Kim, H.-M. *et al.* The genome of the giant Nomura's jellyfish sheds light on the early

evolution of active predation. *BMC Biol.* **17**, 28 (2019).

26. Gold, D. A. *et al.* The genome of the jellyfish Aurelia and the evolution of animal complexity.

Nature Ecology & Evolution **3**, 96 (2019).

27. Shinzato, C. *et al.* Using the *Acropora digitifera* genome to understand coral responses to

environmental change. *Nature* **476**, 320–323 (2011).

28. Wilson, T. J., Lazner, F., Kola, I. & Hertzog, P. J. The Mouse and the Genomic Era. in

Comparative Genomics (ed. Clark, M. S.) 97–121 (Springer US, 2000).

29. Gosse, P. Scolanthus callimorphus. in *World List of Actiniaria,*.

30. Wilson, E. Scolanthus callimorphus Worm anemone. in *Marine Life Information Network:*

Biology and Sensitivity Key Information Reviews, [on-line] (2005).

31. Heger, P., Marin, B., Bartkuhn, M., Schierenberg, E. & Wiehe, T. The chromatin insulator

CTCF and the emergence of metazoan diversity. *Proceedings of the National Academy of*

Sciences **109**, 17507–17512 (2012).

32. Baumgarten, S. *et al.* The genome of Aiptasia, a sea anemone model for coral symbiosis.

Proc. Natl. Acad. Sci. U. S. A. **112**, 11893–11898 (2015).

33. Seppey, M., Manni, M. & Zdobnov, E. M. BUSCO: Assessing Genome Assembly and

Annotation Completeness. in *Gene Prediction: Methods and Protocols* (ed. Kollmar, M.)

227–245 (Springer, 2019).

34. Fredman, D., Schwaiger, M., Rentzsch, F. & Technau, U. Nematostella vectensis

transcriptome and gene models v2.0. Preprint at

https://figshare.com/articles/Nematostella_vectensis_transcriptome_and_gene_models_v2

_0/807696 (2013).

35. Cole, A. G. *et al.* Muscle cell-type diversification is driven by bHLH transcription factor

expansion and extensive effector gene duplications. *Nat. Commun.* **14**, 1747 (2023).

36. Steger, J. *et al.* Single-cell transcriptomics identifies conserved regulators of neuroglandular

lineages. *Cell Rep.* **40**, 111370 (2022).

37. Guo, L. *et al.* An adaptable chromosome preparation methodology for use in invertebrate

research organisms. *BMC Biol.* **16**, 25 (2018).

38. Zacharias, H., Anokhin, B., Khalturin, K. & Bosch, T. C. G. Genome sizes and

chromosomes in the basal metazoan Hydra. *Zoology* **107**, 219–227 (2004).

39. Li, Y. *et al.* Chromosome-level reference genome of the jellyfish Rhopilema esculentum.

Gigascience **9**, (2020).

40. Simakov, O. *et al.* Deeply conserved synteny and the evolution of metazoan chromosomes.

Science Advances **8**, eabi5884 (2022).

41. Simakov, O. *et al.* Deeply conserved synteny resolves early events in vertebrate evolution.

Nature Ecology & Evolution **4**, 820–830 (2020).

42. Dehal, P. & Boore, J. L. Two Rounds of Whole Genome Duplication in the Ancestral

Vertebrate. *PLoS Biol.* **3**, e314 (2005).

43. Pollard, S. L. & Holland, P. W. H. Evidence for 14 homeobox gene clusters in human

genome ancestry. *Curr. Biol.* **10**, 1059–1062 (2000).

44. Braasch, I. *et al.* The spotted gar genome illuminates vertebrate evolution and facilitates

human-teleost comparisons. *Nat. Genet.* **48**, 427–437 (2016).

45. Wang, S. *et al.* Scallop genome provides insights into evolution of bilaterian karyotype and

development. *Nature Ecology & Evolution* **1**, 1–12 (2017).

46. Qu, Z. *et al.* Millipede genomes reveal unique adaptations during myriapod evolution. *PLoS*

Biol. **18**, e3000636 (2020).

47. Kenny, N. J. *et al.* Tracing animal genomic evolution with the chromosomal-level assembly

of the freshwater sponge Ephydatia muelleri. *Nature Communications* **11**, 3676 (2020).

48. Kapli, P. & Telford, M. J. Topology-dependent asymmetry in systematic errors affects

phylogenetic placement of Ctenophora and Xenacoelomorpha. *Science Advances* **6**,

eabc5162 (2020).

49. Butts, T., Holland, P. W. H. & Ferrier, D. E. K. The Urbilaterian Super-Hox cluster. *Trends*

Genet. **24**, 259–262 (2008).

50. Ferrier, D. E. K. Evolution of Homeobox Gene Clusters in Animals: The Giga-Cluster and

Primary vs. Secondary Clustering. *Frontiers in Ecology and Evolution* **4**, (2016).

51. Chourrout, D. *et al.* Minimal ProtoHox cluster inferred from bilaterian and cnidarian Hox

complements. *Nature* **442**, 684–687 (2006).

52. Simakov, O. *et al.* Hemichordate genomes and deuterostome origins. *Nature* **527**, 459–465

(2015).

53. Larroux, C. *et al.* The NK Homeobox Gene Cluster Predates the Origin of Hox Genes. *Curr.*

Biol. **17**, 706–710 (2007).

54. Friedman, N. & Rando, O. J. Epigenomics and the structure of the living genome. *Genome*

Res. **25**, 1482–1490 (2015).

55. Fletcher, C. *et al.* The genome sequence of the starlet sea anemone, Nematostella

vectensis (Stephenson, 1935). *Wellcome Open Res.* **8**, 79 (2023).

56. Rodríguez-Carballo, E. *et al.* The HoxD cluster is a dynamic and resilient TAD boundary

controlling the segregation of antagonistic regulatory landscapes. *Genes Dev.* **31**, 2264–

2281 (2017).

57. Cazet, J. F. *et al.* A chromosome-scale epigenetic map of the *Hydra* genome reveals

conserved regulators of cell state. *Genome Res.* gr.277040.122 (2023).

58. Simakov, O. *et al.* Insights into bilaterian evolution from three spiralian genomes. *Nature*

493, 526–531 (2013).

59. He, S. *et al.* An axial Hox code controls tissue segmentation and body patterning in

Nematostella vectensis. *Science* **361**, 1377–1380 (2018).

60. McGinnis, W. & Krumlauf, R. Homeobox genes and axial patterning. *Cell* **68**, 283–302

(1992).

61. Ryan, J. F. *et al.* Pre-Bilaterian Origins of the Hox Cluster and the Hox Code: Evidence

from the Sea Anemone, Nematostella vectensis. *PLoS One* **2**, e153 (2007).

62. Finnerty, J. R., Pang, K., Burton, P., Paulson, D. & Martindale, M. Q. Origins of Bilateral

Symmetry: Hox and Dpp Expression in a Sea Anemone. *Science* **304**, 1335–1337 (2004).

63. Lazar, N. H. *et al.* Epigenetic maintenance of topological domains in the highly rearranged

gibbon genome. *Genome Res.* **28**, 983–997 (2018).

64. Rowley, M. J. & Corces, V. G. Organizational principles of 3D genome architecture. *Nat.*

Rev. Genet. **19**, 789–800 (2018).

65. Szabo, Q., Bantignies, F. & Cavalli, G. Principles of genome folding into topologically

associating domains. *Science Advances* **5**, eaaw1668 (2019).

66. Sacerdot, C., Louis, A., Bon, C., Berthelot, C. & Roest Crollius, H. Chromosome evolution

at the origin of the ancestral vertebrate genome. *Genome Biol.* **19**, 166 (2018).

67. Clarence, T. *et al.* Robust 3D modeling reveals spatiosyntenic properties of animal

genomes. *iScience* **26**, 106136 (2023).

68. Sinigaglia, C., Busengdal, H., Leclère, L., Technau, U. & Rentzsch, F. The bilaterian head

patterning gene six3/6 controls aboral domain development in a cnidarian. *PLoS Biol.* **11**,

e1001488 (2013).

69. Schwaiger, M. *et al.* An ancestral Wnt-Brachyury feedback loop in axial patterning and

recruitment of mesoderm-determining target genes. *Nat Ecol Evol* **6**, 1921–1939 (2022).

70. Kusserow, A. *et al.* Unexpected complexity of the Wnt gene family in a sea anemone.

Nature **433**, 156–160 (2005).

71. Lengfeld, T. *et al.* Multiple Wnts are involved in Hydra organizer formation and

regeneration. *Dev. Biol.* **330**, 186–199 (2009).

72. Marlétaz, F. *et al.* The little skate genome and the evolutionary emergence of wing-like fin

appendages. *bioRxiv* 2022.03.21.485123 (2022) doi:10.1101/2022.03.21.485123.

73. Crane, E. *et al.* Condensin-driven remodelling of X chromosome topology during dosage

compensation. *Nature* **523**, 240–244 (2015).

74. Kim, J. *et al.* Condensin DC loads and spreads from recruitment sites to create loop-

anchored TADs in C. elegans. *Elife* **11**, (2022).

75. Voolstra, C. R. *et al.* Comparative analysis of the genomes of Stylophora pistillata and

Acropora digitifera provides evidence for extensive differences between species of corals.

Sci. Rep. **7**, 1–14 (2017).

76. Moya, A. *et al.* Whole transcriptome analysis of the coral Acropora millepora reveals

complex responses to CO -driven acidification during the initiation of calcification. *Mol.*

Ecol. **21**, 2440–2454 (2012).

77. Cunning, R., Bay, R. A., Gillette, P., Baker, A. C. & Traylor-Knowles, N. Comparative

1091  analysis of the Pocillopora damicornis genome highlights role of immune system in coral

1092  evolution. *Sci. Rep.* **8**, 16134 (2018).

1093 78. Jiang, J. B. *et al.* A hybrid de novo assembly of the sea pansy (Renilla muelleri) genome.

1094  *Gigascience* **8**, (2019).

1095 79. Jeon, Y. *et al.* The Draft Genome of an Octocoral, Dendronephthya gigantea. *Genome Biol.*

1096  *Evol.* **11**, 949–953 (2019).

1097 80. Marlétaz, F. *et al.* Amphioxus functional genomics and the origins of vertebrate gene

1098  regulation. *Nature* **564**, 64–70 (2018).

1099 81. Fritzenwanker, J. H. & Technau, U. Induction of gametogenesis in the basal cnidarian

1100  Nematostella vectensis (Anthozoa). *Dev. Genes Evol.* **212**, 99–103 (2002).

1101 82. Chen, J. & Dellaporta, S. Urea-based Plant DNA Miniprep. in *The Maize Handbook* (eds.

1102  Freeling, M. & Walbot, V.) 526–527 (Springer, 1994).

1103 83. Sanders, S. M. *et al.* CRISPR/Cas9-mediated gene knockin in the hydroid Hydractinia

1104  symbiolongicarpus. *BMC Genomics* **19**, 649 (2018).

1105 84. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an in vitro method for

1106  long-range linkage. *Genome Res.* **26**, 342–350 (2016).

1107 85. Vurture, G. W. *et al.* GenomeScope: fast reference-free genome profiling from short reads.

1108  *Bioinformatics* **33**, 2202–2204 (2017).

1109 86. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive *k* -mer

1110  weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

1111 87. Roach, M. J., Schmidt, S. A. & Borneman, A. R. Purge Haplotigs: allelic contig

1112  reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460 (2018).

1113 88. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

1114  3100 (2018).

1115 89. Pryszcz, L. P. & Gabaldón, T. Redundans: an assembly pipeline for highly heterozygous

1116  genomes. *Nucleic Acids Res.* **44**, e113–e113 (2016).

90. Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive elements in

eukaryotic genomes. *Mob. DNA* **6**, 11 (2015).

91. Smit, A., Hubley, R. & Green, P. *RepeatMasker Open-4.0.* (2013).

92. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable

element families. *Proceedings of the National Academy of Sciences* **117**, 9451–9457

(2020).

93. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997 [q-bio] (2013).

94. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

95. Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural variant read

extraction. *Bioinformatics* **30**, 2503–2505 (2014).

96. Burton, J. N. *et al.* Chromosome-scale scaffolding of de novo genome assemblies based on

chromatin interactions. *Nat. Biotechnol.* **31**, 1119–1125 (2013).

97. Durand, N. C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps with

Unlimited Zoom. *Cell Systems* **3**, 99–101 (2016).

98. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol.* 9

(2004).

99. Hunt, M. *et al.* REAPR: a universal tool for genome assembly evaluation. *Genome Biol.* **14**,

R47 (2013).

100.Praher, D. *et al.* Conservation and turnover of miRNAs and their highly complementary

targets in early branching animals. *Proceedings of the Royal Society B: Biological Sciences*

288, 20203169 (2021).

101.Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

102.Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-

1143 generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

1144 103. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein

1145 or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

1146 104. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-Seq: reference

1147 generation and analysis with Trinity. *Nat. Protoc.* **8**, (2013).

1148 105. Srivastava, A. *et al.* Alignment and mapping methodology influence transcript abundance

1149 estimation. *bioRxiv* 657874 (2019).

1150 106. Kuo, R. I. *et al.* Illuminating the dark side of the human transcriptome with long read

1151 transcript sequencing. *BMC Genomics* **21**, 751 (2020).

1152 107. Kovaka, S. *et al.* Transcriptome assembly from long-read RNA-seq alignments with

1153 StringTie2. *Genome Biol.* **20**, 278 (2019).

1154 108. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

1155 109. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

1156 features. *Bioinformatics* **26**, 841–842 (2010).

1157 110. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–

1158 D419 (2021).

1159 111. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

1160 112. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for

1161 assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

1162 113. Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:

1163 Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.

1164 *Bioinformatics* **32**, 767–769 (2016).

1165 114. Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: Incorporating Protein

1166 Homology Information into Gene Prediction with GeneMark-EP and AUGUSTUS. 1 (2018).

1167 115. Stanke, M., Schöffmann, O., Morgenstern, B. & Waack, S. Gene prediction in eukaryotes

1168 with a generalized hidden Markov model that uses hints from external sources. *BMC*

Bioinformatics **7**, 62 (2006).

116. Huerta-Cepas, J. *et al.* Fast Genome-Wide Functional Annotation through Orthology

Assignment by eggNOG-Mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

117. El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic Acids Res.* **47**,

D427–D432 (2019).

118. de Mendoza, A. *et al.* Transcription factor evolution in eukaryotes and the assembly of the

regulatory toolkit in multicellular lineages. *Proceedings of the National Academy of*

Sciences **110**, E4858–E4866 (2013).

119. Kent, W. J. BLAT—The BLAST-Like Alignment Tool. *Genome Res.* **12**, 656–664 (2002).

120. Matus, D. Q., Pang, K., Daly, M. & Martindale, M. Q. Expression of Pax gene family

members in the anthozoan cnidarian, Nematostella vectensis: Pax gene expression in

Nematostella vectensis. *Evol. Dev.* **9**, 25–38 (2007).

121. Ryan, J. F. *et al.* The cnidarian-bilaterian ancestor possessed at least 56 homeoboxes:

evidence from the starlet sea anemone, Nematostella vectensis. *Genome Biol.* **7**, R64

(2006).

122. Mazza, M. E., Pang, K., Martindale, M. Q. & Finnerty, J. R. Genomic organization, gene

structure, and developmental expression of three Clustered otx genes in the sea anemone

Nematostella vectensis. *J. Exp. Zool. B Mol. Dev. Evol.* **308B**, 494–506 (2007).

123. Matus, D. Q., Thomsen, G. H. & Martindale, M. Q. Dorso/Ventral Genes Are

Asymmetrically Expressed and Involved in Germ-Layer Demarcation during Cnidarian

Gastrulation. *Curr. Biol.* **16**, 499–505 (2006).

124. Hudry, B. *et al.* Molecular insights into the origin of the Hox-TALE patterning system. *Elife*

3, e01939 (2014).

125. Mazza, M. E., Pang, K., Reitzel, A. M., Martindale, M. Q. & Finnerty, J. R. A conserved

cluster of three PRD-class homeobox genes (homeobrain, rx and orthopedia) in the

Cnidaria and Protostomia. *Evodevo* **1**, 3 (2010).

126. Ying, H. *et al.* The Whole-Genome Sequence of the Coral Acropora millepora. *Genome*

Biol. Evol. **11**, 1374–1379 (2019).

127. Consortium*, The C. elegans Sequencing. Genome Sequence of the Nematode C.

elegans: A Platform for Investigating Biology. *Science* **282**, 2012–2018 (1998).

128. Hoskins, R. A. *et al.* The Release 6 reference sequence of the Drosophila melanogaster

genome. *Genome Res.* **25**, 445–458 (2015).

129. Hydra 2.0 Web Portal. https://research.nhgri.nih.gov/hydra.

130. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

131. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

132. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

133. Sanderson, M. J. r8s: inferring absolute rates of molecular evolution and divergence times

in the absence of a molecular clock. *Bioinformatics* **19**, 301–302 (2003).

134. dos Reis, M. *et al.* Uncertainty in the Timing of Origin of Animals and the Limits of Precision

in Molecular Timescales. *Curr. Biol.* **25**, 2939–2950 (2015).

135. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

136. Dimitrieva, S. & Bucher, P. UCNEbase—a database of ultraconserved non-coding

elements and genomic regulatory blocks. *Nucleic Acids Res.* **41**, D101–D109 (2013).

137. Roth, A. C. J., Gonnet, G. H. & Dessimoz, C. Algorithm of OMA for large-scale orthology

inference. *BMC Bioinformatics* **9**, 518 (2008).

138. Zakas, C., Harry, N. D., Scholl, E. H. & Rockman, M. V. The genome of the poecilogonous

annelid Streblospio benedicti. *bioRxiv* 2021.04.15.440069 (2021).

139. Shingate, P. *et al.* Chromosome-level assembly of the horseshoe crab genome provides

insights into its genome evolution. *Nat. Commun.* **11**, 2322 (2020).

140. Harris, T. W. *et al.* WormBase: a modern Model Organism Information Resource. *Nucleic*

Acids Res. **48**, D762–D767 (2020).

141. Srivastava, M. *et al.* The *Trichoplax* genome and the nature of placozoans. *Nature* **454**,

955–960 (2008).

142. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-

connected communities. *Sci. Rep.* **9**, 5233 (2019).

143. Schultz, D. T. *et al.* A chromosome-scale genome assembly and karyotype of the

ctenophore *Hormiphora californensis. G3 (Bethesda)* **11**, jkab302 (2021).

144. Blaxter, M. *et al.* The genome sequence of the spotted kaleidoscope jellyfish, *Haliclystus*

octoradiatus (James-Clark, 1863). *Wellcome Open Res.* **8**, 6 (2023).

145. Fletcher, C., da Conceicoa, L. P., of Life Consortium, D. T. & Others. The genome

sequence of the starlet sea anemone, *Nematostella vectensis* (Stephenson, 1935).

Wellcome Open Research **8**, 79 (2023).

146. Wood, C. *et al.* The genome sequence of the orange-striped anemone, *Diadumene lineata*

(Verrill, 1869). *Wellcome Open Res.* **7**, 93 (2022).

147. Adams, M. *et al.* One fly-one genome: chromosome-scale genome assembly of a single

outbred *Drosophila melanogaster. Nucleic Acids Res.* **48**, e75 (2020).

148. Kenny, N. J. *et al.* The gene-rich genome of the scallop *Pecten maximus. Gigascience* **9**,

(2020).

149. Mouse Genome Sequencing Consortium *et al.* Initial sequencing and comparative analysis

of the mouse genome. *Nature* **420**, 520–562 (2002).

150. Vietri Rudan, M. *et al.* Comparative Hi-C reveals that CTCF underlies evolution of

chromosomal domain architecture. *Cell Rep.* **10**, 1297–1309 (2015).

151. Zhang, H. *et al.* Fast alignment and preprocessing of chromatin profiles with Chromap. *Nat.*

Commun. **12**, 1–6 (2021).

152. Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome

organization in flies. *Nat. Commun.* **9**, 189 (2018).

153. Abdennur, N. & Mirny, L. A. Cooler: scalable storage for Hi-C data and other genomically

labeled arrays. *Bioinformatics* **36**, 311–316 (2020).

154. Kruse, K., Hug, C. B. & Vaquerizas, J. M. FAN-C: a feature-rich framework for the analysis

and visualisation of chromosome conformation capture data. *Genome Biol.* **21**, 303 (2020).

155. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis

Version 11. *Mol. Biol. Evol.* **38**, 3022–3027 (2021).

156. Seudre, O., Carrillo-Baltodano, A. M., Liang, Y. & Martín-Durán, J. M. ERK1/2 is an

ancestral organising signal in spiral cleavage. *Nat. Commun.* **13**, 2286 (2022).

157. Davidson, P. L. *et al.* Recent reconfiguration of an ancient developmental gene regulatory

network in Heliocidaris sea urchins. *Nat Ecol Evol* **6**, 1907–1920 (2022).

158. Han, W. *et al.* Ancient homomorphy of molluscan sex chromosomes sustained by

reversible sex-biased genes and sex determiner translocation. *Nat Ecol Evol* **6**, 1891–1906

(2022).

159. Zou, Z., Ohta, T., Miura, F. & Oki, S. ChIP-Atlas 2021 update: a data-mining suite for

exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-seq

data. *Nucleic Acids Res.* **50**, W175–82 (2022).

160. Oki, S. *et al.* ChIP-Atlas: a data-mining suite powered by full integration of public ChIP-seq

data. *EMBO Rep.* **19**, (2018).

1268

a

b

c

Actiniaria
Nematostella vectensis
Scolanthus callimorphus
Exaiptasia pallida

Hexacorallia

Acropora digitifera
Acropora millepora
Scleractinia
Pocillopora damicornis
Stylophora pistillata

Anthozoa

Xenia sp.
Octocorallia
Dendronephthya gigantea
Renilla muelleri

Rhopilema esculentum
Nemopilema nomurai
Scyphozoa
Aurelia aurita
Sanderia malayensis

Medusozoa
Hydrozoa
Clytia hemisphaerica
Hydra vulgaris

Evidence

| Assembly | Flow Cytometry |
| FeulgenStaining | K–mer |

Genome Size (MB)
0   200   400   600   800

d *N.vectensis*

log10 Norm. Contact Intensity
0   1   2   3

chr2 chr10 chr5 chr12 chr8 chr3 chr1 chr14 chr4 chr7 chr9 chr6 chr13 chr11 chr15

e *S.callimorphus*

log10 Norm. Contact Intensity
0   1   2

chr1 chr8 chr4 chr3 chr6 chr5 chr9 chr7 chr11 chr15 chr12 chr13 chr14 chr2 chr10