Haploid mouse germ cell precursors from embryonic stem cells reveal *Xist* activation from a single X chromosome

Eishi Aizawa[1], Corinne Kaufmann[1], Sarah Sting[1], Remo Freimann[1], Anton Wutz[1,*]

[1]Institute of Molecular Health Sciences, Swiss Federal Institute of Technology, ETH Zurich, Zurich, Switzerland

*Correspondence: awutz@ethz.ch

Summary

Mammalian haploid cells have applications for genetic screening and substituting gametic genomes. Here we characterize a culture system for obtaining haploid primordial germ cell-like cells (PCGLCs) from haploid mouse embryonic stem cells (ESCs). We find that a haploid genome is maintained in PGCLCs with a high frequency indicating a substantially lower rate of diploidization than somatic cells. Characterization of the differentiating haploid ESCs reveals that *Xist* is activated from the single X chromosome. This observation suggests that X chromosome inactivation is initiated in haploid cells consistent with a model where autosomal blocking factors set a threshold for X-linked activators. The germline segregates from the epiblast and differs from somatic lineages in gene expression and epigenetic mechanisms. The ability of primordial germ cells for repressing *Xist* might contribute to the maintenance of a haploid genome.

Introduction

In mice, the germline derives from cells of the proximal epiblast and primordial germ cells (PGCs) segregate to an extraembryonic location adjacent to the posterior end of the embryo. PGC differentiation has been recapitulated in cultures of mouse epiblast cells (Ohinata et al., 2009). Furthermore, functional gametes have been obtained from primordial germ cell-like cells (PGCLCs) derived from embryonic stem cells (ESCs) in mice (Hayashi et al., 2012; Hayashi et al., 2011). Progress in culture techniques is opening opportunities for studies of the mammalian germline (Hayashi and Saitou, 2013). Germ cells differ from the somatic lineages in several key aspects including gene regulation and epigenetic modification (Hill et al., 2018; Leitch et al., 2013). During migration to the genital ridges, erasure of genomic imprinting occurs in PGCs, and the inactive X chromosome (Xi) is reactivated in female PGCs (Sugimoto and Abe, 2007).

Mammalian dosage compensation is facilitated by inactivation of one of the two X chromosomes in female cells (Lyon, 1961). Therefore, a single transcriptionally active X chromosome (Xa) faces two sets of autosomes in the diploid genome. In mice, X chromosome inactivation (XCI) is initiated by the long noncoding *Xist* RNA, which is expressed from, and accumulates over the inactive X chromosome (Xi) before X-linked gene repression (Galupa and Heard, 2018). *Xist* is first expressed in four-cell stage blastomers and initiates imprinted XCI of the paternally inherited X chromosome (Okamoto et al., 2004; Takagi and Sasaki, 1975). Cells that are destined to form the embryo reactivate the Xi at the blastocyst stage. In the developing female epiblast two Xa are present before embryonic day (E) 5.5, when random XCI is initiated (Mak et al., 2004). Thereafter, the Xi is maintained in all somatic lineages, but in the female germline Xi reactivation is observed.

From E7.0 two Xa are present in PGCs (Ohhata and Wutz, 2013; Sugimoto and Abe, 2007).

These observations show the dynamics of the dosage compensation system.

The regulation of XCI remains to be fully understood. A number of observations

suggest that the X to autosome (X:A) ratio controls the initiation of XCI and *Xist* expression.

One model posits that autosomal blocking factors prevent *Xist* activation for explaining the

observation that a single X chromosome is insufficient for initiation of XCI in male cells

(Barakat et al., 2014; Kay et al., 1993; Pollex and Heard, 2019). In female cells, twice the

number of X-linked activators overcomes an activation threshold for *Xist* leading to the

initiation of XCI. Diffusible X-linked activators and autosomal blocking factors are beginning

to be understood (Barakat et al., 2014; Pollex and Heard, 2019). An alternative model is

based on the observation of paring of the X chromosomes at the X chromosome

inactivation center (*Xic*) (Bacher et al., 2006; Xu et al., 2007; Xu et al., 2006). The *Xic*

encompasses the *Xist* gene and other regulators of XCI including the antisense *Tsix*

transcript. Genetic elements that are required for *Xic* pairing have been identified and

shown to induce XCI when transgenically integrated into autosomes in male ESCs (Augui

et al., 2007). Furthermore, stochastic regulation of *Xist* has been proposed from studies of

tetraploid ESCs, where a variable number of X chromosomes displayed activation of *Xist*

upon entry into differentiation (Monkhorst et al., 2008). Investigating the X counting

mechanism in the context of different autosomal dosage has led to further understanding

of the underlying regulation.

The mechanism of XCI has been extensively studied in ESCs, as these recapitulate

the relevant developmental stages that would be difficult to access in the post-implantation

embryo. In mice, female ESCs initiate XCI at the entry into differentiation. With the

establishment of haploid ESCs (haESCs) the effect of ploidy on regulation of XCI can be

further studied. HaESCs are established from *in vitro* cultured haploid embryos (Elling et

al., 2011; Leeb and Wutz, 2011). HaESCs possess a haploid genome but also show a

tendency towards diploidization, which is strongly enhanced when haESCs enter

differentiation. Diploidization has, thus, far made studies of *Xist* activation difficult to

interpret as haploid cells are in the process of diploidization. Several factors that affect

diploidization have been extensively investigated in somatic lineage differentiation with the

aim to reduce the high rates of diploidization (Freimann and Wutz, 2017; He et al., 2018;

He et al., 2017; Olbrich et al., 2017; Olbrich et al., 2019; Takahashi et al., 2014). A previous

study has reported the germline competence of haESCs *in vivo* by blastocyst injection

(Leeb et al., 2012), but the ploidy of haESCs and their characterization during germ cell

differentiation have been less well studied.

Here, we report the successful differentiation of haESCs into haploid PGCLCs *in vitro*.

We show that a haploid genome is preferentially maintained in PGCLCs compared with

somatic cell lineages. We then use this system to investigate *Xist* activation in haploid cells.

Our data demonstrate that a single X chromosome is sufficient for *Xist* activation in a

haploid genome consistent with a lower threshold of autosomal blocking factors.

Results

Differentiation of haESCs into haploid PGCLCs

To investigate the feasibility of germ cell differentiation from haESCs we followed a

previously established protocol (Hayashi et al., 2012; Hayashi et al., 2011). A mixed

population of haploid and diploid ESCs, containing 16.8% cells with a 1n DNA content

(G0/G1/S-phase haploid), were differentiated to epiblast-like cells (EpiLCs) for 2 days in

the presence of FGF2 and Activin A (Figure 1A, B). Subsequently, we aggregated the

EpiLCs for embryoid body (EB) formation. PGCLCs appeared between day 6 and 8 and

were identified by co-expression of the SSEA1 and integrin β3 surface markers. On day 7,

11.1% of the cells expressed both PGC markers (Figure 1C, D). Analysis of the ploidy

distribution by flow cytometry after DNA staining with Hoechst 33342 showed that 9.3% of

all cells possessed a 1n DNA content corresponding to haploid cells in G0-, G1- and S-

phase, which is approximately half the fraction of haploid ESCs at the beginning of

differentiation (16.8%; Figure 1E, F). To confirm a haploid karyotype in PGCLCs, we sorted

cells expressing SSEA1 and integrin β3 from d7 EBs and prepared chromosome spreads.

As expected, a set of 20 acrocentric chromosomes typical for an intact haploid mouse

genome could be observed (Figure 1G). Transcription analysis of sorted PGCLCs revealed

the upregulation of the PGC-markers *Blimp1*, *Prdm14*, *Tfap2c* and *Stella*, and the

downregulation of *Dnmt3b* in G0/G1/S-phase haploid (1n) as well as in S/G2/M-phase

diploid (4n) PGCLCs relative to EpiLCs and ESCs (Figure 2). These results show that

differentiation of haploid PGCLCs *in vitro* recapitulates gene expression changes that are

anticipated from the development of PGCs *in vivo*.

PGCLCs preferentially maintain a haploid genome

The PGCLC population of d7 EBs appeared to have a notably high content of haploid cells

(Figure 3A, B). We therefore further charcteized the percentage of PGCLCs in windows of

different DNA content. We analyzed the 1n population (G0/G1/S-phase haploid), the 2n

population (G2/M-phase haploid and G0/G1-phase diploid), and 4n population (S/G2/M-

phase diploid). In addition, we also analyzed the combined 2n and 4n population containing

all but the 1n haploid cells. PGCLCs accounted for 32.3% of 1n (G0/G1/S-phase) haploid

cells, while 11.1% of all cells and less than 10% of cell population with higher DNA content

expressed both PGC markers (Figure 3C). Therefore, the 1n haploid population contained

a 3-fold higher percentage of PGCLCs. The apparent enrichment of haploid cells within the

PGCLC population can in part be attributed to high diploidization rates that accompany

differentiation into somatic lineages. Nevertheless, a substantially higher percentage of

haploid cells was observed in PGCLCs (27.5%) compared to overall population of ESCs

(16.8%), EpiLCs (14.3%) and d7 EBs (9.3%; Figure 3D). These results indicate that

PGCLCs preferentially maintain a haploid genome.

Xist is activated from a single X chromosome in a haploid genome

Germline differentiation facilitates an investigation of *Xist* expression in haploid cells

without caveats that arise from diploidization or cell death. We used sorted 1n (G0/G1/S-

phase haploid) and 4n (S/G2/M-phase diploid) cells at different timepoints for *Xist* RNA

FISH analysis. One and two punctate *Xist* signals were observed in haploid and diploid

ESCs, respectively (Figure 4A). Our analysis aimed at the detection of *Xist* clusters and

therefore used double stranded probes that also recognize nascent *Xist* and *Tsix* transcripts

before initiation of XCI as a punctate focal signal. The majority of haploid and diploid ESCs

showed no *Xist* cluster (Figure 4B). After the initiation of differentiation, *Xist* RNA clusters

were observed in 59.0% of diploid EpiLCs as expected (Wutz and Jaenisch, 2000). A clear

Xist cluster was also observed in 42.0% of sorted haploid EpiLCs indicating the initiation of

XCI. *Xist* is activated from the single X chromosome in haploid cells with a similar frequency

as from one of the two X chromosomes in diploid cells. Furthermore, 17.9% of diploid

EpiLCs contained two separate *Xist* clusters. This observation might be explained by

considering that the decision to activate *Xist* may have occurred before diploidization, when

the cell was in a haploid state, and *Xist* expression was maintained on both X chromosomes

after diploidization. Strong upregulation of *Xist* was also confirmed by quantitative RT-PCR

analysis in both haploid and diploid EpiLCs (Figure 2).

With further differentiation, we found that the proportion of cells exhibiting *Xist*

clusters decreased (Figure 4A, B). Only 8.7% and 24.0% of haploid and diploid d7 PGCLCs

showed *Xist* clusters, respectively. This observation might reflect the repression of *Xist* in

germ cell differentiation and cell selection that is expected from the loss of X-linked gene

expression in haploid cells after inactivation of the single X chromosome. The ability to

repress *Xist* may contribute to the ability of the germline to maintain a haploid genome with

a higher frequency than somatic lineages that normally do not express repressors of *Xist*.

Taken together, our data demonstrate that *Xist* is activated from a single X chromosome

without *Xic* pairing in differentiating haESCs.

Discussion

Our observation that haESCs maintain a haploid genome during germ cell differentiation

enabled us to analyze *Xist* activation in the context of a haploid genome. Diploid mouse

ESCs activate *Xist* upon entering differentiation and have been used as a model for

studying the underlying regulation of XCI. Our results are explained by the idea that the

amount of blocking factors produced from a single set of autosomes is insufficient to

counteract activators from a single X chromosome. In contrast, X-linked activators are

titrated by a double dose of blocking factors in diploid male cells preventing *Xist* activation.

Our result therefore supports a model of diffusible X-linked activators and autosomal

blocking factors (Barakat et al., 2014; Pollex and Heard, 2019). We note that haploid cells

have a smaller cell volume than diploid cells (Freimann and Wutz, 2017), and it is not

entirely clear if cell volume could have an influence on XCI. Previous studies have also

linked the activation of *Xist* with *Xic* pairing in differentiating diploid ESCs (Bacher et al.,

2006; Xu et al., 2006). *Xic* pairing cannot occur in haploid cells as only a single X

chromosome is present. Our experiment shows that *Xist* activation does not strictly depend

on *Xic* pairing. This observation does not rule out that *Xic* pairing contributes to XCI in

diploid cells or has a role in ensuring that one X chromosome remains active after the

decision for initiating XCI has been taken in female cells. A recent study has reported on

engineering the *Xic* regions for tethering to the nuclear lamina (Pollex and Heard, 2019).

XCI was initiated in female mouse ESCs despite *Xic* movement was restricted.

Previously, *XIST* expression has been analyzed in human haESCs (Sagi et al., 2016).

XCI in the human embryo is regulated differently from the mouse. *XIST* is initially expressed

from the single and both X chromosomes in male and female embryos, respectively

(Okamoto et al., 2011). In contrast to mouse, *XIST* does not lead to chromosomal silencing

in human preimplantation embryos. The *XACT* gene, which is not present in mice, prevents

silencing and a partial repression of X-linked genes is observed that has been termed

"dampening" (Sahakyan et al., 2017). After implantation *XIST* expression becomes

monoallelic in female embryos and XCI is initiated. Conversely, the post implantation

human male embryo ceases *XIST* expression. Female human ESCs show monoallelic

XIST expression and an Xi. During culture erosion of the Xi has been observed, which is

characterized by a loss of *XIST* expression and reactivation of genes from the Xi to various

extent. Established human haESC lines have been shown to lack *XIST* expression (Sagi

et al., 2016). Although, this observation is expected, it is not entirely clear how an *XIST*

negative state was established. Lack of *XIST* activation, selection of cells with an active X

chromosome, and the erosion of an Xi can be considered. The fact that *XIST* is initially

expressed from all X chromosomes in human diploid preimplantation embryos, suggests

that it is also activated in haploid embryos. It is conceivable that mechanisms that prevent

inactivation including *XACT* could contribute to survival and potentially lead to repression

of *XIST* in human haESCs (Vallot et al., 2017). Irrespectively, it is likely that the lack of *XIST*

expression in established haESC lines does not reflect the state of the embryonic cell of

origin. In contrast, mouse ESCs have not initiated XCI. Capturing of the earliest stages of

XCI upon entry into differentiation permits the study of the X chromosome counting

mechanism in culture.

Our study also reveals that PGCLCs maintain a haploid genome with high frequency.

Therefore, the diploidization rate during germline differentiation is comparable to haESCs

and substantially lower than that of somatic lineages. PGCs have a similar epigenetic state

to ESCs, which might contribute to tolerance of a haploid genome. Firstly, during migration

to the gonads, the Xi becomes reactivated suggesting that dosage compensation is not

essential for germ cell development. Secondly, ESCs and PGCs exhibit genome-wide DNA

hypomethylation (Leitch et al., 2013). Lastly, transcription factors that are associated with

pluripotent cells and germ cells including Oct4 have been implicated as repressors of *Xist*

(Donohoe et al., 2009; Navarro et al., 2010; Nesterova et al., 2011). It is conceivable that

the expression of potential repressors of *Xist* in PGCs might contribute to the maintenance

of a haploid genome consistent with our finding that *Xist* expression is lost at later time

points in differentiating cultures. Oct4 is also expressed in EpiLCs and, thus, cannot explain

repression of *Xist* in ESCs and PGCs. However, other factors have also been implicated in

Xist repression in pluripotent cells. Haploid PGCLCs will be useful for future studies of

ploidy restriction and the genetics of germ cell development.

Experimental Procedures

Derivation and culture of haESC lines

Derivation of haESC lines from 129S6/SvEvTac mice was performed as previously described (Leeb and Wutz, 2011). At passage 5, G0/G1/S-phase haploid cell population of a haESC line was purified by fluorescence-activated cell sorting (FACS) using flow cytometer (MoFlo Astrios EQ, Beckman Coulter) after Hoechst 33342 (Invitrogen) staining. Sorted cells were cultured and maintained on a gelatin-coated plate with irradiated mouse embryonic fibroblasts (MEFs) of DR4 mouse embryos at E12.5 (The Jackson Laboratory, no. 003208). Cells were maintained in Serum+2i+LIF medium, which was mixed with equal volume of Serum+LIF medium without 2i (Postlmayr et al., 2020) and 2i+LIF medium (Hayashi and Saitou, 2013). At passage 10, G0/G1/S-phase haploid cell population of the haESC line was purified by FACS and was maintained on a gelatin-coated plate with MEFs in Serum+2i+LIF medium.

In vitro **germ cell differentiation**

Germ cell differentiation was performed following a published protocol (Hayashi and Saitou, 2013) with a few modification. The haESC line was cultured on an ornithine- and laminin-coated plate without MEFs in 2i+LIF medium from passage 12. At passage 15, EpiLC differentiation was initiated as described in the protocol. After 2 days of EpiLC differentiation, 2.3×10^5 EpiLCs were plated into a well of a Sphericalplate 5D (Kugelmeiers Ltd.) with 1.4 ml of PGCLC differentiation medium not containing BMP8a. After 4 days of PGCLC differentiation, half of the medium was replaced with fresh PGCLC differentiation medium not containing BMP4 and BMP8a.

Flow cytometry analysis and cell sorting

To investigate the cell cycle of haploid and diploid cells and PGCLCs, flow cytometry

analysis was performed to ESCs, EpiLCs and EBs by following procedures. Cells were

harvested from culture vessels as described in a published protocol (Hayashi and Saitou,

2013), followed by staining with 15 µg/ml Hoechst 33342 for 12 min at 37 °C. Subsequently,

PE anti-integrin β3 (BioLegend, no. 104307) and eFluor 660 anti-SSEA1 (eBioscience, no.

50881341) were added to the cell suspension at concentration of 1 µg/ml and 0.12 µg/ml,

respectively, and the cell sample was kept on ice for 12 min. The fluorescence of each dye

was measured by flow cytometry. Cell cycle of haploid and diploid cells was determined

based on peaks of cell population at 1n and 2n DNA contents, corresponding to G0/G1-

phase haploid cells and G2/M-phase haploid and G0/G1-phase diploid cells, respectively,

by measuring the signal of Hoechst 33342. The population of PGCLCs was determined by

measuring the signals of PE and eFluor 660.

Karyotyping

On day 7 of germ cell differentiation, M-phase arrest of EBs was performed by culturing in

PGCLC differentiation medium supplemented with 0.05 mg/ml demecolcine (Merck) for 8

hours. Subsequently, PGCLCs were sorted by FACS. Chromosome counting of PGCLCs

was performed as previously described (Aizawa et al., 2020).

Transcription analysis

During germ cell differentiation of mixed haploid-diploid ESCs, G0/G1/S-phase haploid and

S/G2/M-phase diploid ESCs, EpiLCs and d7 PGCLCs were sorted from ESCs, EpiLCs and

d7 EBs by FACS, respectively. E12.5 gonads were harvested from 129S6/SvEvTac mouse

embryos. Sex of the gonads was determined by their morphology. PGCs were identified as

gonadal cells positive for both SSEA1 and integrin β3 and were sorted by FACS. RNA of

each sample was extracted using the RNeasy Mini Kit (Qiagen) following the

manufacturer's protocol, including an on-column DNA digest using RNase-free DNase

(Qiagen). RNA concentration was determined using a NanoDrop Lite (Thermo Fisher

Scientific). 100 - 500 ng total RNA was reverse transcribed using the PrimeScript RT Master

Mix (Takara) according to the manufacturer's instruction. RT-PCR was performed on a 384

well format with the 480 Lightcycler instrument (Roche) using KAPA SYBR FAST qPCR

KIT (Kapa Biosystems). Fold change expression was calculated using the ΔΔct method.

Gapdh expression of G0/G1/S-phase haploid ESCs was used to normalize the transcription

of G0/G1/S-phase EpiLCs and d7 PGCLCs. *Gapdh* expression of S/G2/M-phase diploid

ESCs was used to normalize the transcription of S/G2/M-phase EpiLCs, d7 PGCLCs and

PGCs. Primers used for transcription analysis are listed in Table S1.

Xist RNA FISH

RNA FISH was performed to analyze *Xist* expression during germ cell differentiation of

mixed haploid-diploid ESCs. FISH probe was prepared from ptetOP-Xist-PA plasmid with

Cy3-dCTP (Amersham Biosciences) as described previously (Wutz and Jaenisch, 2000).

G0/G1/S-phase haploid and S/G2/M-phase diploid ESCs, EpiLCs and d7 PGCLCs were

sorted by FACS, respectively. Sorted cells were mounted onto glass slides using Cytospin

4 (Thermo Scientific) for 3 min at 800 or 900 rpm for haploid or diploid cells, respectively.

Glass slides were immediately rinsed in PBS and washed in CSK buffer (100 nM NaCl, 300

nM sucrose, 3 mM $MgCl_2$, 10 mM PIPES pH 6.8) for 30 sec, in CSK buffer + 0.5% Triton

X-100 for 2 min and again in CSK buffer for 30 sec. Cells were fixed in 4%

paraformaldehyde in PBS for 10 min and washed in 70%, 80%, 95% and 100% EtOH for

2 min each. After airdrying the slides, hybridization was performed by applying probe to the

cells. Probes were covered with a coverslip and sealed with rubber cement. Slides were

then placed in a humidified chamber and incubated overnight at 37°C. Coverslips were

removed and slides were washed in 2X SSC + formamide (50%) for 15 min at 39°C, in 2X

SCC three times for 5 min each at 39°C, in 1X SSC for 10 min and in 4X SSC with a dip

(20X SSC: 3 M NaCl, 0.3 M tri-sodium citrate dihydrate in H_2O). Cellular DNA was

counterstained by incubating slides in 4X SSC + 0.1% Tween + 14 mM DAPI for 1.5 min.

Slides were washed with 4X SSC for 5 min. After washing, the sample was mounted in

Vectashield (Vector Laboratories) and covered with a coverslip. Samples were imaged

under the microscope (Axio Observer Z1, Zeiss) equipped with an ORCA-Flash4.0 camera

(Hamamatsu Photonics K.K.). Images were processed using Zeiss Zen Pro 2.0 software.

Statistical analysis

For comparison of cell population ratio (Figure 3C, D), measurements were analyzed with

the GraphPad Prism 8 software using a two-tailed t-test. A p-value < 0.05 was considered

statistically significant.

Acknowledgments

We thank Dr. Giulio Di Minin for derivation of haESC lines. This study was supported by the Swiss National Science Foundation (grant 31003A_152814/1).

Author Contributions

E.A. and A.W. conceptualized experiments. E.A., C.K., S.S. and R.F. collected the data. E.A. and C.K. analyzed the data. E.A. and A.W. wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

Supplemental Information

Table S1. List of oligos

Figure Legends

Figure 1. HaESCs differentiate to haploid PGCLCs by an in vitro culture.

(A) A scheme of germ cell differentiation of mixed haploid-diploid ESCs. (B) Morphology of ESCs, EpiLCs and d7 EBs derived from mixed haploid-diploid ESCs. Scale bar, 100 µm. (C) A representative flow cytometry analysis of d7 EBs derived from mixed haploid-diploid ESCs. PGCLCs, positive for both SSEA1 and integrin β3, accounted for 15.1% out of all cells. (D, E) The proportion of PGCLCs (D) and G0/G1/S-phase haploid cells (E) out of all cells at the different stages during germ cell differentiation of mixed haploid-diploid ESCs. n = 14 (ESC), 9 (EpiLC), 7 (d6 EB), 18 (d7 EB) and 7 (d8 EB). Data represents the mean value and the standard error of the mean. (F) Flow cytometry analysis of DNA content of diploid ESCs, mixed haploid-diploid ESCs, and d7 EBs derived from mixed haploid-diploid ESCs. Cell cycle phases of haploid and diploid cells (top), and PGCLCs in d7 EBs (green) are indicated. (G) Representative chromosome spread of a haploid d7 PGCLC.

Figure 2. Transcription profile of haploid and diploid cells during germ cell differentiation *in vitro*.

Transcription of PGC markers (*Blimp1*, *Prdm14*, *Tfap2c* and *Stella*), *Dnmt3b* and *Xist* during germ cell differentiation of mixed haploid-diploid ESCs. Gene expression of G0/G1/S-phase haploid and S/G2/M-phase diploid cells of ESCs, EpiLCs and d7 EBs was measured, respectively. Female and male PGCs purified from E12.5 gonads by FACS were used as controls. Gene expression of G0/G1/S-phase haploid and S/G2/M-phase diploid samples was normalized to *Gapdh* expression relative to G0/G1/S-phase haploid and S/G2/M-phase diploid ESCs, respectively. Gene expression of male and female PGCs was

normalized to *Gapdh* expression relative to S/G2/M-phase diploid ESCs. Data represents

relative expression of each sample with the mean value and the standard error of the mean

(n = 4). H, G0/G1/S-phase haploid; D, S/G2/M-phase diploid; F, female; M, male.

Figure 3. PGCLCs preferentially maintain haploidy *in vitro*.

(A) A representative distribution of DNA content in d7 EBs derived from mixed haploid-

diploid ESCs. PGCLCs, positive for SSEA1 and integrin β3, are indicated by green color.

The cell cycle profile of haploid and diploid cells is shown at the top. (B) A representative

flow cytometry analysis of d7 EBs. PGCLCs accounted for 11.2% out of all cells in d7 EBs

derived from mixed haploid-diploid ESCs. PGCLCs accounted for 38.1%, 9.2%, 6.3% and

8.7% out of G0/G1/S haploid cell population, G2/M haploid + G0/G1 diploid cell population,

S/G2/M diploid cell population, and G2/M haploid + all diploid cell population of d7 EBs

derived from mixed haploid-diploid ESCs, respectively. Embryonic female gonads at E12.5

and d7 EBs derived from diploid ESCs were also analyzed as controls. (C) The ratio of

PGCLCs in cell populations of different DNA content in d7 EBs derived from mixed haploid-

diploid ESCs or diploid ESCs. PGCLCs accounted for 32.3% on average out of G0/G1/S

haploid cells in d7 EBs derived from mixed haploid-diploid ESCs. Data represents the mean

and the standard error of the mean. n = 18 (d7 EBs derived from mixed haploid-diploid

ESCs) and n = 12 (d7 EBs derived from diploid ESCs). **** P < 0.0001. (D) Cell cycle

distribution of haploid and diploid cells during germ cell differentiation of mixed haploid-

diploid ESCs. Data represents the mean value and the standard error of the mean. n = 14

(ESC), 9 (EpiLC), 18 (d7 EB) and 18 (d7 PGCLC). **** P < 0.0001.

Figure 4. *Xist* is activated and repressed in both haploid and diploid cells during germ cell differentiation.

(A) Representative images of *Xist* expression during germ cell differentiation of mixed haploid-diploid ESCs detected by RNA FISH using a Cy3-labeled *Xist* probe (orange). No *Xist* cluster but single foci observed in G0/G1/S-phase haploid ESCs and PGCLCs. No cluster but double foci observed in S/G2/M-phase diploid ESCs and PGCLCs. Single *Xist* clusters (arrow) and foci (arrowhead) observed in G0/G1/S-phase haploid EpiLCs. Double clusters (asterisk), single clusters (arrow) and no cluster (arrowhead) observed in each S/G2/M-phase diploid EpiLC. (B) Proportion of *Xist* RNA FISH signals during germ cell differentiation of mixed haploid-diploid ESCs. The number of cells possessing no, single or double *Xist* RNA clusters were counted in G0/G1/S-phase haploid and S/G2/M-diploid cells. Total numbers of counted cells are 138 (H, ESC), 171 (D, ESC), 100 (H, EpiLC), 173 (D, EpiLC), 69 (H, PGCLC) and 121 (D, PGCLC). D, S/G2/M-phase diploid; H, G0/G1/S-phase haploid.

References

Aizawa, E., Dumeau, C.-E., Freimann, R., Di Minin, G., and Wutz, A. (2020). Polyploidy of semi-cloned embryos generated from parthenogenetic haploid embryonic stem cells. PLOS ONE *15*, e0233072.

Augui, S., Filion, G.J., Huart, S., Nora, E., Guggiari, M., Maresca, M., Stewart, A.F., and Heard, E. (2007). Sensing X chromosome pairs before X inactivation via a novel X-pairing region of the Xic. Science (New York, NY) *318*, 1632-1636.

Bacher, C.P., Guggiari, M., Brors, B., Augui, S., Clerc, P., Avner, P., Eils, R., and Heard, E. (2006). Transient colocalization of X-inactivation centres accompanies the initiation of X inactivation. Nature cell biology *8*, 293-299.

Barakat, T.S., Loos, F., van Staveren, S., Myronova, E., Ghazvini, M., Grootegoed, J.A., and Gribnau, J. (2014). The trans-activator RNF12 and cis-acting elements effectuate X chromosome inactivation independent of X-pairing. Molecular cell *53*, 965-978.

Donohoe, M.E., Silva, S.S., Pinter, S.F., Xu, N., and Lee, J.T. (2009). The pluripotency factor Oct4 interacts with Ctcf and also controls X-chromosome pairing and counting. Nature *460*, 128-132.

Elling, U., Taubenschmid, J., Wirnsberger, G., O'Malley, R., Demers, S.P., Vanhaelen, Q., Shukalyuk, A.I., Schmauss, G., Schramek, D., Schnuetgen, F., *et al.* (2011). Forward and reverse genetics through derivation

of haploid mouse embryonic stem cells. Cell Stem Cell *9*, 563-574.

Freimann, R., and Wutz, A. (2017). A fast and efficient size separation method for haploid embryonic stem
cells. Biomicrofluidics *11*, 054117.

Galupa, R., and Heard, E. (2018). X-Chromosome Inactivation: A Crossroads Between Chromosome
Architecture and Gene Regulation. Annual review of genetics *52*, 535-566.

Hayashi, K., Ogushi, S., Kurimoto, K., Shimamoto, S., Ohta, H., and Saitou, M. (2012). Offspring from oocytes
derived from in vitro primordial germ cell-like cells in mice. Science (New York, NY) *338*, 971-975.

Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., and Saitou, M. (2011). Reconstitution of the mouse germ cell
specification pathway in culture by pluripotent stem cells. Cell *146*, 519-532.

Hayashi, K., and Saitou, M. (2013). Generation of eggs from mouse embryonic stem cells and induced
pluripotent stem cells. Nat Protoc *8*, 1513-1524.

He, W., Zhang, X., Zhang, Y., Zheng, W., Xiong, Z., Hu, X., Wang, M., Zhang, L., Zhao, K., Qiao, Z.*, et al.* (2018).
Reduced Self-Diploidization and Improved Survival of Semi-cloned Mice Produced from Androgenetic Haploid
Embryonic Stem Cells through Overexpression of Dnmt3b. Stem cell reports *10*, 477-493.

He, Z.Q., Xia, B.L., Wang, Y.K., Li, J., Feng, G.H., Zhang, L.L., Li, Y.H., Wan, H.F., Li, T.D., Xu, K.*, et al.* (2017).
Generation of Mouse Haploid Somatic Cells by Small Molecules for Genome-wide Genetic Screening. Cell Rep
20, 2227-2237.

Hill, P.W.S., Leitch, H.G., Requena, C.E., Sun, Z., Amouroux, R., Roman-Trufero, M., Borkowska, M., Terragni, J.,
Vaisvila, R., Linnett, S.*, et al.* (2018). Epigenetic reprogramming enables the transition from primordial germ
cell to gonocyte. Nature *555*, 392-396.

Kay, G.F., Penny, G.D., Patel, D., Ashworth, A., Brockdorff, N., and Rastan, S. (1993). Expression of Xist during
mouse development suggests a role in the initiation of X chromosome inactivation. Cell *72*, 171-182.

Leeb, M., Walker, R., Mansfield, B., Nichols, J., Smith, A., and Wutz, A. (2012). Germline potential of
parthenogenetic haploid mouse embryonic stem cells. Development (Cambridge, England) *139*, 3301-3305.

Leeb, M., and Wutz, A. (2011). Derivation of haploid embryonic stem cells from mouse embryos. Nature *479*,
131-134.

Leitch, H.G., McEwen, K.R., Turp, A., Encheva, V., Carroll, T., Grabole, N., Mansfield, W., Nashun, B., Knezovich,
J.G., Smith, A.*, et al.* (2013). Naive pluripotency is associated with global DNA hypomethylation. Nature
structural & molecular biology *20*, 311-316.

Lyon, M.F. (1961). Gene Action in the X-chromosome of the Mouse (Mus musculus L.). Nature *190*, 372-373.

Mak, W., Nesterova, T.B., de Napoles, M., Appanah, R., Yamanaka, S., Otte, A.P., and Brockdorff, N. (2004).
Reactivation of the paternal X chromosome in early mouse embryos. Science (New York, NY) *303*, 666-669.

Monkhorst, K., Jonkers, I., Rentmeester, E., Grosveld, F., and Gribnau, J. (2008). X inactivation counting and
choice is a stochastic process: evidence for involvement of an X-linked activator. Cell *132*, 410-421.

Navarro, P., Oldfield, A., Legoupi, J., Festuccia, N., Dubois, A., Attia, M., Schoorlemmer, J., Rougeulle, C.,
Chambers, I., and Avner, P. (2010). Molecular coupling of Tsix regulation and pluripotency. Nature *468*, 457-
460.

Nesterova, T.B., Senner, C.E., Schneider, J., Alcayna-Stevens, T., Tattermusch, A., Hemberger, M., and
Brockdorff, N. (2011). Pluripotency factor binding and Tsix expression act synergistically to repress Xist in
undifferentiated embryonic stem cells. Epigenetics & chromatin *4*, 17.

Ohhata, T., and Wutz, A. (2013). Reactivation of the inactive X chromosome in development and
reprogramming. Cellular and molecular life sciences : CMLS *70*, 2443-2461.

Ohinata, Y., Ohta, H., Shigeta, M., Yamanaka, K., Wakayama, T., and Saitou, M. (2009). A signaling principle
for the specification of the germ cell lineage in mice. Cell *137*, 571-584.

Okamoto, I., Otte, A.P., Allis, C.D., Reinberg, D., and Heard, E. (2004). Epigenetic dynamics of imprinted X
inactivation during early mouse development. Science (New York, NY) *303*, 644-649.

Okamoto, I., Patrat, C., Thepot, D., Peynot, N., Fauque, P., Daniel, N., Diabangouaya, P., Wolf, J.P., Renard, J.P.,

Duranthon, V.*, et al.* (2011). Eutherian mammals use diverse strategies to initiate X-chromosome inactivation
during development. Nature *472*, 370-374.

Olbrich, T., Mayor-Ruiz, C., Vega-Sendino, M., Gomez, C., Ortega, S., Ruiz, S., and Fernandez-Capetillo, O.
(2017). A p53-dependent response limits the viability of mammalian haploid cells. Proceedings of the
National Academy of Sciences of the United States of America *114*, 9367-9372.

Olbrich, T., Vega-Sendino, M., Murga, M., de Carcer, G., Malumbres, M., Ortega, S., Ruiz, S., and Fernandez-
Capetillo, O. (2019). A Chemical Screen Identifies Compounds Capable of Selecting for Haploidy in
Mammalian Cells. Cell Rep *28*, 597-604 e594.

Pollex, T., and Heard, E. (2019). Nuclear positioning and pairing of X-chromosome inactivation centers are not
primary determinants during initiation of random X-inactivation. Nature genetics *51*, 285-295.

Postlmayr, A., Dumeau, C.E., and Wutz, A. (2020). Cdk8 is required for establishment of H3K27me3 and gene
repression by Xist and mouse development. Development (Cambridge, England) *147*.

Sagi, I., Chia, G., Golan-Lev, T., Peretz, M., Weissbein, U., Sui, L., Sauer, M.V., Yanuka, O., Egli, D., and Benvenisty,
N. (2016). Derivation and differentiation of haploid human embryonic stem cells. Nature *532*, 107-111.

Sahakyan, A., Kim, R., Chronis, C., Sabri, S., Bonora, G., Theunissen, T.W., Kuoy, E., Langerman, J., Clark, A.T.,
Jaenisch, R.*, et al.* (2017). Human Naive Pluripotent Stem Cells Model X Chromosome Dampening and X
Inactivation. Cell Stem Cell *20*, 87-101.

Sugimoto, M., and Abe, K. (2007). X chromosome reactivation initiates in nascent primordial germ cells in
mice. PLoS genetics *3*, e116.

Takagi, N., and Sasaki, M. (1975). Preferential inactivation of the paternally derived X chromosome in the
extraembryonic membranes of the mouse. Nature *256*, 640-642.

Takahashi, S., Lee, J., Kohda, T., Matsuzawa, A., Kawasumi, M., Kanai-Azuma, M., Kaneko-Ishino, T., and Ishino,
F. (2014). Induction of the G2/M transition stabilizes haploid embryonic stem cells. Development (Cambridge,
England) *141*, 3842-3847.

Vallot, C., Patrat, C., Collier, A.J., Huret, C., Casanova, M., Liyakat Ali, T.M., Tosolini, M., Frydman, N., Heard,
E., Rugg-Gunn, P.J.*, et al.* (2017). XACT Noncoding RNA Competes with XIST in the Control of X Chromosome
Activity during Human Early Development. Cell Stem Cell *20*, 102-111.

Wutz, A., and Jaenisch, R. (2000). A shift from reversible to irreversible X inactivation is triggered during ES
cell differentiation. Molecular cell *5*, 695-705.

Xu, N., Donohoe, M.E., Silva, S.S., and Lee, J.T. (2007). Evidence that homologous X-chromosome pairing
requires transcription and Ctcf protein. Nature genetics *39*, 1390-1396.

Xu, N., Tsai, C.L., and Lee, J.T. (2006). Transient homologous chromosome pairing marks the onset of X
inactivation. Science (New York, NY) *311*, 1149-1152.

A

① G0/G1/S haploid

② G2/M haploid + G0/G1 diploid

③ S/G2/M diploid

④ G2/M haploid + all diploid

B

E12.5 Gonad — 18.6%

All in hap-dip d7 EB — 11.2%

G0/G1/S hap in hap-dip d7 EB — 38.1%

G2/M hap + G0/G1 dip in hap-dip d7 EB — 9.2%

S/G2/M dip in hap-dip d7 EB — 6.3%

G2/M hap + all dip in hap-dip d7 EB — 8.7%

All in dip d7 EB — 5.0%

C

(a) All

(b) G0/G1/S hap

(c) G2/M hap + G0/G1 dip

(d) S/G2/M dip

(e) G2/M hap + all dip

D7 EB derived from haploid-diploid mixed ESC

(f) All

D7 EB derived from diploid ESC

****

D

- S/G2/M dip
- G2/M hap + G0/G1 dip
- G0/G1/S hap

****

ESC, EpiLC, D7 EB, D7 PGCLC

A

B

Table S1. List of oligos

Name	Sequence (5' to 3')
Blimp1-F	AGC ATG ACC TGA CAT TGA CAC C
Blimp1-R	CTC AAC ACT CTC ATG TAA GAG GC
Dnmt3b-F	CTC GCA AGG TGT GGG CTT TTG TAAC
Dnmt3b-R	CTG GGC ATC TGT CAT CTT TGC ACC
Gapdh-1	AGG TCG GTG TGA ACG GAT TTG
Gapdh-2	TGT AGA CCA TGT AGT TGA GGT CA
Prdm14-F	ACA GCC AAG CAA TTT GCA CTA C
Prdm14-R	TTA CCT GGC ATT TTC ATT GCT C
Stella-F	AGG CTC GAA GGA AAT GAG TTT G
Stella-R	TCC TAA TTC TTC CCG ATT TTC G
Tfap2c-F	GGG CTT TTC TCT CTT GGC TGG T
Tfap2c-R	TCC ACA CGT CAC CCA CAC AA
Xist-F	GCT GGT TCG TCT ATC TTG TGG GTC
Xist-R	TTG TTC AGA GTA GCG AGG ACT TGA AGA G