# Haploid mouse germ cell precursors from embryonic stem cells reveal *Xist* activation from a single X chromosome

Eishi Aizawa[1], Corinne Kaufmann[1], Sarah Sting[1], Remo Freimann[1], Anton Wutz[1,*]

[1]Institute of Molecular Health Sciences, Swiss Federal Institute of Technology, ETH Zurich, Zurich, Switzerland

*Correspondence: awutz@ethz.ch

## Abstract

In mammals, one of the two X chromosomes of diploid female cells is inactivated for gene dosage compensation between the sexes. *Xist* initiates X chromosome inactivation (XCI) in female cells. Although *Xist* is not expressed in male cells with a single X chromosome, it remains unclear if a single X chromosome in a haploid cell would lead to *Xist* activation. Haploid embryonic stem cells (haESCs) have been derived from several mammalian species [1], and maintain pluripotency in culture. They also show a predisposition for diploidization especially during differentiation. Establishing somatic haploid cells has been reported by inhibiting potential factors for diploidization [2, 3]. Considering high diploidization in somatic lineages, we wondered if specification of germline progenitor cells could be less restrictive to a haploid genome. Here, we report that mouse primordial germ cell-like cells (PGCLCs) preferentially maintain a haploid genome. Using this differentiation system, we find that *Xist* is initially induced in differentiating haESCs but subsides later after germ cell specification. The observation that XCI is initiated in haploid cells

demonstrates *Xist* activation without the requirement of a second X chromosome

inactivation center (*Xic*).

To perform germ cell differentiation we followed a previously established protocol [4].

A mixed population of haploid and diploid ESCs was differentiated to epiblast-like cells

(EpiLCs) for 2 days, followed by aggregation into embryoid bodies (EBs) (Figure 1A, S1A).

In this culture PGCLCs arose between day 6 to 8, and were identified by co-expression of

SSEA1 and integrinβ3 (Figure 1B, S1B). To investigate the ploidy of the cells we stained

the DNA with Hoechst and analyzed the cells by flow cytometry. G0/G1/S-phase haploid

(1n) cells accounted for 9.3% of day 7 EB cells, which is approximately half the fraction at

the beginning of differentiation (16.8%; Figure S1C). We further noted that the 1n haploid

population contained substantial numbers of cells that expressed both SSEA1 and

integrinβ3 (Figure 1B). PGCLCs accounted for 32.3% of G0/G1/S-phase haploid cells.

However, less than 10% of cells with higher DNA content expressed both PGC markers

(Figure 1D, S1D, S1E). Conversely, haploid cells were enriched in the PGCLC population

compared to SSEA1- and integrinβ3-negative cells that likely represent somatic fates

(Figure S1F). We confirmed a set of 20 acrocentric chromosomes in haploid PGCLCs as

expected for a haploid mouse genome (Figure 1C). Furthermore, the PGC markers, *Blimp1*,

*Prdm14*, *Tfap2c* and *Stella*, were upregulated, and *Dnmt3b* was downregulated in sorted

PGCLCs relative to EpiLCs and ESCs (Figure S2A). Our results show that haploid PGCLCs

can be obtained *in vitro* whereby a haploid genome is preferentially maintained in PGCLCs

over somatic lineages.

We next used this differentiation system to address the question if *Xist* is activated

from a single X chromosome upon differentiation of haESCs. We sorted G0/G1/S-phase

haploid (1n) and S/G2/M-phase diploid (4n) cells at different timepoints during germ cell

differentiation of a mixed haploid-diploid ESC culture and performed RNA FISH for

detecting *Xist* RNA. At the start of differentiation, we observed one and two foci in haploid

and diploid ESCs, respectively (Figure S2B), which correspond to nascent *Xist* and *Tsix*

expression before initiation of XCI. In EpiLCs, *Xist* clusters were observed indicating the

activation of *Xist*. A single *Xist* cluster was observed in 42% and 59% of haploid and diploid

EpiLCs, respectively (Figure 1E, F). Surprisingly, 17.9% of diploid EpiLCs had two *Xist*

clusters. This observation is unexpected and might be explained by a potential deregulation

of XCI when the cell was in a haploid state. We confirmed a strong upregulation of *Xist* by

quantitative RT-PCR analysis in both haploid and diploid EpiLCs (Figure S2A). Taken

together, our data demonstrate that *Xist* can be activated from a single X chromosome at

the onset of haESC differentiation. With further differentiation we observed that the

proportion of cells exhibiting *Xist* clusters decreased in haploid and diploid d7 PGCLCs.

How this decrease is achieved is unclear, but repression of *Xist* in germ cell differentiation

or death of *Xist* expressing cells are possible mechanisms.

The persistence of haploid cells during germ cell differentiation enabled us to perform

an analysis of *Xist* activation in the context of a haploid genome. Previous studies reported

*Xist* activation in genomes of different ploidies [5]. Tetraploid cells transiently activate *Xist*

on a variable number of X chromosomes and a stochastic regulation has been proposed.

In contrast, observations on pairing of X chromosomes at their *Xic* loci has been reported

and implicated in the activation of *Xist* [6, 7]. Furthermore, elements that are required for

*Xic* pairing have been identified and transgenically shown to induce XCI [8]. In haploid cells,

*Xic* pairing cannot occur as only a single X chromosome is present similar to male cells. In

contrast to diploid male cells, we find that a single X chromosome is sufficient for *Xist*

activation in haploid cells. Therefore, *Xist* activation can occur independent of *Xic* pairing.

Our finding does not rule out that *Xic* pairing could have a role in ensuring that one X

chromosome remains active in female cells. We further note that our interpretation is

consistent with a recent study that has used tethering of the *Xic* regions to the nuclear

lamina to prevent *Xic* pairing [9]. Although it is hard to rule out in this study that *Xic* pairing

had occurred at some point, the report that XCI could be initiated when *Xic* movement was

constraint to the lamina is consistent with our observation.

We also find that PGCLCs are obtained from haESCs with higher frequency than

somatic lineages. This observation might reflect a similar epigenetic state of PGCs and

ESCs. In mice, PGCs initiate reactivation of the inactive X chromosome and genome-wide

DNA demethylation during migration to the gonads. A lack of dosage compensation and

low DNA methylation level are also associated with naive ESC and the early epiblast [10].

Therefore, it is conceivable that ESCs and PGCs are less restricted for ploidy and relative

X-linked gene expression. Haploid PGCLCs will be useful for future studies of the genetics

of germ cell development.

## Acknowledgments

89   We thank Dr. Giulio Di Minin for derivation of haESC lines. This study was supported by

90   the Swiss National Science Foundation (grant 31003A_152814/1).

## Author Contributions

E.A. and A.W. conceptualized experiments. E.A., C.K., S.S. and R.F. collected the data.

E.A. and C.K. analyzed the data. E.A. and A.W. wrote the manuscript.

## Declaration of Interests

The authors declare no competing interests.

## Figure Legends

**Figure 1. Preferential maintenance of haploidy in PGCLCs and *Xist* activation during germ cell differentiation.** (A) A scheme of germ cell differentiation of mixed haploid-diploid ESCs. (B) Flow cytometry analysis of DNA content of diploid ESCs, mixed haploid-diploid ESCs, and d7 EBs derived from mixed haploid-diploid ESCs. Cell cycle phases (top), and PGCLCs in d7 EBs, positive for SSEA1 and integrinβ3, are indicated (green). (C) Representative chromosome spread of a haploid d7 PGCLC. (D) The ratio of PGCLCs in cell populations of different DNA content in d7 EBs derived from mixed haploid-diploid ESCs or diploid ESCs. PGCLCs accounted for 32.3% on average out of G0/G1/S haploid cells in d7 EBs derived from mixed haploid-diploid ESCs. Data represents the mean and the standard error of the mean. n = 18 (d7 EBs derived from mixed haploid-diploid ESCs) and n = 12 (d7 EBs derived from diploid ESCs). **** P < 0.0001. (E) Representative images of *Xist* expression in G0/G1/S-phase haploid EpiLCs and PGCLCs detected by RNA FISH using a Cy3-labeled *Xist* probe (orange). Single *Xist* clusters (arrow) and foci (arrowhead) were observed in G0/G1/S-phase haploid EpiLCs. Single foci but no cluster were observed in G0/G1/S-phase haploid PGCLCs. (F) Proportion of *Xist* RNA FISH signals during germ

cell differentiation of mixed haploid-diploid ESCs. The number of cells possessing no,

single or double *Xist* RNA clusters were counted in G0/G1/S-phase haploid and S/G2/M-

diploid cells. Total numbers of counted cells are 138 (H, ESC), 171 (D, ESC), 100 (H,

EpiLC), 173 (D, EpiLC), 69 (H, PGCLC) and 121 (D, PGCLC). D, S/G2/M-phase diploid;

dip, diploid; H, G0/G1/S-phase haploid; hap, haploid

## Supplemental Information

Supplemental Information includes experimental procedures, two figures and one table.

## References

1.    Leeb, M., and Wutz, A. (2011). Derivation of haploid embryonic stem cells from mouse embryos. Nature *479*, 131-134.

2.    He, Z.Q., Xia, B.L., Wang, Y.K., Li, J., Feng, G.H., Zhang, L.L., Li, Y.H., Wan, H.F., Li, T.D., Xu, K., et al. (2017). Generation of Mouse Haploid Somatic Cells by Small Molecules for Genome-wide Genetic Screening. Cell Rep. *20*, 2227-2237.

3.    Takahashi, S., Lee, J., Kohda, T., Matsuzawa, A., Kawasumi, M., Kanai-Azuma, M., Kaneko-Ishino, T., and Ishino, F. (2014). Induction of the G2/M transition stabilizes haploid embryonic stem cells. Development *141*, 3842-3847.

4.    Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., and Saitou, M. (2011). Reconstitution of the mouse germ cell specification pathway in culture by pluripotent stem cells. Cell *146*, 519-532.

5.    Monkhorst, K., Jonkers, I., Rentmeester, E., Grosveld, F., and Gribnau, J. (2008). X inactivation counting and choice is a stochastic process: evidence for involvement

of an X-linked activator. Cell *132*, 410-421.

6.   Bacher, C.P., Guggiari, M., Brors, B., Augui, S., Clerc, P., Avner, P., Eils, R., and

Heard, E. (2006). Transient colocalization of X-inactivation centres accompanies

the initiation of X inactivation. Nat. Cell Biol. *8*, 293-299.

7.   Xu, N., Tsai, C.L., and Lee, J.T. (2006). Transient homologous chromosome pairing

marks the onset of X inactivation. Science *311*, 1149-1152.

8.   Augui, S., Filion, G.J., Huart, S., Nora, E., Guggiari, M., Maresca, M., Stewart, A.F.,

and Heard, E. (2007). Sensing X chromosome pairs before X inactivation via a

novel X-pairing region of the Xic. Science *318*, 1632-1636.

9.   Pollex, T., and Heard, E. (2019). Nuclear positioning and pairing of X-chromosome

inactivation centers are not primary determinants during initiation of random X-

inactivation. Nat. Genet. *51*, 285-295.

10.   Leitch, H.G., McEwen, K.R., Turp, A., Encheva, V., Carroll, T., Grabole, N.,

Mansfield, W., Nashun, B., Knezovich, J.G., Smith, A., et al. (2013). Naive

pluripotency is associated with global DNA hypomethylation. Nat. Struct. Mol. Biol.

*20*, 311-316.