Were Neoarchean atmospheric methane hazes and early Paleoproterozoic

glaciations driven by the rise of oxygen in surface environments?

Anshuman Swain[1]*, Alan J Kaufman[2], Marcin Kalinowski[3], Stephanie A Yarwood[4], and

William F Fagan[1]

[1]Department of Biology, University of Maryland, College Park, MD 20742, USA

[2]Department of Geology and Earth System Science Interdisciplinary Center, University

of Maryland, College Park, MD 20742, USA

[3]Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland

[4]Department of Environmental Science and Technology, University of Maryland,

College Park, MD, 20742, USA

Abstract

Geochemical evidence suggests methane was the predominant greenhouse gas in the

Archean and early Proterozoic eons. Consequently, fluctuations in methane

concentration in Earth's atmosphere (i.e., 'methane hazes') would have contributed to

climate change and influenced the flux of UV radiation reaching surface environments. If

correct, understanding what factors (e.g., O_2 and/or resource concentration) drove the

biological methane cycle might shed light on the repetition of biological, atmospheric and

climatic events preserved in the sedimentary rock record from ~2.8 to 2.0 BYA. To explore

these interdependencies, we developed a novel dynamical model of microbial ecological

interactions to investigate the conditions under which methane is preferentially released

to the atmosphere. We found that the interplay between resource and O_2 availability

results in complex cyclic methane dynamics unrelated to the functional groups or

efficiencies of microbial communities, to initial conditions, or to other model constraints.

Based on these results, we propose that the cyclicity of methane haze events and glacial

episodes in the Neoarchean and early Paleoproterozoic may have been linked to the

progressive increase in oceanic and atmospheric O_2 through the interval.

Keywords: Methane haze, methanogenesis, microbial interaction, oxygen, oscillations

*Corresponding author: answain@terpmail.umd.edu; Address: 3205, Biology-

Psychology Bldg., University of Maryland, College Park, MD 20742, USA

Introduction

Like Titan, Saturn's orange-shrouded moon, the early Earth may have been

episodically enveloped in a thick blanket of atmospheric methane (CH_4), a powerful

greenhouse gas produced by archaeal methanogens in anoxic environments. A methane

haze would have formed if the CH_4/CO_2 ratio was >0.2 (Trainer et al., 2004; Domagal-

Goldman et al., 2008) contrasting with the value of 0.005 in the modern oxygenated

atmosphere. Although the depth and distribution of the CH_4 haze during such events likely

varied across the planet due to vertical atmospheric motion and mixing (Wolf and Toon,

2010), the biogenic gas would nonetheless have warmed the troposphere, and through

attenuation of the solar UV flux, protected surface environments from high energy

radiation.

Carbon isotope evidence from Neoarchean (2.8 to 2.5 BYA) organic matter

preserved in sedimentary rocks (Eigenbrode and Freeman, 2006) suggests a prolonged

increase in the production of CH_4 and its consumption by methanotrophs to form biomass.

Depending on the leak rate of the greenhouse gas from anoxic sediments, CH_4 may have

accumulated in sufficient concentrations in the atmosphere to form periodic hazes. The

extent and depth of such hazes is likely to have been modulated by the photosynthetic

production and accumulation of molecular oxygen (O_2) in the oceans and atmosphere.

Due to the rapid oxidation of CH_4 to CO_2 in the presence of O_2 (Saunois et al.,

2016), the atmospheric concentration of CH_4 is generally thought to represent a balance

between source and sink reactions. Tipping the scales between the two would thus

modulate the size of the atmospheric reservoir of CH_4. On the source side, the production

of CH_4 could have been affected by competition for critical metabolic substrates between

methanogens and sulfate reducers, or by the lack of key metal cofactors, like nickel,

required for methanogenic enzymes (Anbar and Knoll, 2002; Konhauser et al., 2009). In

this analysis, however, we consider ecological factors and biological interactions that may

have increased the CH_4 flux from surface environments that are directly associated with

the release of photosynthetic O_2. Such a scenario would be particularly relevant in the

Neoarchean when periodic CH_4 hazes are suggested from geochemical measurements

of sedimentary rocks and photochemical models (Zerkle et al., 2012; Farquhar et al.,

2013; Izon et al., 2015; 2017, Williford et at., 2016).

Our dynamic resource-driven model for CH_4 generation and release is based upon

the metabolic repercussions of the Warburg Effect in microbial functional groups (see

Swain and Fagan, 2018) – which was initially proposed to explain uncontrolled cellular

proliferation during cancer growth (Warburg, 1956) – to an ecological scenario

appropriate to Neoarchean and early Paleoproterozoic surface environments and

atmospheric fluctuations. The Warburg Effect in microorganisms describes a shift in

metabolic respiration where the energetic process of oxidative phosphorylation (OP),

which produces CO_2 as an end product, is replaced by glycolysis that instead generates

products that can be incorporated into biomass but yields 10 times less energy. This shift

in respiration mechanisms is particularly pronounced during the rapid growth of microbial

cells that are not resource limited – even in the presence of oxygen. This metabolic

phenomenon has been observed in a wide variety of organisms such as bacteria (where

it is termed "overflow metabolism"; Basan et al., 2015, Schuster et al., 2015; Zhuang et

al., 2011), yeast (the "Crabtree effect"; Deken, 1966), and a diversity of human cells,

including Kupffer and microglial cells (specialized macrophages in the liver and brain,

respectively), lymphocytes (white blood cells), and fast twitch muscle fibers (Schuster et

al., 2015; Vander Heiden et al., 2009). Especially in microorganisms, this phenomenon is

known to have cascading effects on the stoichiometry and community dynamics in natural

ecosystems (see Manzoni et al., 2012; Carlson et al., 2018; Kreft et al., 2020). While the

Warburg Effect may seem energetically counterintuitive, one can observe that as cells

grow rapidly, the kinetic advantage of glycolysis that occurs in the cytoplasm results in a

two-fold energy advantage over OP, which is restricted to the cell membrane (Voet and

Voet, 2004). Past models thus describe a metabolic arms race between OP and glycolysis

that includes: i) efficiency vs. rate (OP is more efficient, but glycolysis is faster), ii) surface

area vs. volume (glycolysis occurs in the cell interior whereas OP is constrained to the

cell surface), and iii) biomass production (glycolysis produces organic substrates that can

be used in biomass production directly, which is generally not true for OP) (see Swain

and Fagan, 2018; Schuster et al., 2015). Results from these models suggest that large

inputs of resources (i.e., any carbon substrate that allows the microbes to grow) with even

small inputs of oxygen can stimulate glycolysis over the more energetic process of OP,

and therefore, can affect ecological processes, specifically methanogenesis, that use

these glycolytic overflow products (e.g., formate, lactate, and acetate).

To understand the dynamics of how resource and O_2 availability can affect the

production and consumption of CH_4, we present a dynamical model of microbial

ecological interactions, using the concept of overflow metabolism associated with the

Warburg Effect. The ecological interactions described in this ordinary differential equation

(ODE) model may be relevant to geological episodes where pulsed resource inputs or the

availability of O_2 can alter the CH_4 production rate. Our ODE model might then apply to

climatic and environmental perturbations of the early Earth, including the cyclic patterns

of Neoarchean CH_4 haze events (Pavlov et al., 2001; Zerkle et al., 2012; Izon et al., 2015;

2017) and early Paleoproterozoic glacial episodes (e.g., Bekker and Kaufman, 2007).

In our model, we focus on five major functional groups of microbes: obligate

aerobic heterotrophic bacteria (A), methanogens (B), aerobic methanotrophs (C),

generalized fermenters (F), anaerobic methanotrophs (G; also known as the anaerobic

oxidation of methane (AOM) consortia), and anaerobic heterotrophs (H; such as iron and

sulfur reducers). All of these microbial functional groups are involved in the production

and/or consumption of generalized resources (R), glycolytic/fermentation products (FP),

O_2, CH_4, and CO_2 (**Fig. 1**). Here, the generalized resources are an abstract representation

of a variety of materials that organisms might use as a carbon source, or in an indirect

assimilation of carbon source and growth. Through this generalization of resources, we

allow that some carbon-based compounds come from oxygenic photosynthesis. For

example, anoxygenic photosynthesis (which, in turn, depended on and was limited by

electron acceptors; see Ward et al., 2019) appears to have provided carbon sources for

microbes, especially in Neoarchean and early Paleoproterozoic sedimentary

environments. As a consequence of this generalization of resources, our model considers

a decoupled system of R and O_2. In addition, non-carbon resources can vary in their

stoichiometry in organisms (see Sterner and Elser, 2002); therefore, this decoupling of R

and O_2 also helps to decode the effects of the continental nutrient influx to the oceans, or

related phenomena.

Figure 1: *Flow diagram representation of the model system. Here (in blue) A, B, C, F, G*

and H are functional groups of heterotrophic bacteria, methanogens, aerobic

methanotrophs, fermenters, anaerobic methanotrophs, and anaerobic heterotrophs,

respectively. The flux arrows indicate processes that produce (in red) either metabolic

reactants or gases cycled among the microbial functional groups. Ordinary differential

equations with detailed explanations representing this dynamical system appear in

Methods.

Key relationships within the flow diagram (Fig. 1) can be understood as follows.

Methanogens depend upon fermentation products (FP) for their methane production; FP

are simple organic substrates such as formate, lactate, and acetate that are produced by

Fermenters (F). Aerobic heterotrophic bacteria (A) also produce fermentation products

(FP) through overflow metabolism, insofar as this functional group responds to increasing

R by producing more FP than CO_2, and thereby stimulating methanogenesis (see

Methods). Anaerobic heterotrophs (H) compete with methanogens (B) for FP. Methane

is consumed by aerobic methanotrophs (C) that utilize O_2 to produce CO_2, and by

anaerobic methanotrophs (G), in consortia with various other groups of microbes, to

produce either alkalinity or CO_2 through the anaerobic oxidation of methane (AOM).

Changes in the size of microbial functional groups are modelled as functions of the

availability of R and the efficiency of their utilization (energy production), as well as their

basal energy requirements necessary to maintain homeostasis (energy requirement). We

assume methanogens (B) represent both acetoclastic and hydrogenotrophic

methanogenesis with respect to reduced FP compounds. Acetoclastic methanogens use

primary FP, and hydrogenotrophic methanogens depend on secondary fermentation for

H_2 and formate. We did not include the hydrogenotrophic methanogens as a sink for CO_2,

however, because this can cause unintended model estimation errors that are important

for the decoupling between R and O_2. As we are assuming a decoupling between CO_2,

O_2, and R (given the unknown extent of oxygenic photosynthesis), we can either leave

CO_2 as a sink (as modeled) or couple CO_2 back to R as a function of oxygenic and

anoxygenic autotrophy. In the latter case, we would include an explicit CO_2 flux to the

hydrogenotrophic methanogens. In the former (as chosen), we assume only that there is

enough CO_2 to fulfill all necessary demands, and therefore the substrate is not a limiting

reagent.

The main rationale behind constructing this coupled ODE-based ecological model

is to understand the response of microbial functional groups to the availability of R and

O_2, through mutual interactions and usage of substrates. Changes in the abundance of

resources, metabolic products (FP, CH_4, and CO_2), and O_2 associated with consortia

interactions are functions of efficiencies of i) overflow metabolism, ii) respiration, and iii)

biomass production and maintenance, as well as the rate of addition of both resources

and O_2. To explore the model, we considered both driven and non-driven states. First, for

non-driven dynamics, we explore the basic dependence of metabolic parameters and

species interactions in closed systems wherein microbes and resources start from an

initial condition, and are then closed off to external changes. Second, we consider driven

dynamics in open systems, in which there are constant rates of addition of R and O_2.

Numerical exploration of the model suggest that the interplay between the two results in

complex, but robust, cyclic methane dynamics. We relate this new dynamic perspective

to the episodic nature of Neoarchean methane haze events and early Paleoproterozoic

glaciation.

Results

For the case of no external forcing (non-driven) where the only initial inputs were

R and O_2, we ran 500,000 model simulations to steady-state (**Figure S1**). Across these

simulations, we varied key model parameters (the efficiency of various metabolic

reactions, including methanogenesis, AOM, and fermentation) and initial conditions

(starting populations of different microbial functional groups, and original abundance of R

and O_2) using Latin Hypercube Sampling (LHS; Carnell, 2020) to explore their impacts

on final gas (O_2, CO_2, and CH_4) concentrations. We gauged the importance of the different

parameters and conditions using partial rank correlation coefficients (PRCC) (see

Methods), and found that steady-state CH_4 and CO_2 concentrations varied the most

based upon the metabolic parameters of methanogens (these included resource

incorporation efficiency (β), resource utilization efficiency (κ) and energy requirement (u))

(**Figure S2B,C**). The model further reveals that steady state CO_2 concentration varied

strongly as a function of the metabolic parameters that controlled anaerobic

methanotrophy (or AOM, G) (**Figure S2B**). Steady state O_2 concentrations varied as

functions of the metabolic parameters of aerobic methanotrophs (C), aerobic

heterotrophic bacteria (A), and methanogens (B) (**Figure S2A**). None of the initial

conditions that we varied (i.e., starting populations of different microbial functional groups

and initial concentrations of R and O_2) had a high PRCC for any of the stable state gas

concentrations **(Figure S3)**, except for the dependence of the final CO_2 concentration on

the initial value of O_2 concentration **(Figure S3B)**.

Although the steady state gas concentrations varied quantitatively with certain

parameters, the qualitative trends that they depicted were relatively invariant across the

Latin Hypercube sweep of parameter space and initial conditions. **Figure 2** depicts a

specific case, where we plot the steady state variations of the proportions of various gases

in a closed, non-driven system (initialized with a biologically-relevant base

parametrization) due to changes in the methane production efficiency coefficient (β, see

Eqs. 4, 9, and 11 in **Methods**), which was set at 1%, 10%, and 50%. With increasing

efficiency, the system became more anoxic (lower O_2) and less acidic (lower CO_2), while

CH_4 concentrations increased, especially at low initial O_2 and high initial R inputs. We

recorded similar model responses for variations of all other parameters (See **Figures**

S4A-O). However, we noted lower than expected CO_2 concentrations at low initial O_2

inputs for low efficiencies of parameters governing the incorporation and utilization of

resources by the aerobic heterotrophs (see Eqs. 2, 3, 8, and 10 in **Methods**; and **Figs.**

S4D and S4H); these can be attributed to the PRCC dependence of CO_2 on initial O_2

concentration. We also noted a minor deviation in CH_4 concentration at a high initial O_2

input in the Ω (resource utilization efficiency for anaerobic methanotrophs, G; see

Methods) simulations (see Eqs. 7 and 10 in **Methods**; **Fig. S4K**); this was attributable to

the formation of CO_2 and its dependence on metabolic parameters of G. These non-

linearities arose in cases where the efficiency of usage, or of incorporation, is either

biologically unrealistic, or are at the extreme end of what is possible for biological systems

(e.g., a very low O_2 affinity for aerobic heterotrophs or a high CH_4 usage efficiency in

anaerobic methanotrophs [see Morris and Schmidt 2013; Bowles et al., 2019]). Overall,

these results signify that, on average, the qualitative outputs of our ODE computations

remained consistent across the parametrizations and initial conditions in the non-driven

setting.

We further tested the stability and the sensitivity of the modeled system to the

addition of new microbial functional groups to the main structure by comparing with results

of a model lacking H (anaerobic heterotrophs) that compete directly with B

(methanogens). The comparison shows no significant differences among the two models

in terms of their partial rank correlation coefficients (see Figure S2).

Figure 2: *Dynamic variations in O_2, CO_2, and CH_4 proportions (in columns A-C) run to*

steady state at set O_2 and R input values, with the coefficient of CH_4 production (β) set

at 1, 10, and 50% in rows 1-3, respectively. Note that the resource (R) axis begins at 2;

this is due to constraints posed by the minimal energy requirements for each microbial

functional group, below which all functional groups diminish.

<u>External forcing</u>

In contrast with the no external forcing scenario above, microbial systems may also be forced, particularly by influxes of metabolizable organic substrates (resource, R) and O_2 that are linked through oxygenic photosynthesis, wherein R is synthesized and O_2 is released as a byproduct of the metabolic activity. Model runs with only forced R inputs were qualitatively simple. As illustrated in **Fig. S5**, initiating the model with a constant rate of addition of R depletes O_2 rapidly regardless of its initial abundance. Under these conditions, the Warburg Effect in oxidative heterotrophs does not play a major role in the long-term dynamics of the microbial system. In this scenario, the Warburg Effect only affects the system initially, so that most FP is derived from general fermenters (F) for the subsequent production of CH_4 through methanogenesis. Most notably, simulations with lower initial O_2 (**Fig. S5B**) resulted in an oscillatory behavior over longer periods of CH_4 generation. These results contrast with simulations with higher initial O_2 (**Fig. S5A**), which resulted in rapid damping of the oscillatory behavior. Note, however, that all simulations with the same parameter values reached the same steady state irrespective of the initial conditions, as one can extrapolate from **Figure S5** (and PRCC values from steady state dynamics). Thus, these simulations revealed that differences in the initial O_2 input only affected the relative timescale of events, as the saturation values (amplitudes) and profiles of the CH_4, CO_2 and R concentrations remained the same. Higher initial O_2 stabilized the system more rapidly to steady state condition, while higher initial R under low O_2 resulted in sustained oscillations over longer intervals. If the initial R input is held constant, the dynamic solution to forcing the microbial system with a fixed rate of addition

of O_2 is that CH_4 will be produced briefly associated with the initial concentration of

resource, and O_2 will fall as long as methane is produced and aerobic methanotrophs

consume it (**Fig. S6**). After the CH_4 resource is depleted, the continued input of O_2

overwhelms the system and kills the anaerobes, thereby allowing O_2 to increase at a

linear rate. Different starting concentrations of R have no apparent effect on system-level

dynamics.

Truly complex dynamics occur when both R and O_2 are supplied at constant rates

(the input rate per unit time, denoted as $\sigma_1(t)$ for R and $\sigma_2(t)$ for O_2; see Eqs. 8 and 12 in

Methods) (**Fig. 3).** Low oxygen supply rates relative to resource supply rates (i.e., high

$\sigma_2(t)$ / $\sigma_1(t)$) or conversely low $\sigma_2(t)$ / $\sigma_1(t)$) result in large amplitude, slow frequency

oscillations in methane and oxygen concentrations in the system, a phenomenon that

differs from the model behavior with only resource addition (see **Figure S5; Figure 3A**).

With decreasing $\sigma_2(t)$ / $\sigma_1(t)$), these oscillations reduce in amplitude and increase in

frequency until a threshold value (of relative O_2 input), where the system reaches a non-

oscillatory stable state (**Figure 3A-J; Figure 4**). The transitions into and out of the periodic

behavior appear smooth, and the amplitude and periodicity of the CH_4 peaks within the

dynamic interval decline with increasing O_2 input rate. Moreover, the ratio of resource to

oxygen input rates at which oscillatory behavior transitions to a stable, non-oscillatory

pattern remains constant across the parameter variations and initial conditions (i.e. the

ratio of R to O_2 input remained at 5, or, $\sigma_2(t)$ / $\sigma_1(t)$~5)(**Figure 4B**). **Figure 4A** further

illustrates the relationship between the O_2 input rate (σ_2, varied from 10 to 20% of σ_1) on

the amplitude and periodicity of CH_4 peaks. Clearly the greatest amplitudes and periods

in CH_4 concentration occur at low O_2 input rates, and these both decline rapidly as oxygen

overwhelms the system, leading to a phase transition when σ_2 is ~ 20% of σ_1 (**Figures 3**

and 4).

We conduct a similar robustness test for the dynamics of a reduced model (lacking

the H functional group of anaerobic heterotrophs) with the dynamics of the full model, and

find negligible differences between the two (e.g., Figure S7). This signifies that the model

is mostly affected by the relative addition rates of oxygen to resources, and not by other

parameters or initial conditions (see Figure S2, S3, S7) for large qualitative changes.

Figure 3: *Dynamic variations in O_2 and CH_4 for the externally forced model with constant*

rates of addition of R and O_2. The O_2 input rate (σ_2) varied between 1% to 50% relative to

the resource input rate across panels (σ_1), but in all cases the R input rate (σ_1) was held

constant at 1.

Figure 4: *(A) shows the dependence of amplitude (in red) and time period (in black) of the CH_4 oscillations in Fig. 3 as a function of the O_2 input rate (σ_2). (B) demarcates the resource input (σ_1) -oxygen input (σ_2) space on the basis of oscillatory behavior for our standard set of parameters. (C) depicts the dependence of the slope of the line separating the space between oscillatory and non-oscillatory behavior in σ_1-σ_2 space on specific parameters (for description of parameters, see **Methods**; Table 1) over a wide range of parameter combinations through partial rank correlation coefficient (PRCC).*

We also perform a sensitivity analysis of the slope demarcating the oscillatory and non-oscillatory behavior in a variety of parameter combination (selected through LHS)

with the results illustrated in Figure 4C. We find that the space demarcation is affected

most by the resource utilization efficiency of methanogens (β), anaerobic methanotrophs

(Ω) and anaerobic heterotrophs ( ), as well as resource incorporation efficiency of aerobic

methanotrophs (λ) and methanogens (κ). Increased conversion efficiency of carbon

substrates to methane (represented by β) has been hypothesized to result in severe biotic

crises and methanogenic bursts in the past (Rothman et al., 2014). The role of aerobic

and anaerobic methanotrophs (represented by Ω and λ, *respectively*) is usually to control

the large-scale methane flux, and that of anaerobic heterotrophs (through) is to compete

for carbon substrates with methanogens (Lyu et al., 2018).

Discussion

Our dynamic, resource-driven ecological model based on the Warburg Effect

reveals clues to the production of CH_4 in response to the availability of molecular O_2 when

applied to a consortium of aerobic and anaerobic microbial functional groups. By

progressively increasing the input rate of O_2 from 10-20% with respect to the resource

input rate, the overflow metabolism of the heterotrophic bacteria provides the resources

necessary to produce CH_4 in greater proportions than it is consumed by both aerobic and

anaerobic methanotrophs.

Applied to potentially coupled biological and environmental events across the

Neoarchean-early Paleoproterozoic transition, this dynamic ecological model may help to

explain how the evolution of oxygenic photosynthesis drove fluctuations in the oxidation

state of surface environments, as well as the pulsed releases of CH_4 to the atmosphere

that formed periodic methane hazes. Carbon isotope compositions of organic matter

preserved in sedimentary rocks deposited between ~2.8 to 2.5 billion years ago reveal

unprecedented levels of ^{12}C enrichment that are most parsimoniously attributed to the

incorporation of biomass formed by methanotrophs (Hayes, 1994; Eigenbrode and

Freeman, 2006). Under aerobic conditions, CH_4 is oxidized with molecular O_2 to form

biomass through the serine or ribulose monophosphate pathways, and CO_2

(Chistoserdova et al., 2005), where the minimum concentrations of CH_4 and O_2 required

to drive the microbial process are estimated at 5-10 μM and 12-20 μM, respectively

(Harris and Hanson, 1980; Lidstrom and Somers, 1984). Under anaerobic conditions NO_3^-

, NO_2^-, SO_4^{2-}, Fe^{3+}, or Mn^{2+} may also be used to oxidize CH_4 to form simple organic

compounds, like acetate (Yan et al., 2018). If photosynthetically derived resources were

high — as expected by the pervasive occurrence of thick organic-rich black shales and

carbonates in Neoarchean sedimentary successions worldwide (e.g. Zhelezinskaia et al.,

2014) — then almost any local increase in O_2 could have initiated overflow metabolism in

extant heterotrophic bacteria. Through microbial community dynamics, this could have

resulted in the enhanced release of CH_4 to the atmosphere, as described in the model

above. Distributed globally, oceanic O_2 oases could then have been the source of

Neoarchean CH_4 hazes.

Independent evidence for the presence of repetitious Neoarchean CH_4 haze

events comes from multiple sulfur isotope measurements of pyrite from the same samples

where organic matter was isolated and analyzed. In Neoarchean marginal marine

sedimentary successions on multiple continents, carbon isotope evidence links enhanced

methanotrophic biomass inputs in organic-rich sedimentary rocks to the presence of a

CH_4 haze that would have attenuated the UV flux and isotopically altered the products of

atmospheric SO_2 photochemistry (see Domagal-Goldman et al., 2008), predominantly

sulfate aerosols and elemental sulfur (Kasting, 2001). The photochemical reactions

resulted in mass-independent fractionation, or redistribution, of the sulfur isotope masses

(S-MIF) in both of the products that rained down and accumulated in surface

environments. Notable deviations in the ratio of multiple rare sulfur isotope abundances

($\Delta^{33}S/\delta^{34}S$ and $\Delta^{36}S/\Delta^{33}S$) (see **Methods**) relative to normal distribution of Archean S-

MIF data (typically referred to as Archean Reference Arrays, or ARAs) coincide in several

discrete stratigraphic intervals with negative excursions in carbon isotope values

suggestive of methanotrophic inputs of organic carbon (Zerkle et al., 2012; Farquhar et

al., 2013; Izon et al., 2015; 2017, Williford et at., 2016). The most recent of these studies

(Izon et al., 2017) used available radiometric constraints to estimate the duration of a

single haze event at ~1 million years. Through accelerated H_2 loss to space, the haze

events are postulated by these authors to have hastened planetary oxidation.

Atmospheric models of Neoarchean hazes suggest that CH_4 concentrations may have

been at the level of 20-30 ppm (e.g. Zerkle et al., 2012), as opposed to ~2 ppm in the

modern atmosphere.

Photochemical experiments with SO_2 have shown that isotope effects and

resulting sulfur isotope fractionations depend on both reaction condition and the

wavelength of incident radiation. Recent results (Whitehill et al. 2012; Endo et al., 2016)

show that shallow positive slopes for $\Delta^{33}S/\delta^{34}S$ (<0.1) are produced by photolysis

reactions at wavelengths shorter than ~220 nm, and steeper slopes (>3) are produced

when longer wavelengths (240 – 340 nm) cause photoexcitation. Similarly, photolysis of

SO_2 yields steeper negative $\Delta^{36}S/\Delta^{33}S$ (<-2), while photoexcitation reactions yield positive

$\Delta^{36}S/\Delta^{33}S$ values (e.g. Whitehill et al., 2012). Applying these observations to the

Neoarchean and early Paleoproterozoic atmosphere, shifts to shallower $\Delta^{33}S/\delta^{34}S$ and

steeper $\Delta^{36}S/\Delta^{33}S$ slopes would be anticipated with the transmission of shorter-

wavelength UV light resulting in greater proportional photolysis reactions. This may result

from the shielding of UV wavelengths by a photochemical CH_4 haze (e.g. Wolf and Toon,

2010), although quantitative details are lacking.

Pertinent to our model, it appears possible that anoxygenic photosynthesis may

have been dominant during intervals when a CH_4 haze blanketed the planet. The

presence of a methane haze would have shielded surface environments from harmful UV

radiation that might then allow ammonia (a strong greenhouse gas) to accumulate in the

lower atmosphere, and thus have shifted upward the peak wavelength of solar radiation

reaching the oceans (Sagan and Chyba, 1996; Wolf and Toon, 2010). Insofar as a

Neoarchean CH_4 haze would scatter and absorb UV and visible wavelengths (Wolf and

Toon, 2010), it may have influenced photosynthetic reactions and primary productivity. In

this regard, the pigments used during anoxygenic photosynthesis differ from those

associated with oxygenic photosynthesis in their molecular details and peak wavelength

of light absorbed. Notably the peak absorption wavelength for chlorophyll-a used by

oxygen-producing plants and cyanobacteria is 100-200 nm less than the peak absorption

wavelength used by bacteriochlorophylls during anoxygenic photosynthesis. The

development of a significant methane haze may thus have shifted primary production of

organic matter towards anoxygenic photosynthesis where the electrons necessary to

drive the metabolic activity ultimately come from the oxidation of H_2S or Fe (II) (rather

than from H_2O in oxygenic photosynthesis). If correct, the dominance of the anaerobic

metabolism over oxygenic photosynthesis would have decoupled inputs of R and O_2,

matching a key assumption of our model.

While our resource-driven ecological model is neither calibrated for amount nor

time, the consistent, robust periodic dynamics observed in the system under some

parameter regimes nonetheless accord with the episodic release of CH_4 from surface

environments assuming a constant, but not overwhelming, source of O_2. According to our

ecological model, if this oxidant was globally distributed at low concentrations, and if

resources were high, a CH_4 haze would likely result. Izon et al. (2017) suggest that the

coupled carbon/sulfur isotope anomalies were associated with biospheric responses to

increased nutrients, with the CH_4 flux controlled by the availability of organic carbon and

sulfate. Increased loading of the oceans with nutrients (e.g., nitrogen, phosphorus) is

most often associated with uplift, mountain building and weathering of exposed

continental masses, all of which could erode, disrupt, or accelerate sedimentation on

marginal marine platforms. In the relatively continuous sedimentary successions where

these episodic coupled anomalies have been recorded, there is no physical evidence for

any temporal sedimentary disruption in the form of unconformities or major hiatal

surfaces. As an alternative hypothesis, we propose that if nutrients (including ferrous iron,

which would have been soluble in the ocean under anoxic conditions) were widely

available in the Neoarchean ocean, then the dynamic release of CH_4 and the episodic

development of CH_4 hazes in the atmosphere could simply have been the result of

constant low level (or gradually increasing) inputs of photosynthetic O_2 to anaerobic

environments where overflow metabolism was dominant.

Geochemically documented CH_4 haze events are currently known only from the

Neoarchean, but the episodic and potentially globe-encompassing ice ages of the

succeeding early Paleoproterozoic Era may similarly be understood in terms of

fluctuations of O_2 and greenhouse gases in the ancient atmosphere (Pavlov et al, 2000;

Bekker et al., 2005; Kopp et al., 2005; Goldblatt et al., 2006; Claire et al., 2006; Bekker

and Kaufman, 2007). Insofar as the early Paleoproterozoic glacial epoch is related to the

transition from the anoxic CH_4-rich atmosphere to an oxic CO_2-rich atmosphere, the

multiplicity of ice ages (including at least three discrete events) in this interval is arguably

related to oscillations in O_2 and CH_4 production and utilization. Related to the dynamic

ecological model presented here, the rise of photosynthetic O_2 prior to each of these

Paleoproterozoic glaciations would have progressively drawn down atmospheric CH_4

levels resulting in surface refrigeration. Primary productivity would then be severely

limited beneath ice-covered oceans, thereby constraining the production of both

fermentative products (FP) and O_2. Atmospheric CO_2 levels during the glaciations likely

rose due to unabated volcanic fluxes and limited chemical weathering of exposed

continental rocks (e.g., Hoffman et al., 1998). However, once threshold concentrations of

CO_2 were reached, the glaciations would have ended rapidly, resulting in high rates of

silicate weathering on land, and the enhanced delivery of sediments, oxidants (including

sulfate and nitrate), nutrients, and alkalinity to the oceans. Melting glaciers and sea ice

coupled with higher post-glacial temperatures would have flooded continental margins

and stimulated both oxygenic photosynthesis and methanogenesis.

Support for such an environmental scenario comes from carbon isotope analyses

of drill core samples collected from across the oldest recorded ice age deposit (the

Ramsay Lake diamictite) in southern Ontario, which reveal strongly negative $\delta^{13}C$ values

(with a nadir of ca. -41‰) in organic matter from sediments that accumulated in the glacial

aftermath (Bekker and Kaufman, 2007). The magnitude and duration of ^{13}C depletion in

the post-glacial Pecors Formation (expressed over 100 meters of sedimentary strata) is

consistent with long-term methanogenic production and methanotrophic oxidation of CH_4

in a shallow oceanic setting. Multiple rare sulfur isotope abundances in pyrite extracted

from the same Pecors samples reveal even more profound negative deviations in

$\Delta^{36}S/\Delta^{33}S$ values relative to the ARA (Wing et al., 2002; 2004), which may similarly record

atmospheric processes during this (and other) post-glacial early Paleoproterozoic CH_4

haze events. Considering our model results, we suggest that the end of the early

Paleoproterozoic glacial epoch around 2.2 billion years ago likely marked the end of

extended methane-dominated atmospheres of the early Earth. This repetitive series of

ice ages was followed by a remarkably long-lived (>100 million years) interval of highly

positive carbon isotope compositions in carbonates known as the Lomagundi-Jatuli event

(Karhu and Holland, 1996; Bachan and Kump, 2015; Mänd et al., 2020). Insofar as

significant ^{13}C enrichments in marine carbonates are driven by the burial and

sequestration of ^{12}C-rich organic matter (Broecker, 1970; Hayes, 1983), the attendant

photosynthetic production and release of O_2 at this time would most likely have marked

the transition to an oxygenated atmosphere with carbon dioxide acting as the major

greenhouse gas thereafter.

Episodically enhanced CH_4 release and buildup in the atmosphere later in Earth

history may be related to the tectonic reorganization of the mobile continents, which

modulate the delivery of reduced gases from spreading ridges and weathering fluxes from

uplifted mountains. These variable fluxes would have impacted the photosynthetic

production of organic matter and degree of ocean oxygenation during discrete climatic

and biological events. Insofar as our model results indicate that these factors control CH_4

fluxes to the atmosphere, we speculate that younger ice ages of the Neoproterozoic and

Paleozoic eras, as well as some Phanerozoic mass extinctions, may be linked to the

dynamic microbial processes modeled here.

Conclusion

Patterns of CH_4 production and release in both driven and non-driven versions of

a novel microbial ecological model, based upon the metabolic repercussions of the

Warburg Effect, suggest that resources and O_2 play important roles in defining the

dynamics of microbial ecosystems over both short and long-time scales. We demonstrate

that forced low level inputs of O_2 result in pulsed releases of CH_4 via overflow metabolism,

a pattern that may explain periodic CH_4 haze events and episodic glaciation deep in

Earth's history. Such pulsed methane releases also occur in modern-day wetlands where

tidal flooding repeatedly tips the balance between oxic and anoxic conditions (see Zhu et

al., 2017), suggesting similar mechanisms may be at work. The qualitative behavior of

this ecosystem model provides a platform for understanding the dynamics of mixed

microbial communities in ancient, modern, and future ecosystems. More detailed models

that blend empirical gas-release data with calibrated input parameters should shed further

light on the links between microbial communities and the timing of atmospheric dynamics.

Methods

Modeling Microbial System Dynamics

We model the dynamics of the interacting microbial system as a system of ordinary

differential equations (ODEs). First, we define functions pertaining to the Warburg Effect

(eq. 1-2) and then describe the equations for each of the microbial functional groups.

We begin by defining two functions that describe Warburg Effect dynamics in

aerobic heterotrophic bacteria (A):

$$f([R], A) = \begin{cases} \alpha\, A\, ([R] - R_0) & \text{for } [R] > R_0 \\ 0 & \text{for } [R] < R_0 \end{cases}$$ Eq. 1 (overflow metabolism of A)

$$g([R], A) = \begin{cases} \varepsilon\, A\, R_1 & \text{for } [R] > R_1 \\ \varepsilon\, A\, [R] & \text{for } [R] < R_1 \end{cases}$$ Eq. 2 (respiratory function of A)

The first function $f(R, A)$ denotes the process of overflow metabolism, where, despite the

presence of ample oxygen, heterotrophic aerobic bacteria (A) produce fermentation

products when resources are abundant. In this function, at resource concentrations higher

than base levels (represented by R_0), there is a linear increase in overflow products (see

Basan et al., 2015, Swain and Fagan, 2018). The second function $g(R, A)$ denotes the

respiratory function (see Eq. 3 below) for these organisms (with the maximal resource

usage condition represented by resource concentration of R_1). Both α and ϵ are

proportionality constants that denote the efficiency of overflow metabolism and

respiration, respectively, and fall in the range [0, 1].

The differential equations below pertain to changes through time of the dominance

of metabolic functional groups as the different microbial functional groups (A, B, C, F, and

G as defined in Fig. 1) compete for available substrates under variable environmental

conditions.

$\frac{dA}{dt} = (1 - \mu)\, [O_2]\, g([R], A) - s\, A$ Eq. 3

$\frac{dB}{dt} = (1 - \kappa)\,\beta\,[\text{FP}]\,B - u\,B$ Eq. 4

$\frac{dC}{dt} = (1 - \lambda)\,\gamma\,[O_2][CH_4]\,C - v\,C$ Eq. 5

$\frac{dF}{dt} = (1 - \xi)\,\omega\,[R]\,F - a_1\,F$ Eq. 6

$\frac{dG}{dt} = (1 - \psi)\,\Omega\,[CH_4]\,G - a_2\,G$ Eq. 7

$\frac{dH}{dt} = (1 - \rho)\,\tau\,[\text{FP}]\,H - m\,H$ Eq. 8

The first term on the right-hand side (RHS) of each differential equation is positive and

denotes the process of assimilation of substrates needed for growth (increase in biomass)

and energy production for the metabolic class. The second RHS term in each equation is

negative and denotes the energy requirement for homeostasis (see Table 1). In equations

3-8, the terms governing resource incorporation efficiency, generically denoted $(1 - X)$

with X replaced by the consortium-specific parameters $(\mu, \kappa, \lambda, \xi \text{ and } \psi)$, specify the

proportion of the required substrate in the extracellular environment that is incorporated

into cells, which can then be used for making biomass by those microbes. Variation in

these efficiencies is assumed to be modulated by biochemical and physical constraints

of the different microbial functional groups. The remaining reservoir of resources outside

of the cell either leave the system as gases or are available for later incorporation. All

twelve coefficients of efficiency $(\mu, \kappa, \lambda, \xi, \psi, \rho, \varepsilon, \beta, \gamma, \omega, \Omega, \text{ and } \tau)$ lie in the range [0, 1]. The

coefficients related to homeostasis $(s, u, v, a_1, a_2, \text{or } m)$ represent the basal metabolic

requirement values for microbial functional groups A, B, C, F, G, and H respectively. All

parameters and their interpretations are summarized in Table 1.

Functional group	Resource incorporation efficiency	Resource utilization efficiency	Energy requirements for homeostasis
A (aerobic heterotrophs)	$(1 - \mu)$	ε (for R)	s
B (methanogens)	$(1 - \kappa)$	β (for FP)	u
C (aerobic methanotrophs)	$(1 - \lambda)$	γ (for O_2 and CH_4)	v
F (fermenters)	$(1 - \xi)$	ω (for R)	a_1
G (anaerobic methanotrophs)	$(1 - \psi)$	Ω (for CH_4)	a_2
H (anaerobic heterotrophs)	$(1- \rho)$	τ (for FP)	m

Table 1: *Differential equation parameters governing the dynamics of a system of five*

microbial functional groups. Parameters summarize the efficiency of resource

incorporation, the efficiency of resource utilization, and the energy requirements for

homeostasis.

The following set of differential equations reflect dynamic changes in the

abundance of resources or metabolic products in the five microbial functional groups

through time:

$\frac{d[R]}{dt} = -f([R], A) - g([R], A) - \xi \, \omega \, [R] \, F + \sigma_1(t)$ Eq 9

$$\frac{d[\text{FP}]}{dt} = f([R], A) - \beta B [\text{FP}] + \xi \omega R F - \tau [\text{FP}] H \qquad \text{Eq 10}$$

$$\frac{d[\text{CO}_2]}{dt} = \mu [O_2] g([R], A) + \gamma \lambda [O_2] [\text{CH}_4] C + \psi \Omega [\text{CH}_4] G + \rho \tau [\text{FP}] H \qquad \text{Eq 11}$$

$$\frac{d[\text{CH}_4]}{dt} = \kappa \beta B [\text{FP}] - \gamma [O_2] [\text{CH}_4] C - \Omega [\text{CH}_4] G \qquad \text{Eq 12}$$

$$\frac{d[O_2]}{dt} = -\{\mu [O_2] g([R], A) + \gamma \lambda [O_2] [\text{CH}_4] C\} + \sigma_2(t) \qquad \text{Eq 13}$$

In equations 8-12 the RHS positive terms indicate either input or production of the

resources, while negative terms indicate metabolic resource consumption. All the terms

involve interaction of the resources with one or more of the microbial functional groups,

which are modelled in equations 3-8, except for the 'forcing terms' $\sigma_1(t)$ and $\sigma_2(t)$. These

terms (see Eqs. 9 and 13) denote the time-dependent input rates of R and O_2 that are

controlled externally. In this model, CO_2 is considered a sink that evades to the

atmosphere and is decoupled from the autotrophic processes. This assumption makes R

an independent variable, and further ensures that substrate production is not

photoautotrophic (the case in many scenarios). Moreover, as developed here, the

ecological model is not spatially compartmented in order to make the results more general

in nature.

There exist many methanogenic systems where oxic and anoxic parts are

heterogeneously mixed, albeit at generally small scales or temporally, such as in wetlands

(Kirsche et al., 2013). In this ecological model we have simplified the process of

methanogenesis in several ways: i) FPs can be one of many substrates, including

formate, acetate, and lactate, all of which can be used for methane production by various

methanogens; and ii) for hydrogen (H_2) + CO_2 methane production, we assume that

reduced reactants are unlimited, and insofar as CO_2 is a sink and evades the system as

a gas, it is unlimited as well (i.e., it is not used for autotrophic processes that deliver new

resources to the community). For the anaerobic oxidation of methane by functional group

G, we lump all of the potential anaerobic methanotrophs here, but recognizing that both

CO_2 and carbonate alkalinity may be formed, we include a conversion efficiency term (Ω)

to calculate the relative concentrations of CO_2 alone.

This coupled differential equation system was solved through Wolfram

Mathematica v. 12.1 and R v. 3.6.2. A regular parameter sweep was first performed for

the non-driven case (no external input: $\sigma_1, \sigma_2 = 0$) using Latin Hypercube sampling (LHS)

for 500,000 different scenarios using R packages *deSolve* (Soetaert et al., 2010) and *lhs*

(Carnell, 2020). We then calculated the partial rank correlation coefficients (PRCCs) to

explore the effect of each parameter and initial condition on the concentrations of gases

(CH_4, O_2 and CO_2) through R package *sensitivity* (Iooss et al., 2020).

To further explore the trends in these gas emissions, we calculated the relative

proportions of these gases under a set of base parameters that depict biologically

possible scenarios (according to data from Payne and Wiebe, 1978; Geyer et al., 2016;

Bowles et al., 2019; and Morris and Schmidt, 2013). All the efficiencies were set at half

maximum for the base case. We then varied one parameter at a time to look at the

dependence on that specific variable on gas emissions as a function of resource and

oxygen initial conditions. In doing so, we collated the data for different initial conditions

(such as different initial ratios of the microbial functional groups) and plotted the averaged

values of the equilibrium state as a function of initial resource and oxygen inputs. For

driven dynamics, we followed the same procedure of parameter dependence, but now

including the two R and O_2 external input parameters, σ_1 and σ_2. To calculate the period

and amplitude of the driven dynamics, values were numerically estimated based on

dynamic results from the first few oscillations. Periodicity statistics for the dynamics

resulting from both high and low extreme values of σ_2 are estimates given the difficulty of

estimating them for very long or extremely short oscillations that occur at these values of

σ_2.

For looking at the sensitivity of the model to addition or removal of microbial

functional groups, we ran the same tests for a model without H (anaerobic heterotrophs).

To measure sensitivity of the demarcation of the oscillatory phase in the space of resource

and oxygen addition ($\sigma_1 - \sigma_2\ space$), across a wide variety of parameter combinations, we

first used LHS to sample 100,000 combinations through the parameters space. We

calculated, for each set of parameters, twenty points on the $\sigma_1 - \sigma_2$ space, estimated the

relative value for which there is a phase change from oscillatory behavior to no

oscillations, and from these values calculated the slope of the line which demarcates the

space for each set of LHS parameters. We performed PRCC analysis to find the

dependence of the slope value on the parameters.

Notation for Sulfur Isotope Compositions

Sulfur-isotope ratios are conventionally reported in delta (δ) notation and reflect

the permil (‰) deviation of the ratio of the least abundant isotope (^{33}S, ^{34}S, or ^{36}S) to the

most abundant isotope (^{32}S), relative to the same ratio in an international reference

standard (Vienna Canyon Diablo Troilite, V-CDT). For example, δ^{34}S =

[(^{34}S/^{32}S)$_{sample}$/(^{34}S/^{32}S)$_{V\text{-}CDT}$ − 1] x 1000. The majority of terrestrial processes fractionate

S-isotopes mass-dependently, whereby δ^{33}S ≈ 0.515 * δ^{34}S and δ^{36}S ≈ 1.91 * δ^{34}S.

Departure from mass-dependent behavior, or mass-independent fractionation (S-MIF), is

expressed in capital-delta (Δ) notation as either non-zero $\Delta^{33}S = [(^{33}S/^{32}S)_{sample}/(^{33}S/^{32}S)_{V\text{-}CDT}$

$_{CDT} - (^{34}S/^{32}S)_{sample}/(^{34}S/^{32}S)_{V\text{-}CDT}]^{0.515}$ or $\Delta^{36}S = [(^{36}S/^{32}S)_{sample}/(^{36}S/^{32}S)_{V\text{-}CDT} -$

$(^{34}S/^{32}S)_{sample}/(^{34}S/^{32}S)_{V\text{-}CDT}]^{1.9}$.

Acknowledgements

This research was supported in part by NSF award EAR-1338810 to AJK. AS would like

to thank the COMBINE program at the University of Maryland (DGE-1632976) for

technical and academic support.

Code and data availability

All the codes that were used to explore the ODE model and generate the relevant figures

are available publicly at the following github repository:

https://github.com/anshuman21111/ch4_haze . All extra figures and simulation data,

above and exceeding the supplementary materials are also found on this repository.

References

1. Anbar, A.D. and Knoll, A.H., 2002. Proterozoic ocean chemistry and evolution: a

bioinorganic bridge?. science, 297(5584), pp.1137-1142.

2. Bachan, A. and Kump, L.R., 2015. The rise of oxygen and siderite oxidation

during the Lomagundi Event. Proceedings of the National Academy of Sciences,

112(21), pp.6562-6567.

3. Basan, M., Hui, S., Okano, H., Zhang, Z., Shen, Y., Williamson, J.R. and Hwa,

T., 2015. Overflow metabolism in Escherichia coli results from efficient proteome

allocation. Nature, 528(7580), pp.99-104.

4. Bekker, A., Kaufman, A.J., Karhu, J.A. and Eriksson, K.A., 2005. Evidence for

Paleoproterozoic cap carbonates in North America. Precambrian Research,

137(3-4), pp.167-206.

5. Bekker, A. and Kaufman, A.J., 2007. Oxidative forcing of global climate change:

a biogeochemical record across the oldest Paleoproterozoic ice age in North

America. Earth and Planetary Science Letters, 258(3-4), pp.486-499.

6. Carlson, R.P., Beck, A.E., Phalak, P., Fields, M.W., Gedeon, T., Hanley, L.,

Harcombe, W.R., Henson, M.A. and Heys, J.J., 2018. Competitive resource

allocation to metabolic pathways contributes to overflow metabolisms and

emergent properties in cross-feeding microbial consortia. Biochemical Society

Transactions, 46(2), pp.269-284.

7. Carnell, R., 2020. lhs: Latin Hypercube Samples. R package version 1.0.2.

https://CRAN.R-project.org/package=lhs

8. Chistoserdova, L., Vorholt, J.A. and Lidstrom, M.E., 2005. A genomic view of

methane oxidation by aerobic bacteria and anaerobic archaea. Genome biology,

6(2), p.208.

9. Eigenbrode, J.L. and Freeman, K.H., 2006. Late Archean rise of aerobic

microbial ecosystems. Proceedings of the National Academy of Sciences,

103(43), pp.15759-15764.

10. Farquhar, J., Cliff, J., Zerkle, A.L., Kamyshny, A., Poulton, S.W., Claire, M.,

Adams, D. and Harms, B., 2013. Pathways for Neoarchean pyrite formation

constrained by mass-independent sulfur isotopes. Proceedings of the National

Academy of Sciences, 110(44), pp.17638-17643.

11. Halverson, G.P., Hoffman, P.F., Schrag, D.P. and Kaufman, A.J., 2002. A major

perturbation of the carbon cycle before the Ghaub glaciation (Neoproterozoic) in

Namibia: Prelude to snowball Earth?. Geochemistry, Geophysics, Geosystems,

Vol. 3(6)

12. Hayes, J.M., 1994. Early life on Earth. In Nobel symposium (Vol. 84, pp. 220-36).

13. Iooss, B., Da Veiga, S., Janon, A., Pujol, G., et al., 2020. sensitivity: Global

Sensitivity Analysis of Model Outputs. R package version 1.20.0. https://CRAN.R-

project.org/package=sensitivity

14. Izon, G., Zerkle, A.L., Zhelezinskaia, I., Farquhar, J., Newton, R.J., Poulton,

S.W., Eigenbrode, J.L. and Claire, M.W., 2015. Multiple oscillations in

Neoarchaean atmospheric chemistry. Earth and Planetary Science Letters, 431,

pp.264-273.

15. Izon, G., Zerkle, A.L., Williford, K.H., Farquhar, J., Poulton, S.W. and Claire,

M.W., 2017. Biological regulation of atmospheric chemistry en route to planetary

oxygenation. Proceedings of the National Academy of Sciences, 114(13),

pp.E2571-E2579.

16. Kasting, J.F., 2001. The rise of atmospheric oxygen. Science, 293(5531),

pp.819-820.

17. Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J.G., Dlugokencky,

E.J., Bergamaschi, P., Bergmann, D., Blake, D.R., Bruhwiler, L. and Cameron-

Smith, P., 2013. Three decades of global methane sources and sinks. Nature

geoscience, 6(10), pp.813-823.

18. Konhauser, K.O., Pecoits, E., Lalonde, S.V., Papineau, D., Nisbet, E.G., Barley,

M.E., Arndt, N.T., Zahnle, K. and Kamber, B.S., 2009. Oceanic nickel depletion

and a methanogen famine before the Great Oxidation Event. Nature, 458(7239),

pp.750-753.

19. Kopp, R.E., Kirschvink, J.L., Hilburn, I.A. and Nash, C.Z., 2005. The

Paleoproterozoic snowball Earth: a climate disaster triggered by the evolution of

oxygenic photosynthesis. Proceedings of the National Academy of Sciences,

102(32), pp.11131-11136.

20. Kreft, J.U., Griffin, B.M. and González-Cabaleiro, R., 2020. Evolutionary causes

and consequences of metabolic division of labour: why anaerobes do and

aerobes don't. Current opinion in biotechnology, 62, pp.80-87.

21. Lyu, Z., Shao, N., Akinyemi, T. and Whitman, W.B., 2018. Methanogenesis.

Current Biology, 28(13), pp.R727-R732.

22. Manzoni, S., Taylor, P., Richter, A., Porporato, A. and Ågren, G.I., 2012.

Environmental and stoichiometric controls on microbial carbon-use efficiency in

soils. New Phytologist, 196(1), pp.79-91.

23. Pavlov, A.A., Kasting, J.F., Eigenbrode, J.L. and Freeman, K.H., 2001. Organic

haze in Earth's early atmosphere: Source of low-13C Late Archean kerogens?.

Geology, 29(11), pp.1003-1006

24. Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J.G.,

Dlugokencky, E.J., Etiope, G., Bastviken, D., Houweling, S. and Janssens-

Maenhout, G., 2016. The global methane budget 2000–2012. *Earth System*

Science Data, *8*(2), pp.697-751.

25. Soetaert, K., Petzoldt, T. and Setzer, R.W., 2010. Package deSolve: solving

initial value differential equations in R. J Stat Softw, 33(9), pp.1-25.

26. Swain, A. and Fagan, W.F., 2018. A mathematical model of the Warburg Effect:

Effects of cell size, shape and substrate availability on growth and metabolism in

bacteria. Math. Biosci. Eng, 16, pp.168-186.

27. Voet, D. and Voet, J.G., 2004. Biochemistry. Hoboken. John Wiley & Sons

28. Warburg, O., 1956. On the origin of cancer cells. Science, 123(3191), pp.309-

314.

29. Ward, L.M., Rasmussen, B. and Fischer, W.W., 2019. Primary productivity was

limited by electron donors prior to the advent of oxygenic photosynthesis. Journal

of Geophysical Research: Biogeosciences, 124(2), pp.211-226.

30. Williford, K.H., Ushikubo, T., Lepot, K., Kitajima, K., Hallmann, C., Spicuzza,

M.J., Kozdon, R., Eigenbrode, J.L., Summons, R.E. and Valley, J.W., 2016.

Carbon and sulfur isotopic signatures of ancient life and environment at the

microbial scale: Neoarchean shales and carbonates. Geobiology, 14(2), pp.105-

128.

31. Wing, B.A., Bekker, A., Brabson, L., Farquhar, J., Kaufman, A.J. and Rumble, D.,

2004, November. Atmospheric chemistry of SO2 during the earliest Huronian

glaciation. In Abstr. Progr. Geol. Soc. Am (Vol. 36).

32. Wing, B.A., Brabson, E., Farquhar, J., Kaufman, A.J., Rumble, D. and Bekker, A.,

2002, August. Delta S-33, delta S-34 and delta C-13 constraints on the

Paleoproterozoic atmosphere during the earliest Huronian glaciation. In

Geochimica et Cosmochimica Acta Vol. 66, No. 15 A, pp. A840-A840

33. Wolf, E.T. and Toon, O.B., 2010. Fractal organic hazes provided an ultraviolet

shield for early Earth. Science, 328(5983), pp.1266-1268.

34. Yan, Z., Joshi, P., Gorski, C.A. and Ferry, J.G., 2018. A biochemical framework

for anaerobic oxidation of methane driven by Fe (III)-dependent respiration.

Nature communications, 9(1), pp.1-9.

35. Zerkle, A.L., Claire, M.W., Domagal-Goldman, S.D., Farquhar, J. and Poulton,

S.W., 2012. A bistable organic-rich atmosphere on the Neoarchaean Earth.

Nature Geoscience, 5(5), pp.359-363.

36. Zhelezinskaia, I., Kaufman, A.J., Farquhar, J. and Cliff, J., 2014. Large sulfur

isotope fractionations associated with Neoarchean microbial sulfate reduction.

Science, 346(6210), pp.742-744.

735